

Title	PRE-MELTING MODE IN THE DISORDERED KNO <sub>3</sub> CRYSTAL OBSERVED BY THE LOW-FREQUENCY RAMAN SCATTERING(Session I : Cross-Disciplinary Physics, The 1st Tohwa University International Meeting on Statistical Physics Theories, Experiments and Computer Simulations)
Author(s)	Sanada, Mizuho; Yagi, Toshirou
Citation	物性研究 (1996), 66(3): 384-385
Issue Date	1996-06-20
URL	<a href="http://hdl.handle.net/2433/95839">http://hdl.handle.net/2433/95839</a>
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

PRE-MELTING MODE IN THE DISORDERED KNO<sub>3</sub> CRYSTAL OBSERVED BY THE LOW-FREQUENCY RAMAN SCATTERING

Mizuho Sanada\* and Toshirou Yagi\*\*

\*Department of Electrical Engineering, Kyushu Kyoritsu University, Kitakyushu 807, Japan

\*\*Research Institute for Electronic Science, Hokkaido University, Sapporo 060, Japan

On heating, potassium nitrate, KNO<sub>3</sub> undergoes a structural phase transition from the orthorhombic structure (phase II) to the trigonal structure (phase I) at 403 K ( $T_c$ ) and melts at 606 K ( $T_m$ ). The central peak in low-frequency Raman scattering spectra of KNO<sub>3</sub> was studied from room temperature up to 130 K above the melting point  $T_m$ . A drastic growth was observed with increasing temperature below and above  $T_m$ . The spectra were numerically analyzed and the results revealed that the central peak of KNO<sub>3</sub> consists of two spectral components; one is common between the solid state phase I and the molten state, and the other is sensitive for melting process.

Experiment

A single crystal of KNO<sub>3</sub> was grown from aqueous solution. The crystal axes were determined from the crystal habit. After cut and polished, the specimen was placed in the scattering




Figure 1: Numerical fit of the central peak of KNO<sub>3</sub> in the solid state phase I (a), and in the molten state (b). Upper figures, dots are observed spectra corrected by Bose-Einstein thermal factor  $n(\omega) + 1$ . (a) Two damped-harmonic oscillator functions,  $O_i = A_i \omega_i^2 \Gamma_i \omega / [(\omega_i^2 - \omega^2)^2 + \Gamma_i^2 \omega^2]$ ,  $i = 1, 2$ . (b) One damped-harmonic oscillator function  $O_1$  and a Gaussian function  $G = A_g \exp(-a_g \omega^2)$ . Lower figures indicate the residue of the fit.




Figure 2: Temperature dependence of the components  $O_1$  and  $O_2$  obtained from numerical analysis.

cell with a temperature controller. An  $\text{Ar}^+$  ion laser was used as light source with 514.5 nm at 100 mW. The light scattered at 90 degree from the specimen was analyzed with an 80 cm double monochromator.

The low-frequency Raman spectra were observed in  $a(cc)b, a(ca)b, a(ba)b$  and  $a(bc)b$  geometries, which is denoted by the orthorhombic (phase II) axes. The spectra shows good agreement with previous studies.<sup>1,2)</sup> The central peak, in any geometry, shows drastic growth with increasing temperature. In particular, the polarized central peak grows extremely in the  $a(cc)b$  geometry.

#### Numerical Analysis

We tried to fit the observed spectra using several functions. As shown in Figure 1, the best fit was obtained for the spectra in solid state phase I with a set of two damped-harmonic oscillator functions ( $O_i = A_i \omega_i^2 \Gamma_i \omega / [(\omega_i^2 - \omega^2)^2 + \Gamma_i^2 \omega^2]$ ,  $i = 1, 2$ ), and for the spectra in molten state with a set of a Gaussian function ( $G = A_g \exp(-a_g \omega^2)$ ) and a damped-harmonic oscillator function ( $O_1$ ).<sup>3,4)</sup>

Temperature dependence of the damped harmonic oscillator functions  $O_i$  is shown in Figure 2. The component  $O_1$  is characteristic of the molten state even in the solid state, or it exists as a common mode. The other one  $O_2$  becomes unstable near  $T_m$ , which indicates the relation to the melting process, thus it should be termed as a melting mode in the solid state. These modes should be coupled with each other in the solid state, because of the similar temperature dependence of frequency.

#### References

- 1) M. H. Brooker, *Can. J. Chem.*, **55**(1977), 1242.
- 2) K. Akiyama, Y. Morioka and I. Nakagawa, *J. Phys. Soc. Jpn.*, **48**(1980)898.
- 3) G. Neumann and H. Vogt, *Phys. Stat. Sol. (b)*, **85**(1978)179.
- 4) W. Hayes and R. Loudon, *Scattering of Light by Crystals* (John Wiley: New York, 1978).