

Title	Bounds for Global Solutions of Some Semilinear Parabolic Equations
Author(s)	OTANI, Mitsuharu
Citation	数理解析研究所講究録 (1989), 698: 36-50
Issue Date	1989-06
URL	http://hdl.handle.net/2433/101456
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

Bounds for Global Solutions of Some Semilinear Parabolic Equations

By

Mitsuharu ÔTANI (大谷光春)

Department of Mathematics, Faculty of Science, Tokai University

1. Introduction. In this note we are concerned with the asymptotic behavior of global solutions of the initial boundary value problem $(E)_N$ (or $(E)_D$) for the semilinear parabolic equation :

$$(E) \quad \begin{cases} (1) & u_t(x,t) - \Delta u(x,t) = f(x,u(x,t)), \quad (x,t) \in \Omega \times [0, \infty), \\ (2) & u(x,0) = u_0(x), \quad x \in \Omega, \end{cases}$$

with the Neumann boundary condition

$$(3) \quad \partial u(x,t) / \partial n = 0, \quad (x,t) \in \partial \Omega \times [0, \infty),$$

(or the Dirichlet boundary condition

$$(4) \quad u(x,t) = 0, \quad (x,t) \in \partial \Omega \times [0, \infty).)$$

Here Ω is a bounded domain in \mathbb{R}^N with smooth boundary $\partial \Omega$ and f is a continuous function from $\Omega \times \mathbb{R}^1$ to \mathbb{R}^1 . It is well known that if f is superlinear in u , then (E) has solutions which blow up in finite time. So it would be natural to ask whether (E) has a global solution which blows up at ∞ or not.

This kind of problem was first studied by [7,8] for an abstract equation of the form $u_t + \partial \varphi^1(u(t)) - \partial \varphi^2(u(t)) = 0$ in a real Hilbert space \hat{H} , where $\partial \varphi^i$ are subdifferentials of lower semi-continuous convex and homogeneous functionals φ^i ($i=1,2$) on H . As an application of a result of [7], it is shown in [8] that

every global solution of $(E)_D$ with $f(x,u) = |u|^{p-2}u$ is uniformly bounded in $H_0^1(\Omega)$ with respect to time t , provided that $p < 2^*$, $2^* = \infty$ for $N=1,2$; $2^* = 2N/(N-2)$ for $N \geq 3$. Ni-Sacks-Tavantzis [6] also studied $(E)_D$ for the case where Ω is convex and $f(x,u) = |u|^{p-2}u$ and showed that if $2 < p < 2 + 2/N$, then every positive global solution of $(E)_D$ is uniformly bounded in $L^\infty(\Omega)$ with respect to t and that if $2^* \leq p$ ($N \geq 3$), then $(E)_D$ has a global solution u such that $\|u(t)\|_{L^\infty} \rightarrow \infty$ as $t \rightarrow +\infty$.

Cazenave-Lions [2] treated more general nonlinear terms $f(x,u) = f_0(u)$ satisfying

$$(f)_0 \begin{cases} (o) & f_0 \in C^1(\mathbb{R}^1; \mathbb{R}^1), \\ (i) & |f_0(u)| \leq C_1|u| + C_2|u|^{p-1} \quad \forall u \in \mathbb{R}^1, \quad p < 2^*, \\ (ii) & u f_0(u) \geq (2+\varepsilon) F_0(u), \quad \forall u \in \mathbb{R}^1, \quad \varepsilon > 0, \quad F_0(u) = \int_0^u f_0(t) dt, \end{cases}$$

and showed that every global solution of $(E)_D$ is bounded in $L^\infty(\Omega)$ uniformly in $[t, \infty)$ for any $t > 0$. Furthermore this bound depends only on t and the $H_0^1(\Omega)$ -norm of u_0 , provided that $2 < p < 2_*$, $2_* = \infty$ for $N = 1$; $2_* = 2 + 12/(3N - 4)$ for $N \geq 2$. Giga [4] removed this restriction on p for positive solutions, i.e., he showed that if $p < 2^*$, then the $L^\infty(\Omega)$ bound for every positive global solution depends only on the $L^\infty(\Omega)$ -norm of u_0 .

For all these studies, it seems that there is no result for the Neumann problem $(E)_N$ in this direction in spite of its importance. In studying $(E)_N$, it must be noted that the methods in [2,4,6] rely much on the Dirichet boundary condition, so do not work for $(E)_N$.

The main purpose of this note is to show that "phase plane" method introduced in [7,8] works also for $(E)_N$. However, in order to apply this method to $(E)_N$, we have to remove two major restrictions in [7,8], i.e., conditions that $\partial\varphi^1$ is coercive and $\partial\varphi^2$ is homogeneous. As a matter of course, in carrying out this, we need much more careful consideration in the phase space than in [7,8]. In this note we are concerned with the following nonlinearity of f :

$$(f) \left\{ \begin{array}{l} f(.,.) \text{ is a continuous function from } \Omega \times \mathbb{R}^1 \text{ into } \mathbb{R}^1 \\ \text{and there exist constants } K_i (i=0,1,2,3) \text{ and numbers} \\ p \in (2, 2^*), \delta > 0 \text{ and } \xi > 0 \text{ such that} \\ (i) \quad |f(x,u)| \leq K_0 (1 + |u|^{p-1}) \quad \forall (x,u) \in \Omega \times \mathbb{R}^1, \\ (ii) \quad F(x,u) = \int_0^u f(x,t) dt \geq K_1 |u|^{2+\delta} - K_2 \quad \forall (x,u) \in \Omega \times \mathbb{R}^1, \\ (iii) \quad u f(x,u) \geq (2 + \xi) F(x,u) - K_3 \quad \forall (x,u) \in \Omega \times \mathbb{R}^1. \end{array} \right.$$

Then our main results are stated as follows.

Theorem I. Let (f) be satisfied and u be a global solution of $(E)_N$ (or $(E)_D$) such that $u \in V \equiv W_{loc}^{1,2}([0,\infty); L^2(\Omega)) \cap L_{loc}^2([0,\infty); H^2(\Omega))$. Then there exists a positive constant $C_0 = C_0(|u_0|_{H^1}, K_0, K_1, K_2, K_3, \delta, \xi, |\Omega|)$ such that

$$(5) \quad \sup_{t \geq 0} |u(t)|_{L^2} \leq C_0,$$

$$(6) \quad \sup_{t \geq 0} |u(t)|_{H^1} < +\infty,$$

$$(7) \quad \text{There exists a number } T_1 \text{ such that } \sup_{t \geq T_1} |u(t)|_{H^1} \leq C_0,$$

$$(8) \quad \sup_{t \geq 0} |u(t)|_{H^1} \leq C_0, \text{ provided that } p \in (2, 2_*), 2_* = \infty \text{ for } N=1 \\ \text{and } 2_* = 2 + 12/(3N - 4).$$

Theorem II. Let (f) be satisfied and u be a global solution of $(E)_N$ (or $(E)_D$) such that $u_0 \in L^\infty(\Omega)$ and $u \in L_{loc}([0, \infty); L^\infty(\Omega)) \cap W_{loc}^{1,2}((0, \infty); L^2(\Omega)) \cap L_{loc}^2((0, \infty); H^2(\Omega))$. Then there exists a positive constant $C_1 = C_1(|u_0|_{L^\infty}, K_0, K_1, K_2, K_3, \delta, \varepsilon, |\Omega|)$ such that

$$(9) \quad \sup_{t \geq 0} |u(t)|_{L^\infty} < \infty,$$

$$(10) \quad \text{There exists a number } T_1 \text{ such that } \sup_{t \geq T_1} |u(t)|_{L^\infty} \leq C_1,$$

$$(11) \quad \sup_{t \geq 0} |u(t)|_{L^p} \leq C_1, \text{ provided that } p \in (2, 2_*).$$

Remark 1. (1) Assertions in Theorems I and II hold true also for Robin problem $(E)_R$, i.e., (E) with more general boundary condition:

$$(3)' \quad \frac{\partial u}{\partial n}(x, t) + \sigma(x) u(x, t) = 0, \quad (x, t) \in \partial\Omega \times [0, \infty), \quad \sigma \in L^\infty(\partial\Omega).$$

The following arguments for $(E)_N$ can be applied for $(E)_R$ with slight modifications.

(2) Conditions $(f)_0$ does not allow f to contain linear or sublinear parts, but condition (f) allow it. For example, $f(x, u) = |u|^{q-2}u + |u|^{p-2}u$ with $1 < q \leq 2 < p$ satisfies (f) but not $(f)_0$.

(3) If f satisfies condition (f), then so does $g(x, u) = f(x, u) + u$. Indeed it is clear that g satisfies (i)-(ii) of (f), and since

$$u g(x, u) = u^2 + u f(x, u) \geq u^2 + (2 + \varepsilon) F(x, u) - K_3,$$

$$(2 + \varepsilon/2)(u^2/2 + F(x, u)) + \varepsilon F(x, u)/2 - \varepsilon u^2/4 - K_3,$$

$$(2 + \varepsilon/2) \int_0^u g(x, t) dt + \varepsilon(K_1 |u|^{2+\delta} - K_2 - u^2/2)/2 - K_3,$$

and $K_1 |u|^{2+\delta} - K_2 - |u|^2/2$ is bounded below, $g(x,u)$ also satisfies (iii) of (f) with ε and K_3 replaced by $\varepsilon/2$ and some K'_3 .

2. Proofs of Theorems. We shall give here proofs only for Neumann problem $(E)_N$, which are more complicated than those for $(E)_D$ and also valid for $(E)_D$.

Instead of (1), let us here consider its equivalent :

$$(1)' \quad u_t - \Delta u + u = u + f(x,u) = g(x,u) \quad (x,t) \in \Omega \times [0,\infty).$$

As was seen in (3) of Remark 1, g also satisfies (f). We use the same K_i , δ and ε for g as before. In what follows, (\cdot, \cdot) and $|\cdot|$ denote the inner product and norm of $L^2(\Omega)$. We also denote by $|\cdot|_r$ and $\|\cdot\|$ the $L^r(\Omega)$ norm and $H^1(\Omega)$ norm respectively. Now we introduce several functionals on $H^1(\Omega)$

$$- \quad A(u) = \frac{1}{2} (|\nabla u|^2 + |u|^2) = \frac{1}{2} \|u\|^2$$

$$- \quad G(u) = \int_{\Omega} \int_0^{u(x)} g(x,t) dt dx,$$

$$- \quad J(u) = A(u) - G(u),$$

$$- \quad j(u) = (g(\cdot, u(\cdot)), u(\cdot)) - 2A(u)$$

and subsets of $H^1(\Omega)$:

$$- \quad S_a = \{ u \in H^1(\Omega) ; j(u) = a \}, \quad a \in \mathbb{R}^1.$$

Then, by virtue of (ii) and (iii) of (f), we obtain

$$(12) \quad G(u) \geq K_1 |u|_{2+\delta}^{2+\delta} - K_2 |\Omega| \quad \forall u \in H^1(\Omega), \quad (|\Omega| \text{ is the volume of } \Omega.)$$

$$(13) \quad j(u) \geq \varepsilon G(u) - 2J(u) - K_3 \quad \forall u \in H^1(\Omega),$$

$$(14) \quad A(u) \geq (1 + \varepsilon/2)G(u) - K_3/2 - a/2 \quad \forall u \in S_a.$$

Furthermore, by (i) of (f) and Sobolev's inequality :

$|u|_p \leq C_b \|u\|$, there exists a constant d_0 such that $d_0 \geq K_3$ and $G(u) \leq d_0 (A(u)^{p/2} + 1)$, whence follows

$$(15) \quad A(u) \geq ([G(u)/d_0 - 1]^+)^{2/p}, \quad \text{where } [a]^+ = \max(a, 0).$$

(Note that d_0 is a constant depending only on $K_0, K_3, p, |\Omega|$ and embedding constant C_b .) Introducing a new parameter

$\alpha = (2 + \varepsilon)d_0 - K_3 > 0$ and taking account of (14) and (15),

we can draw the following Fig. 1 which illustrates how S_0, S_α and lines $J(u) = J(u_0)$ and $J(u) = -d_0$ are located in the $(G(u), A(u))$ -phase plane.

We here claim the following proposition.

Proposition 1. Let u be a global solution of $(E)_N$ belonging to V . Then we have

(i) $J(u(t))$ is monotone decreasing in t ,

(ii) $J(u(t)) \geq -d_0 \quad \forall t \geq 0$,

(iii) $\int_0^\infty |u_t(t)|^2 dt \leq J(u_0) + d_0$,

(iv) $|u(t)| \leq C_0 \quad \forall t \geq 0$,

(v) $\int_t^{t+1} G(u(s))^2 ds \leq C_0 \quad \forall t \geq 0$,

(vi) $\int_t^{t+1} \|u(s)\|^4 ds \leq C_0 \quad \forall t \geq 0$,

where C_0 is a constant depending only on $K_i (i=0,1,2,3), \varepsilon, \delta, d_0$ and $J(u_0)$.





Fig. 1  In this region there is no element of $H^1(\Omega)$.

$$\alpha = (2 + \epsilon) d_0 - K_3 > 0$$

$$d_1 = (2 + \epsilon)(J(u_0) + d_0) / \epsilon$$

$$d_2 = (2J(u_0) + (2 + \epsilon)d_0) / \epsilon$$

$$d_0 = d_0(K_0, K_3, p, |\Omega|, C_b)$$

Proof of Proposition 1. Multiplying (1)' by u_t and u , we have

$$(16) \quad |u_t(t)|^2 + \frac{d}{dt} J(u(t)) = 0 \quad \text{for a.e. } t \in [0, \infty),$$

$$(17) \quad \frac{1}{2} \frac{d}{dt} |u(t)|^2 = j(u(t)) \quad \text{for a.e. } t \in [0, \infty).$$

Then assertion (i) is a direct consequence of (16), and it follows from (12), (13), (17) and (i) that

$$(18) \quad \begin{aligned} \frac{1}{2} \frac{d}{dt} |u(t)|^2 &\geq \varepsilon G(u(t)) - 2J(u_0) - K_3 \\ &\geq \varepsilon K_1 |\Omega|^{-\delta/2} |u(t)|^{2+\delta} - \varepsilon K_2 |\Omega| - 2J(u_0) - K_3. \end{aligned}$$

Suppose that $J(u(t_0)) < -d_0$ for some t_0 , then in view of (i) and Fig. 1, we find that $j(u(t)) \geq \alpha > 0$ for all $t \geq t_0$. Hence (17) and (18) assure that there exists $t_1 > 0$ such that

$$(19) \quad \frac{d}{dt} |u(t)|^2 \geq \varepsilon K_1 |\Omega|^{-\delta/2} |u(t)|^{2+\delta} \quad \text{for a.e. } t \geq t_1,$$

which implies that $|u(t)|$ blows up in a finite time. This is a contradiction. Thus (ii) is verified. Consequently integration of (16) over $[0, \infty)$ gives (iii). Suppose now that there exists a $t_1 > 0$ such that

$$\varepsilon K_1 |\Omega|^{-\delta/2} |u(t_1)|^{2+\delta} / 2 \geq \varepsilon K_2 |\Omega| + 2J(u_0) + K_3,$$

then by (18), $|u(t)|$ is monotone increasing in the neighbourhood of t_1 . Consequently (19) holds for a.e. $t \geq t_1$, this again leads to a contradiction. Thus we obtain a priori bound:

$$(20) \quad \sup_{t \geq 0} |u(t)| \leq K_4 = \left[\{4 \varepsilon K_2 |\Omega| + 4J(u_0) + 2K_3\} |\Omega|^{\delta/2} / \varepsilon K_1 \right]^{1/(2+\delta)}$$

Since $d|u(t)|^2/dt = 2(u(t), u_t(t)) \leq 2|u(t)||u_t(t)|$, assertion (v) is derived from (iii), (iv) and integration of (18) over $[t, t+1]$. Assertion (vi) follows from (v) and the fact that $J(u(t)) \leq J(u_0)$ or $A(u(t)) \leq B(u(t)) + J(u_0)$. Q.E.D.

Before proceeding to the proof of Theorem I, we prepare the following lemmas.

Lemma 1. Let $r, q \in [1, \infty]$ and m be a non-negative integer.

Then

$$|u|_s \leq C |u|_{W^{m,r}}^a |u|_q^{1-a} \quad \forall u \in W^{m,r}(\Omega) \cap L^q(\Omega)$$

holds for any numbers $a \in [0, 1]$ and $s \in [1, \infty]$ satisfying

$$1/s = a(1/r - m/N) + (1-a)/q,$$

where C is a constant depending only on Ω, r, q, m and a .

For a proof of this lemma, see Friedmann [3].

Lemma 2. Let (i) of (f) be satisfied and u be a global solution of $(E)_N$ belonging to V . Then there exists a positive monotone decreasing function $T(\cdot)$ such that

$$(21) \quad \|u(t)\| \leq \|u(t_0)\| + 1 \quad \text{for all } t_0 \text{ and } t \in [t_0, t_0 + T(\|u(t_0)\|)].$$

Proof. First of all, we note that there exists a number $\lambda \in (0, 2]$ such that

$$(22) \quad |u|_{2(p-1)}^{2(p-1)} \leq C |u|_{H^2}^{2-\lambda} \|u\|^{2p-4+\lambda} \quad \forall u \in H^2(\Omega).$$

Indeed, for the case $N=1, 2$ or $N \geq 3$ and $2(p-1) \leq 2N/(N-2)$, we can take $\lambda = 2$. For the other case, we have only to apply Lemma 1 with $s = 2(p-1), m = r = 2$ and $q = 2N/(N-2)$ and use the

fact that $H^1(\Omega)$ is continuously embedded in $L^{2N/(N-2)}(\Omega)$.

This inequality implies that there exists a monotone increasing function $M(\cdot)$ such that

$$(23) \quad |g(\cdot, u)|^2 \leq \frac{1}{2} |\Delta u|^2 + M(\|u\|) \quad \forall u \in H^2(\Omega).$$

Multiplying (1)' by $-\Delta u(t) + u(t)$, we get

$$\frac{1}{2} \frac{d}{dt} \|u(t)\|^2 + |\Delta u(t)|^2 + |u(t)|^2 + 2 |\nabla u(t)|^2 \leq |g(\cdot, u)| |\Delta u(t)|.$$

Then, by (23), we obtain

$$\frac{1}{2} \frac{d}{dt} \|u(t)\|^2 \leq M(\|u(t)\|) \quad \text{for a.e. } t \in [0, \infty),$$

by which we can easily verify (21) by taking $T(r) = 1/2M(r+1)$.

Q.E.D.

Now we proceed to the proof of Theorem I.

Proof of (6) and (7). By virtue of (iii) of Proposition 1, there exists a positive number T_0 such that

$$(24) \quad \int_{T_0}^{\infty} |u_t(t)|^2 dt \leq \left(\frac{\alpha}{K_4}\right)^2 T(d_1).$$

Then we have

$$(25) \quad \|u(t)\| \leq d_1 + 1 \quad \text{for all } t \geq T_1 \equiv T_0 + \frac{K_4^2}{2\alpha}.$$

Suppose that this does not hold. Then there exists a $t_1 \geq T_1$ such that $\|u(t_1)\| > d_1 + 1$. Hence, by Fig. 1, $j(u(t_1)) > \alpha$.

Therefore there exists a $t_0 < t_1$ such that $j(u(t_0)) = \alpha$ and $j(u(t)) > \alpha$ for all $t \in (t_0, t_1]$. Then integration of (17) over $[t_0, t_1]$ and (20) give $t_1 - t_0 \leq K_4^2 / 2\alpha$, i.e., $t_0 \geq T_0$.

Again integrating (17) over $[t_0, t_1]$, we find

$$\begin{aligned} \alpha (t_1 - t_0) &\leq \int_{t_0}^{t_1} j(u(t)) dt \leq \int_{t_0}^{t_1} |u_t(t)| |u(t)| dt \\ &\leq K_4 (t_1 - t_0)^{1/2} \left(\int_{T_0}^{\infty} |u_t(t)|^2 dt \right)^{1/2}. \end{aligned}$$

Hence, by (24), $t_1 - t_0 \leq T(d_1)$. Since $\|u(t_0)\| \leq d_1$, Lemma 2 assures that $\|u(t_1)\| \leq d_1 + 1$. This is a contradiction. Thus (5)-(7) of Theorem I are verified.

Proof of (8). As was seen above, $u(t)$ can not stay in the region $\{u \in H^1(\Omega); j(u) > \alpha\}$ longer than $\gamma \equiv K_4^2 / 2\alpha$. Then for any $t \geq 0$, there exists $t_0 \in [\max(0, t-\gamma), t]$ such that $\|u(t_0)\| \leq d_3 \equiv \max(\|u_0\|, d_1)$. Therefore, in order to prove (8), it suffices to show that $\|u(t)\| \leq C_0$ for all $t \in I \equiv [t_0, t_0 + \gamma]$. For this purpose, we prepare several results on u .

Lemma 3. If $u \in L^r(I; L^r(\Omega))$, $r < 2^*$, then

$$\|u\|_{L^\infty(I; L^{r/2+1}(\Omega))} \leq C \left(\|u\|_{L^r(I; L^r(\Omega))}, d_0, d_3, r \right).$$

Proof. We note that

$$\begin{aligned} \frac{1}{s} \frac{d}{dt} |u(t)|_s^s &= (u_t(t), |u|^{s-2} u(t)) \leq |u_t(t)| |u(t)|_r^{s-1} \\ &\leq \frac{1}{2} (|u_t(t)|^2 + |u(t)|_r^r), \quad s = \frac{r}{2} + 1. \end{aligned}$$

Then integration of this over I and (iii) of Proposition 1 assure the assertion. Q.E.D.

Proposition 2. If $u \in L^\infty(I; L^r(\Omega))$ with $N(p-2)/2 < r < 2^*$, then

$$\|u\|_{L^\infty(I; H^1(\Omega))} \leq C \left(\|u_0\|_{L^\infty(I; L^r(\Omega))}, C_0 \right).$$

Proof. Multiplying (1)' by $|u|^{\ell-2}u(t)$ ($2 < \ell < 2^*$), we obtain, by (i) of (f),

$$(26) \quad \frac{1}{\ell} \frac{d}{dt} |u(t)|_{\ell}^{\ell} + \frac{4(\ell-1)}{\ell^2} \| |u(t)|^{\ell/2} \|^2 \leq K_0 (2 |u(t)|_{\frac{p+\ell-2}{p+\ell-2}}^{p+\ell-2} + |\Omega|).$$

For all $\ell > (p-2)N/2$, Lemma 1 with $s = 2(p+\ell-2)/\ell$, $m = 1$, $r = q = 2$ and u replaced by $|u|^{\ell/2}$ gives

$$(27) \quad |u|_{\frac{p+\ell-2}{p+\ell-2}}^{p+\ell-2} \leq C |u|^{p+\lambda\ell-2} \| |u|^{\ell/2} \|^2, \quad \lambda > 0.$$

Then, by integration of (26) over I and (27) with $\ell = r$, we get

$$(28) \quad \| |u|^{r/2} \|_{L^2(I; H^1(\Omega))} \leq C (\|u\|_{L^\infty(I; L^r(\Omega))}^r, \|u_0\|)$$

Again by Lemma 1 with $s = 2(N+2)/N$, $m = 1$, $r = q = 2$ and u replaced by $|u|^{r/2}$, we now deduce

$$(29) \quad |u|_{\frac{(2+N)r/N}{(2+N)r/N}}^{(2+N)r/N} \leq C |u|_r^{2/N} \| |u|^{r/2} \|^2.$$

Then it follows from (28) and (29) that $u \in L^{(2+N)r/N}(I; L^{(2+N)r/N}(\Omega))$. Hence, by Lemma 3, $u \in L^\infty(I; L^{(N+2)r/2N+1}(\Omega))$. Repeating this procedure, we observe that $u \in L^\infty(I; L^{r_i}(\Omega))$, where r_i are defined by the recurrence formula

$$r_{i+1} = \frac{N+2}{2N} r_i + 1, \quad i = 1, 2, \dots, \quad r_1 = r.$$

Since $r_i \rightarrow 2^*$ as $i \rightarrow \infty$ and $p < 2^*$, we can show by finite steps that $|u|_{L^\infty(I; L^p(\Omega))} \leq C (\|u\|_{L^\infty(I; L^r(\Omega))}^r, C_0)$. Thus the assertion follows from (ii) of Proposition 1. Q.E.D.

Proposition 3. For $N=1,2,3$ or 4 , we have

$$|u|_{L^\infty(I;L^q(\Omega))} \leq C_0 \quad \text{for all } q < q^*,$$

where $q^* = \infty$ for $N=1$ and $q^* = 2 + 8/(3N-4)$ for $N=2,3,4$.

Proof. Let $q_1 = 2$ and $q_{i+1} = (4-N)q_i/2N + (6N+8)/2N$, $i = 1,2,\dots$, and apply Lemma 1 with $s = q_{i+1}$, $m=1$, $r=2$ and $q = (2+q_i)/2$. Then we

have

$$|u|_{q_{i+1}}^{q_{i+1}} \leq C |u|_{(2+q_i)/2}^{q_{i+1}-4} \|u\|^4.$$

Therefore, by virtue of (vi) of Proposition 1 and Lemma 3,

$u \in L^{q_i}(I;L^{q_i}(\Omega))$ for all q_i . Since $q_i \rightarrow q^*$ as $i \rightarrow \infty$, we can derive the assertion by finite steps. Q.E.D.

Proposition 4. For $N \geq 5$, we have

$$|u|_{L^\infty(I;L^q(\Omega))} \leq C_0 \quad \text{for all } q < q_* = 3 + (N-4)(2-p)/4.$$

Proof. Since $p < 2^* \leq 4$ for $N \geq 4$ and (vi) of Proposition 1 assures that $|u|_{L^4(I;L^{2^*}(\Omega))} \leq C_0$, we get $|u|_{L^p(I;L^p(\Omega))} \leq C_0$.

Let $p_1 = p$, $p_{i+1} = (N-4)p_i/(N-2) + [8+(N-4)(2-p)]/(N-2)$, and

$s_i = 2 + p_i - p$, $i = 1,2,\dots$. Suppose that $u \in L^{p_i}(I;L^{p_i}(\Omega))$,

then by (26) with $\ell = s_i$, $|u|_{s_i/2}^{s_i/2} \in L^2(I;H^1(\Omega))$. Moreover

Lemma 1 with $s = 2p_{i+1}/s_i$, $m=1$, $r=2$, $q = 2 \cdot 2^*/s_i$ and u replaced by $|u|^{s_i/2}$ yields

$$|u|_{p_{i+1}}^{p_{i+1}} \leq C |u|_{2^*}^{\lambda_1} \| |u|^{s_i/2} \|^{\lambda_2} \quad \text{with } \lambda_1/2 + \lambda_2 \leq 2,$$

whence follows $|u|_{L^{p_{i+1}}(I;L^{p_{i+1}}(\Omega))} \leq C_0$. Since $p_i \rightarrow q_*$ as

$i \rightarrow \infty$, the assertion can be derived by finite steps. Q.E.D.

In order to prove (8), we have only to combine Propositions 2 with 3 (for the case $N \leq 4$) and 2 with 4 (for the case $N \geq 5$). In fact, $N(p-2)/2 < q^*$ or $N(p-2)/2 < q_*$ holds if and only if $p < 2_*$. Q.E.D.

Proof of Theorem II. We shall rely on Moser's iteration scheme to obtain L^∞ bound via H^1 bound. The following lemma plays an important role in this procedure.

Lemma 4. Let $w \in W_{loc}^{1,1}([0, \infty); L^2(\Omega)) \cap L_{loc}^\infty([0, \infty); L^\infty(\Omega) \cap H^1(\Omega))$

satisfy

$$(30) \quad \frac{d}{dt} |w(t)|_r^r + C_1 r^{-\theta_1} \| |w(t)|^{r/2} \|^2 \leq C_2 r^{\theta_2} (|w(t)|_r^{r+1}) \quad \text{a.e. } t \in [0, \infty)$$

for all $r \geq 2$, where $C_1 > 0$ and $C_2, \theta_1, \theta_2 \geq 0$.

Then there exist constants a, b, c, d such that

$$\sup_{t \geq 0} |w(t)|_\infty \leq a 2^{\theta_2 + (\theta_1 + \theta_2) b} M_0,$$

where $M_0 = \max(1, c |u_0|_\infty, \sup_{t \geq 0} |w(t)|^d)$.

Proof. When w belongs to $H^1(\Omega)$ for a.e. t , this is proved by Nakao [5] (See Lemma 3.1). By using Lemma 1 instead of Gagliardo-Nirenberg's inequality, one can prove this lemma by the same argument as in [5]. Q.E.D.

Put $\lambda_1 = r(1 - N(p-2)/2p)$, $\lambda_2 = p - 2$, $\lambda_3 = Nr(p-2)/2p$ and $\theta = Nr/(N-2)$. Then, by Hölder's inequality, we get

$$|u|_{p+r-2}^{p+r-2} \leq |u|_r^{\lambda_1} |u|_p^{\lambda_2} |u|_\theta^{\lambda_3}.$$

Applying Sobolev's embedding theorem and Young's inequality, we deduce

$$\begin{aligned}
 (31) \quad |u(t)|_{p+r-2}^{p+r-2} &\leq C |u(t)|_r^{\lambda_1} \| |u(t)|^{r/2} \|^{2\lambda_3/r} \\
 &\leq \frac{2(r-1)}{r^2} \| |u(t)|^{r/2} \|^2 + C r^{\frac{N(p-2)}{2p-(p-2)N}} |u(t)|_r^r,
 \end{aligned}$$

where C is a general constant depending on $\sup\{|u(t)|_p; t \geq 0\}$.

Then (31) and (26) with $\ell = r$ imply that $u(t)$ satisfies (30) with $C_1 = 1$, $\theta_1 = 0$ and $\theta_2 = 2p/(2p-(p-2)N)$. Thus (9) is verified by Lemma 4.

On the other hand, it is easy to show that there exists a positive number T_0 depending only on $|u_0|_\infty$ such that $|u(t)|_\infty \leq |u_0|_\infty + 1$ for all $t \in [0, 2T_0]$ and $\|u(T_0)\| \leq C(|u_0|_\infty)/T_0$. Hence, (10) and (11) follows from (7) and (8) respectively.

Q.E.D.

References

- 1 Alikakos, N.D., L^p bounds of solutions of reaction-diffusion equations, Comm. Partial Differential Equations 4 (1979), 827-868.
- 2 Cazenave, T. and Lions, P.L., Solutions globales d'équations de la chaleur semi linéaires, *ibid.*, 9 (1984), 955-978.
- 3 Friedmann, A., Partial Differential Equations, Holt, Rinehart and Winston, New York (1969).
- 4 Giga, Y., A bound for global solutions of semilinear heat equations, Comm. Math. Phys., 103 (1986), 415-421.
- 5 Nakao, M., L^p -estimate of solutions of some nonlinear degenerate diffusion equations, J. Math. Soc. Japan, 37 (1985), 41-64.
- 6 Ni, W.M., Sacks, P.E., and J. Tavantzis, On the asymptotic behavior of solutions of certain quasilinear equation of parabolic type, J. Differ. Equations 54 (1984), 97-120.
- 7 Ôtani, M., Existence and asymptotic stability of strong solution of nonlinear evolution equations with a difference term of subdifferentials, Colloquia Math. Soc. Janos Bolyai, 30, Qualitative theory of differential equations, North Holland, 1980.
- 8 Ôtani, M., 寄微分作用素の差の項をもつ発展方程式の解の漸近挙動について, 京都大学教理解析研究所 講究録 (1980)