


scientific reports



OPEN

Chitosan increases *Pinus pinaster* tolerance to the pinewood nematode (*Bursaphelenchus xylophilus*) by promoting plant antioxidative metabolism

Marta Nunes da Silva^{1,3}, Carla S. Santos^{1,3}, Ana Cruz¹, Adrián López-Villamor^{1,2} & Marta W. Vasconcelos¹

The pine wilt disease (PWD), for which no effective treatment is available at the moment, is a constant threat to *Pinus* spp. plantations worldwide, being responsible for significant economic and environmental losses every year. It has been demonstrated that elicitation with chitosan increases plant tolerance to the pinewood nematode (PWN) *Bursaphelenchus xylophilus*, the causal agent of the PWD, but the biochemical and genetic aspects underlying this response have not been explored. To understand the influence of chitosan in *Pinus pinaster* tolerance against PWN, a low-molecular-weight (327 kDa) chitosan was applied to mock- and PWN-inoculated plants. Nematode population, malondialdehyde (MDA), catalase, carotenoids, anthocyanins, phenolic compounds, lignin and gene expression related to oxidative stress (thioredoxin 1, *TRX*) and plant defence (defensin, *DEF*, and α -farnesene synthase, *AFS*), were analysed at 1, 7, 14, 21 and 28 days post-inoculation (dpi). At 28 dpi, PWN-infected plants elicited with chitosan showed a sixfold lower nematode population when compared to non-elicited plants. Higher levels of MDA, catalase, carotenoids, anthocyanins, phenolic compounds, and lignin were detected in chitosan-elicited plants following infection. The expression levels of *DEF* gene were higher in elicited plants, while *TRX* and *AFS* expression was lower, possibly due to the disease containment-effect of chitosan. Combined, we conclude that chitosan induces pine defences against PWD via modulation of metabolic and transcriptomic mechanisms related with plant antioxidant system.

Bursaphelenchus xylophilus, commonly known as the pinewood nematode (PWN), is the etiological agent of the pine wilt disease (PWD), which affects several *Pinus* spp., particularly the maritime pine *P. pinaster*¹. The PWD leads to important economic losses in the timber and wood industries, and increases the costs of disease management procedures and control, along with their inherent environmental impacts^{2–4}. Currently, large areas of forests damaged by PWD are located in Japan, South Korea, China, and southern Europe, and these are estimated to increase by 50% in the next 50 years⁵.

Following infection, nematodes move and feed on plant resin canals leading to loss of water conductivity in stems and, subsequently, decreased transpiration and photosynthesis in leaves, resulting in generalized oxidative damage^{6,7}. The production of hydrogen peroxide (H₂O₂) and superoxide ion radical (O²⁻), for example, has been associated with increased *B. xylophilus* pathogenicity⁸. To counteract the harmful effects of these molecules, plants activate several antioxidant enzymes, such as superoxide dismutase, peroxidases and catalase. However, if excessive oxidative stress occurs, plant cells may accumulate malondialdehyde (MDA), a secondary product of cell wall lipid peroxidation⁹, whose content was already demonstrated to increase in plants inoculated with PWN¹. In addition to the production of these metabolites, soluble phenolic compounds and lignin synthesis have also been implicated in plant defence against pathogens^{10,11}. Phenolic compounds production

¹Universidade Católica Portuguesa, CBQF - Centro de Biotecnologia e Química Fina – Laboratório Associado, Escola Superior de Biotecnologia, Rua de Diogo Botelho 1327, 4169-005 Porto, Portugal. ²Misión Biológica de Galicia (MBG-CSIC), Carballeira 8, Salcedo, 36143 Pontevedra, Spain. ³These authors contributed equally: Marta Nunes da Silva and Carla S. Santos. ✉email: masilva@porto.ucp.pt

and accumulation are associated with the browning of the leaf tissues injured by the PWN^{12–14}, but the role of these secondary metabolites is still not fully understood. In addition, lignin biosynthesis was found to occur in stem tissues during the advanced stages of the PWD, and constitutive lignin was suggested to be a mechanical barrier against nematode invasion, conferring higher plant tolerance to the pathogens^{5,16}. These enzymatic and other biochemical responses occur as soon as a few hours after infection, and are activated when a group of genes related to plant resistance recognizes pathogen effectors, initiating a resistance response¹⁷. In fact, genes related to secondary metabolite biosynthesis (α -farnesene synthase, *AFS*), defence against pathogens (defensins, *DEF*) and oxidative stress (thioredoxin, *TRX*) were found to be highly expressed in pine trees within a few hours after PWN infection¹⁸, especially when infected with virulent strains of *B. xylophilus*¹⁶.

Prevention is undoubtedly the best approach to reduce PWD incidence and different strategies have been suggested to avoid or treat the disease by: targeting the nematode itself, the vector, the host, or a combination of all three. Several synthetic compounds have already been developed to control the PWD, but many of them are toxic to the environment, labour intensive to apply, and expensive¹⁹. Previous studies suggested that chitosan can be used to enhance plant defence against bacteria^{20,21}, fungi²² and nematodes^{23,24}. Chitosan acts as a plant growth promoter, stimulating responses associated with both primary and secondary metabolism, including: carbon and nitrogen metabolism, primary photochemistry and photosynthesis, the tricarboxylic acid cycle, and terpenoid and phenolic compounds biosynthesis²⁵. For example, carotenoids, one of the most important plant terpenoids, were found to be increased in basil and strawberry tissues following chitosan-elicitation^{26,27}. Phenolic compounds, which play very important functions during pathogen infection (by e.g., providing mechanical strength through cell wall lignification processes) and in preventing oxidative stress, were also demonstrated to increase following elicitation with chitosan²⁸. In fact, total phenolic content, and anthocyanins in particular, have been found to accumulate with chitosan treatment in leaf tissues of lemon balm and basil, and in cell cultures of grape vines^{29–31}. Chitosan application also enhances the accumulation of several enzymatic antioxidants, including catalase, decreasing malondialdehyde (MDA) levels in leaf tissues thus improving plant antioxidant status^{30,32}.

The use of chitosan in plant protection against pests and pathogens presents various advantages compared with the currently employed control compounds, as it is physically and biologically functional, biodegradable and biocompatible with tissues and cells^{33,34}. Khalil et al.²³ described the nematocidal activity of chitosan with different molecular weights (MW) against *Meloidogyne incognita* in tomato seedlings, reporting that low MW had the highest efficacy in controlling pathogen progression. Previous studies on the effect of chitosan as a control agent against the PWN demonstrated that low MW components had the highest nematocidal effect, reducing nematode density inside *P. pinaster* tissues up to 24 days post-inoculation (dpi)²⁴. More recently, it has been demonstrated that *P. pinaster* and *P. pinea* supplemented with diazotrophic bacteria and a chitosan-producing fungus, *Cunninghamella elegans*, showed a 36-fold reduction of nematode colonization when compared to non-supplemented plants, as well as improved photosynthetic pigments, water content and phenolic compounds biosynthesis³⁵. Despite the promising pieces of evidence that chitosan may be a useful tool in the control of the PWD, the physiological, metabolic and genetic mechanisms induced by chitosan application to PWN-infected *Pinus* spp. have not been addressed yet. The lack of knowledge regarding the regulatory mechanisms involved in the crosstalk between *Pinus* spp. plants and PWN after plant elicitation with chitosan greatly hinders the efforts to develop evidence-based, sustainable and affordable control methods against the PWD.

As such, in the present work, we hypothesized that the protective action of chitosan in PWN-infected plants might be due to systemic acquired resistance-related responses associated with a readjustment of the plant's oxidative status. To test this hypothesis, the potential of low MW chitosan against the PWD in one-year-old *P. pinaster* plants was analysed, through the evaluation of the effect of chitosan on (1) nematode population dynamics, (2) plant antioxidative system and secondary metabolites accumulation, and (3) gene expression related to oxidative stress and defence responses.

Results

Nematode population in infected plants. Total nematode count inside stem tissues of inoculated plants without chitosan elicitation significantly increased by 44-fold from 1 to 28 dpi, indicating a successful infection and multiplication of PWN inside plant tissues over time (Fig. 1). In contrast, with chitosan elicitation the number of nematodes was low and statistically identical over the entire experimental period. In fact, only about 0.14-fold of the number of nematodes found in non-elicited plants were recovered from plants inoculated in the presence of chitosan, being the effect of the chitosan treatment significant ($P < 0.0001$) (Table 1).

Oxidative stress-related mechanisms. Plants inoculated with PWN displayed a progressive and significant increase in MDA concentration throughout the experimental period, regardless of chitosan elicitation (Fig. 2A), with a significant effect of the chitosan treatment and the DPI (Table 1). In fact, at the end of the experimental period, PWN-inoculated plants elicited with chitosan had significantly higher MDA concentration than non-elicited plants (by 1.8-fold). Similarly to MDA, during the first 14 days of infection, catalase activity was not significantly affected in PWN-infected plants, regardless of chitosan elicitation, which remained similar to catalase levels in water-inoculated plants (Fig. 2B). Non-elicited inoculated plants presented a progressive increase in catalase levels and, at 28 dpi, these were 4.5-fold significantly higher than at 1 dpi. In chitosan-elicited plants, a more drastic and significant increase was registered after 14 dpi, with catalase activity increasing by 25-fold at 21 dpi, resulting in a significant interaction between chitosan treatment and DPI (Table 1).

Plants behaved similarly regarding the accumulation of carotenoids (Fig. 3A) and anthocyanins (Fig. 3B). From 1 to 28 dpi PWN-infected plants without chitosan elicitation showed a progressive and significant increase of 15- and sevenfold in carotenoid and anthocyanin concentrations, respectively. In chitosan-elicited plants, carotenoid and anthocyanin concentrations did not vary significantly between control and PWN-infected plants.




Figure 1. Nematode population in plant tissues. Nematode density in *P. pinaster* stem tissues 1, 7, 14, 21 and 28 days post-inoculation (dpi). Plants were inoculated with the pinewood nematode (PWN) *B. xylophilus* 14 days following the application of 0.5% acetic acid (without chitosan, non-elicited controls) or a chitosan solution to plants' substrate. Data represents the difference between the averages of PWN-inoculated and water-inoculated plants \pm SE, and different letters indicate statistically different means at $P < 0.05$.

	Chitosan treatment		DPI		Interaction	
	F _(1,32)	P value	F _(4,32)	P value	F _(4,32)	P value
Nematode population	32.60	<0.0001	19.75	<0.0001	13.42	<0.0001
MDA	F _(1,26)	P value	F _(4,26)	P value	F _(4,26)	P value
	13.94	0.0009	14.89	<0.0001	0.7780	0.5496
Catalase activity	F _(1,40)	P value	F _(4,40)	P value	F _(4,40)	P value
	37.18	<0.0001	34.58	<0.0001	12.16	<0.0001
Carotenoids	F _(1,35)	P value	F _(4,35)	P value	F _(4,35)	P value
	0.013	0.9116	13.02	<0.0001	2.906	0.0355
Anthocyanins	F _(1,30)	P value	F _(4,30)	P value	F _(4,30)	P value
	0.324	0.5737	25.21	<0.0001	4.488	0.0058
Total Polyphenols	F _(1,33)	P value	F _(4,33)	P value	F _(4,33)	P value
	19.24	0.0001	14.25	<0.0001	1.219	0.3215
Lignin	F _(1,31)	P value	F _(4,31)	P value	F _(4,31)	P value
	7.172	0.012	11.45	<0.0001	2.415	0.0699
Gene expression	F _(1,20)	P value	F _(4,20)	P value	F _(4,20)	P value
<i>TRX1</i>	34.37	<0.0001	72.75	<0.0001	42.92	<0.0001
<i>DEF</i>	7.522	0.0125	10.83	<0.0001	15.53	<0.0001
<i>AFS</i>	10.10	0.0047	29.26	<0.0001	20.77	<0.0001

Table 1. Effect of chitosan treatment, time-points of analysis (days post-inoculation, DPI) and their interaction in the physiological, biochemical and gene expression analyses of control and inoculated *P. pinaster* plants, following elicitation with chitosan.

Hence, no significant effect was found for the chitosan treatment for both analytes (Table 1). However, their concentration abruptly increased at 28 dpi by 21-fold in carotenoids concentration and 13-fold in anthocyanins concentration (when compared to the 21 dpi time-point), with a significant effect of the DPI and its interaction with the chitosan treatment (Table 1).

For all time-points, total phenolic concentration was about two times higher in PWN-infected plants with chitosan elicitation ($P < 0.0001$), when compared to non-elicited plants (Fig. 4A, Table 1). At 14 dpi, plants treated with chitosan presented the highest levels of total phenolic compounds. Moreover, in both elicited and non-elicited plants, polyphenols concentration was drastically reduced in infected plants at the end of the assay (28 dpi), returning to concentration levels similar to non-infected plants. In general, lignin concentration was




Figure 2. Lipid peroxidation and antioxidant activity. Malondialdehyde concentration and catalase activity in *P. pinaster* leaf tissues 1, 7, 14, 21 and 28 days post-inoculation (dpi) with the pinewood nematode (PWN) *B. xylophilus*. Plants were inoculated 14 days following the application of 0.5% acetic acid (without chitosan, non-elicited controls) or a chitosan solution to plants' substrate. Data represents the difference between the averages of PWN-inoculated and water-inoculated plants \pm SE, and different letters indicate statistically different means at $P < 0.05$.




Figure 3. Carotenoids and anthocyanins. Carotenoids and anthocyanins concentrations in *P. pinaster* leaf tissues 1, 7, 14, 21 and 28 days post-inoculation (dpi) with the pinewood nematode (PWN) *B. xylophilus*. Plants were inoculated 14 days following the application of 0.5% acetic acid (without chitosan, non-elicited controls) or a chitosan solution to plants' substrate. Data represents the difference between the averages of PWN-inoculated and water-inoculated plants \pm SE, and different letters indicate statistically different means at $P < 0.05$.




Figure 4. Total phenolic compounds and lignin. Total phenolic compounds and lignin concentrations in *P. pinaster* stem tissues 1, 7, 14, 21 and 28 days post-inoculation (dpi) with the pinewood nematode (PWN) *B. xylophilus*. Plants were inoculated 14 days following the application of 0.5% acetic acid (without chitosan, non-elicited controls) or a chitosan solution to plants' substrate. Data represents the difference between the averages of PWN-inoculated and water-inoculated plants \pm SE, and different letters indicate statistically different means at $P < 0.05$.

similar between PWN-infected plants regardless of chitosan elicitation (Fig. 4B). However, at 28 dpi, inoculated plants showed increased lignin concentration (as compared with water-inoculated control plants), particularly with chitosan elicitation (by ca. twofold). No significant interaction between chitosan treatment and DPI was found for both total phenolic compounds and lignin accumulation (Table 1).

Relative expression of stress-related genes. Regarding the expression of *TRX1* gene (Fig. 5A), non-elicited PWN-infected plants showed an increase by more than twofold at 7 dpi, which was followed by an eightfold increase at 14 dpi, when compared to 1 dpi. The expression decreased to less than half of the values seen at 14 dpi, and remained at this level until the end of the experiment. For elicited PWN-infected plants, *TRX1* relative expression was also significantly increased at 7 dpi, however, it immediately decreased in the following time-points and was generally lower than the expression values in non-elicited plants. Following PWN infection, *DEF* expression (Fig. 5B) was differently modulated throughout the experiment depending on the chitosan treatment ($P < 0.05$). In non-elicited plants, from 1 to 7 dpi it was significantly reduced to almost null values and maintained at this level until 21 dpi, until its expression increased again, by about twofold, at 28 dpi. In elicited plants, there was a significant decrease in *DEF* expression at 7 dpi, but at 14 dpi its expression increased again, remaining unaltered until the end of the experimental period. Chitosan elicitation significantly impacted the expression pattern of *AFS* (Fig. 5C), with non-elicited PWN-infected plants having a progressive increase in its expression over time, reaching the highest value at 28 dpi (ninefold higher than at 1 dpi). Elicited plants showed constant and relatively low *AFS* expression throughout the experiment, which was twofold higher in PWN-infected plants than in controls. Chitosan treatment and DPI had a significant effect in the expression of all genes, also having a significant interaction among them (Table 1).

Discussion

When the PWN infects susceptible plant species, there is a primary phase of the disease (early phase), where the nematode population remains low, without producing any visual symptoms. When the second phase develops (advanced stage), PWN population increases exponentially, beginning a more destructive phase for the plant due to nematode feeding and reproduction on the epithelial cells^{6,7,36}. Previous histological and biochemical works on the PWN population dynamics demonstrated that the PWD progresses strongly between 7 and 14 days after infection^{37,38}, which is in accordance to what was observed in the present work in the absence of chitosan elicitation. It has been previously shown that chitosan application prevents nematode multiplication in plant tissues over time²⁴, and reduces the number of juveniles in soil³⁹. The present work seems to attest for the promising role of chitosan elicitation of *P. pinaster* plants against the PWD, as the nematode population remained at low levels throughout the entire experimental period, indicating that this compound prevented, or, at least, slowed disease progression.

In susceptible *Pinus* spp. host plants, PWN damages xylem and phloem parenchyma and cortex cells, which can compromise water transport, lead to the peroxidation of the unsaturated lipids of cellular membranes and to ROS formation, culminating in leaf necrosis and often in plant death^{7,40}. MDA, a secondary compound of lipid peroxidation reactions, is frequently used as an indicator of cell damage induced by free radicals⁴¹. Interestingly, despite preventing nematode reproduction in plants tissues, chitosan did not prevent cellular damage, leading to increased MDA accumulation. This could be a result of the abrupt induction of plant defence mechanisms upon PWN infection, with consequent formation of ROS⁴². To counteract the negative effects of intracellular ROS levels, increased or de novo synthesis of several antioxidant enzymes occur. One of those enzymes, catalase, can capture H_2O_2 and convert it in water and oxygen molecules, which do not have a damaging impact to plant cells^{43,44}. Here, following PWN inoculation, catalase activity was significantly induced throughout the progression of the disease, although more drastically in chitosan-elicited plants, putatively representing the activation of plants' antioxidant defences. Hydrogen peroxide-mediated oxidative burst has been observed in several plant species supplemented with chitosan, and it is generally acknowledged that it is required for chitosan-induced defence responses^{25,45}. Increased H_2O_2 content in chitosan-elicited plants in generally accompanied by the accumulation of several antioxidant enzymes, among which catalase and peroxidase seem to be the most important^{46–48}. The application of chitosan may, therefore, lead to a stronger, and possibly more efficient, antioxidant response against PWN infection, which may contribute to an increased tolerance against the pathogen.

Carotenoids and anthocyanins are metabolites involved in plant protection by preventing photosystem damage, participating in signalling, ROS scavenging and inhibiting the growth of noxious microorganisms^{49–51}. It appears that their concentration increased in a similar manner to the PWN population, MDA concentration and catalase activity, as they gradually increased in inoculated plants (as compared to the controls), in both elicited and non-elicited plants. Previous studies have reported that chitosan-elicitation promotes plant defences through increased activity of several defence-related enzymes, e.g. phenylalanine ammonia-lyase, involved in the biosynthesis of phenolic compounds and terpenoids (which include, among several others, carotenoids and anthocyanins)^{52,53}. We hypothesise that increased concentrations of carotenoids and anthocyanins in PWN-infected tissues, especially by the end of the experimental period, may be a coping mechanism to prevent further cellular damage caused by the pathogen⁵⁴. As observed for catalase, chitosan elicitation seems to promote the accumulation of these metabolites, thus corroborating their potential in improving plants' antioxidant defences. Along with carotenoids and anthocyanins biosynthesis, phenolics and lignin are also involved in plant resistance to biotic stress^{15,55,56}. In general, total phenolic compounds concentration increased at 14 dpi (with statistical significance in PWN-infected elicited plants), which seems to correspond to the beginning of the advanced stage of the PWD. This is concordant with a previously reported positive relationship found between nematode migration and phenolics concentration⁵⁷. In the case of lignin, chitosan-elicited plants had higher accumulation than non-elicited ones, which may indicate an intensified tissue lignification after pathogen infection, possibly




Figure 5. Stress-related gene expression. Fold expression of genes *thioredoxin H1* (TRX1) (A), *defensin* (DEF) (B) and *α -farnesene synthase* (AFS) (C) in plants inoculated with *B. xylophilus* relative to their expression in water-inoculated plants, 1, 7, 14, 21 and 28 days post-inoculation (dpi). Plants were inoculated 14 days following the application of 0.5% acetic acid (without chitosan, non-elicited controls) or a chitosan solution to plants' substrate. Data represents the difference between the averages of PWN-inoculated and water-inoculated plants \pm SE, and different letters indicate statistically different means at $P < 0.05$.

increasing plant tolerance to the pathogen⁵⁸. Chitosan also induces a generalized H₂O₂-mediated hypersensitive response, involved in cell wall lignification processes that confer protection against pathogen-induced mechanical damage^{30,59}. The higher accumulation of carotenoids, anthocyanins, phenolics and lignin observed in the present study, particularly during the later stages of PWN infection and with chitosan elicitation, are in agreement in these previous works, and attest the potential of chitosan elicitation in promoting *Pinus* spp. defences against the pathogen.

In plants, thioredoxins act as signalling molecules through their reduction-regulatory properties in response to pathogen attacks^{60,61}. In the present work, significant overexpression of *TRX1* was observed in non-elicited plants at 14 dpi, corroborating the hypothesis that the advanced stage of the PWD was triggered around this time-point. In PWN-infected elicited plants, *TRX1* expression levels were not induced at this time-point, which is coherent with the lower nematode population and increased antioxidative activity previously discussed. Plant defensins, on the other hand, are usually associated with antimicrobial and antifungal activity, constituting the first defensive line against pathogens^{62,63}. It is interesting to note that, here, *DEF* expression pattern was very distinct between chitosan-elicited and non-elicited plants following PWN infection. It seems that, without chitosan elicitation, plants were not able to activate the expression of this gene as the PWN population progressed, whereas in elicited plants *DEF* expression was significantly increased along time, possibly contributing to the lower extent of disease progression observed. Also involved in plant defence is α -farnesene (*AFS*), an herbivore-induced plant volatile, usually increased in plants suffering pathogen attacks, being key in disease resistance^{18,64}. Here, non-elicited plants which showed increased nematode population over time, also displayed higher levels of *AFS* expression, particularly at 28 dpi, when the highest number of nematodes were registered. Contrastingly, chitosan-elicited plants had lower levels of *AFS* expression, possibly due to the low nematode progression and multiplication within their tissues.

In general, chitosan increased *P. pinaster* tolerance to the PWN, resulting in decreased nematode density, increased accumulation of metabolites involved in antioxidant activity, and differential expression of stress-related gene expression. This further supports the potential of chitosan in the prevention and/or treatment of the PWD in *Pinus* spp., which should be further explored and considered for the development of sustainable disease control strategies.

Methods

Plant maintenance and chitosan elicitation. One-year-old (40–50 cm height) *P. pinaster* plants were maintained in a growth chamber (Fitoclima 10 000 EHF; Aralab, Rio de Mouro, Portugal) under a 16 h light/8 h darkness photoperiod at 25/18 °C, respectively, and 80% relative humidity. Photon flux density during the day was 380 $\mu\text{mol m}^{-2} \text{s}^{-1}$. Plants were kept in a commercial substrate (COMPO SANA universal substrate; Compo GmbH, Münster, Germany) composed of (mg L^{-1}): 200–450 N; 200–500 P_2O_5 ; 300–550 K_2O , pH 5.0–6.5. Plants were watered weekly until field capacity.

An acidic solution (0.5% of acetic acid, pH 6.0) of 4.4% chitosan (molecular weight 327 kDa, deacetylation degree $\geq 75\%$, Ref.: 448869, Sigma-Aldrich, Missouri, USA) was prepared and left shaking for 24 h before use. A single chitosan treatment was applied to a group of 75 plants by adding 80 mL the chitosan solution to the substrate of each plant. An additional group of 75 plants served as non-elicited control, where a 0.5% solution of acetic acid (no chitosan) was added to the soil.

Nematode culture and plant inoculation. Fourteen days following plant elicitation, a virulent strain of PWN (65 GO, isolated from Góis, Portugal) was inoculated into plant stems. Nematodes were maintained in agar plates with *Botrytis cinerea* mycelia for 14 days, after which they were extracted from the growing media using the Baermann funnel technique⁶⁵. Plant inoculation was performed with 2 000 nematodes at *ca.* 3 cm from the top of each plant as described by Nunes da Silva et al.²⁴.

Experimental design and plant sampling. A total of 150 *P. pinaster* plants were used in this experiment. Chitosan was applied to a group of 75 plants, in which 25 were mock-inoculated with water (non-inoculated controls) and 50 were inoculated with PWN. An additional group of 75 plants served as non-elicited control, where a 0.5% solution of acetic acid (without chitosan) was added to the substrate: 25 plants were mock-inoculated with water (non-inoculated controls) and 50 were inoculated with PWN. At 1, 7, 14, 21 and 28 days post-inoculation (dpi) five biological replicates of each treatment were sampled. At each time-point, leaves were recovered for the quantification of lipid peroxidation (MDA), catalase, carotenoids and anthocyanins, and stems were used for total soluble phenolics and lignin quantification and gene expression analysis. Five additional PWN-inoculated plants were sampled at the same time-points for whole-stem PWN quantification.

Nematode quantification in plant tissues. In each time-point, five inoculated plants from each treatment were randomly selected for PWN quantification. Leaves were removed, and the entire stem was cut into small portions (*ca.* 0.5 cm) and used for nematode quantification using the Baermann funnel technique for 24 h at 25 °C. Nematode density in stem tissues was estimated taking into consideration whole stem fresh weight.

Lipid peroxidation and catalase. Quantification of MDA, a sub-product of lipid peroxidation, was performed by mixing vigorously 100 mg of leaf tissue previously homogenized in liquid nitrogen with 10 mL of 0.5% thiobarbituric acid in 20% of trichloroacetic acid and incubated at 100 °C for 30 min. After the incubation period, the reaction was terminated by transferring the tubes into ice. Samples were centrifuged during 10 min at 5 000 g and the supernatant was filtered, after which absorbances were measured at 450, 532 and 600 nm and MDA estimated as described by Yang et al.⁶⁶.

Quantification of catalase activity was performed following the protocol by Ruley et al.⁶⁷, adding 1.5 mL of phosphate buffer 1 M to 100 mg of leaf tissue previously homogenized with liquid nitrogen. Samples were vigorously mixed for 2 min, centrifuged at 5 000 g for 10 min at 4 °C and the supernatant was recovered and diluted 3 times with phosphate buffer. Three hundred and forty-four microlitres of a 73 mM H_2O_2 solution (in 0.5 M Tris/HCl pH 7.0) were then added to 666 μL of sample extract. Enzyme activity was measured for 3 min at 240 nm. One unit of enzyme corresponded to a decrease of 0.001 in absorbance.

Plant secondary metabolites. For carotenoids and anthocyanins quantification, leaves sampled as previously described (100 mg) were homogenized with liquid nitrogen and extracted with 10 mL of cold acetone/Tris buffer solution at 1 M (80:20 v/v, pH 7.8). Samples were incubated at 4 °C for 72 h and absorbances were recorded at 470, 537, 647 and 663 nm. Metabolite concentration was calculated as described by Sims and Gamon⁶⁸.

Total soluble phenolic compounds were quantified as in Azevedo et al.⁶⁹. Lyophilized leaf tissues previously homogenized with liquid nitrogen (100 mg) were extracted with 5 mL of methanol for 24 h in the dark at 4 °C. One hundred µL of sample extract were mixed with 5 mL of ultrapure water and 0.5 mL of Folin-Denis reagent, stirred vigorously and incubated at room temperature for 5 min. After incubation, 1.5 mL of sodium carbonate at 35% (w/v) was mixed with each sample and let to react in the dark for 2 h, after which 2.9 mL of ultrapure water were added and absorbances were determined at 760 nm. Soluble phenolic compounds were determined using a quercetin calibration curve.

Lignin was determined by the acetyl bromide method⁷⁰ using the leaf biomass used for phenolics extraction, which was subjected to sequential 24 h extractions with water, acetone and hexane. Samples were dried at 60 °C for 48 h, and 10 mg of biomass were mixed with 500 µL of glacial acetic acid and 500 µL of acetyl bromide 25%. The mixture was digested at 50 °C for 2 h with vigorous stirring, after which samples were centrifuged for 10 min at 15 000 g. One hundred microlitre of sample were recovered to a new microcentrifuge tube and homogenized with 200 µL of glacial acetic acid, 150 µL of NaOH 0.3 M, 50 µL of hydroxylamine hydrochloride 0.5 M and 500 µL of glacial acetic. Absorbances were recorded at 280 nm and lignin concentration in each sample was determined through a standard calibration curve.

Gene expression analysis. From the five biological replicates from each treatment, three were randomly selected for gene expression analysis. After stem tissue homogenization with liquid nitrogen, RNA extraction was performed using the RNeasy Mini Kit (QIAGEN GmbH, Hilden, Germany) following the manufacturer's protocol. RNA quality and quantity were determined spectrophotometrically and single-strand cDNA was synthesised using iScript cDNA Synthesis Kit (Bio-Rad Laboratories, Inc, California, USA) following the manufacturer's instructions. The transcript levels of genes encoding proteins related to oxidative stress (thioredoxin 1, *TRX*) and plant defence (defensin, *DEF*; α-farnesene synthase, *AFS*) were analysed. Primer sequences were retrieved from available published literature¹⁸. Reactions of RT-qPCR were carried out in a thermal cycler CFX96 Touch Deep Well Real-Time PCR Detection System using iQ SYBR Green Supermix (Bio-Rad Laboratories, Inc, California, USA) and data visualized with the software Bio-Rad CFX Manager 3.1. The amplification protocol cycle was 95 °C for 3 min and 40 cycles at 95 °C for 15 s, 30 s at 54 °C and 71 °C for 30 s. The comparative CT method ($\Delta\Delta CT$) was used for the relative quantification of gene expression values using the geometric mean of the expression of the control transcripts ubiquitin and 18S ribosomal RNA genes, and the plants inoculated with water (non-inoculated controls) as reference sample⁷¹. For each sample and target gene, two technical replicates were analysed.

Statistical analysis. Data were analysed using GraphPad Prism v6.0 software (GraphPad Software, California, USA) and significant differences between chitosan treatments were determined using two-way analysis of variance (ANOVA) followed by Tukey's post hoc test with $P=0.05$. Due to the large number of datasets, and to facilitate result viewing and discussion, results concerning MDA, catalase, carotenoids, anthocyanins, phenolic compounds and lignin are presented as the difference between the averages of PWN- and water-inoculated plants. Bars showing positive values represent an increase in PWN-inoculated plants relatively to water-inoculated plants, whereas negative results represent a decrease. Propagation of uncertainty was used to calculate the standard error of each value.

Received: 11 September 2020; Accepted: 1 February 2021

Published online: 12 February 2021

References

- Nunes da Silva, M., Solla, A., Sampedro, L., Zas, R. & Vasconcelos, M. Susceptibility to the pinewood nematode (PWN) of four pine species involved in potential range expansion across Europe. *Tree Physiol.* **35**(9), 987–999 (2015).
- Mamiya, Y. Pine wilt disease. In *Pine Wilt Disease in Japan* (eds Mota, M. & Vieira, P.) 9–20 (Springer, New York, 2004).
- Yang, L., Wang, Z., Hou, Y., Han, R. & Sun, Y. Effects of Cu²⁺ on wheat seedlings exposed to enhanced ultraviolet-b radiation. *Am. J. Plant Sci.* **5**, 3060–3065 (2014).
- Shimazu, M. Current status on research and management of pine wilt disease in Japan. Current status on research and management of pine wilt disease, International Symposium, October 20. Korea Forest Research Institute, Seoul, Korea: 1–18 (2006).
- Hirata, A. et al. Potential distribution of pine wilt disease under future climate change scenarios. *PLoS ONE* **12**, e0182837. <https://doi.org/10.1371/journal.pone.0182837> (2017).
- Kuroda, K. Pine wilt disease. In *Physiological Incidences Related to Symptom Development and Wilting Mechanism* (eds Zhao, B. G. et al.) 204–222 (Springer, New York, 2008).
- Yamada, T. Pine wilt disease. In *Biochemical Responses in Pine Trees Affected by Pine Wilt Disease* (eds Zhao, B. G. et al.) 223–234 (Springer, New York, 2008).
- Zhang, W. et al. Enhancement of oxidative stress contributes to increased pathogenicity of the invasive pine wood nematode. *Philos. Trans. R. Soc. Lond. B* **374**, 20180323. <https://doi.org/10.1098/rstb.2018.0323> (2019).
- Heath, R. L. & Packer, L. Photoperoxidation in isolated chloroplasts. I. Kinetics and toichiometry of fatty acid peroxidation. *Arch. Biochem. Biophys.* **125**, 189–198 (1968).
- Treutter, D. Significance of flavonoids in plant resistance: A review. *Environ. Chem. Lett.* **4**, 147–157 (2006).
- Liu, Q., Luo, L. & Zheng, L. Lignins: Biosynthesis and biological functions in plants. *Int. J. Mol. Sci.* **19**(2), 335. <https://doi.org/10.3390/ijms19020335> (2018).
- Mamiya, Y. Inoculation of the first year pine (*Pinus densiflora*) seedlings with *Bursaphelenchus lignicolus* and the histopathology of diseased seedlings. *J. Jap. For. Soc.* **62**, 176–183 (1980).

13. Myers, R. F. Cambium destruction in conifers caused by pinewood nematodes. *J. Nematol.* **18**, 398–402 (1986).
14. Futai, K. Abnormal metabolites in pine wood nematode-inoculated Japanese black pine. *Jap. J. Nematol.* **33**, 45–56 (2003).
15. Kawaguchi, E. Relationship between the anatomical characteristics of cortical resin canals and migration of *Bursaphelenchus xylophilus* in stem cuttings of *Pinus thunbergii* seedlings. *J. Jap. Soc. Hort. Sci.* **88**, 240–244 (2006).
16. Lee, I. H. *et al.* Comparative transcriptome analysis of *Pinus densiflora* following inoculation with pathogenic (*Bursaphelenchus xylophilus*) or non-pathogenic nematodes (*B. thailandae*). *Sci. Rep.* **9**, 12180. <https://doi.org/10.1038/s41598-019-48660-w> (2019).
17. Schiffer, R. *et al.* Tissue dependence and differential cordycep sensitivity of race-specific resistance responses in the barley-powdery mildew interaction. *Mol. Plant-Microbe Interact.* **10**, 830–839 (1997).
18. Santos, C. S., Pinheiro, M., Silva, A., Egas, C. & Vasconcelos, M. W. Searching for resistance genes to *Bursaphelenchus xylophilus* using high throughput screening. *BMC Genom.* **13**, 599. <https://doi.org/10.1186/1471-2164-13-599> (2012).
19. Kamata, N. Pine wilt disease. In *Integrated Pest Management of Pine Wilt Disease in Japan: Tactics and Strategies* (eds Zhao, B. G. *et al.*) 304–322 (Springer, New York, 2008).
20. Tikhonov, V. E. *et al.* Bactericidal and antifungal activities of a low molecular weight chitosan and its N-(2(3)-(dodec-2-enyl)succinoyl)-derivatives. *Carbohydr. Polym.* **64**, 66–72 (2006).
21. Rabea, E. I. & Steurbaut, W. Chemically modified chitosans as antimicrobial agents against some plant pathogenic bacteria and fungi. *Plant Protect. Sci.* **4**, 149–158 (2010).
22. Trotel-Aziz, P., Couderchet, M., Vernet, G. & Aziz, A. Chitosan stimulates defense reactions in grape vine leaves and inhibits development of *Botrytis cinerea*. *Eur. J. Plant Pathol.* **114**, 405–413 (2006).
23. Khalil, M. S. & Badawy, M. E. I. Nematicidal activity of a biopolymer chitosan at different molecular weights against root-knot nematode *Meloidogyne incognita*. *Plant Protect. Sci.* **48**(4), 170–178 (2012).
24. Nunes da Silva, M. *et al.* Chitosan as a biocontrol agent against the pinewood nematode (*Bursaphelenchus xylophilus*). *For. Pathol.* **44**(5), 420–423 (2014).
25. Mukhtar Ahmed, K. B., Khan, M. M., Siddiqui, H. & Jahan, A. Chitosan and its oligosaccharides, a promising option for sustainable crop production: A review. *Carbohydr. Polym.* **227**, 115331. <https://doi.org/10.1016/j.carbpol.2019.115331> (2020).
26. Pirbalouti, A. G., Malekpoor, F., Salimi, A. & Golparvar, A. Exogenous application of chitosan on biochemical and physiological characteristics, phenolic content and antioxidant activity of two species of basil (*Ocimum ciliatum* and *Ocimum basilicum*) under reduced irrigation. *Sci. Hort.* **217**, 114–122 (2017).
27. Rahman, M. *et al.* Chitosan biopolymer promotes yield and stimulates accumulation of antioxidants in strawberry fruit. *PLoS ONE* **13**(9), e0203769. <https://doi.org/10.1371/journal.pone.0203769> (2018).
28. Mehregan, M., Mehrafarin, A., Labbafi, M. R. & Naghdi Badi, H. Effect of different concentrations of chitosan biostimulant on biochemical and morphophysiological traits of stevia plant (*Stevia rebaudiana* Bertoni). *J. Med. Plants Res.* **2**(62), 169–181 (2017).
29. Cai, Z., Kastell, A., Mewis, I., Knorr, D. & Smetanska, I. Polysaccharide elicitors enhance anthocyanin and phenolic acid accumulation in cell suspension cultures of *Vitis vinifera*. *Plant Cell Tissue Organ Cult.* **108**, 401–409 (2011).
30. Hawrylak-Nowak, B., Dresler, S., Rubinowska, K. & Matraszek-Gawron, R. Eliciting effect of foliar application of chitosan lactate on the phytochemical properties of *Ocimum basilicum* L. and *Melissa officinalis* L. *Food Chem.* **342**, 128358. <https://doi.org/10.1016/j.foodchem.2020.128358> (2020).
31. Singh, R. K., Martins, V., Soares, B., Castro, I. & Falco, V. Chitosan application in vineyards (*Vitis vinifera* L. cv. Tinto Cão) induces accumulation of anthocyanins and other phenolics in berries, mediated by modifications in the transcription of secondary metabolism genes. *Int. J. Mol. Sci.* **21**(1), 306. <https://doi.org/10.3390/ijms21010306> (2020).
32. Zong, H., Liu, S., Xing, R., Chen, X. & Li, P. Protective effect of chitosan on photosynthesis and antioxidative defense system in edible rape (*Brassica rapa* L.) in the presence of cadmium. *Ecotoxicol. Environ. Saf.* **138**, 271–278 (2017).
33. Badawy, M. E. I. & Rabea, E. I. A biopolymer chitosan and its derivatives as promising antimicrobial agents against plant pathogens and their applications in crop protection. *Int. J. Carbohydr. Chem.* **2011**, 460381. <https://doi.org/10.1155/2011/460381> (2011).
34. Rendina, N., Nuzzaci, M., Scopa, A., Cuypers, A. & Sofo, A. Chitosan-elicited defense responses in Cucumber mosaic virus (CMV)-infected tomato plants. *J. Plant Physiol.* **234–235**, 9–17 (2019).
35. Nunes da Silva, M., Pintado, M., Sarmiento, B., Stamford, N. P. & Vasconcelos, M. W. A biofertilizer with diazotrophic bacteria and a filamentous fungus increases *Pinus pinaster* tolerance to the pinewood nematode (*Bursaphelenchus xylophilus*). *Biol. Control* **132**, 72–80 (2019).
36. Myers, R. F. Pathogenesis in pine wilt caused by pinewood nematode, *Bursaphelenchus xylophilus*. *J. Nematol.* **20**, 236–244 (1988).
37. Son, J., Komatsu, M., Matsushita, N. & Hogetsu, T. Migration of pine wood nematodes in the tissues of *Pinus thunbergii*. *J. For. Res.* **15**(3), 186–193 (2009).
38. Trindade, C.M.P.P. Avaliação da expressão de genes relacionados com a susceptibilidade a *Bursaphelenchus xylophilus*, agente causal da doença da murchidão dos pinheiros (pine wilt disease) em *Pinus pinaster* Ait e *Pinus yunnanensis* Franch. Master's thesis in Cellular and Molecular Biotechnology, University of Lisbon, Faculty of Sciences (Lisbon, Portugal, 2012).
39. El-Sayed, S. M. & Mahdy, M. E. Effect of chitosan on root-knot nematode, *Meloidogyne javanica* on tomato plants. *Int. J. Chem. Tech. Res.* **7**(4), 1985–1992 (2015).
40. Apel, K. & Hirt, H. Reactive oxygen species: Metabolism, oxidative stress, and signal transduction. *Annu. Rev. Plant Biol.* **55**, 373–399 (2004).
41. Santos, C. S. *et al.* Understanding the role of the antioxidant system and the tetrapyrrole cycle in iron deficiency chlorosis. *Plants* **8**(9), 348. <https://doi.org/10.3390/plants8090348> (2019).
42. Agrawal, G. K. *et al.* Chitosan activates defense/stress response(s) in the leaves of *Oryza sativa* seedlings. *Plant Physiol. Biochem.* **40**, 1061–1069 (2002).
43. Tavares, J. T. Q., Silva, C. L., Carvalho, L. A., Silva, M. A. & Santos, C. M. G. Estabilidade do ácido ascórbico em suco de laranja submetido a diferentes tratamentos. *Magistra* **12**, 1–2 (2000).
44. Barreiros, L. B. S., David, J. M. & David, J. P. Estresse oxidativo: relação entre geração de espécies reativas e defesas do organismo. *Quím. Nova* **29**, 113–123 (2006).
45. Zhao, T., Wang, J., Wang, Y., Sun, J. W. & Cao, Y. Effects of reactive oxygen species metabolic system on soybean (*Glycine max*) under exogenous chitosan to ozone stress. *Bull. Environ. Contam. Toxicol.* **85**, 59–63 (2010).
46. Ferri, M. *et al.* Chitosan treatment induces changes of protein expression profile and stilbene distribution in *Vitis vinifera* cell suspensions. *Proteomics* **9**, 610–624 (2009).
47. Povero, G. *et al.* Transcript profiling of chitosan-treated *Arabidopsis* seedlings. *J. Plant Res.* **124**, 619–629 (2010).
48. Pongprayoon, W., Roytrakul, S., Pichayangkura, R. & Chadchawan, S. The role of hydrogen peroxide in chitosan-induced resistance to osmotic stress in rice (*Oryza sativa* L.). *Plant Growth Regul.* **70**, 159–173 (2013).
49. Schaefer, H. M., Rentzsch, M. & Breuer, M. Anthocyanins reduce fungal growth in fruits. *Nat. Prod. Commun.* **3**(8), 1267–1272 (2008).
50. Nishino, A., Yasui, H. & Maoka, T. Reaction of paprika carotenoids, capsanthin and capsorubin, with reactive oxygen species. *J. Agric. Food Chem.* **64**, 4786–4792 (2016).
51. Elkhouni, A. *et al.* Structural and functional integrity of *Sulla carnosa* photosynthetic apparatus under iron deficiency conditions. *Plant Biol.* **20**, 415–425 (2018).
52. Kim, H. J., Chen, F., Wang, X. & Rajapakse, N. C. Effect of chitosan on the biological properties of sweet basil (*Ocimum basilicum* L.). *J. Agric. Food Chem.* **53**(9), 3696–3701 (2005).

53. Pichyangkura, R. & Chadchawan, S. Biostimulant activity of chitosan in horticulture. *Sci. Hortic.* **196**, 49–65 (2015).
54. Nakajima, J., Tanaka, I., Seo, S., Yamazaki, M. & Saito, K. LC/PDA/ESI-MS profiling and radical scavenging activity of anthocyanins in various berries. *J. Biomed. Biotechnol.* **2004**(5), 241–247 (2004).
55. Kuroda, H., Goto, S., Kazumi, E. & Kuroda, K. The expressed genes of Japanese red pine (*Pinus densiflora*) involved in the pine wilt disease severity. *BMC Proc.* **5**(7), 92. <https://doi.org/10.1186/1753-6561-5-S7-P92> (2011).
56. Gaspar, D. *et al.* Expression profiling in *Pinus pinaster* in response to infection with the pine wood nematode *Bursaphelenchus xylophilus*. *Forest* **8**(8), 279. <https://doi.org/10.3390/f8080279> (2017).
57. Zas, R. *et al.* Intraspecific variation of anatomical and chemical defensive traits in Maritime pine (*Pinus pinaster*) as factors in susceptibility to the pinewood nematode (*Bursaphelenchus xylophilus*). *Trees* **29**, 663–673 (2014).
58. Raes, J., Rohde, A., Christensen, J. H., Van de Peer, Y. & Boerjan, W. Genome-wide characterization of the lignification toolbox in arabidopsis. *Plant Physiol.* **133**(3), 1051–1071 (2003).
59. Lucini, L. *et al.* Chitosan treatment elicited defence mechanisms, pentacyclic triterpenoids and stilbene accumulation in grape (*Vitis vinifera* L.) bunches. *Phytochem.* **156**, 1–8 (2018).
60. Jacquot, J. P., Lancelin, J. M. & Meyer, Y. Thioredoxins: Structure and function in plant cells. *New Phytol.* **136**, 543–570 (1997).
61. Laloi, C., Mestres-Ortega, D., Marco, Y., Meyer, Y. & Reichheld, J. The arabidopsis cytosolic thioredoxin h5 gene induction by oxidative stress and its W-box-mediated response to pathogen elicitor. *Plant Physiol.* **134**, 1006–1016 (2004).
62. Kovaleva, V., Krynytskyy, H., Gout, I. & Gout, R. Recombinant expression, affinity purification and functional characterization of Scots pine defensin 1. *Appl. Microbiol. Biotechnol.* **89**, 1093–1101 (2011).
63. Zamany, A., Liu, J. J., Ekramoddoullah, A. & Sniezko, R. Antifungal activity of a *Pinus monticola* antimicrobial peptide 1 (Pm-AMP1) and its accumulation in western white pine infected with *Cronartium ribicola*. *Can. J. Microbiol.* **57**, 667–679 (2011).
64. Wang, X. *et al.* Formation of α -farnesene in tea (*Camellia sinensis*) leaves induced by herbivore-derived wounding and its effect on neighbouring tea plants. *Int. J. Mol. Sci.* **20**, 4151. <https://doi.org/10.3390/ijms20174151> (2019).
65. Baermann, G. Eine einfache methode zur auffindung von Anchylostomum-(Nematoden)-larven in erdproben. *Geneeskundig Tijdschrift Nederlands Indië* **57**, 131–133 (1917).
66. Yang, B. Pine wilt disease. In *The history, dispersal and potential threat of pine wood nematode in China* (eds Mota, M. & Vieira, P.) 21–24 (Springer, New York, 2004).
67. Ruley, A. T., Sharma, N. C. & Sahi, S. V. Antioxidant defence in a lead accumulating plant, *Sesbania drummondii*. *Plant Physiol.* **42**, 899–906 (2004).
68. Sims, D. A. & Gamon, J. A. Relationships between leaf pigment content and spectral reflectance across a wide range of species, lead structures and development stages. *Remote Sens. Environ.* **81**, 337–354 (2002).
69. Azevedo, H.A.Q.P. Contributions to the study of the *Pinus pinaster*-*Botrytis cinerea* interaction. Ph.D. dissertation in Biology, Minho University (Braga, Portugal, 2005).
70. Fukushima, R. S. & Hatfield, R. D. Extraction and isolation of lignin for utilization as a standard to determine lignin concentration using the acetyl bromide spectrophotometric method. *J. Agric. Food Chem.* **49**(7), 3133–3139 (2001).
71. Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ method. *Methods* **25**, 402–408 (2001).

Acknowledgements

This article was supported by the project “POINTERS - Interactions between nematodes and host pine trees: the discovery of sustainable alternatives for the management of the pine wilt disease”, funded by the Competitiveness and Internationalization Operational Program (POCI-01-0145- FEDER-031999) and by Fundação para a Ciência e a Tecnologia under its OE component (PTDC/ASP-SIL/31999/2017). The authors would like to thank Dr. Manuel Mota for providing the nematode strain HF. This work was also supported by National Funds from Fundação para a Ciência e a Tecnologia through the scientific collaboration under the Project UID/Multi/50016/2019, and through A.L.V.’s scholarship funded by the Galician government (Ref.: 530 IN606B).

Author contributions

M.N.S. and C.S.S. designed the experimental setup and M.W.V. coordinated the funding project of this study. M.N.S., C.S.S. and A.C. maintained and inoculated the plants and conducted the analytical procedures. All authors participated in result analysis, critical discussion and writing, and approved the submitted version of the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to M.N.d.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2021