University of New Hampshire

University of New Hampshire Scholars' Repository

NHAES Bulletin

New Hampshire Agricultural Experiment Station

10-1-1963

Growth and feed standards for broilers -- 1963, Station Bulletin, no.478

Wabeck, C. J.

Skoglund, W. C.

New Hampshire Agricultural Experiment Station

Follow this and additional works at: https://scholars.unh.edu/agbulletin

Recommended Citation

Wabeck, C. J.; Skoglund, W. C.; and New Hampshire Agricultural Experiment Station, "Growth and feed standards for broilers – 1963, Station Bulletin, no.478" (1963). *NHAES Bulletin*. 440. https://scholars.unh.edu/agbulletin/440

This Text is brought to you for free and open access by the New Hampshire Agricultural Experiment Station at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in NHAES Bulletin by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact nicole.hentz@unh.edu.

tronives

E21

October 1963

Growth and Feed Standards for Broilers—1963

Ву

C. J. WABECK and W. C. SKOGLUND

Agricultural Experiment Station University of New Hampshire Durham, New Hampshire



Growth and Feed Standards for Broilers—1963

By

C. J. WABECK

anu

W. C. SKOGLUND

Growth and Feed Standards for Broilers—1963

By C. J. Wabeck and W. C. Skoglund*

BROILER growth and feed standards provide a means whereby the commercial broiler grower may compare his flock against a standard for growth, feed consumption, and feed conversion at different age levels. In order to meet competition, every effort should be made to meet or exceed these standards.

The Department of Poultry Science, at the University of New Hampshire, has published two bulletins, in the past, Station Bulletin 401 in 1952 and Station Bulletin 466 in 1959, to provide up-to-date information on growth and feed standards. Due to rapid improvements in feed, management recommendations, and strains of broilers, such standards are outdated and new up-to-date figures are needed.

Data was obtained from four commercially available strains of broilers during the period of 1962-63. The birds were assigned randomly by strain to pens of 140 birds each, at the rate of one square foot per bird. A total of eight pens were used for each trial. The chicks were brooded by a central hot water heating system. A four foot watering trough in each pen provided approximately 0.8 inches per chick of watering space. Feeder capacity was increased proportionately with age until four weeks of age when approximately 3.1 inches of feeder space was provided for each bird for the duration of the trial. A high energy broiler feed in crumble form was fed for the first three weeks of age, and pellets were fed for the remainder of the trial. A total of 4,480 birds were reared in four different trials during all seasons of the year.

Table I presents the average weekly weights and gains for the males, females, and mixed sexes.

Table II presents the weekly and cumulative feed consumption and feed conversion figures.

Table III shows the comparative figures for 1953, 1959 and 1963 for broiler weight, feed consumption and conversion.

^{*} Mr. Wabeck is a Graduate Research Assistant, New Hampshire Agricultural Experiment Station. Dr. Skoglund is Professor of Poultry Science and Poultry Scientist, New Hampshire Agricultural Experiment Station.

Table 1. Weekly Average Weight and Gain for Commercial Broilers, 1962-63

	Males		Fem	ales	Mixed Sexes		
Week	Avg. Wgt.	Gain/Week	Avg. Wgt.	Gain/Week	Avg. Wgt.	Gain/Weel	
Initial	.09		.09		.09		
1	.23	.14	.22	.13	.23	.14	
2	.48	.25	.45	.23	.47	.24	
3	.87	.39	.77	.32	.82	.35	
4	1.32	.45	1.14	.37	1.23	.41	
5	1.86	.54	1.58	.44	1.72	.49	
6	2.52	.66	2.06	.48	2.29	.57	
7	3.23	.71	2.61	.55	2.92	.63	
8	3,90	.67	3.13	.52	3.52	.60	
9	4.61	.71	3.64	.51	4.13	.61	

Table 2. Weekly and Cumulative Feed Consumption and Feed Conversion for Mixed Sex Commercial Broilers, 1962-63

Feed	Consumption per	Feed Conversion*			
Week	Weekly	Cumulative	Weekly	Cumulative	
1	.16	.16	1.23	1.23	
2	.36	.52	1.49	1.38	
3	.60	1.12	1.68	1.53	
4	.82	1.94	1.94	1.67	
5	1.01	2.95	2.01	1.79	
6	1.28	4.23	2.14	1.87	
7	1.44	5.67	2.35	1.95	
8	1.64	7.31	2.77	2.10	
9	1.78 9.09		2.91	2.23	

^{*} Pounds of feed required to produce one pound of live weight.

Comparative Figures 1953¹, 1959², and 1963, for Broiler Weights, Feed Consumption and Conversion.

10	9	8	7	6	5	4	သ	2	_	Initial	Week	
3.26		2.35		1.51		.91		.37		.10	1953	Av
4.05	3.57	3.01	2.50	1.99	1.51	1.04	.65	.37	.19	.09	1959	For Mixed Sexe
	4.13	3.52	2.92	2.29	1.72	1.23	.82	.47	.23	.09	1963	ight es
8.70		5.71		3,38		1.65		.47			1953	
9.39	7.84	6.28	4.84	3.59	2.53	1.61	.93	.44	.14		1959	Cumulative Feed Consumption
	9.09	7.31	5.67	4.23	2.95	1.94	1.12	.52	.16	1	1963	
2.67		2.43		2.24	1	1.81	İ	1.26	1		1953	Çı
2.32	2.20	2.09	1.94	1.80	1.68	1.55	1.43	1.19	.74		1959	umulative Feed Conversion
	2.23	2.10	1.95	1.87	1.79	1.67	1.53	1.38	1.23		1963	

¹ Potter, L. M., and R. C. Ringrose, 1953. Growth and Feed Standards for New Hampshire. Station Bulletin 401, N. H. Agricultural Experiment Station.

²Reed, Willis S., and W. C. Skoglund, 1959. Growth and Feed Standards for Broilers Station Bulletin 466, N. H. Agricultural Experiment Station.



Table 3. Comparative Figures 1953¹, 1959², and 1963, for Broiler Weights, Feed Consumption and Conversion.

9 10	8 7	6 5	ယ္န	2	Initial	Week -	
3.26	2.35	1.51	.91	.37	.10	1953	Αv
3.57 4.05	2.50 3.01	1.51 1.99	.65 1.04	.19	.09	For Mixed Sexe	erage Body We
4.13	2.92 3.52	1.72 2.29	.82 1.23	.47	.09	1963	ight
8.70	5.71	3,38	1.65	.47		1953	
7.84 9.39	4.84 6.28	2.53 3.59	.93 1.61	.14 .44		Consumption 1959	Cumulative Feed
9.09	5.67 7.31	2.95 4.23	1.12 1.94	.16 .52		1963	
2.67	2.43	2.24	1.81	1.26		1953	Cu
2.20 2.32	2.09	1.68	1.43	.74 1.19		Conversion 1959	mulative Feed
2.23	1.95 2.10	1.79	1.53	1.23 1.38		1963	

¹ Potter, L. M., and R. C. Ringrose, 1953. Growth and Feed Standards for New Hampshire. Station Bulletin 401, N. H. Agricultural Experiment Station.

²Reed, Willis S., and W. C. Skoglund, 1959. Growth and Feed Standards for Broilers Station Bulletin 466, N. H. Agricultural Experiment Station.