

Title	Colloid-polymer mixtures and the limits of criticality
Author(s)	Royall, Paddy; Tanaka, Hajime; Aarts, Dirk
Citation	物性研究 (2006), 87(1): 147-147
Issue Date	2006-10-20
URL	<a href="http://hdl.handle.net/2433/110604">http://hdl.handle.net/2433/110604</a>
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

## Colloid-polymer mixtures and the limits of criticality

Institute of Industrial Science, University of Tokyo. Paddy Royall<sup>1</sup> & Hajime Tanaka  
Ecole Normale Supérieure, Paris. Dirk Aarts

Since the industrial revolution, boiling and condensation have come to be one of the best recognised phase transitions of condensed matter. Approaching the critical point, a liquid becomes indistinguishable from its vapour, the interfacial thickness diverges and the system is entirely dominated by long-wavelength density fluctuations so microscopic properties such as chemical structure may be safely neglected. This universality concept assumes a large separation between the lengthscales of the critical density fluctuations and the constituent particles. However, the connection between these mesoscopic and microscopic regimes remains a long-standing open question. Here we access this “missing link” for the first time by using a mesoscopic analogue of simple liquids, colloid-polymer mixtures.

Direct visualization of both the individual colloidal particles and critical density fluctuations provides full access to all lengthscales. We reveal the first particle-level images of the critical clusters and liquid-gas interface. It turns out that the lengthscale of both is identical. Surprisingly, we find that the behaviour can be very well described by critical scaling almost to the single-particle level, as shown by the correlation length, critical cluster fractal dimension and free energy, all of which we directly measure. However, when the correlation length falls to 2-3 times the constituent particle size, we reveal a clear crossover from critical to non-critical behaviour. Our results provide a framework in which to unify the disparate particle- and correlation- lengthscales, and bring new insight into the nature of the liquid-gas interface and the limit of the critical regime.

---

<sup>1</sup>E-mail: paddy@iis.u-tokyo.ac.jp