

Title	Perception-Based Extension of Probability Models and its Application to Finance (Information and mathematics of non-additivity and non-extensivity : non-additivity and convex analysis)
Author(s)	Yoshida, Yuji
Citation	数理解析研究所講究録 (2009), 1630: 96-105
Issue Date	2009-02
URL	<a href="http://hdl.handle.net/2433/140382">http://hdl.handle.net/2433/140382</a>
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

## Perception-Based Extension of Probability Models and its Application to Finance

北九州市立大学・経済学部 吉田 祐治 (Yuji Yoshida)  
Faculty of Economics and Business Administration  
University of Kitakyushu

### 1 Introduction

The risk allocation is an important topic in asset management under uncertainty, and in finance the portfolio is one of the most important methods for the risk allocation. Recently, *value-at-risk* ( $VaR$ ) is used widely in financial trading to estimate the risk of worst-scenarios. In this paper, we deal with *conditional value-at-risk* ( $CVaR$ ), which is derived from  $VaR$ , and a  $CVaR$ -portfolio problem. From the viewpoint of risk theory, it is also known that  $VaR$  is not a coherent risk measure ([1]), however  $CVaR$  is not easy to find the correspondence with parameters in finance.  $VaR$  and  $CVaR$  have a merit and a demerit as a risk allocation tool. This paper discusses them in a fuzzy and probabilistic environment from the viewpoint of risk measures.

Estimation of uncertain quantities is important in decision making ([22, 15, 16]). To represent uncertainty in this portfolio model, we use *fuzzy random variables* which have two kinds of uncertainties, i.e. randomness and fuzziness. In this paper, randomness is used to represent the uncertainty regarding the belief degree of frequency, and fuzziness is applied to linguistic imprecision of data because of a lack of knowledge regarding the current stock market. We extend the  $CVaR$  for real random variables to one regarding fuzzy random variables from the viewpoint of perception-based approach in Yoshida [20]. We formulate the  $CVaR$  portfolio problem with fuzzy random variables, and we discuss the fundamental properties of the extended  $CVaR$  using the results in Yoshida [21]. Recently, Yoshida [17, 19] introduced the mean, the variance and the measurement of fuzziness of fuzzy random variables, using *evaluation weights* and  $\lambda$ -mean functions. This paper estimates fuzzy numbers/fuzzy random variables by the probabilistic expectation and these criteria, which are characterized by *possibility/necessity criteria* for subjective estimation and a *pessimistic-optimistic index* for subjective decision. These parameters are decided by the investor and are based on the degree of his certainty regarding the current information in the market. In this portfolio model, we use triangle-type fuzzy numbers/fuzzy random variables for computation in actual models, and we analyze mathematically the  $CVaR$  portfolio problem under some regularity condition.

### 2 A portfolio model under stochastic and fuzzy environment

In this paper, we consider a portfolio model with  $n$  stocks as risky assets, where  $n$  is a positive integer. We assume *small investors hypothesis* such that an investor's actions do not have any impact on the stock market ([9]). Let a positive integer  $T$  denote an expiration date, and let  $\mathbb{R}$  denote the set of all real numbers. Let  $(\Omega, P)$  be a probability

space, where  $P$  is a non-atomic probability measure on a sample space  $\Omega$ . For an asset  $i = 1, 2, \dots, n$ , a *stock price process*  $\{S_t^i\}_{t=0}^T$  is given by *rates of return*  $R_t^i$  at time  $t$  as follows. Let a stock price  $S_t^i := S_{t-1}^i(1 + R_t^i)$  for time  $t = 1, 2, \dots, T$ , where  $\{R_t^i\}_{t=1}^T$  is assumed to be a sequence of integrable real random variables. In this paper, we discuss a portfolio model where stock prices  $S_t^i$  take fuzzy values using fuzzy random variables, taking into account from linguistic imprecision of data because of a lack of knowledge regarding the current stock market. Mathematical notations of fuzzy random variables are introduced later. Hence, we deal with a portfolio with *portfolios given by portfolio weight vectors*  $w = (w^1, w^2, \dots, w^n)$  such that  $w^1 + w^2 + \dots + w^n = 1$  and  $w^i \geq 0$  ( $i = 1, 2, \dots, n$ ). The rate of return for the portfolio  $w = (w^1, w^2, \dots, w^n)$  is given by

$$R_t := w^1 R_t^1 + w^2 R_t^2 + \dots + w^n R_t^n. \quad (1)$$

This paper assumes that  $R_t^i$  ( $i = 1, 2, \dots, n$ ) has a normal distribution ([8, 23, 24]).

Next, we introduce fuzzy numbers/fuzzy random variable and we give a portfolio model under uncertainty. A fuzzy number is denoted by its membership function  $\tilde{a} : \mathbb{R} \mapsto [0, 1]$  which is normal, upper-semicontinuous and quasi-concave and has a compact support ([15, 16, 25]).  $\mathcal{R}$  denotes the set of all fuzzy numbers. In this paper, we identify fuzzy numbers with their corresponding membership functions. The  $\alpha$ -cut of a fuzzy number  $\tilde{a} (\in \mathcal{R})$  is given by  $\tilde{a}_\alpha := \{x \in \mathbb{R} \mid \tilde{a}(x) \geq \alpha\}$  ( $\alpha \in (0, 1]$ ) and  $\tilde{a}_0 := \text{cl}\{x \in \mathbb{R} \mid \tilde{a}(x) > 0\}$ , where  $\text{cl}$  denotes the closure of an interval. We write the closed intervals as  $\tilde{a}_\alpha := [\tilde{a}_\alpha^-, \tilde{a}_\alpha^+]$  for  $\alpha \in [0, 1]$ . Hence we also introduce a partial order  $\succeq$ , so called the *fuzzy max order*, on fuzzy numbers  $\mathcal{R}$  ([4]). An addition, a subtraction and a scalar multiplication for fuzzy numbers are defined by Zadeh's extension principle ([15, 16, 25]).

A fuzzy-number-valued map  $\tilde{X} : \Omega \mapsto \mathcal{R}$  is called a *fuzzy random variable* if the maps  $\omega \mapsto \tilde{X}_\alpha^\pm(\omega)$  are measurable for all  $\alpha \in (0, 1]$ , where  $\tilde{X}_\alpha(\omega) = [\tilde{X}_\alpha^-(\omega), \tilde{X}_\alpha^+(\omega)] = \{x \in \mathbb{R} \mid \tilde{X}(\omega)(x) \geq \alpha\}$  ([6, 10]). We need to introduce expectations of fuzzy random variables in order to describe a portfolio model. A fuzzy random variable  $\tilde{X}$  is said to be integrably bounded if  $\omega \mapsto \tilde{X}_\alpha^\pm(\omega)$  are integrable for all  $\alpha \in (0, 1]$ . Let  $\tilde{X}$  be an integrably bounded fuzzy random variable. The expectation  $E(\tilde{X})$  of the fuzzy random variable  $\tilde{X}$  is defined by a fuzzy number  $E(\tilde{X})(x) := \sup_{\alpha \in [0, 1]} \min\{\alpha, 1_{E(\tilde{X})_\alpha}(x)\}$ , where  $E(\tilde{X})_\alpha := [\int_\Omega \tilde{X}_\alpha^-(\omega) dP(\omega), \int_\Omega \tilde{X}_\alpha^+(\omega) dP(\omega)]$  for  $\alpha \in (0, 1]$  ([5, 10, 14]).

Now we deal with a case where the rate of return  $\{R_t^i\}_{t=1}^T$  has some imprecision ([19]). In this paper, we use triangle-type fuzzy random variables for computation, however we can apply the similar approach to general fuzzy random variables. We define a *rate of return process with imprecision*  $\{\tilde{R}_t^i\}_{t=0}^T$  by a sequence of triangle-type fuzzy random variables

$$\tilde{R}_t^i(\cdot)(x) = \begin{cases} 0 & \text{if } x < R_t^i - c_t^i \\ \frac{x - R_t^i + c_t^i}{c_t^i} & \text{if } R_t^i - c_t^i \leq x < R_t^i \\ \frac{x - R_t^i - c_t^i}{-c_t^i} & \text{if } R_t^i \leq x < R_t^i + c_t^i \\ 0 & \text{if } x \geq R_t^i + c_t^i, \end{cases} \quad (2)$$

where  $c_t^i$  is a positive number. We call  $c_t^i$  a *fuzzy factor* for asset  $i$  at time  $t$ . Hence we can represent  $\tilde{R}_t^i$  by the sum of the real random variable  $R_t^i$  and a fuzzy number  $\tilde{a}_t^i$ :

$$\tilde{R}_t^i(\omega)(\cdot) := 1_{\{R_t^i(\omega)\}}(\cdot) + \tilde{a}_t^i(\cdot) \quad (3)$$

for  $\omega \in \Omega$ , where  $1_{\{\cdot\}}$  denotes the characteristic function of a singleton and  $\tilde{a}_t^i$  is a triangle-type fuzzy number defined by

$$\tilde{a}_t^i(x) = \begin{cases} 0 & \text{if } x < -c_t^i \\ \frac{x+c_t^i}{c_t^i} & \text{if } -c_t^i \leq x < 0 \\ \frac{x-c_t^i}{-c_t^i} & \text{if } 0 \leq x < c_t^i \\ 0 & \text{if } x \geq c_t^i. \end{cases} \quad (4)$$

For assets  $i = 1, 2, \dots, n$ , we define *stock price processes*  $\{\tilde{S}_t^i\}_{t=0}^T$  by the *rates of return with imprecision*  $\tilde{R}_t^i$  as follows:  $\tilde{S}_0^i := S_0^i$  is a positive number and

$$\tilde{S}_t^i = \tilde{S}_0^i \prod_{s=1}^t (1 + \tilde{R}_s^i) \quad (5)$$

for  $t = 1, 2, \dots, T$  ([16]). Hence, we present a portfolio with *trading strategies* given by *portfolio weight vectors*  $w = (w^1, w^2, \dots, w^n)$  such that  $w^1 + w^2 + \dots + w^n = 1$  and  $w^i \geq 0$  ( $i = 1, 2, \dots, n$ ). For the portfolio  $w = (w^1, w^2, \dots, w^n)$ , the rate of return with imprecision for the portfolio is given by a linear combination of fuzzy random variables

$$\tilde{R}_t := w^1 \tilde{R}_t^1 + w^2 \tilde{R}_t^2 + \dots + w^n \tilde{R}_t^n. \quad (6)$$

In Section 4, we discuss a CVaR model regarding (6).

### 3 An extension of CVaR for fuzzy random variables

In this section, we introduce a conditional value-at-risk for fuzzy random variables and we apply it to the rate of return (6). Let  $\mathcal{X}$  be the set of all integrable real random variables  $X$  on  $\Omega$  with a continuous distribution function  $x \mapsto F_X(x) := P(X < x)$  for which there exists a non-empty open interval  $I$  such that  $F_X(\cdot) : I \mapsto (0, 1)$  is a strictly increasing and onto. Then there exists a strictly increasing and continuous inverse function  $F_X^{-1} : (0, 1) \mapsto I$ . We put  $F_X(\inf I) := \lim_{x \downarrow \inf I} F_X(x) = 0$  and  $F_X(\sup I) := \lim_{x \uparrow \sup I} F_X(x) = 1$ . Then, the *value-at-risk* (*VaR*) at a risk probability  $p$  is given by the percentile of the distribution function  $F_X$ . Define

$$\text{VaR}_p(X) := \sup\{x \in I \mid F_X(x) \leq p\} \quad (7)$$

if  $0 < p < 1$ ,  $\text{VaR}_p(X) := 0$  if  $p = 0$  and  $\text{VaR}_p(X) := 1$  if  $p = 1$ . Then we have  $\text{VaR}_p(X) = F_X^{-1}(p)$  for  $0 < p < 1$ . The *conditional value-at-risk* (*CVaR*) at a probability level  $p$  (*Expected Shortfall* with at a confidence probability level  $1 - p$ ) is given by

$$\text{CVaR}_p(X) := \frac{1}{p} \int_0^p \text{VaR}_q(X) dq \quad (8)$$

if  $0 < p \leq 1$  and  $\text{CVaR}_p(X) := \inf I$  if  $p = 0$  ([11]) It is known that CVaR has the following properties, which implies CVaR is a *coherent risk measure*.

**Lemma 1.** *Let  $X, Y \in \mathcal{X}$  and let  $p$  be a positive probability. Then the conditional value-at-risk  $\text{CVaR}_p$  defined by (8) has the following properties:*

- (i) *If  $X \leq Y$ , then  $\text{CVaR}_p(X) \leq \text{CVaR}_p(Y)$ . (monotonicity)*
- (ii)  *$\text{CVaR}_p(\zeta X) = \zeta \text{CVaR}_p(X)$  for  $\zeta > 0$ . (positively homogeneity)*
- (iii)  *$\text{CVaR}_p(X + \theta) = \text{CVaR}_p(X) + \theta$  for  $\theta \in \mathbb{R}$ . (translation invariance)*
- (iv)  *$\text{CVaR}_p(X + Y) \geq \text{CVaR}_p(X) + \text{CVaR}_p(Y)$ . (super-additivity)*
- (v) *Let  $\{X_n\}_n (\subset \mathcal{X})$  be a monotone sequence of real random variables with a limit  $X (\in \mathcal{X})$ . Then  $\lim_{n \rightarrow \infty} \text{CVaR}_p(X_n) = \text{CVaR}_p(X)$  (continuity).*
- (vi)  *$\text{CVaR}_p(X + Y) = \text{CVaR}_p(X) + \text{CVaR}_p(Y)$  if  $X$  and  $Y$  are comonotonic (comonotonically additive).*

**Remark.**

- (a) Lemma 1(ii) and (iv) imply the convexity, which is an important property in risk theory.
- (b) Regarding Lemma 1(iv), we note that the super-additivity for the value-at-risk

$$\text{VaR}_p(X + Y) \geq \text{VaR}_p(X) + \text{VaR}_p(Y),$$

$(X, Y \in \mathcal{X})$  does not hold in general ([1]).

Let  $\tilde{\mathcal{X}}$  be the set of all fuzzy random variables  $\tilde{X}$  on  $\Omega$  such that their  $\alpha$ -cuts  $\tilde{X}_\alpha^\pm$  are integrable and  $\lambda \tilde{X}_\alpha^- + (1 - \lambda) \tilde{X}_\alpha^+ \in \mathcal{X}$  for all  $\lambda \in [0, 1]$  and  $\alpha \in [0, 1]$ . Hence, from (8) we introduce a CVaR for a fuzzy random variable  $\tilde{X} (\in \tilde{\mathcal{X}})$  at a positive risk probability  $p$  as follows.

$$\text{CVaR}_p(\tilde{X})(x) := \sup_{X \in \mathcal{X} : \text{CVaR}_p(X) = x} \inf_{\omega \in \Omega} \tilde{X}(\omega)(X(\omega)),$$

$x \in \mathbb{R}$ . Yoshida [20] has studied *perception-based estimations* extending the concept of the expectations in Kruce and Meyer [5]. This definition is an extension from the CVaR on real random variables to the CVaR on fuzzy random variables. Hence, the CVaR on fuzzy random variables is characterized by the following representation ([20]).

**Theorem 1.** *Let  $\tilde{X} \in \tilde{\mathcal{X}}$  be a fuzzy random variable and let  $p$  be a positive probability. Then the conditional value-at-risk  $\text{CVaR}_p(\tilde{X})$  is a fuzzy number whose  $\alpha$ -cuts are*

$$\text{CVaR}_p(\tilde{X})_\alpha = [\text{CVaR}_p(\tilde{X}_\alpha^-), \text{CVaR}_p(\tilde{X}_\alpha^+)], \quad (9)$$

for  $\alpha \in (0, 1]$ .

The CVaR on fuzzy random variables has the following properties similar to Lemma 1 for the CVaR (8). Theorem 2 shows that CVaR is a *coherent risk measure on the fuzzy random variables*.

**Theorem 2.** *Let  $\tilde{X}, \tilde{Y} \in \tilde{\mathcal{X}}$  be fuzzy random variables and let  $p$  be a positive probability. Then the conditional value-at-risk  $\text{CVaR}_p$  on fuzzy random variables has the following properties:*

- (i) *If  $\tilde{X} \preceq \tilde{Y}$ , then  $\text{CVaR}_p(\tilde{X}) \preceq \text{CVaR}_p(\tilde{Y})$ . (monotonicity)*
- (ii)  *$\text{CVaR}_p(\zeta \tilde{X}) = \zeta \text{CVaR}_p(\tilde{X})$  for  $\zeta > 0$ . (positively homogeneity)*
- (iii)  *$\text{CVaR}_p(\tilde{X} + \tilde{a}) = \text{CVaR}_p(\tilde{X}) + \tilde{a}$  for a fuzzy number  $\tilde{a} \in \mathcal{R}$ . (translation invariance)*
- (iv)  *$\text{CVaR}_p(\tilde{X} + \tilde{Y}) \succeq \text{CVaR}_p(\tilde{X}) + \text{CVaR}_p(\tilde{Y})$ . (super-additivity)*
- (v) *Let  $\{\tilde{X}_n\}_n \subset \tilde{\mathcal{X}}$  be a monotone sequence of fuzzy random variables with a limit  $\tilde{X} \in \tilde{\mathcal{X}}$ . Then  $\lim_{n \rightarrow \infty} \text{CVaR}_p(\tilde{X}_n) = \text{CVaR}_p(\tilde{X})$  (continuity).*
- (vi)  *$\text{CVaR}_p(\tilde{X} + \tilde{Y}) = \text{CVaR}_p(\tilde{X}) + \text{CVaR}_p(\tilde{Y})$  if  $\tilde{X}$  and  $\tilde{Y}$  are comonotonic (comonotonically additive).*

Next we need to evaluate the fuzziness of fuzzy numbers/fuzzy random variables since the conditional value-at-risk  $\text{CVaR}_p(\tilde{R}_t)$  for the rate of return (6) with portfolio is a fuzzy number. There are many studies regarding the defuzzification of fuzzy numbers. Here we adopt the evaluation method of fuzzy numbers/fuzzy random variables, which is given by possibility/necessity criteria ([3, 15, 16]). In the rest of this section we introduce the definitions from [17, 18, 19], and in the next section we estimate the CVaR regarding the rate of return (6) by the evaluation method. Yoshida [17, 19] has studied an evaluation of fuzzy numbers by *evaluation weights* which are induced from fuzzy measures to evaluate a confidence degree that a fuzzy number takes values in an interval. With respect to fuzzy random variables, the randomness is evaluated by the probabilistic expectation and the fuzziness is estimated by the evaluation weights and the following function. Let  $g^\lambda : \mathcal{I} \mapsto \mathbb{R}$  be a map such that

$$g^\lambda([x, y]) := \lambda x + (1 - \lambda)y \quad (10)$$

for  $[x, y] \in \mathcal{I}$ , where  $\lambda$  is a constant satisfying  $0 \leq \lambda \leq 1$  and  $\mathcal{I}$  denotes the set of all bounded closed intervals. This scalarization is used for the estimation of fuzzy numbers to give a mean value of the interval  $[x, y]$  with a weight  $\lambda$ , and  $g^\lambda$  is called a  $\lambda$ -*mean function* and  $\lambda$  is called a *pessimistic-optimistic index* which indicates the pessimistic degree of attitude in decision making ([2]). Let a fuzzy number  $\tilde{a} \in \mathcal{R}$ . A mean value of the fuzzy

number  $\tilde{a}$  with respect to  $\lambda$ -mean functions  $g^\lambda$  and an evaluation weight  $w(\alpha)$ , which depends only on  $\tilde{a}$  and  $\alpha$ , is given as follows ([17, 18]):

$$\tilde{E}(\tilde{a}) := \int_0^1 g^\lambda(\tilde{a}_\alpha) w(\alpha) d\alpha / \int_0^1 w(\alpha) d\alpha, \quad (11)$$

where  $\tilde{a}_\alpha = [\tilde{a}_\alpha^-, \tilde{a}_\alpha^+]$  is the  $\alpha$ -cut of the fuzzy number  $\tilde{a}$ . In (11),  $w(\alpha)$  indicates a *confidence degree that the fuzzy number  $\tilde{a}$  takes values in the interval  $\tilde{a}_\alpha$  at each level  $\alpha$* . Hence, an evaluation weight  $w(\alpha)$  is called the *possibility evaluation weight  $w^P(\alpha)$*  if  $w^P(\alpha) := 1$  for  $\alpha \in [0, 1]$ , and  $w(\alpha)$  is called the *necessity evaluation weight  $w^N(\alpha)$*  if  $w^N(\alpha) := 1 - \alpha$  for  $\alpha \in [0, 1]$ . Especially, for a fuzzy number  $\tilde{a} \in \mathcal{R}$ , the mean  $\tilde{E}^P(\tilde{a})$  in the possibility case and the mean  $\tilde{E}^N(\tilde{a})$  in the necessity case are represented as follows ([17, 18]):

$$\tilde{E}^P(\tilde{a}) = \int_0^1 g^\lambda(\tilde{a}_\alpha) d\alpha, \quad (12)$$

$$\tilde{E}^N(\tilde{a}) = \int_0^1 g^\lambda(\tilde{a}_\alpha) (2 - 2\alpha) d\alpha. \quad (13)$$

The mean  $\tilde{E}$  has the following natural properties of the linearity and the monotonicity regarding the fuzzy max order.

**Lemma 2** ([17, 18, 19]). *Let  $\lambda \in [0, 1]$ . For fuzzy numbers  $\tilde{a}, \tilde{b} \in \mathcal{R}$  and real numbers  $\theta, \zeta$ , the following (i) – (iv) hold.*

- (i)  $\tilde{E}(\tilde{a} + 1_{\{\theta\}}) = \tilde{E}(\tilde{a}) + \theta$ , where  $1_{\{\theta\}}$  is the characteristic function of a set.
- (ii)  $\tilde{E}(\zeta\tilde{a}) = \zeta\tilde{E}(\tilde{a})$  if  $\zeta \geq 0$ .
- (iii)  $\tilde{E}(\tilde{a} + \tilde{b}) = \tilde{E}(\tilde{a}) + \tilde{E}(\tilde{b})$ .
- (iv) If  $\tilde{a} \succeq \tilde{b}$ , then  $\tilde{E}(\tilde{a}) \geq \tilde{E}(\tilde{b})$ .

For a fuzzy random variable  $\tilde{X}$ , the mean of the expectation  $E(\tilde{X})$  is a real number

$$E(\tilde{E}(\tilde{X})) := E \left( \int_0^1 g^\lambda(\tilde{X}_\alpha) w(\alpha) d\alpha / \int_0^1 w(\alpha) d\alpha \right).$$

From Lemma 2, we obtain the following results regarding fuzzy random variables.

**Lemma 3** ([17, 18, 19]). *Let  $\lambda \in [0, 1]$ . For a fuzzy number  $\tilde{a} \in \mathcal{R}$ , integrable fuzzy random variables  $\tilde{X}, \tilde{Y}$ , an integrable real random variable  $Z$  and a nonnegative real number  $\zeta$ , the following (i) – (v) hold.*

- (i)  $E(\tilde{E}(\tilde{X})) = \tilde{E}(E(\tilde{X}))$ .
- (ii)  $E(\tilde{E}(\tilde{a})) = \tilde{E}(\tilde{a})$  and  $E(\tilde{E}(Z)) = E(Z)$ .

- (iii)  $E(\tilde{E}(\zeta\tilde{X})) = \zeta E(\tilde{E}(\tilde{X}))$ .
- (iv)  $E(\tilde{E}(\tilde{X} + \tilde{Y})) = E(\tilde{E}(\tilde{X})) + E(\tilde{E}(\tilde{Y}))$ .
- (v) If  $\tilde{X} \succeq \tilde{Y}$ , then  $E(\tilde{E}(\tilde{X})) \geq E(\tilde{E}(\tilde{Y}))$ .

#### 4 A CVaR portfolio model under stochastic and fuzzy environment

In this section, we discuss portfolio problems under uncertainty. First we estimate the rate of return with imprecision for a portfolio. Let the mean, the variance and the covariance of the rate of return  $R_t^i$  by

$$\begin{aligned}\mu_t^i &:= E(R_t^i), \\ (\sigma_t^i)^2 &:= E((R_t^i - \mu_t^i)^2), \\ \sigma_t^{ij} &:= E((R_t^i - \mu_t^i)(R_t^j - \mu_t^j))\end{aligned}$$

for  $i, j = 1, 2, \dots, n$ . We assume that the determinant of the variance-covariance matrix  $[\sigma_t^{ij}]$  is not zero and there exists its inverse matrix. For a portfolio  $w = (w^1, w^2, \dots, w^n)$  satisfying  $w^1 + w^2 + \dots + w^n = 1$  and  $w^i \geq 0$  ( $i = 1, 2, \dots, n$ ), we calculate the expectation and the variance regarding  $\tilde{R}_t = w^1 \tilde{R}_t^1 + w^2 \tilde{R}_t^2 + \dots + w^n \tilde{R}_t^n$ . From Lemma 3, the expectation  $\bar{\mu}_t := E(\tilde{E}(\tilde{R}_t))$  follows

$$\bar{\mu}_t = \sum_{i=1}^n w^i \bar{\mu}_t^i, \quad (14)$$

where  $\bar{\mu}_t^i := E(\tilde{E}(\tilde{R}_t^i))$  for  $i = 1, 2, \dots, n$ . On the other hand, regarding this model, in Yoshida [19] we can find that the variance  $(\bar{\sigma}_t)^2$  equals to the variance  $(\sigma_t)^2 := E((R_t - \mu_t)^2)$  of  $R_t$ :

$$(\bar{\sigma}_t)^2 = (\sigma_t)^2 = \sum_{i=1}^n \sum_{j=1}^n w^i w^j \sigma_t^{ij}. \quad (15)$$

Hence, applying Lemmas 2 and 3 to (3), we obtain the following lemma regarding CVaR of the rates of return  $\tilde{R}_t^i$ .

**Lemma 4.** *Let  $p$  be a positive probability. The following (i) and (ii) hold:*

- (i)  $\bar{\mu}_t^i = \mu_t^i + \tilde{E}(\bar{a}_t^i)$  for  $i = 1, 2, \dots, n$ .
- (ii) *The mean of  $\text{CVaR}_p(\tilde{R}_t)$  is evaluated by*

$$\tilde{E}(\text{CVaR}_p(\tilde{R}_t)) = \sum_{i=1}^n w^i \bar{\mu}_t^i - \kappa \sqrt{\sum_{i=1}^n \sum_{j=1}^n w^i w^j \sigma_t^{ij}}$$



where  $\kappa := \frac{1}{p} \int_0^p \kappa(q) dq$  for  $\kappa(q)$  defined by

$$\bar{E}(\text{VaR}_q(\tilde{R}_t)) = \sum_{i=1}^n w^i \tilde{\mu}_t^i - \kappa(q) \sqrt{\sum_{i=1}^n \sum_{j=1}^n w^i w^j \sigma_t^{ij}}.$$

Now we discuss the following CVaR portfolio without allowance for short selling. The following form (16) comes from the conditional value-at-risk  $\bar{E}(\text{CVaR}_p(\tilde{R}_t))$  in Lemma 4.

**CVaR-portfolio problem (P):** Maximize the conditional value-at-risk

$$\sum_{i=1}^n w^i \tilde{\mu}_t^i - \kappa \sqrt{\sum_{i=1}^n \sum_{j=1}^n w^i w^j \sigma_t^{ij}} \quad (16)$$

with respect to portfolios  $w = (w^1, w^2, \dots, w^n)$  satisfying  $w^1 + w^2 + \dots + w^n = 1$  and  $w^i \geq 0$  for  $i = 1, 2, \dots, n$ .

Let  $\tilde{\mu}$  be the vector whose elements are  $\tilde{\mu}^i = \mu_t^i + \bar{E}(\tilde{a}_t^i)$  ( $i = 1, 2, \dots$ ), and let  $\mathbf{1}$  be the vector whose elements are 1. Let

$$\Sigma := \begin{bmatrix} \sigma_t^{11} & \sigma_t^{12} & \dots & \sigma_t^{1n} \\ \sigma_t^{21} & \sigma_t^{22} & \dots & \sigma_t^{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_t^{n1} & \sigma_t^{n2} & \dots & \sigma_t^{nn} \end{bmatrix},$$

$$A := \mathbf{1}^T \Sigma^{-1} \mathbf{1}, \quad B := \mathbf{1}^T \Sigma^{-1} \tilde{\mu}, \quad C := \tilde{\mu}^T \Sigma^{-1} \tilde{\mu} \quad \text{and} \quad \Delta := AC - B^2.$$

Hence, in a similar way as the proof in Yoshida [21, Theorem 4.2], we arrive at the following analytical solutions regarding CVaR-portfolio problem for  $\kappa = \frac{1}{p} \int_0^p \kappa(q) dq$ .

**Theorem 3.** *Let  $A$  and  $\Delta$  be positive. Let  $\kappa$  satisfy  $\kappa^2 > C$ . Then the following (i) and (ii) hold.*

(i) *The solution of CVaR-portfolio problem (P) is given by*

$$w^* := \xi \Sigma^{-1} \mathbf{1} + \eta \Sigma^{-1} \tilde{\mu} \quad (17)$$

*and then the corresponding CVaR is*

$$v^* := \frac{B - \sqrt{A\kappa^2 - \Delta}}{A} \quad (18)$$

*at the expected rate of return*

$$\gamma^* := \frac{B}{A} + \frac{\Delta}{A\sqrt{A\kappa^2 - \Delta}}, \quad (19)$$

where  $\xi := \frac{C - B\gamma^*}{\Delta}$  and  $\eta := \frac{A\gamma^* - B}{\Delta}$ .

- (ii) Further, if  $\Sigma^{-1}\mathbf{1} \geq \mathbf{0}$  and  $\Sigma^{-1}\bar{\mu} \geq \mathbf{0}$ , then the portfolio (17) satisfies  $w^* \geq \mathbf{0}$ , i.e. the portfolio  $w^*$  is a portfolio without allowance for short selling. Here,  $\mathbf{0}$  denotes the zero vector.

## 5 Conclusion

In this paper, we have discussed the following terms:

- Extension of CVaR for fuzzy random variable, and its coherence as a risk measure.
- A CVaR-portfolio model under randomness and fuzziness.
- An optimality portfolio for this model.

VaR is directly related to the falling rate of the asset prices, and it is used widely in real finance. On the other hand, CVaR is not easy to find a direct relation with parameters in real finance, however CVaR is a coherence risk measure. The coherence is a necessary property as a criterion from the viewpoint of axiomatic approach for risk measures.

## References

- [1] P.Arznner, F.Delbaen, J.-M.Eber and D.Heath (1999). Coherent measures of risk, *Mathematical Finance*, volume 9, pages 203-228.
- [2] P. Fortemps and M. Roubens (1996). Ranking and defuzzification methods based on area compensation, *Fuzzy Sets and Systems*, volume 82, pages 319-330.
- [3] M.Inuiguchi and T.Tanino (2000). Portfolio selection under independent possibilistic information, *Fuzzy Sets and Systems*, volume 115, pages 83-92.
- [4] G.J.Klir and B.Yuan (1995). *Fuzzy Sets and Fuzzy Logic: Theory and Applications*, Prentice-Hall, London.
- [5] R.Kruse and K.D.Meyer (1987). *Statistics with Vague Data*, Riedel Publ. Co., Dordrecht.
- [6] H.Kwakernaak (1978). Fuzzy random variables – I. Definitions and theorem, *Inform. Sci.*, volume 15, pages 1-29.
- [7] H.Markowitz (1990). *Mean-Variance Analysis in Portfolio Choice and Capital Markets*, Blackwell, Oxford.
- [8] A.Meucci (2005). *Risk and Asset Allocation*, Springer-Verlag, Heidelberg.
- [9] S.R.Pliska (1997). *Introduction to Mathematical Finance: Discrete-Time Models*, Blackwell Publ., New York.
- [10] M.L.Puri and D.A.Ralescu (1986). Fuzzy random variables, *J. Math. Anal. Appl.*, volume 114, pages 409-422.

- [11] R.T.Rockafellar and S.P.Uryasev (2000). Optimization of conditional value-at-risk, *Journal of Risk*, volume 2, pages 21-42.
- [12] S.M.Ross (1999). *An Introduction to Mathematical Finance*, Cambridge Univ. Press, Cambridge.
- [13] H.Tanaka, P.Guo and L.B.Türksen (2000). Portfolio selection based on fuzzy probabilities and possibilistic distribution, *Fuzzy Sets and Systems*, volume 111, pages 397-397.
- [14] R.R.Yager (1981). A procedure for ordering fuzzy subsets of the unit interval. *Inform. Sciences*, volume 24, pages 143-161.
- [15] Y.Yoshida (2003). The valuation of European options in uncertain environment, *European J. Oper. Res.*, volume 145, pages 221-229.
- [16] Y.Yoshida (2003). A discrete-time model of American put option in an uncertain environment, *European J. Oper. Res.*, volume 151, pages 153-166.
- [17] Y.Yoshida (2004). A mean estimation of fuzzy numbers by evaluation measures, in: M.Ch.Negoita, R.J.Howlett and L.C.Jain, eds., *Knowledge-Based Intelligent Information and Engineering Systems, Part II*, LNAI 3214, Springer, pages 1222-1229.
- [18] Y.Yoshida, M.Yasuda, J.Nakagami and M.Kurano (2006). A new evaluation of mean value for fuzzy numbers and its application to American put option under uncertainty, *Fuzzy Sets and Systems*, volume 157, pages 2614-2626.
- [19] Y.Yoshida (2006). Mean values, measurement of fuzziness and variance of fuzzy random variables for fuzzy optimization, *Proceedings of SCIS & ISIS 2006*, Tokyo, pages 2277-2282, Sept. 2006.
- [20] Y.Yoshida (2008). Perception-based estimations of fuzzy random variables: Linearity and convexity, *International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems (IJUFKS)*, 16, suppl. 1 (2008) 71-87.
- [21] Y.Yoshida, A value-at-risk portfolio under uncertainty, submitted.
- [22] Z.Zmeškal (2001). Application of the fuzzy-stochastic methodology to appraising the firm value as a European call option, *European J. Oper. Res.*, volume 135, pages 303-310.
- [23] Z.Zmeškal (2005). Value at risk methodology of international index portfolio under soft conditions (fuzzy-stochastic approach), *Intern. Rev. of Financial Analysis*, volume 14, pages 263-275.
- [24] Z.Zmeškal (2005). Value at risk methodology under soft conditions approach (fuzzy-stochastic approach), *European J. Oper. Res.*, volume 161, pages 337-347.
- [25] L.A.Zadeh (1965). Fuzzy sets, *Inform. and Control*, volume 8, pages 338-353.