

Title	Diverse molecular forms of plasma B-type natriuretic peptide in heart failure.
Author(s)	Nishikimi, Toshio; Minamino, Naoto; Nakao, Kazuwa
Citation	Current heart failure reports (2011), 8(2): 140-146
Issue Date	2011-06
URL	<a href="http://hdl.handle.net/2433/142513">http://hdl.handle.net/2433/142513</a>
Right	The final publication is available at <a href="http://www.springerlink.com">www.springerlink.com</a>
Type	Journal Article
Textversion	author

Editorial Manager(tm) for Current Heart Failure Reports  
Manuscript Draft

Manuscript Number: HF48

Title: Diverse Molecular Forms of Plasma B-type Natriuretic Peptide in Heart Failure

Article Type: Review Article

Corresponding Author: Toshio Nishikimi, M.D., Ph.D.

Corresponding Author's Institution: Kyoto University Graduate School of Medicine

First Author: Toshio Nishikimi, M.D., Ph.D.

Order of Authors: Toshio Nishikimi, M.D., Ph.D.; Naoto Minamino, Ph.D.; Kazuwa Nakao, M.D., Ph.D.

# Diverse Molecular Forms of Plasma B-type Natriuretic Peptide in Heart Failure

Running head: Molecular forms of BNP in heart failure

Toshio Nishikimi, MD, PhD<sup>1)</sup>, Naoto Minamino, PhD<sup>2)</sup>, and Kazuwa Nakao, MD, PhD<sup>1)</sup>

<sup>1)</sup>Department of Medicine and Clinical Science, Kyoto University Graduate School of Medicine, 54, Shogoin-Kawara-cho, Sakyo-ku, Kyoto 606-8507, Japan, and <sup>2)</sup>Department of Molecular Pharmacology, National Cerebral and Cardiovascular Center Research Institute, Fujishirodai, Suita, Osaka 565-8565, Japan

**Total Words counts:** 3844 words

**Key words:** brain natriuretic peptide; atrial natriuretic peptide; heart failure; molecular form; atrial overload; ventricular overload; proBNP-108; BNP-32; processing; secretion; gel-filtration; *O*-glycosylation; natriuretic peptide receptor-A; cGMP,

Address correspondence to:

Toshio Nishikimi, MD, PhD

Department of Medicine and Clinical Science

Kyoto University Graduate School of Medicine

54, Shogoin-Kawara-cho, Sakyo-ku, Kyoto 606-8507, Japan

E-mail: nishikim@kuhp.kyoto-u.ac.jp

TEL: 81-75-751-4287; FAX: 81-75-771-9452

## Abstract

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Recent studies have shown that not only plasma brain natriuretic peptide (BNP)-32, but also plasma proBNP-108 are increased in heart failure (HF) and that current BNP-32 assay kit crossreacts proBNP-108. We also showed that both BNP-32 and proBNP-108 were higher in HF than in normal. The proBNP-108/total BNP (BNP-32+proBNP-108) ratio was widely distributed and HF with ventricular overload had higher proBNP-108/ total BNP ratio than HF with atrial overload. Consistent with this finding, proBNP-108 was the major molecular form in ventricular tissue, and BNP-32 was the major molecular form in atrial tissue. In addition, proBNP-108 was the major molecular form of BNP in pericardial fluid. The proBNP-108/total BNP ratio increased with deterioration of HF and decreased with improvement of HF. Thus, not only BNP-32, but also proBNP-108 is increased in HF and the proBNP/total BNP ratio also rises in association with pathophysiological conditions such as ventricular overload. A new hypothesis that *O*-glycosylation at Thr-71 in a region close to the cleavage site impairs proBNP-108 processing was proposed. In the future, the precise mechanism of increased proBNP-108 in HF should be elucidated.

## Introduction

Atrial natriuretic peptide (ANP) and brain natriuretic peptide (BNP) is a cardiac hormone and a member of the natriuretic family. (1) Both ANP and BNP preferentially bind to natriuretic receptors-A (NPR-A), whereas C-type natriuretic peptide (CNP) specifically binds to NPR-B. The most abundantly expressed in the heart are ANP and BNP, and they are mainly secreted from the cardiac atrium and ventricle, respectively. ANP and BNP play an important role in the regulation of blood pressure and body fluid homeostasis. In addition these peptides have pleiotropic effects, because NPR-A is widely distributed throughout the body (e.g., in the kidneys, vessels, adrenal glands, central nervous system, bone, adipocytes, and heart).

BNP has a strikingly similar to ANP with regard to both its amino acid sequence and its pharmacologic profile (2). Stress to the ventricle such as pressure overload, volume overload, myocardial ischemia, neurohumoral factor and cytokine stimulate BNP gene expression, and proBNP[1-108] (proBNP-108) is produced (3,4,5). Thus, plasma BNP is increased in heart failure and it has been used as a useful biochemical marker for heart failure. When proBNP-108 is secreted from the ventricular myocyte (4,5), it has been thought to be cleaved to proBNP[77-108] (BNP-32) and N-terminal proBNP[1-76] (NT-proBNP-76) in an equimolar fashion by furin, corin or other currently unknown proteases. However, recent studies have shown that not only BNP-32 and NT-proBNP-76, but also proBNP-108 circulates in human plasma and that the proBNP-108 level is also increased in heart failure (6,7,8). Other studies have found that the present assay kit for BNP-32 recognizes proBNP-108 at high cross-reactivity in addition to authentic BNP-32 (9,10).

However, the reason why proBNP-108 is secreted without specific conversion into BNP-32 and NT-proBNP-76 remains unknown. The proBNP-108 levels and relative ratio of proBNP-108 to BNP-32 in plasma from patients with heart failure is not well investigated. In addition, plasma proBNP-108 is very

1 recently reported to be heavily *O*-glycosylated in its N-terminal peptide (11,12).  
2 Thus, it is considered to be important to examine the ratio of proBNP-108 to  
3 BNP-32 in the plasma and its relation to pathophysiological conditions of the  
4 patients with heart failure.  
5  
6  
7

8 We review the current status of BNP studies regarding its molecular forms  
9 and pathophysiology of heart failure, primarily based on our recent report (13).  
10  
11  
12

### 13 **Method for the measurement of the proBNP/total BNP ratio in normal and** 14 **heart failure** 15 16 17

18 First, we compared plasma BNP-32 and proBNP-108 levels in control  
19 subjects, patients with lone atrial fibrillation, and patients with heart failure.  
20 Blood samples (3 mL) were withdrawn from all subjects via the antecubital vein.  
21 Blood was immediately transferred into chilled glass tube containing disodium  
22 EDTA (1 mg/mL) and aprotinin (500 U/mL). Blood was centrifuged immediately  
23 at 4°C and the plasma was frozen and stored at -80° C. Plasma sample was  
24 extracted with Sep-Pak C18 cartridges (Waters, Milford, MA, USA) as previously  
25 reported method (13). The eluate was lyophilized and dissolved in 30%  
26 acetonitrile containing 0.1% TFA. An aliquot of the solution was subjected to gel  
27 filtration high performance liquid chromatography (HPLC) on a TSK gel  
28 G2000SWXL column (7.8 x 300 mm, Tosoh) as previously reported (13). Each  
29 fraction was measured by immunofluorescent BNP assay kit (TOSO, Japan).  
30 ProBNP-108/total BNP ratio was calculated based on the summation of a high  
31 molecular weight (MW) immunoreactive (IR-) proBNP-108 and a low MW  
32 IR-BNP-32 with the following formula: proBNP-108/total BNP ratio =  
33 IR-proBNP-108/(IR-proBNP-108+IR-BNP-32).  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

53 Since very recent studies have revealed that glycosylated pro-BNP of MW  
54 about 35K is circulating in plasma (11,12), we examined their elution positions of  
55 recombinant proBNP-108 (HyTest, Finland), glycosylated proBNP-108 (HyTest,  
56 Finland), and synthetic BNP-32 (Peptide Institute, Japan) in the gel filtration. To  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 evaluate the cross-reactivity of proBNP-108 and glycosylated proBNP-108 in the  
2 immunofluorescent assay (FIA) kit for BNP-32, an aliquot of each peptide was  
3 subjected to acid hydrolysis at 110C for 22 h to estimate peptide contents by  
4 amino acid analysis (L-8500 analyzer, Hitachi, Tokyo, Japan), and another aliquot  
5 was submitted to the FIA for BNP-32, after desalting purchased peptides. In this  
6 system, we were able to accurately evaluate the proBNP/total BNP ratio because  
7 we used one assay system almost equivalently recognizing BNP-32 and  
8 proBNP-108. Some studies evaluated the proBNP/total BNP ratio by using a  
9 distinct assay system for proBNP or BNP-32. In such a study, different affinities  
10 of antibodies against proBNP, BNP-32 and their related peptides in each system  
11 hamper accurate evaluation of the proBNP/total BNP ratio.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

### 26 **Plasma concentrations of IR-BNP-32, IR-proBNP-108 and proBNP/total** 27 **BNP ratios in control, atrial fibrillation, and heart failure.**

28 As shown in Figures 1-A and 1-B, two peaks of IR-BNP were constantly  
29 observed in all cases. The first peak was observed in fractions #9-16 of MW larger  
30 than 13K, and the second peak was in fractions #18-21 of MW 3.5K  
31 corresponding to BNP-32. In this gel filtration HPLC, recombinant proBNP-108  
32 and glycosylated proBNP-108 were eluted mainly in fraction #14 and fraction  
33 #15, being not separable of glycosylated proBNP-108 and non-glycosylated  
34 proBNP-108 with each other. In the following part the term IR-proBNP-108  
35 indicates high MW IR-BNP, which includes both proBNP-108 and glycosylated  
36 proBNP-108, while IR-BNP-32 indicates low MW IR-BNP that is mainly  
37 composed of BNP-32. The proBNP-108/total BNP ratio was narrowly distributed  
38 in control and atrial fibrillation; however, the proBNP-108/total BNP ratio was  
39 widely distributed in patients with heart failure (Figure 1-C). Thus, the mean  
40 proBNP-108/total BNP ratio in heart failure was significantly lower in heart  
41 failure than in control or atrial fibrillation.  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58

59 It is thought that proBNP-108 is cleaved to BNP-32 and N-terminal  
60  
61  
62  
63  
64  
65

1 proBNP-76 in an equimolar fashion by a processing enzyme, when proBNP-108  
2 is secreted from the ventricular myocyte (1,5). However, recent studies have  
3 shown that not only BNP-32, but also proBNP-108 circulates in human plasma  
4 and they are increased in patients with heart failure (7-9). However, there were  
5 few studies, which measured both BNP-32 and proBNP-108 using the same assay  
6 kit. From these results, we confirmed that high and low MW forms of BNP,  
7 BNP-32 and proBNP-108, were present in plasma from control subjects and  
8 patients with atrial fibrillation and heart failure, suggesting that a significant  
9 percentage of proBNP-108 is secreted from the heart without processing.  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20

### 21 **Comparison of plasma ProBNP/total BNP ratio of ventricular overload with** 22 **that of atrial overload** 23 24

25  
26 Next, we measured plasma BNP-32 and proBNP-108 in patients with heart  
27 failure with atrial overload and heart failure with ventricular overload. Heart  
28 failure with atrial overload included mitral stenosis, mitral regurgitation, and  
29 atrial septal defect, whereas heart failure with ventricular overload included aortic  
30 regurgitation and aortic stenosis. We compared the ratio of proBNP-108/total  
31 BNP in heart failure with atrial overload to that in heart failure with ventricular  
32 overload. Two peaks of IR-BNP were observed in both groups. Interestingly,  
33 IR-BNP-32 peak was more dominant than IR-proBNP-108 peak in heart failure  
34 with atrial overload; in contrast, IR-proBNP-108 and IR-BNP-32 peaks were  
35 nearly equivalent in heart failure with ventricular overload. As a result, the mean  
36 proBNP-108/total BNP ratio was higher in heart failure with ventricular  
37 overload than that in heart failure with atrial overload (Figure 2-A). These results suggest  
38 that ventricular overload alters processing of proBNP-108 (13).  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53

### 54 **Tissue concentrations of proBNP and BNP-32 in ventricle and atrium** 55 56

57 To elucidate the mechanism for the higher proBNP-108/total BNP ratio in  
58 ventricular overload than atrial overload, we analyzed molecular forms of BNP in  
59  
60  
61  
62  
63  
64  
65



1 ventricular and atrial tissues. Resected samples of left atrial tissues and left  
2 ventricles were frozen in liquid nitrogen, and stored at  $-80^{\circ}\text{C}$ . The atrial and  
3 ventricular tissues were boiled in 10 volumes of 1 mol/L acetic acid to inhibit the  
4 endogenous peptidase activity (14). The tissues were homogenized, centrifuged,  
5 lyophilized, and subjected to gel filtration HPLC on a TSK gel G2000SWXL  
6 column as described above for plasma. Then, each fraction was submitted to the  
7 FIA for BNP-32 similar to the plasma samples. Two IR-BNP peaks corresponding  
8 to BNP-32 and proBNP-108 were also observed in atrial and ventricular tissue.  
9 Interestingly, in atrial tissue, the low MW IR-BNP peak corresponding to BNP-32  
10 was a dominant molecular form of IR-BNP; in contrast, the high MW IR-BNP  
11 peak, corresponding to proBNP-108 was a dominant molecular form in  
12 ventricular tissue. Consequently, the mean proBNP-108/total BNP ratio was  
13 much higher in ventricular tissue than in atrial tissue (Figure 2-B).  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27

28 There were few studies, which measured molecular forms of BNP in  
29 cardiac tissue. A previous study showed that two molecular forms of IR-BNP of  
30 MW 4K and MW 13-15K were present in human atrial tissue (15), which is  
31 consistent with the present results. However, to our best of knowledge, no study  
32 has previously examined the molecular form of BNP in human ventricular tissue.  
33 Although the exact meaning of dominant presence of proBNP-108 in ventricular  
34 tissue remains unknown at present, these results suggest that alteration of proBNP  
35 processing may occur when some stress is loaded onto ventricle.  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46

### 47 **Pericardial fluid concentrations of IR-BNP-32 and IR-proBNP-108 in** 48 **patients with heart failure** 49

50  
51 To confirm the molecular form of BNP produced and secreted from the  
52 ventricular tissue, we characterized IR-BNP-32 and IR-proBNP-108 in  
53 pericardial fluid and plasma in patients who underwent cardiac surgery. Two  
54 IR-BNP peaks corresponding to BNP-32 and proBNP-108 in plasma and  
55 pericardial fluid in both patients with mitral regurgitation and with aortic stenosis  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 were present. In plasma, IR-BNP-32 was a dominant molecular form in patients  
2 with mitral regurgitation, while IR-proBNP-108 was a dominant molecular form  
3 in patients with aortic stenosis; however, IR-proBNP-108 was exclusively the  
4 dominant molecular form in pericardial fluid in both cases. Consequently, the  
5 mean proBNP-108/total BNP ratio was greater in pericardial fluid than in the  
6 plasma (Figure 2-C).  
7  
8  
9  
10  
11  
12

13 Pericardial fluid is known to contain abundant levels of various bioactive  
14 substances produced in the heart (16,17) and its composition is known to be close  
15 to the interstitial fluid in the ventricle (18). In addition, the concentrations of  
16 bioactive substances, such as adrenomedullin, BNP, ANP, basic fibroblast growth  
17 factor, and vascular endothelial growth factor, have been reported to be higher in  
18 pericardial fluid than in plasma (16-19). Most IR-BNP was present as  
19 IR-proBNP-108, irrespective of the type of heart failure. These results support the  
20 hypothesis again that proBNP-108 is the major molecular form of BNP in the  
21 ventricle and that most of proBNP-108 is secreted from ventricle without  
22 proteolytic processing.  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

### 36 **The post-treatment or natural course of plasma concentrations of** 37 **IR-BNP-32 and IR-proBNP-108 in patients with heart failure** 38 39

40 To investigate whether the pathophysiological status of heart failure affects  
41 the molecular form of BNP in plasma, plasma proBNP-108 and BNP-32 levels  
42 were measured in patients with heart failure before and after their symptom had  
43 improved in response to treatments. These peptides levels were also measured in  
44 patients with heart failure before and after their symptom had deteriorated.  
45 Elevated plasma IR-BNP levels decreased after the treatments, accompanied by a  
46 reduction in the proBNP-108/total BNP ratio. In the cases that heart failure  
47 deteriorated during the observation, IR-BNP levels increased concomitantly with  
48 an increase in the proBNP-108/total BNP ratio.  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59

60 Thus, it is concluded that the proBNP-108/total BNP ratio is altered  
61  
62  
63  
64  
65

1 depending on the pathophysiological status of heart failure. An increased  
2 proBNP-108/total BNP ratio in severe heart failure may be explained in part by  
3 the increased production and secretion of proBNP-108 from the ventricle.  
4 Another possibility is that gene expression of proteolytic processing enzyme is  
5 not increased in parallel with the increase in gene expression of BNP precursor in  
6 severe heart failure, which results in the reduced proteolytic conversion of  
7 proBNP-108 into BNP-32.  
8  
9  
10  
11  
12  
13  
14  
15  
16

### 17 **The pathophysiological significance of increased proBNP in heart failure**

19 In vitro studies have shown remarkably reduced cGMP producing ability of  
20 proBNP-108 in vascular smooth muscle and endothelial cells compared with that  
21 of BNP-32 [11]. The plasma cGMP level is increased in proportion to the severity  
22 of mild to moderate HF, and levels of plasma cGMP correlates with those of  
23 plasma BNP. However, increased cGMP levels are attenuated in severe heart  
24 failure [20], of which mechanism has not been fully elucidated. A definite  
25 increase in the amount of less hormonally active proBNP-108 observed in severe  
26 heart failure might explain this phenomenon. Indeed, the proBNP-108/BNP-32  
27 ratio is increased in decompensated heart failure and that medical therapy for such  
28 patients decreases plasma BNP and patients' symptoms in concert with a  
29 reduction in the proBNP-108/BNP-32 ratio [13]. Therefore, elucidation of the  
30 mechanism underlying the increased ratio of proBNP-108/BNP-32 should help to  
31 clarify the pathogenesis of heart failure and/or pave the way towards novel  
32 therapies.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

### 51 **The mechanism of proBNP secretion without processing: new hypothesis of** 52 **processing of proBNP**

53 Like the increasing levels of proBNP-108, BNP-32 and NT-proBNP-76 in  
54 heart failure, proBNP-108 is also *O*-glycosylated according to heart failure  
55 severity [12,22]. Thus, understanding the clinical relevance of proBNP-108  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

glycosylation is a matter of obvious importance.

Pressure overload, volume overload, ischemia and other conditions stimulate BNP gene transcription. The BNP mRNA is translated in the endoplasmic reticulum to produce preproBNP. Removal of its signal peptide generates proBNP-108 that can be posttranslationally glycosylated to varying degrees at several sites (Ser36, Thr37, Thr44, Thr48, Thr53, Ser58, and Thr71) of its N-terminal region throughout the Golgi network [22]. *O*-glycosylated (glyco-) proBNP-108 is transported to the trans-Golgi network, where it is cleaved to BNP-32 and NT-proBNP-76 probably by furin [22,23]. Both BNP-32 and NT-proBNP-76 are thought to be secreted via a constitutive pathway without storing in secretory granules (Figure 3).

Levels of glyco-proBNP-108, but not of proBNP-108, are increased in the plasma of patients with severe heart failure [24,25]. Western blot analyses have revealed the presence of high MW (20-22 K) IR-BNP and low MW (4 K) IR-BNP species in the plasma of such patients. Digestion with deglycosylation enzymes for 8 h reduced the 20-22K BNP species to about 12K, which is the predicted size of pro-BNP-108, whereas the 4K form, corresponding to BNP-32, remained unchanged [11]. Thus, glyco-proBNP-108 definitely circulates in the plasma of patients with heart failure and is a major molecular form of proBNP.

How glyco-proBNP-108 is secreted without processing under conditions of severe heart failure is not fully understood. A recent study has shown that *O*-glycosylation at Thr71 in a region close to the cleavage site impairs proBNP-108 processing by furin in HEK293 cell, a specific cell line originally derived from human embryonic kidney cells [22,23] (Figure 3). Since the effect of *O*-glycosylation was only evaluated with furin in that vitro study, the roles of other possible processing enzymes, such as corin, remains unclear. It has not been examined whether this finding is also valid in the in the cardiac myocytes of atrium and ventricle in vitro, either. Therefore, further studies using cardiac myocytes are required to elucidate the exact mechanism of proBNP-108

processing and the effect of *O*-glycosylation.

## Conclusion

We review the recent studies regarding complex molecular forms of BNP, in plasma and tissue, primarily based on our recent results. The presence of proBNP-108 in plasma in patients with heart failure has almost been established. Recent studies also showed that the proBNP/total BNP ratio increases in association with pathophysiological conditions such as ventricular overload. An interesting hypothesis that *O*-glycosylation at Thr71 in a region close to the cleavage site impairs proBNP-108 processing by furin is proposed based on the experiments in HEK293 cell (Figure 3). Whether this hypothesis is true in cardiac myocytes should be investigated in the near future.

## ACKNOWLEDGMENTS

We thank Dr. Masashi Ikeda for helpful advice. We thank Mr. Kazumi Akimoto, Ms. Masako Matsubara, Ms. Keiko Ishikawa, Ms. Masako Minato, Ms. Kyoko Tabei, and Ms. Machiko Sakata for technical assistance.

**Funding:** This work was supported in part by Scientific Research Grants-in-Aid [‘18590787’ and ‘20590837’] from the Ministry of Education, Culture, Sports, Science and Technology of Japan, by the Fund from the Suzuken Memorial Foundation, by the Science Research Promotion Fund from the Promotion and Mutual Aid Corporation for Private Schools of Japan; and by the Intramural Research Fund of National Cerebral and Cardiovascular Center of Japan.

**Competing interests:** None.

## FIGURE LEGENDS

### **Figure 1: ProBNP-108/total BNP ratios in plasma in the control, atrial fibrillation and heart failure.**

(A, B) Gel filtration HPLC profile of the venous plasma extracts from control (A) and heart failure (B). Two peaks of high MW and low MW IR-BNPs corresponding to proBNP-108 and BNP-32 were observed. (C) The proBNP-108/total BNP ratios in plasma in the control, atrial fibrillation (Af) and heart failure (HF). (modified Ref. 13)

### **Figure 2: ProBNP-108/total BNP ratios in plasma in heart failure patients with atrial overload and ventricular overload, cardiac tissue, and pericardial fluid**

(A) The proBNP-108/total BNP ratio in heart failure patients with atrial overload and those with ventricular overload. (B) The proBNP-108/total BNP ratio in the atrial and ventricular tissue in patients with heart failure who underwent cardiac surgery. (C) The proBNP-108/total BNP ratios in pericardial fluid and plasma from the same patients who underwent cardiac surgery. (modified Ref. 13)

### **Figure 3. Schematic representation of processing and glycosylation of proBNP in cardiomyocytes.**

Some of the biosynthesized proBNP-108 is *O*-glycosylated in Golgi apparatus. ProBNP-108 is converted into BNP-32 and NT-proBNP-76 by processing enzyme in trans-Golgi network, if *O*-glycosylation does not occur at Thr71. Then, BNP-32 and NT-proBNP-76 is secreted in an equimolar fashion via a constitutive pathway. If proBNP is *O*-glycosylated at Thr71 and is not cleaved by processing enzyme, non-cleaved glycosylated proBNP-108 is secreted into circulation.

**REFERENCES**

1. Minamino N, Horio H, Nishikimi T: Chapter 165. Natriuretic peptides in the cardiovascular system. In: Kastin AJ, ed. *THE HANDBOOK OF BIOLOGICALLY ACTIVE PEPTIDES*. Academic Press, 2006. P. 1217-1225
2. Sudoh T, Kangawa K, Minamino N, Matsuo H: A new natriuretic peptide in porcine brain. *Nature*. 1988;**332**:78-81.
3. Nishikimi T, Maeda N, Matsuoka H: The role of natriuretic peptides in cardioprotection. *Cardiovasc Res*. 2006;**69**:318-28.
4. Daniels LB, Maisel AS: Natriuretic peptides. *J Am Coll Cardiol*. 2007;**50**:2357-68.
5. Weber M, Hamm C: Role of B-type natriuretic peptide (BNP) and NT-proBNP in clinical routine. *Heart*. 2006;**92**:843-9.
6. Lam CS, Burnett JC, Jr., Costello-Boerrigter L, et al.: Alternate circulating pro-B-type natriuretic peptide and B-type natriuretic peptide forms in the general population. *J Am Coll Cardiol*. 2007;**49**:1193-202
7. Waldo SW, Beede J, Isakson S, et al.: Pro-B-type natriuretic peptide levels in acute decompensated heart failure. *J Am Coll Cardiol*. 2008;**51**:1874-82.
8. Seferian KR, Tamm NN, Semenov AG, et al.: The brain natriuretic peptide (BNP) precursor is the major immunoreactive form of BNP in patients with heart failure. *Clin Chem*. 2007;**53**:866-73
9. Shimizu H, Masuta K, Asada H, et al.: Characterization of molecular forms of probrain natriuretic peptide in human plasma. *Clin Chim Acta*. 2003;**334**:233-9.
10. Nishikimi T, Minamino N, Horii K, Matsuoka H: Do commercially available assay kits for B-type natriuretic peptide measure Pro-BNP1-108, as well as BNP1-32? *Hypertension*. 2007;**50**:e163
11. Liang F, O'Rear J, Schellenberger U, et al.: Evidence for functional heterogeneity of circulating B-type natriuretic peptide. *J Am Coll Cardiol*. 2007;**49**:1071-8.
12. Schellenberger U, O'Rear J, Guzzetta A, et al.: The precursor to B-type



1 natriuretic peptide is an O-linked glycoprotein. *Arch Biochem Biophys.*  
2 2006;**451**:160-6.  
3

4 13. \*\*Nishikimi T, Minamino N, Masashi I, et al.: Diversity of Molecular Forms  
5 of Plasma Brain Natriuretic Peptide in Heart failure -Different proBNP-108 to  
6 BNP-32 Ratios in Atrial and Ventricular Overload- *Heart.* 2010; **96**:432-439  
7  
8 *This study first shows the relation between increase of proBNP-108 and*  
9 *ventricular overload in patients with heart failure.*  
10

11 14. Yoshihara F, Nishikimi T, Sasako Y, et al.: Plasma atrial natriuretic peptide  
12 concentration inversely correlates with left atrial collagen volume fraction in  
13 patients with atrial fibrillation: plasma ANP as a possible biochemical marker to  
14 predict the outcome of the maze procedure. *J Am Coll Cardiol.* 2002;**39**:288-94.  
15

16 15. Hino J, Tateyama H, Minamino N, et al.: Isolation and identification of human  
17 brain natriuretic peptides in cardiac atrium. *Biochem Biophys Res Commun.*  
18 1990;**167**:693-700.  
19

20 16. Nishikimi T, Shibasaki I, Iida H, et al.: Molecular forms of adrenomedullin in  
21 pericardial fluid and plasma in patients with ischaemic heart disease. *Clin Sci*  
22 (Lond). 2002;**102**:669-77.  
23

24 17. Nishikimi T, Asakawa H, Iida H, et al.: Different secretion patterns of two  
25 molecular forms of cardiac adrenomedullin in pressure- and volume-overloaded  
26 human heart failure. *J Card Fail.* 2004;**10**:321-7.  
27

28 18. Fujita M, Ikemoto M, Kishishita M, et al.: Elevated basic fibroblast growth  
29 factor in pericardial fluid of patients with unstable angina. *Circulation.*  
30 1996;**94**:610-3.  
31

32 19. Tambara K, Fujita M, Nagaya N, et al.: Increased pericardial fluid  
33 concentrations of the mature form of adrenomedullin in patients with cardiac  
34 remodelling. *Heart.* 2002;**87**:242-6.  
35

36 20. Tsutamoto T, Wada A, Maeda K, et al.: Attenuation of compensation of  
37 endogenous cardiac natriuretic peptide system in chronic heart failure: prognostic  
38 role of plasma brain natriuretic peptide concentration in patients with chronic  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 symptomatic left ventricular dysfunction. *Circulation*. 1997;96:509-16.

2 21. Hammerer-Lercher A, Halfinger B, Sarg B, et al.: Analysis of circulating  
3 forms of proBNP and NT-proBNP in patients with severe heart failure. *Clin Chem*.  
4 2008;54:858-865.  
5  
6

7 22. \*\*Semenov AG, Postnikov AB, Tamm NN, et al.: Processing of pro-Brain  
8 natriuretic peptide is suppressed by O-glycosylation in the region close to the  
9 cleavage site. *Clin Chem*. 2009;55:493-498. *This study first shows the*  
10 *O-glycosylation at Thr-71 in a region close to the cleavage site impairs*  
11 *proBNP-108 processing by furin in HEK293 cell.*  
12  
13  
14  
15  
16  
17  
18

19 23. \*Semenov AG, Tamm NN, Seferian KR, et al.: Processing of pro-B-type  
20 natriuretic peptide: furin and corin as candidate convertases. *Clin Chem*.  
21 2010;56:1166-1176. *This study shows that both furin and corin are involved in*  
22 *the proBNP processing pathway.*  
23  
24  
25  
26

27 24. Luckenbill KN, Christenson RH, Jaffe AS, et al.: Cross-reactivity of BNP,  
28 NT-proBNP, and proBNP in commercial BNP and NT-proBNP assays:  
29 preliminary observations from the IFCC Committee for Standardization of  
30 Markers of Cardiac Damage. *Clin Chem*. 2008;54:619-621  
31  
32  
33  
34  
35

36 25. Seferian KR, Tamm NN, Semenov AG, et al.: Immunodetection of  
37 glycosylated NT-proBNP circulating in human blood. *Clin Chem*.  
38 2008;54:866-873.  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65





