

Title	Grazing on Microcystis (Cyanophyceae) by testate amoebae with special reference to cyanobacterial abundance and physiological state
Author(s)	Mizuta, Seiko; Imai, Hiroyuki; Chang, Kwang-Hyeon; Doi, Hideyuki; Nishibe, Yuichiro; Nakano, Shin-ichi
Citation	Limnology (2011), 12(3): 205-211
Issue Date	2011-12
URL	<a href="http://hdl.handle.net/2433/151697">http://hdl.handle.net/2433/151697</a>
Right	The final publication is available at <a href="http://www.springerlink.com">www.springerlink.com</a>
Type	Journal Article
Textversion	author

1 Grazing on *Microcystis* (Cyanophyceae) by testate amoebae with  
2 special reference to cyanobacterial abundance and physiological  
3 state

4  
5 Seiko Mizuta<sup>1</sup>, Hiroyuki Imai<sup>1</sup>, Chang Kwang-Hyeon<sup>1,2</sup>, Hideyuki Doi<sup>1,3</sup>, Yuichiro  
6 Nishibe<sup>1,4</sup> and Shin-ichi Nakano<sup>1,5\*</sup>

7  
8 <sup>1</sup> LAFWEDY, Ehime University, Tarumi 3-5-7, Matsuyama 790-8566, Ehime, Japan

9 <sup>2</sup> Department of Environmental Science and Engineering, Kyung Hee University,  
10 Seochen-dong 1, Giheung-gu, Yongin-Si, Gyeonggi-Do 446-701, Republic of  
11 Korea

12 <sup>3</sup> Institute for Chemistry and Biology of the Marine Environment, Carl-von-Ossietzky  
13 University Oldenburg, Schleusenstrasse 1 26382 Wilhelmshaven, Germany

14 <sup>4</sup> Tohoku National Fisheries Research Institute, Shinhamacho 3-27-5, Shiogama  
15 985-0001, Miyagi, Japan

16 <sup>5</sup> Center for Ecological Research, Kyoto University, Hirano 2-509-3, Otsu 520-2101,  
17 Shiga, Japan

18  
19 \*Corresponding author: [nakano@ecology.kyoto-u.ac.jp](mailto:nakano@ecology.kyoto-u.ac.jp)

20

21

22

23

1 **Abstract**

2 We examined the growth of testate amoebae preying on *Microcystis* whose  
3 physiological states were different in laboratory experiments and a hypertrophic pond.  
4 We prepared three experimental systems using water samples dominated by *Microcystis*  
5 *aeruginosa*: light incubation (control), dark incubation (dark), and light incubation with  
6 addition of nitrogen and phosphorus (+NP). In all the systems, colony density of *M.*  
7 *aeruginosa* decreased slightly during incubation. Physiological activity of  
8 phytoplankton as determined by chlorophyll fluorescence was high and almost constant  
9 in the control and +NP systems, whereas that decreased in the dark system. Cell  
10 densities of testate amoebae increased in the control and +NP systems, whereas in the  
11 dark system they remained low. Thus, growth of the amoebae was low in the systems  
12 where physiological activity of *Microcystis* was low. In a hypertrophic pond, cell  
13 density of testate amoebae increased and remained high when *M. aeruginosa*  
14 predominated. Cell density of testate amoebae increased remarkably, simultaneously  
15 with the increases in *M. aeruginosa* colony density and phytoplankton physiological  
16 activity. We also found a significant correlation between densities of *M. aeruginosa*  
17 colonies and testate amoebae. We suggested that the physiological activity of  
18 *Microcystis* is one important factor affecting the growth of testate amoebae grazing on  
19 *Microcystis*.

1 Keywords: *Microcystis*, testate amoebae, grazing, physiological state

2

### 3 **Introduction**

4 Blooms of cyanobacteria are notorious symptoms of eutrophication in  
5 freshwaters all over the world, deteriorating water quality as well as the health of human  
6 and natural resources. The genus *Microcystis* is the most frequently found in  
7 cyanobacterial blooms. We already have numerous reports on the physiological and  
8 ecological characteristics of *Microcystis*, and their bloom-forming mechanisms have  
9 been clarified (Reynolds et al. 1981; Oliver and Ganf 2000; Nakano et al. 2001a).  
10 However, loss processes of *Microcystis* populations are not yet fully understood.

11 *Microcystis* abundance is influenced by the usual biological interactions such as  
12 competition, grazing and infection, of which grazing may be the most important loss  
13 process controlling *Microcystis* abundance. Previous studies have reported as possible  
14 grazers of *Microcystis*: protists (Cole and Wynne 1974; Dryden and Wright 1987),  
15 rotifers (Snell 1980; Fulton and Pearl 1987), crustacean zooplankton (Hanazato and  
16 Yasuno 1984; Jarvis et al. 1987), fish (Moriarty 1973, Kawanabe and Mizuno 1989;  
17 Miura 1990). There are only a limited number of rotifers, crustaceans and fish which  
18 graze on *Microcystis* but various protistan species have been shown to do so (Zhang et

1 al. 1996; Nishibe et al. 2002 and 2004; Kim et al. 2006; Wilken et al. 2010). Indeed,  
2 grazing on *Microcystis* by protists occasionally dominates in the collapse of *Microcystis*  
3 blooms (Dryden and Wright 1987). Thus, it is possible that the wax and wane of a  
4 *Microcystis* bloom is dependent on grazing by protists.

5 Among such protistan grazers, rhizopods, including both naked and testate  
6 amoebae, are frequently found to be abundant when significant decreases in *Microcystis*  
7 abundance are detected in lakes, and grazing on *Microcystis* by some rhizopod species  
8 has been demonstrated in laboratory experiments (Yamamoto 1981; Yamamoto and  
9 Suzuki 1984; Nishibe et al. 2004) and field observation (Whitton 1973; Nishibe et al.  
10 2004). Unfortunately, we still have limited eco-physiological information about the  
11 rhizopods which graze on *Microcystis*. Rodriguez-Zaragoza (1994) has reported that  
12 excessive nutrients and elevated water temperatures may be beneficial to common  
13 rhizopod species because such environmental conditions favor bacterial growth, which  
14 in turn feed rhizopods. Nishibe et al. (2004) reported that the abundance of the testate  
15 amoebae *Penardochlamys* sp. which grazes on *Microcystis* was high when *Microcystis*  
16 was abundant in a hypereutrophic pond. High rhizopod abundance with high  
17 *Microcystis* abundance may be reasonable, since the relationship of consumption by a  
18 grazer on various densities of prey follows the Michaelis-Menten equation. However,

1 not only quantity but also the quality of prey is also important for growth of the grazer.  
2 For the Excavata amoebae, Liu et al. (2006) examined the food selection mechanism  
3 and the digestion process of a Vahlkampfiid amoebae *Naegleria* sp. using several  
4 cyanobacterial strains and found that *Microcystis* was inappropriate food for the  
5 amoebae even when the cyanobacteria were heat-killed. By contrast, we still do not  
6 have any information about the effects of prey quality on the growth of testate amoeba  
7 (the Amoebozoa, Unikonts). In addition, no studies have so far examined the  
8 importance of the physiological state of *Microcystis* for the growth of rhizopods until  
9 now.

10 In the present study, we hypothesized that the rhizopods which grazed on  
11 healthy *Microcystis* would grow actively. To examine this hypothesis, we conducted  
12 laboratory experiments in which we fed testate amoebae using *Microcystis* with  
13 different physiological states. We also conducted field monitoring in a hypereutrophic  
14 pond to collect information about seasonal changes in abundance of testate amoebae  
15 together with the abundance and physiological state of the *Microcystis*. This is the first  
16 study which reports the importance of physiological state of *Microcystis* for growth of  
17 rhizopods.

18

## 1 **Materials and methods**

### 2 Laboratory experiment

3           We conducted laboratory experiments using twelve liters of water collected on  
4 30 June and 27 September 2005, from Furuike Pond (33°49'N, 132°48'E), Matsuyama  
5 City, Ehime Prefecture, Japan using a water column sampler to collect an integrated  
6 water sample from the whole water column. The pond is hypertrophic due to  
7 anthropogenic loading from the watershed, and its physical and chemical characteristics  
8 have been described in our previous studies (Nakano et al. 1998 and 2001b; Manage et  
9 al. 1999 and 2001; Nishii et al. 2001). *Microcystis* species usually become dominant in  
10 this pond from May to October (Nakano et al. 1998; Manage et al. 2001; Nishii et al.  
11 2001).

12           We prepared three experimental systems, each in duplicate. A 1.5-liter portion of  
13 the mixed water sample was poured into a 3-liter flask, and  $\text{KH}_2\text{PO}_4$  and  $\text{KNO}_3$  were  
14 added at  $5 \mu\text{mol P L}^{-1}$  and  $80 \mu\text{mol N L}^{-1}$  respectively (+NP system). For the control  
15 system, a 1.5 liter portion of the water sample was poured into a 3 liter flask. These two  
16 systems were then incubated at 25 °C at a photon flux density of  $60 \mu\text{mol m}^{-2} \text{s}^{-1}$  under a  
17 12:12 hour light:dark cycle with daily shaking. It is likely that the light intensity used in  
18 the present study was appropriate, since most species of phytoplankton have the light

1 intensity of saturation in the range of 60 and 100  $\mu\text{E m}^{-2} \text{s}^{-1}$  (Harris 1978). The  
2 remaining 1.5 liters of the water sample were poured into a 3 liter flask and incubated in  
3 the dark at 25 °C with daily shaking. We took a 100 ml subsample from each system  
4 every day and followed changes in density of *Microcystis* colonies, cell density of  
5 amoebae and physiological activity of the phytoplankton.

6 For enumeration of *Microcystis* colonies and amoeba cells, a 50 ml portion of  
7 the water sample was fixed with acidified Lugol's solution at a final concentration of  
8 1% and concentrated by natural sedimentation. *Microcystis* colonies and amoeba cells  
9 were counted in a haematocytometer (Burker-Turk) under a light microscope at a  
10 magnification of  $\times 400$  at least 3 times.

11 A 50 ml portion of the water sample was used to measure the physiological  
12 activity of the phytoplankton using a Water-PAM Chlorophyll Fluorometer  
13 (Heinz-Walz). PAM fluorescence measurements are based on the determination of the  
14 ground fluorescence;  $F_0$  which is measured in weak, constant irradiation of a  
15 dark-adapted sample (all reaction centers in the open state). The maximal fluorescence,  
16  $F_m$ , is measured in a saturation pulse light (all reaction centers in the close state). The  
17 variable fluorescence,  $F_v$ , is calculated as the difference between  $F_0$  and  $F_m$ . The  
18 efficiency of photochemistry of open reaction centers of photosystem II ( $F_v/F_m$ ) was



1 calculated as follows:

$$2 \quad F_v / F_m = (F_m - F_0) \times F_m^{-1}$$

3 Immediately after taking the water samples, a 20 ml subsample was placed in  
4 the dark for 20 min and then we measured the minimum ( $F_0$ ) and maximum ( $F_m$ )  
5 fluorescence yield.

6

7 Field monitoring

8 Weekly field monitoring was conducted from 16 May to 23 November 2006 in Furuike  
9 Pond. Surface water temperature was determined using a thermistor (ABT-1, ALEC  
10 Electronics Co. Ltd.). Water samples were taken as described previously.

11 To determine chlorophyll *a* concentration, 10 ml of each water sample were  
12 filtered through a 0.2  $\mu$ m Nuclepore filter (25 mm in diameter, CORNING Nuclepore)  
13 under negative pressure at 0.05 MPa to retain seston. The filter was then transferred into  
14 a glass tube containing 8 ml of *N,N*-dimethylformamide to extract chlorophyll *a* and  
15 kept in a freezer at -20 °C. The amount of chlorophyll *a* was determined using a  
16 fluorometer (Turner Designs, 10-AU) (Moran and Porath 1980).

17 For enumeration of phytoplankton and testate amoebae, 300 ml of the water  
18 sample was fixed with acid Lugol's solution at a final concentration of 1%.

1 Enumeration of cells of phytoplankton and amoebae were conducted as explained  
2 above.

3         Physiological activity of phytoplankton was determined using another 50 ml  
4 portion of the water sample as described previously.

5

## 6 **Results**

### 7 Laboratory experiment

8 The experiments started from 30 June and 27 September had high reproducibility.

9 During the experimental period, densities of *Microcystis* colonies decreased in all the  
10 systems (Figs. 1A). Almost no difference was found between *Microcystis* colony  
11 densities in the controls and the +NP systems, although those in the dark system were  
12 the lowest (Figs. 1A). *Microcystis aeruginosa* predominated throughout the  
13 experimental period in the control, +NP and dark systems.

14         Physiological activity of the phytoplankton was almost constant in the control and  
15 +NP systems (Figs. 1B). By contrast, physiological activity in the dark system gradually  
16 decreased (Figs. 1B).

17         The dominant amoeba found in the present study was a testate amoeba which  
18 belonged to the genus *Penardochlamys*. We also counted naked amoebae, but their

1 densities were very low compared with those of the testate amoebae (data not shown).  
2 Cell densities of the testate amoeba gradually increased in the control and +NP systems  
3 (Figs. 1C) from 550 cells ml<sup>-1</sup> (0 day) to 3.2 × 10<sup>4</sup> cells ml<sup>-1</sup> (9 day) and from 330 cells  
4 ml<sup>-1</sup> (0 day) to 3.8 × 10<sup>4</sup> cells ml<sup>-1</sup> (9 day), respectively. The testate amoeba grew in the  
5 dark system (Fig. 1C). However, its growth was negligible, ranging between 660 cells  
6 ml<sup>-1</sup> (0 day) and 9500 cells ml<sup>-1</sup> (6 day) (Fig. 1C).

7

#### 8 Field monitoring

9 Chlorophyll *a* concentration in Furuike Pond increased from May to June, reaching their  
10 maximum (723 μg l<sup>-1</sup>) on 13 June, remained relatively high in July and August with  
11 fluctuations and then gradually decreased in September and October (Fig. 2A).  
12 Physiological activity of the phytoplankton showed cyclic oscillations from May to  
13 October, ranging between 0.194 and 0.528 (Fig. 2B). Dominant phytoplankton species  
14 during the study period were *Microcystis aeruginosa* and *M. wesenbergii*.

15 Densities of *M. aeruginosa* colonies fluctuated widely between May and June,  
16 remained low in July and August (Fig. 2C), then increased to high densities recorded in  
17 September and October, followed by a decrease in November (Fig. 2C). *M. wesenbergii*  
18 colony density increased from 16 May to 24 July and became almost stable from 31

1 July onwards, although a relatively high density ( $15.4 \times 10^3$  colonies  $\text{ml}^{-1}$ ) was found on  
2 17 November (Fig. 2C).

3 The increase in cell density of testate amoebae was slow between 16 May (37 cells  
4  $\text{ml}^{-1}$ ) and 21 August (222 cells  $\text{ml}^{-1}$ ), followed by a rapid increase to 30 August (1037  
5 cells  $\text{ml}^{-1}$ ) (Fig. 3A). The maximum density was recorded on 21 September (1593 cells  
6  $\text{ml}^{-1}$ ), and then the cell density decreased from 27 September onwards (Fig. 3A).

7 The percentage of testate amoebae attached to *Microcystis* colonies was relatively  
8 high between June and July, and between September and October, but in August was  
9 negligible (Fig. 3B). Relatively high densities of testate amoebae were found on *M.*  
10 *wesenbergii* colonies between June and July, whereas the densities of amoebae attached  
11 to *M. aeruginosa* colonies between August and October were higher than those on *M.*  
12 *wesenbergii* colonies (Fig. 3C).

13 Pearson Correlation Analysis showed that there was no significant relationship  
14 between cell density of testate amoebae and concentration of chlorophyll *a*, or  
15 physiological activity of the phytoplankton (Table 1). We did find a significant  
16 correlation between the densities of testate amoebae and *M. aeruginosa* ( $r = 0.7664$ ,  $n =$   
17  $25$ ,  $p < 0.001$ ), but the correlation between the densities of testate amoebae and *M.*  
18 *wesenbergii* ( $r = 0.1562$ ,  $n = 25$ ) was insignificant (Table 1).

1

## 2 **Discussion**

### 3 Laboratory experiment

4       During the present study, colony density of *Microcystis* decreased in all our  
5 experimental systems, despite the fact that we added large amounts of N and P to the  
6 +NP systems (see Materials and methods). The patterns of decrease in *Microcystis*  
7 colony density in the +NP systems were similar to those in the control systems (Fig. 1A),  
8 indicating that the *Microcystis* in the present study was not subjected to N or P  
9 limitation. Thus, it is likely that another element(s) was responsible for the decrease in  
10 *Microcystis* colony density, since light was available in the +NP and the control systems.  
11 All experiments in the present study were conducted in batch cultures where no  
12 additional nutrients were supplied after the beginning of the experiment. This was not  
13 the case for CO<sub>2</sub>, because CO<sub>2</sub> would be supplied by mixing in each system when we  
14 took subsamples. However, we did not bubble-mix the systems, and CO<sub>2</sub> supply in our  
15 experiments might have been insufficient for *Microcystis* growth. Thus, we think that  
16 the decrease in *Microcystis* colony density in our experimental systems was due to  
17 carbon limitation. Even if carbon was limiting for *Microcystis*, the decrease in colony  
18 density was small in the +NP and control systems, and we had a variety of physiological

1 activities in our systems. We therefore believe that carbon limitation on *Microcystis* in  
2 the present study does not affect our interpretation of the results.

3 The dominant amoeba found in the present study belonged to the testate amoeba  
4 *Penardochlamys*. It has been reported that *Microcystis* is the only prey for  
5 *Penardochlamys* (Nishibe et al. 2004), suggesting that the abundance of amoebae  
6 depend on that of the cyanobacteria. However, in the dark system, growth of the testate  
7 amoeba was low (Figs. 1C) in spite of high *Microcystis* colony densities (Fig. 1A). Thus,  
8 the low growth of the testate amoeba could not be explained by *Microcystis* abundance.  
9 In the dark system, physiological activity of the phytoplankton predominated by  
10 *Microcystis* decreased (Figs. 1B), although the *Microcystis* colony density was high  
11 (Figs. 1A). Thus, growth of the testate amoebae was low in the systems where the  
12 physiological activity of *Microcystis* was low which suggests that the physiological  
13 activity of *Microcystis* is responsible for changes in growth of the testate amoebae.

14

#### 15 Field monitoring

16 In our field monitoring we found a significant logarithmic correlation between  
17 chlorophyll *a* concentration and *Microcystis* colony density (*M. aeruginosa* plus *M.*  
18 *wesenbergii*) (n = 25, r = 0.464, p<0.05). Thus, the phytoplankton physiological activity

1 shown in Fig. 2B can be regard as that of *Microcystis*. However, we did not find any  
2 significant correlations between concentration of chlorophyll *a* and cell density of  
3 testate amoebae, or between physiological activity of phytoplankton and cell density of  
4 testate amoebae (Table 1). These results suggest the importance of prey species for  
5 growth of testate amoebae. There was a clear succession of dominant *Microcystis*  
6 species (Fig. 2C), and *M. aeruginosa* predominated during September (Fig. 2C) when  
7 cell density of testate amoebae increased and remained high (Fig. 3A). In addition, a  
8 significant correlation between densities of *M. aeruginosa* colonies and testate amoebae  
9 (Table 1) and higher densities of testate amoebae attached to *M. aeruginosa* colonies  
10 (Fig. 3C) suggest that prey availability is important for the growth of this testate  
11 amoebae, and that the food linkage between *M. aeruginosa* and testate amoebae in the  
12 present study is to some extent species-specific.

13       However, this is contrary to the results of our previous study (Nishibe et al. 2004)  
14 where the food linkage between *Microcystis* and testate amoebae was not  
15 species-specific. Indeed, also in the present study, percentages of amoebae attached to  
16 *M. wesenbergii* colonies (Fig. 3C) were high relative to those on *M. aeruginosa* colonies  
17 during the dominance of *M. wesenbergii* (Fig. 2C), although overall cell density of  
18 testate amoebae was low (Fig. 3A). The dominant testate amoebae species described in

1 Nishibe et al. (2004) was *Penardochlamys* sp., and this might be the case in the present  
2 study. Some previous studies have reported that some rhizopods collected from natural  
3 waters seemed to have strong feeding selectivity on specific prey (Cook et al. 1974;  
4 Becares and Romo 1994), but others have shown that some rhizopods have a wide range  
5 of prey within cyanobacterial species (Ho and Alexander 1974; Yamamoto and Suzuki  
6 1984; Laybourn-Parry et al. 1987). We need to collect more information about  
7 abundance and composition of amoebae attached to *Microcystis* colonies In order to  
8 understand which amoebae species are important as grazers of the cyanobacteria.

9 In early and mid-August, colony density and physiological activity of both *M.*  
10 *aeruginosa* and *M. wesenbergii* remained low (Fig. 2), and cell density of testate  
11 amoebae was also low (Fig. 3A). Moreover, almost no amoebae were attached to  
12 *Microcystis* colonies during that period (Fig. 3B). Cell density of testate amoebae  
13 markedly increased from late August (Fig. 3A), simultaneously with the increases in  
14 density of *M. aeruginosa* colonies (Fig. 2C) and phytoplankton physiological activity  
15 (Fig. 2B). Thus, our field monitoring suggested that not only prey availability but also  
16 prey quality are important for growth of testate amoebae grazing on *Microcystis*,  
17 although we did not find any significant correlation between physiological activity of  
18 phytoplankton and cell density of testate amoebae (Table 1).



1

## 2 **Conclusions**

3 Rhizopods have long been considered to be of minor importance in the food webs  
4 of freshwater and marine systems. However, food linkage between rhizopods and  
5 *Microcystis* may provide another important role for amoebae in aquatic food webs.  
6 From the results obtained in the present study, we concluded that physiological activity  
7 of *Microcystis* is another important factor which affects the growth of testate amoebae  
8 that graze on *Microcystis*. The chemical composition of *Microcystis* probably varies,  
9 depending on the physiological state of these cyanobacteria. Thus, changes in growth of  
10 testate amoebae may be affected by the chemical composition or nutritional value of  
11 their prey. Unfortunately, we did not examine the chemical composition of the  
12 *Microcystis* in the present study. Further studies are required to elucidate changes in the  
13 growth of amoebae in relation to changes in the chemical composition or nutritional  
14 value of their prey. In addition, not only are biological but also physico-chemical  
15 variables most likely to be responsible for growth of amoebae, and further studies to  
16 examine which environmental factors affect the abundance and composition of amoebae  
17 are needed to elucidate their ecology.

18

1 **Acknowledgments** We thank Dr. Kotaro Takayama, Ehime University for teaching  
2 us the PAM technique. Thanks are also due to the members of LAFWEDY, Ehime  
3 University, for their advice, discussions and encouragement throughout the study. We  
4 thank Dr. M. J. Morris for correction of our English and constructive comments on the  
5 manuscript. The present study was partly supported by the Center of Excellence (COE)  
6 Program at the “Global Center of Excellence for Interdisciplinary Studies on  
7 Environmental Chemistry”, by the Espec Foundation for Global Environment Research  
8 and Technology and by the Environment Research and Technology Development Fund  
9 (D-0905) of the Ministry of Environment, Japan.

10

1   **References**

- 2   Becares E, Romo S (1994) Selective predation of *Thecamoeba sphaeronucleolus*  
3       (Greeff, 1891) on filamentous algae in natural conditions. J Gen Appl Microbiol  
4       40:15–21
- 5   Cole GT and Wynne MJ (1974) Endocytosis of *Microcystis aeruginosa* by *Ochromonas*  
6       *danica*. J Phycol 10:397-410
- 7   Cook WL, Ahearn DG, Reinhardt DJ, Reiber RJ (1974) Blooms of an algophorous  
8       amoeba associated with *Anabeba* in a freshwater lake. Water Air Soil Pollut  
9       3:71-80
- 10   Dryden RC, Wright SJL (1987) Predation of cyanobacteria by protozoa. Can J  
11       Microbiol 33:471-482
- 12   Fulton RS, Paerl HW (1987) Toxic and inhibitory effects of the blue-green alga  
13       *Microcystis aeruginosa* on herbivorous zooplankton. J Plankton Res 9: 837-855
- 14   Hanazato T, Yasuno M (1984) Growth, reproduction and assimilation of *Moina*  
15       *macrocopa* fed on *Microcystis* and/or *Chlorella*. Jpn J Limnol 34: 195-202
- 16   Harris GP (1978) Photosynthesis, productivity and growth: the physiological ecology of  
17       phytoplankton. Arch Hydrobiol Beih Ergebn Limnol 10: 1-177
- 18   Ho TS, Alexander M (1974) The feeding of amoebae on the filamentous blue-green

- 1 algae. Bot Bull Acad Sin 23:63-70
- 2 Jarvis AC, Hart RC, Combrink S (1987) Zooplankton feeding on size fractionated  
3 *Microcystis* colonies and *Chlorella* in a hypertrophic lake (Hartbeespoort Dam,  
4 South Africa): implications to resource utilization and zooplankton succession. J  
5 Plankton Res 9: 1231-1249
- 6 Kawanabe H, Mizuno T (1989) Freshwater fishes in Japan. Yama-to-Keikoku-sha Co.  
7 Ltd. Tokyo
- 8 Kim BR, Nakano S, Kim BH, Han MS (2006) Growth and grazing of the heterotrophic  
9 nanoflagellate, *Diphyllia rotans* on the cyanobacterium *Microcystis aeruginosa*.  
10 Aquat Microb Ecol 45: 163-170
- 11 Laybourn-Parry J, Jones K, Holdich JP (1987) Grazing by *Mayorella* sp. (Protozoa:  
12 Sarcodina) on cyanobacteria. Funct Ecol 1:99–104
- 13 Liu X, Shi M, Liao Y, Gao Y, Zhang Z, Wen D (2006) Feeding Characteristics of an  
14 Amoeba (*Lobosea: Naegleria*) Grazing Upon Cyanobacteria: Food Selection,  
15 Ingestion and Digestion Progress. Microb Ecol 51, 315–325
- 16 Manage PM, Kawabata Z, Nakano S (1999) Seasonal changes in densities of  
17 cyanophage infectious to *Microcystis aeruginosa* in a hypereutrophic pond.  
18 Hydrobiologia 411:211-216

- 1 Manage, P. M., Z. Kawabata and S. Nakano (2001) Dynamics of cyanophages and  
2 algicidal bacteria causing *Microcystis aeruginosa* mortality. *Limnology* 2: 73-78
- 3 Miura T (1990) The effects of planktivorous fishes on the plankton community in a  
4 eutrophic lake. *Hydrobiologia* 200/210: 567-579
- 5 Moran R, Porath D (1980) Chlorophyll determination in intact tissues using N,  
6 N-dimethylformamide. *Plant Physiol* 65: 478-479
- 7 Moriarty DJW (1973) The physiology of digestion of blue-green algae in the cichlid  
8 fish, *Tilapia nilotica*. *J Zool* 171: 25-39
- 9 Nakano S, Ishii N, Manage PM, Kawabata Z (1998) Trophic roles of heterotrophic  
10 nanoflagellates and ciliates among planktonic organisms in a hypereutrophic  
11 pond. *Aquat Microb Ecol* 16:153-161
- 12 Nakano S, Hayakawa K, Frenette JJ, Nakajima T, Jiao C, Tsujimura S, Kumagai M  
13 (2001a) Cyanobacterial blooms in a shallow lake: a large-scale enclosure assay  
14 of the importance of diurnal stratification. *Arch Hydrobiol* 150:491-509
- 15 Nakano S, Manage PM, Nishibe Y, Kawabata Z (2001b) Trophic linkage among  
16 heterotrophic nanoflagellates, ciliates and metazoan zooplankton in a  
17 hypereutrophic pond. *Aquat Microb Ecol* 25:259-270
- 18 Nishibe Y, Kawabata Z, Nakano S (2002) Grazing on *Microcystis aeruginosa* by the

- 1 heterotrophic flagellate *Collodictyon triciliatum* in a hypertrophic pond. *Aquat*  
2 *Microb Ecol* 29:173-179
- 3 Nishibe Y, Manage PM, Kawabata Z, Nakano S (2004) Trophic coupling of testate  
4 amoeba and *Microcystis* species in a hypertrophic pond. *Limnology* 5:71-76
- 5 Nishii K, Nakano S, Tamada M, Manage PM, Nishibe S, Kawabata Z (2001) Microbial  
6 decomposition of dissolved organic matter in a hypertrophic pond. *Limnology*  
7 2:207-212
- 8 Oliver RL, Ganf GC (2000) Freshwater blooms. In: *Ecology of Cyanobacteria: Their*  
9 *Diversity in Time and Space* (Eds B.A Whitton & M. Potts) pp.149-194, Kluwer  
10 Academic Publishers, Dordrecht.
- 11 Reynolds CS, Jaworski GHM, Cmiench HA, Leedale GF (1981) On the annual cycle of  
12 the blue-green alga *Microcystis aeruginosa* Kutz. emend. Elenkin. *Phil Trans R*  
13 *Soc Lond Ser B* 293:419-477
- 14 Rodriguez-Zaragoza S (1994) Ecology of free-living amoebae. *Crit Rev Microbiol*  
15 20:225-241
- 16 Snell TW (1980) Blue-green algae and selection in rotifer populations. *Oecologia* 46:  
17 343-346
- 18 Yamamoto Y (1981) Observation on the occurrence of microbial agents which cause

- 1 lysis of blue-green algae in lake Kasumigaura. Jap J Limnol 42:20-27
- 2 Yamamoto Y, Suzuki K (1984) Light and electron microscope observations and prey  
3 specificities of an algophorous amoeba from japanese freshwater. J Gen Appl  
4 Microbiol 31:411-417
- 5 Whitton BA (1973) Interactions with other organism. In: Carr NG, Whitton BA (eds)  
6 The biology of blue-green algae. Blackwell, Oxford, pp 415-433
- 7 Wilken S, Wiezer S, Huisman J, Van Donk E (2010) Microcystins do not provide  
8 anti-herbivore defence against mixotrophic flagellates. Aquat Microb Ecol  
9 59:207-216
- 10 Zhang X and Watanabe MM, Inouye I (1996) Light and electron microscopy of grazing  
11 by *Poteroochromonas malhamensis* (Chrysophycease) on a range of  
12 phytoplankton taxa. J Phycol 32:486-492

13

1

Table 1 Pearson Correlation Analysis between testate amoebae density and phytoplankton variables.

	r	Significance	Relationship
Chlorophyll <i>a</i> concentration	0.116	Not significant	Negative
Fv / Fm	0.076	Not significant	Positive
Colony density of <i>M. aeruginosa</i>	0.766	p<0.001	Positive
Colony density of <i>M. wesenbergii</i>	0.156	Not significant	Negative

2




1 Figure captions

2 Fig. 1 Changes in colony density of *Microcystis aeruginosa* (A), physiological  
3 activity of *M. aeruginosa* (B) and cell density of testate amoebae (C) in the  
4 experiment started on 30 June 2005. Vertical bars that indicate differences  
5 between duplicates are shown when they exceeded the size of the symbol.

6 Fig. 2 Seasonal changes in chlorophyll *a* concentration (A), phytoplankton  
7 physiological activity (B) and colony density of *Microcystis aeruginosa* and *M.*  
8 *wesenbergii* (C) in Furuike Pond between May and November 2006.

9 Fig. 3 Seasonal changes in cell density of testate amoebae (A), the percentage of  
10 testate amoebae attached to *Microcystis* colonies (B) and the percentage of  
11 testate amoebae attached to *Microcystis aeruginosa* colonies (white part) or to *M.*  
12 *wesenbergii* colonies (gray part) (C) in Furuike Pond between May and  
13 November 2006.


14



1


2 Fig. 1 Mizuta et al.

1



2

3 Fig. 2 Mizuta et al.



1

2 Fig. 3 Mizuta et al.