

| | |
|-------------|--------------------------------------------------------------------------------------------------------------------|
| Title | Nonparametric model reconstruction for stochastic differential equations from discretely observed time-series data |
| Author(s) | Ohkubo, Jun |
| Citation | Physical Review E (2011), 84(6) |
| Issue Date | 2011-12 |
| URL | http://hdl.handle.net/2433/152425 |
| Right | ©2011 American Physical Society |
| Type | Journal Article |
| Textversion | publisher |

Nonparametric model reconstruction for stochastic differential equations from discretely observed time-series data

Jun Ohkubo*

Graduate School of Informatics, Kyoto University, Yoshida Hon-machi, Sakyo-ku, Kyoto-shi, Kyoto 606-8501, Japan

(Received 21 September 2011; revised manuscript received 1 November 2011; published 14 December 2011)

A scheme is developed for estimating state-dependent drift and diffusion coefficients in a stochastic differential equation from time-series data. The scheme does not require to specify parametric forms for the drift and diffusion coefficients in advance. In order to perform the nonparametric estimation, a maximum likelihood method is combined with a concept based on a kernel density estimation. In order to deal with discrete observation or sparsity of the time-series data, a local linearization method is employed, which enables a fast estimation.

DOI: [10.1103/PhysRevE.84.066702](https://doi.org/10.1103/PhysRevE.84.066702)

PACS number(s): 05.10.Gg, 05.45.Tp, 87.18.Tt

I. INTRODUCTION

Recently, it has been clarified that stochastic nature in small systems such as cells plays an important role in dynamics and behavior of biological systems [1–3]. In addition, due to recent experimental developments such as single-molecule spectroscopy, it becomes possible to obtain time-series data for various stochastic phenomena. From a theoretical point of view, it is important to develop methods for analysis of the time-series data, and actually there are many studies for the analysis of the single-molecule time series (for example, see Refs. [4–6]).

Parameter estimations from observed time-series data are also important research topics. If one obtains experimental data for a specified biochemical system, parameters in the specified biochemical system can be estimated from the experimental data. The biochemical system could be modeled by using a master equation or a stochastic differential equation (a Langevin equation). The master equation or the stochastic differential equation for the specified biochemical system has some parameters (e.g., reaction rates). If the reaction rates are estimated from the experimental data, we will obtain a reconstructed model, which would reproduce the experimental data adequately. The reconstructed model enables us to perform more detailed numerical simulations and to have deep insights for the phenomenon. In recent years, a discrete property or sparsity in observations has attracted much attention; it would be difficult to completely observe the phenomenon and to obtain detailed time-series data, and then we would perform the estimation from discretely observed time-series data. For example, estimation procedures based on Markov chain Monte Carlo methods [7–9] and variational methods [10,11] have been proposed for the problem of the discrete observations. (In addition, there is a recent review article [12] for the estimation problem.)

Here, we consider the following situation: We know that time-series data can be modeled with a stochastic differential equation, but specific forms of drift and diffusion coefficients of the stochastic differential equation are unknown in advance. That is, there is no prior knowledge about time-series data, except for some basic properties such as a memoryless property. While there are some works for the parametric

estimation based on a maximum likelihood method [13,14], simple applications of these parametric estimations are not suitable for our problem here, and a nonparametric estimation scheme is needed. For example, we here focus on a bimodal distribution of a chemical substance. Relations between the bimodal distributions and stochasticity have been discussed experimentally [15,16] and theoretically [17,18]. Although one may consider that the bimodal distribution is produced from a double-well potential system, it has been known that the bimodal distribution can also be produced from state-dependent noise [19,20]. In addition, a recent study indicates that such noise-induced bimodality may play an important role in decision making in a noisy environment [21,22]. In these situations, it is necessary to judge whether bimodal distribution is produced from a double-well potential system or a state-dependent noise, and it is enough to estimate how the drift and diffusion coefficients of the stochastic differential equation depend on the state. For the nonparametric estimations, there are many studies in various research fields. For example, in Ref. [23], a method based on estimations of Kramers-Moyal coefficients has been proposed. However, in the Kramers-Moyal coefficients estimation, adjoint Fokker-Planck equations should be solved numerically, which needs additional computational costs. In order to perform nonparametric estimations for complicated systems, fast algorithms are required.

In the present paper, we develop a nonparametric model-reconstruction method for a stochastic differential equation from discretely observed time-series data. A kind of local estimations is employed in order to extract the state dependency in the nonparametric estimation. In order to perform the local estimations efficiently, we propose a maximum likelihood method combined with a concept based on a kernel density estimation, which has been studied a lot and widely used for nonparametric density estimation [24]. The difficulty caused from the discrete observations is dealt with a local linearization method [25,26], which enables us to approximate a nonlinear stochastic differential equation by a locally linear stochastic differential equation. In addition, a useful second-order form suitable for the local linearization method is proposed. We demonstrate that the combination of these ideas enables us to estimate the state-dependent drift and diffusion coefficients from only a small set of discretely observed time-series data.

*ohkubo@i.kyoto-u.ac.jp

The present paper is constructed as follows. In Sec. II, we give problem settings and an example of time-series data. In Sec. III, we briefly review a kernel density estimation, and the scheme is reformulated from a different point of view; we show that a maximum likelihood method reproduces the kernel density estimation adequately. Section IV is the main part in the present paper; an explicit estimation scheme based on the local linearization method is explained. Examples of estimation results for the problem introduced in Sec. II are given in Sec. V. Section VI is the conclusion.

II. PROBLEM SETTINGS

The aim of our estimation problem is to reconstruct a stochastic differential equation only from observed time-series data. In order to demonstrate the estimation, we here use the following Ito-type stochastic differential equation as a toy model:

$$dx_t = f^{\text{true}}(x_t)dt + g^{\text{true}}(x_t)dW_t, \quad (1)$$

where W_t is a Wiener process, $f^{\text{true}}(x)$ and $g^{\text{true}}(x)$ are state-dependent drift and state-dependent diffusion coefficients defined as

$$f^{\text{true}}(x) = -4x^3 + 4x, \quad g^{\text{true}}(x) = 0.2 \sin(\pi x), \quad (2)$$

respectively. This model has state-dependent drift and diffusion coefficients. In addition, this form of the stochastic differential equation suggests that the potential landscape is a double-well form, and there are two stable points around $x = 1$ and $x = -1$. Using the Euler-Maruyama scheme [27], we generate an original path for the stochastic differential equation (1). In the generation of the original path, we employ a time interval of $\Delta t = 0.001$. After the generation of the original path, we sampled data points discretely, which corresponds to discrete observations. Although the time interval of the observation can be varied, we here take data points at equally spaced time interval $\Delta t = 0.05$ for simplicity. The generated time-series data are depicted in Fig. 1. Hereafter, we must forget the stochastic differential equation (1) and the state dependency of the drift and diffusion coefficients in Eq. (2), and only the time-series data in Fig. 1 is focused and analyzed.

Our aim is to analyze the time-series data in Fig. 1 under a situation that we do not have any information about the original model. A first guess may be as follows: It seems that there are two stable points around $x \sim 1.0$ and $x \sim -1.0$, and the time spent around $x \sim -1.0$ seems to be a little longer than that around $x \sim 1.0$. Is the difference caused by a potential landscape, or other reasons? A simple way to solve this problem is to reconstruct an explicit model that reproduces the time-series data. Hence, the problem settings are as follows: Estimate $\hat{f}(x)$ and $\hat{g}(x)$ in

$$dx_t = \hat{f}(x_t)dt + \hat{g}(x_t)dW_t, \quad (3)$$

from time-series data $\{(X_i, T_i) | i = 1, \dots, N\}$, where each data point X_i is observed at time T_i , and N is the total number of the observed data points. Note that there is no prior knowledge about the state dependencies of $\hat{f}(x)$ and $\hat{g}(x)$, except that these coefficients are time independent. It could be possible to




FIG. 1. (a) Observed time-series data. There are totally 2000 points observed discretely. (b) Enlarged figure of (a). Each circle is an observation with $\Delta t = 0.05$, and the solid thin line is an original path which is produced using the Euler-Maruyama approximation with $\Delta t = 0.001$.

deal with time-dependent cases, but it is beyond the scope of the present paper.

III. KERNEL DENSITY ESTIMATION

In order to develop a scheme to estimate the drift and diffusion coefficients in Eq. (3), we first explain a kernel density estimation, which gives us many insights for our final aim.

A. Brief review

A first and simple step to analyze time-series data is to construct a probability density $p(x)$ for the observed data. A histogram, in which the x coordinate is split into several bins and the numbers of data points within the bins are counted, is a simple method to estimate the probability density $p(x)$. However, the histogram is based on a discrete approximation. One of the most widely used methods for nonparametric density estimation is a kernel density estimation [24,28]. The kernel density estimation has been recently studied in the context of biophysics [29], in which applications for the forced unfolding and unbinding data for proteins are discussed.

In the kernel density estimation, a non-negative real function $K(x)$ (i.e., a kernel function) is used. The kernel function satisfies the normalization condition, $\int_{-\infty}^{\infty} K(x)dx = 1$, and it has a zero first moment, $\int_{-\infty}^{\infty} xK(x)dx = 0$, and a finite second moment $\int_{-\infty}^{\infty} x^2K(x)dx < \infty$. In the present paper, a parameter h , a so-called bandwidth, is introduced explicitly in the kernel function, and we write the kernel function as




FIG. 2. Probability density function $p(x)$. The solid line corresponds to an exact solution of Eq. (1). The estimated density from the observed time-series data is depicted with the dashed line. The bimodality of the density is reproduced, but heights of the peaks are different from the exact one because of the small size of the data in Fig. 1.

$K_h(x) \equiv K(x/h)/h$. Using the kernel function, the probability density is estimated as

$$\hat{p}(x) = \frac{1}{N} \sum_{i=1}^N K_h(x - X_i). \quad (4)$$

There are some kernel functions, and in the present paper, we consider only a Gaussian kernel $K_h(x) = (1/\sqrt{2\pi}h) \exp[-x^2/(2h^2)]$, which has been widely used.

The remaining task for the kernel density estimation is the choice of the bandwidth h of the kernel function. Various choices have been studied, and a famous data-driven method is a method based on cross validation [30]. In the cross-validation method, the following risk function is minimized with respect to the bandwidth h for the Gaussian kernel:

$$\hat{Q} = A + B \sum_{i < j} [\exp(-\Delta_{ij}^2/4) - C \exp(-\Delta_{ij}^2/2)], \quad (5)$$

where

$$A = (2Nh\sqrt{\pi})^{-1}, \quad B = (N^2h\sqrt{\pi})^{-1}, \\ C = 2\sqrt{2}N/(N-1), \quad \Delta_{ij} = (X_i - X_j)/h.$$

Figure 2 shows the estimated probability density. Here, we used $h = 0.06476$, which is selected based on the risk function (5). The solid line corresponds to an exact solution obtained from Eq. (1). Because of the small size of the data in Fig. 1, there are differences between the estimated probability density and the exact solution. We note that if there is longer time-series data, better estimate results are obtained.

B. Kernel density estimation through maximum likelihood estimation

The estimation of the probability density is not the aim in the present paper, but it is helpful for us to reformulate the kernel density estimation from the viewpoint of a maximum likelihood estimation; this discussion gives us a way to reconstruct a stochastic differential equation without any prior knowledge.

The maximum likelihood estimation enables us to estimate parameters in a statistical model (for example, see Ref. [31]). In the context of the density estimation, the probability density plays a role as the parameters, and we seek probable probability density function $\hat{p}(x)$ from observed data.

Due to the concept of the kernel function, a contribution from one observed data point should be distributed according to the kernel function. For example, assume that we observe a data point $X = 1.0$. Although we have only one data point, here we introduce many replicas for the observation. Each replica has a different pseudo-observation. For instance, when there are four replicas and the replicas have pseudo-data-points $X = 1.0, 1.2, 0.7, 1.2$, the probability with which we observe the total replicas is

$$p(X = 1.0)(p(X = 1.2))^2 p(X = 0.7).$$

There are four pseudo-observations for only one real observation, and then one may consider the fourth root,

$$[p(X = 1.0)]^{1/4} [p(X = 1.2)]^{2/4} [p(X = 0.7)]^{1/4}.$$

Note that the frequency of the pseudo-observation of X is the power index of each probability. Extending this discussion and using the kernel function instead of the frequency of pseudo-observation, we construct the kernelized likelihood function, as follows. Using a discretization of the x coordinate as $x = j\Delta x$, a probability with which we observe a distributed data point X_i is written as

$$\lim_{\Delta x \rightarrow 0} \prod_{j=-\infty}^{\infty} p(j\Delta x)^{K_h(j\Delta x - X_i)}.$$

Note that if we use a δ -like function as the kernel function, only a contribution from $p(X_i)$ remains, and a usual interpretation without the kernel function is recovered. Using a notation $\mathbf{X} = \{X_1, \dots, X_N\}$, a likelihood function $L[\hat{p}(x)|\mathbf{X}]$ is written as

$$L[\hat{p}(x)|\mathbf{X}] = \lim_{\Delta x \rightarrow 0} \prod_{i=1}^N \prod_{j=-\infty}^{\infty} \hat{p}(j\Delta x)^{K_h(j\Delta x - X_i)}, \quad (6)$$

where the hat of \hat{p} means that this is just the parameter to be estimated. Hence, the log-likelihood function is

$$l(\hat{p}(x)|\mathbf{X}) = \sum_{i=1}^N \int_{-\infty}^{\infty} dx_i [\ln \hat{p}(x_i)] K_h(x_i - X_i). \quad (7)$$

In order to obtain the maximum likelihood estimates for $\hat{p}(x)$, we take a functional derivative of Eq. (7) with respect to $\hat{p}(x)$ under a constraint $\int dx \hat{p}(x) = 1$. A Lagrange multiplier λ is introduced and we consider a maximization of the following function:

$$\sum_{i=1}^N \int_{-\infty}^{\infty} dx_i [\ln \hat{p}(x_i)] K_h(x_i - X_i) + \lambda \left(\int dx \hat{p}(x) - 1 \right).$$

Taking the functional derivative with respect to $\hat{p}(x)$ and setting the functional derivative is equal to zero, we obtain

$$\sum_{i=1}^N \frac{1}{\hat{p}(x)} K_h(x - X_i) + \lambda = 0, \quad (8)$$

and therefore

$$\hat{p}(x) = \frac{1}{\lambda} \sum_{i=1}^N K_h(x - X_i). \quad (9)$$

Inserting Eq. (9) into the constraint condition, we have

$$1 = \frac{1}{\lambda} \int_{-\infty}^{\infty} dx \sum_{i=1}^N K_h(x - X_i) = \frac{N}{\lambda}. \quad (10)$$

Hence, $\lambda = N$ and we recover Eq. (4).

The above discussion indicates that the concept of distributed data points in pseudo-observation corresponds to the kernel function. It would be expected that the combination of the distributed data points and the maximum likelihood method gives us a more flexible estimation scheme than the usual one.

It is straightforward to extend the above discussions to an estimation of a conditional density. Given $\mathcal{D} = \{(X_1, Y_1), \dots, (X_N, Y_N)\}$, a sample of independent observations from the distribution of (X, Y) , we want to obtain the estimation of the conditional density $\hat{p}(y|x)$. In this case, the log-likelihood function should be set as

$$\begin{aligned} l(\hat{p}(y|x)|\mathcal{D}) \\ = \sum_{i=1}^N \int_{-\infty}^{\infty} dy_i [\ln \hat{p}(y_i|x)] \tilde{K}_w(y_i - Y_i) K_W(x - X_i), \end{aligned} \quad (11)$$

where $\tilde{K}_w(y)$ is a kernel function for the y coordinate, and $K_W(x)$ is that for the x coordinate. Using the similar discussion as Eq. (7), we obtain the following conditional density estimator

$$\hat{p}(y|x) = \frac{\sum_{i=1}^N K_W(x - X_i) \tilde{K}_w(y - Y_i)}{\sum_{i=1}^N K_W(x - X_i)}, \quad (12)$$

which is the same as a conventional conditional density estimator [32,33].

Figure 3 shows a schematic illustration of the kernel conditional density estimator. We here consider a conditional density $p(y|x_0)$. The kernel functions \tilde{K}_w with bandwidth w are distributed according to the observations. The conditioning $x = x_0$ is carried out by another kernel function in the x coordinate. The kernel function K_W has a bandwidth W , and the center of the kernel function is at $x = x_0$. The estimation of the conditional density is performed by summing the N kernel functions in the y coordinate, weighted by the kernel function in the x coordinate.

For $\{(X_1, Y_1), \dots, (X_N, Y_N)\}$ being random samples from a population having a density $p(x, y)$, there are many studies for estimating conditional densities and properties of the conditional densities [34–36]. The problem in the present paper is different from these studies; $\{(X_1, Y_1), \dots, (X_N, Y_N)\}$ are not generated from an identical density (see the next section). The maximum likelihood method developed in this section enables us to deal with the nonparametric estimation for a stochastic differential equation, and we will propose the method in the next section.




FIG. 3. A schematic illustration of the kernel density estimate $p(y|x_0)$. Each data point has a weight, which is smoothly distributed according to the kernel function. The conditioning on $x = x_0$ is carried out by another kernel function in the x coordinate.

IV. PROPOSED METHOD

A. Basics

We are now ready for constructing a method to estimate drift and diffusion coefficients in a stochastic differential equation from time-series data.

Firstly, the observed data $\{(X_i, T_i) | i = 1, \dots, N\}$ is converted to a slightly different form $\mathcal{D} = \{(X_i, \Delta X_i, \Delta t_i) | i = 1, \dots, N - 1\}$, where $\Delta X_i = X_{i+1} - X_i$ and $\Delta t_i = T_{i+1} - T_i$. This conversion means that when a current coordinate is X_i , we have an amount of change ΔX_i during the time interval Δt_i . For each i , the time interval can be varied in general, and then the amount of change ΔX_i depends on Δt_i ; $\{(X_i, \Delta X_i)\}$ is not from an identical density. Due to an assumption that the time series has a Markov property, $\{(X_i, \Delta X_i)\}$ are independent of each other.

Secondly, we consider a conditional probability density $p(\Delta x|x, \theta_x)$, which has a set of parameters θ_x . The parameters θ_x depend on a specific coordinate x , and the dependency of θ_x on x is unknown in advance. Note that the parametrization of the conditional probability density is not related to the parametrization of drift and diffusion coefficients in the stochastic differential equation in Sec. II. Using the localized parameters θ_x , the drift and diffusion coefficients only for a specific coordinate x could be estimated.

Thirdly, we consider the following log-likelihood function:

$$\begin{aligned} l(\theta_x|\mathcal{D}) = \sum_{i=1}^{N-1} \int_{-\infty}^{\infty} d(\Delta x_i) [\ln p(\Delta x_i|x, \theta_x)] \\ \times \tilde{K}_w(\Delta x_i - \Delta X_i) K_W(x - X_i). \end{aligned} \quad (13)$$

Maximizing the above log-likelihood function with respect to the parameters θ_x , it is possible to estimate the localized parameter θ_x adequately.

A remaining task is to specify the conditional probability density $p(\Delta x|x, \theta_x)$. If a stochastic differential equation is linear, the conditional probability density is expressed as a

normal distribution (i.e., ΔX_i obeys a normal distribution with mean E_i and variance V_i , where E_i and V_i depend on X_i and Δt_i). For a nonlinear stochastic differential equation, the description based on the normal distribution is impossible in general. However, if the conditional probability density cannot be written as the normal distribution, the calculation scheme would become very complicated. Hence, we here assume that the conditional probability density is written as the normal distribution. The restriction with the normal distribution seems to be severe, but we will discuss a method to approximate the nonlinear stochastic differential equation to a locally linear stochastic differential equation. The assumption of the Gaussian form gives the conditional probability density,

$$\ln p(\Delta x_i | x, \theta_x) = -\frac{1}{2} \left\{ \frac{(\Delta x_i - E_i)^2}{V_i} + \ln(2\pi V_i) \right\}, \quad (14)$$

where E_i and V_i depend on X_i , Δt_i , and the parameters θ_x .

B. Simple estimation

We here discuss the most simple case (i.e., $E_i = \mu_{x_0} \Delta t_i$, $V_i = \sigma_{x_0}^2 \Delta t_i$) and $\theta_{x_0} = \{\mu_{x_0}, \sigma_{x_0}\}$. This means that for the estimation at $x = x_0$, we assume a stochastic differential equation with constant drift and diffusion coefficients. Note that this constant property is assumed only for the estimation at $x = x_0$, and we do not assume that $\hat{f}(x)$ and $\hat{g}(x)$ in Eq. (3) are constant for all x . Repeating the estimation of μ_{x_0} and σ_{x_0} for various points $x = x_0$, we obtain the estimated drift and diffusion coefficients as $\hat{f}(x) = \mu_x$ and $\hat{g}(x) = \sigma_x$, respectively.

The above procedure is enough for the estimation, but here we give a discussion for the choice of the kernel bandwidth. For simplicity, we here assume that the kernel \tilde{K}_w is the Gaussian kernel, and that $\Delta t_i = 1$ for all i . In addition, we consider a simple case in which $X_i = x_0$ for all i (i.e., all current coordinates are at x_0). Hence, the kernel function K_W is constant for all i . In this case, the following log-likelihood function is obtained:

$$\begin{aligned} l(\theta_{x_0} | \mathcal{D}) &\propto -\frac{1}{2} \sum_{i=1}^{N-1} \int_{-\infty}^{\infty} d(\Delta x_i) \left\{ \frac{(\Delta x_i - \mu_{x_0})^2}{\sigma_{x_0}^2} + \ln(2\pi \sigma_{x_0}^2) \right\} \\ &\quad \times \tilde{K}_w(\Delta x_i - \Delta X_i) \\ &= -\frac{1}{2} \sum_{i=1}^{N-1} \left\{ \frac{(\Delta X_i - \mu_{x_0})^2}{\sigma_{x_0}^2} + \frac{w^2}{\sigma_{x_0}^2} + \ln(2\pi \sigma_{x_0}^2) \right\} \end{aligned} \quad (15)$$

The maximum likelihood method gives

$$\mu_{x_0} = \frac{1}{N-1} \sum_{i=1}^{N-1} \Delta X_i \quad (16)$$

and

$$\sigma_{x_0}^2 = \frac{1}{N-1} \sum_{i=1}^{N-1} (\Delta X_i - \mu_{x_0})^2 + w^2. \quad (17)$$

Here, note that the unbiased estimator for the variance is given as

$$\bar{\sigma}_{x_0}^2 = \frac{1}{N-2} \sum_{i=1}^{N-1} (\Delta X_i - \mu_{x_0})^2. \quad (18)$$

Comparing Eq. (17) with Eq. (18), it is clear that the bandwidth w should be taken small enough for large N . In our problem settings in the present paper, it is possible to consider that $w \simeq 0$. Hence, the kernel $\tilde{K}_w(\Delta x_i - \Delta X_i)$ is replaced as a Dirac δ function $\delta(\Delta x_i - \Delta X_i)$ hereafter.

Combining the above all discussions, we obtain the following simple estimation scheme.

Algorithm 1 (simple method)

- (i) Set a bandwidth W .
- (ii) For a point x_0 , maximize the following log-likelihood function with respect to $\theta_{x_0} = \{\mu_{x_0}, \sigma_{x_0}\}$:

$$\begin{aligned} l(\theta_{x_0} | \mathcal{D}) &= -\frac{1}{2} \sum_{i=1}^{N-1} \left\{ \frac{(\Delta X_i - \mu_{x_0} \Delta t_i)^2}{\sigma_{x_0}^2 \Delta t_i} + \ln(2\pi \sigma_{x_0}^2) \right\} \\ &\quad \times K_W(x - X_i), \end{aligned} \quad (19)$$

i.e., calculate the following quantities:

$$\mu_{x_0} = \frac{\sum_{i=1}^{N-1} \Delta X_i K_W(x - X_i)}{\sum_{i=1}^{N-1} \Delta t_i K_W(x - X_i)}, \quad (20)$$

$$\sigma_{x_0}^2 = \frac{\sum_{i=1}^{N-1} (\Delta X_i - \mu_{x_0} \Delta t_i)^2 K_W(x - X_i) / \Delta t_i}{\sum_{i=1}^{N-1} K_W(x - X_i)}. \quad (21)$$

- (iii) Repeat (ii) for various x_0 .
- (iv) Estimate the drift and diffusion coefficients as $\hat{f}(x) = \mu_x$ and $\hat{g}(x) = \sigma_x$.

C. Local linearization method and second-order approximation

In Sec. IV B, the local drift and diffusion coefficients have simple forms, so that we easily solve the stochastic differential equation explicitly. Note that if we have nonlinear drift and diffusion coefficients, we cannot obtain an explicit solution for the stochastic differential equation exactly in general. However, using a local linearization method [25,26], it is possible to obtain the approximate solution with a Gaussian form for the nonlinear stochastic differential equation. Hence, the estimation scheme developed in Sec. IV A is available even for the nonlinear cases. We can assume arbitrary drift and diffusion coefficients, and a second-order approximation, which will be introduced soon, is one of the tractable schemes.

We first note that the diffusion coefficient in the stochastic differential equation must be positive for all x ; this fact needs an additional constraint for the optimization procedures. We, therefore, use the following second-order approximation; $\hat{f}(x)$ and $\hat{g}(x)$ around x_0 is approximated as

$$\hat{f}_{x_0}(x) = \mu_{x_0}^{(0)} + \mu_{x_0}^{(1)}(x - x_0) + \frac{1}{2} \mu_{x_0}^{(2)}(x - x_0)^2, \quad (22)$$

$$\hat{g}_{x_0}(x) = \exp(s_{x_0}^{(0)} + s_{x_0}^{(1)}(x - x_0) + \frac{1}{2} s_{x_0}^{(2)}(x - x_0)^2), \quad (23)$$

and $\theta_{x_0} = \{\mu_{x_0}^{(0)}, \mu_{x_0}^{(1)}, \mu_{x_0}^{(2)}, s_{x_0}^{(0)}, s_{x_0}^{(1)}, s_{x_0}^{(2)}\}$. Due to the exponential form in Eq. (23), the diffusion coefficient $\hat{g}_{x_0}(x)$ is positive for all x .

We comment that a final estimation for the drift and diffusion coefficients should be performed as $\hat{f}(x) = \mu_x^{(0)}$, and $\hat{g}(x) = \exp[s_x^{(0)}]$, respectively, as discussed in Sec. IV B.

A stochastic differential equation with the state-dependent drift coefficient $\hat{f}_{x_0}(x)$ and the state-dependent diffusion coefficient $\hat{g}_{x_0}(x)$ is nonlinear, and the local linearization method gives an analytical solution in a Gaussian form. As a result, the conditional probability density $p(\Delta x_i | x, \theta_x)$ can be written as a Gaussian distribution. We briefly explain the local linearization method in the Appendix, and only the consequence is shown here. We note that the kernel function \tilde{K}_w is replaced with the Dirac δ function according to the discussion in Sec. IV B. The estimation scheme based on the local linearization method is as follows:

Algorithm 2 (LL method)

- (i) Set a bandwidth W .
- (ii) For a point x_0 , maximize the following log-likelihood function with respect to $\theta_{x_0} = \{\mu_{x_0}^{(0)}, \mu_{x_0}^{(1)}, \mu_{x_0}^{(2)}, s_{x_0}^{(0)}, s_{x_0}^{(1)}, s_{x_0}^{(2)}\}$:

$$l(\theta_{x_0} | \mathcal{D}) = -\frac{1}{2} \sum_{i=1}^{N-1} \left\{ \frac{(\phi_{x_0} - E_i)^2}{V_i} + \ln(2\pi V_i) - s_{x_0}^{(0)} - s_{x_0}^{(1)}(X_i - x_0) - \frac{s_{x_0}^{(2)}}{2}(X_i - x_0)^2 \right\} \times K_W(x - X_i), \quad (24)$$

where

$$\phi_{x_0} = \int_{X_i - x_0}^{X_i + \Delta X_i - x_0} du \exp\left(-s_{x_0}^{(0)} - s_{x_0}^{(1)}u - \frac{1}{2}s_{x_0}^{(2)}u^2\right), \quad (25)$$

and

$$F_i = \frac{\hat{f}_{x_0}(X_i)}{\hat{g}_{x_0}(X_i)} - \frac{\hat{g}_{x_0}(X_i)}{2} [s_{x_0}^{(1)} + s_{x_0}^{(2)}(X_i - x_0)], \quad (26)$$

$$L_i = \mu_{x_0}^{(1)} + \mu_{x_0}^{(2)}(X_i - x_0) - \hat{f}_{x_0}(X_i) [s_{x_0}^{(1)} + s_{x_0}^{(2)}(X_i - x_0)] - [(\hat{g}_{x_0}(X_i))^2 \left(\frac{s_{x_0}^{(2)}}{2} + \frac{[s_{x_0}^{(1)} + s_{x_0}^{(2)}(X_i - x_0)]^2}{2} \right)], \quad (27)$$

$$M_i = \frac{\hat{g}_{x_0}(X_i)}{2} \left\{ \mu_{x_0}^{(2)} - s_{x_0}^{(2)} \hat{f}_{x_0}(X_i) - [\mu_{x_0}^{(1)} + \mu_{x_0}^{(2)}(X_i - x_0)] [s_{x_0}^{(1)} + s_{x_0}^{(2)}(X_i - x_0)] - 2[\hat{g}_{x_0}(X_i)]^2 s_{x_0}^{(2)} [s_{x_0}^{(1)} + s_{x_0}^{(2)}(X_i - x_0)] - [\hat{g}_{x_0}(X_i)]^2 [s_{x_0}^{(1)} + s_{x_0}^{(2)}(X_i - x_0)]^3 \right\}, \quad (28)$$

$$E_i = \frac{F_i}{L_i} (e^{L_i \Delta t_i} - 1) + \frac{M_i}{L_i^2} (e^{L_i \Delta t_i} - 1 - L_i \Delta t_i), \quad (29)$$

$$V_i = \frac{e^{2L_i \Delta t_i} - 1}{2L_i}. \quad (30)$$

(iii) Repeat (ii) for various x_0 .

(iv) Estimate the drift and diffusion coefficients as $\hat{f}(x) = \mu_x^{(0)}$ and $\hat{g}(x) = \exp(s_x^{(0)})$.

For step (ii), various standard numerical maximization or minimization algorithms are available. We note that the function ϕ_{x_0} can be written using an error function or an imaginary error function.

V. ESTIMATION RESULTS

We apply the two algorithms in Sec. IV to the discretely observed data in Sec. II. In the numerical experiments, we use the Gaussian kernel.

We should first set the kernel bandwidth W . It would be possible to select the kernel bandwidth W using some criteria, for example, using a cross-validation method. However, here we set W heuristically. A simple choice for W is the optimized bandwidth for the kernel density estimator discussed in Sec. III. In Figs. 4(a) and 4(d), we show the results obtained from the simple estimation (Algorithm 1) and the estimation based on the local linearization (Algorithm 2), using $W = 0.06476$. Because the bandwidth is narrow and the number of data points, N , is small, the estimation results have large fluctuations, as shown in Figs. 4(a) and 4(b).

In the estimation based on the local linearization, the second-order approximation is used, and then the bandwidth W could be taken larger than that of the kernel density estimator; intuitively, it would be reasonable to select a bandwidth three or five times as large as the optimized bandwidth for the kernel density estimator. Larger bandwidth W enables us to use a larger number of effective data points for the estimation. Of course, if large W is employed, the second-order approximation becomes invalid, and hence the estimation results would be worse. We here set $W = 0.3$ ad hoc. Figures 4(c) and 4(d) are the estimated results. As expected, the estimation results become smooth because a larger number of effective data points is available. Due to the simple assumption for Algorithm 1, the estimated results are not good; the usage of the larger bandwidth W gives a kind of averaging effect. In contrast, the estimation based on the local linearization (Algorithm 2) gives good estimates.

The above results are only for one time-series data in Fig. 1. Next, we generated 200 time-series data with the same parameters, and checked the validity of our proposed method. Figures 4(e) and 4(f) show results of averages of the drift and diffusion coefficients for 200 trajectories, respectively. Here, we used $W = 0.3$ for the estimations. The error bars in Figs. 4(e) and 4(f) are standard deviations of the results. From the results, we see that the estimation method based on the local linearization (Algorithm 2) works well, especially for the estimation of the drift coefficients.

Based on results on other numerical experiments, we comment on the choice of the bandwidth W as follows: The choice of the bandwidth W should be small, but if the number of data points is not enough, a slightly larger bandwidth would be better. One may intuitively estimate the bandwidth from the probability density $\hat{p}(x)$. Although a method of the choice of the bandwidth is beyond the scope of the present paper, the above choice would be enough in a practical sense.




FIG. 4. Estimation results for the drift coefficients $\hat{f}(x)$ [panels (a), (c), and (e)], and for the diffusion coefficients $\hat{g}(x)$ [panels (b), (d), and (f)]. Panels (a)–(d) show results for only one trajectory in Fig. 1, and panels (e) and (f) are results of averages for 200 trajectories. The filled circle (Simple) corresponds to the results for the simple Euler scheme, and the empty boxes (LL) are those for the local linearization method. In (a) and (b), the bandwidth of the kernel is $W = 0.06476$, and in (c), (d), (e), and (f), $W = 0.3$. Solid curves in left panels [panels (a), (c), and (e)] and in right panels [panels (b), (d), and (f)] correspond to $f^{\text{true}}(x)$ and $g^{\text{true}}(x)$ in Eq. (2), respectively. Error bars in panels (e) and (f) are the standard deviations.

VI. CONCLUSION

In the present paper, we developed a nonparametric estimation scheme for a stochastic differential equation from discretely observed time-series data. In order to make the estimation scheme, a concept based on a kernel density estimation was extended and a kernelized likelihood function was derived. The word, kernelized, means that an observed point is distributed with a bandwidth. In addition, we employ a local linearization method in order to deal with the discrete property or sparsity of the observations. A second-order approximation was introduced, in which the diffusion coefficient is restricted to be positive naturally. This avoids adding any constraint for the maximization or minimization for the log-likelihood

function in the algorithm. Using a toy model, we demonstrated that the estimation method based on the local linearization method works well.

Although results are not shown, we applied the estimation schemes to several different models, and confirmed that they work well. In addition, we performed numerical experiments for cases with larger Δt_j . If we have larger Δt_j , results become worse. This is because that the approximation of the local linearization is inadequate for the larger time interval. For the large-interval cases, the Kramers-Moyal coefficients estimation would be available [23], as explained in Sec. I. However, as stated before, some additional computational costs are needed. On the other hand, our estimation scheme is

based on an approximated analytical solution of a stochastic differential equation, and then the computational time is largely reduced. For example, a computational time for one point x_0 in Algorithm 2 is a few seconds in a laptop computer. In this sense, our estimation method is a complementary one with previous works.

Finally, we comment that the estimation scheme for stochastic differential equations could be extended to multivariate cases straightforwardly, because the local linearization method has already been formulated for multivariate cases [25]. In addition, estimations in the presence of strong measurement noise [37–39] should be considered in future works.

ACKNOWLEDGMENTS

The author thanks T. J. Kobayashi for motivating this work. This work was supported in part by grant-in-aid for scientific research (Grants No. 20115009 and No. 21740283) from the Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan.

APPENDIX: LOCAL LINEARIZATION METHOD

The local linearization method [25,26] is one of the useful approximations for nonlinear stochastic differential equations. The basic concept of the local linearization method is that the nonlinear stochastic differential equation is approximated locally as a linear stochastic differential equation. It would be possible to obtain the same consequence using a Fokker-Planck equation. In Refs. [25,26], an approximation for the stochastic differential equation is explicitly given, and for the reader’s convenience, we briefly review the local linearization method. For details, see Refs. [25,26].

For simplicity, we here consider a one-dimensional stochastic process x_t satisfying

$$dx_t = A(x_t)dt + B(x_t)dB_t, \tag{A1}$$

where $A(x_t)$ is twice continuously differentiable with respect to x_t , $B(x_t)$ is a continuously differentiable function of x_t , and B_t is a standard Brownian motion. The above stochastic differential equation can be transformed into a more tractable equation as follows:

$$dz_t = \left(A \frac{d\phi}{dx} + \frac{B^2}{2} \frac{d^2\phi}{dx^2} \right) dt + dB_t, \tag{A2}$$

where $z_t = \phi(x_t)$ and $\phi(x_t)$ satisfies an ordinary differential equation $B \frac{d\phi}{dx} = 1$. Ito’s formula immediately gives Eq. (A2).

Hence, we here only consider the following stochastic differential equation with a constant diffusion coefficient:

$$dx_t = A(x_t)dt + dB_t. \tag{A3}$$

In the local linearization method, the drift term $A(x_t)$ is locally approximated by a linear function of x_t . Using Ito’s formula, we have

$$dA = \frac{1}{2} \frac{\partial^2 A}{\partial x^2} dt + \frac{\partial A}{\partial x} dx. \tag{A4}$$

Here, an assumption that both coefficients in Eq. (A4) are constant for a small interval $[s, t)$ gives

$$A(x_t) - A(x_s) = \frac{1}{2} \frac{\partial^2 A}{\partial x^2} \Big|_{x=x_s} (t - s) + \frac{\partial A}{\partial x} \Big|_{x=x_s} (x_t - x_s). \tag{A5}$$

Hence, we obtain the drift coefficient $A(x_t)$ as

$$A(x_t) = L_s x_t + M_s t + N_s, \tag{A6}$$

where

$$L_s = \frac{\partial A}{\partial x} \Big|_{x=x_s}, \quad M_s = \frac{1}{2} \frac{\partial^2 A}{\partial x^2} \Big|_{x_s},$$

$$N_s = A(x_s, s) - \frac{\partial A}{\partial x} \Big|_{x=x_s} x_s - \frac{1}{2} \frac{\partial^2 A}{\partial x^2} \Big|_{x=x_s} s.$$

Finally, we obtain a linear stochastic differential equation as follows:

$$dx_t = (L_s x_t + M_s t + N_s)dt + dB_t. \tag{A7}$$

The linear stochastic differential equation can be solved analytically, and the solution is

$$x_t = x_s + \frac{A(x_s)}{L_s} (e^{L_s(t-s)} - 1) + \frac{M_s}{L_s^2} (e^{L_s(t-s)} - 1 - L_s(t-s)) + \int_s^t e^{L_s(t-u)} dB_u, \tag{A8}$$

where the fourth term follows the Gaussian distribution with mean 0 and variance $\{\exp[2L_s(t-s)] - 1\}/(2L_s)$. As a result, $x_t - x_s$ follows the Gaussian distribution with mean

$$\frac{A(x_s)}{L_s} (e^{L_s(t-s)} - 1) + \frac{M_s}{L_s^2} [e^{L_s(t-s)} - 1 - L_s(t-s)]$$

and variance $\{\exp[2L_s(t-s)] - 1\}/(2L_s)$.

Combining the above results, the variable transformation $z_t = \phi(x_t)$, and its Jacobian, we finally obtain the conditional probability used in Eq. (24)

[1] C. V. Rao, D. M. Wolf, and A. P. Arkin, *Nature (London)* **420**, 231 (2002).
 [2] M. B. Elowitz, A. J. Levine, E. D. Siggia, and P. S. Swain, *Science* **297**, 1183 (2002).
 [3] M. Kærn, T. C. Elston, W. J. Blake, and J. J. Collins, *Nature Rev. Genetics* **6**, 451 (2005).
 [4] A. Baba and T. Komatsuzaki, *Proc. Natl. Acad. Sci. USA* **104**, 19297 (2007).
 [5] C.-B. Li, H. Yang, and T. Komatsuzaki, *J. Phys. Chem. B* **113**, 14732 (2009).
 [6] M. Miyazaki and T. Harada, *J. Chem. Phys.* **134**, 085108 (2011).
 [7] A. Golightly and D. J. Wilkinson, *Biometrics* **61**, 781 (2005).

- [8] R. J. Boys, D. J. Wilkinson, and T. B. L. Kirkwood, *Stat. Comput.* **18**, 125 (2008).
- [9] Y. Wang, S. Christley, E. Mjolsness, and X. Xie, *BMC Syst. Biol.* **4**, 99 (2010).
- [10] A. Ruttor and M. Opper, *Phys. Rev. Lett.* **103**, 230601 (2009).
- [11] M. D. Vrettas, D. Cornford, M. Opper, and Y. Shen, *Neurocomputing* **73**, 1186 (2010).
- [12] R. Friedrich, J. Peinke, M. Sahimi, and M. R. R. Tabar, *Phys. Rep.* **506**, 87 (2011).
- [13] D. Kleinhans, R. Friedrich, A. Nawrothb, and J. Peinkeb, *Phys. Lett. A* **346**, 42 (2005).
- [14] D. Kleinhans and R. Friedrich, *Phys. Lett. A* **368**, 194 (2007).
- [15] M. Acar, A. Becskei, and A. van Oudenaarden, *Nature (London)* **435**, 228 (2005).
- [16] Y. Toyooka, D. Shimosato, K. Murakami, K. Takahashi, and H. Niwa, *Development* **135**, 909 (2008).
- [17] M. N. Artyomov, J. Das, M. Kardar, and A. K. Chakraborty, *Proc. Natl. Acad. Sci. USA* **104**, 18958 (2007).
- [18] L. M. Bishop and H. Qian, *Biophys. J.* **98**, 1 (2010).
- [19] Y. Togashi and K. Kaneko, *Phys. Rev. Lett.* **86**, 2459 (2001).
- [20] J. Ohkubo, N. Shnerb, and D. A. Kessler, *J. Phys. Soc. Jpn.* **77**, 044002 (2008).
- [21] T. J. Kobayashi, *Phys. Rev. Lett.* **104**, 228104 (2010).
- [22] T. J. Kobayashi, *Phys. Rev. Lett.* **106**, 228101 (2011).
- [23] C. Honisch and R. Friedrich, *Phys. Rev. E* **83**, 066701 (2011).
- [24] B. W. Silverman, *Density Estimation for Statistics and Data Analysis* (Chapman & Hall, Boca Raton, 1998).
- [25] R. Biscay, J. C. Jimenez, J. J. Riera, and P. A. Valdes, *Ann. Inst. Statist. Math.* **48**, 631 (1996).
- [26] I. Shoji and T. Ozaki, *Stoch. Anal. Appl.* **16**, 733 (1998).
- [27] P. E. Kloeden and E. Platen, *Numerical Solution of Stochastic Differential Equations, Corrected Third Printing* (Springer, Berlin, 1999).
- [28] M. Rosenblatt, *Ann. Math. Stat.* **27**, 832 (1956).
- [29] E. E. Bura, A. Zhmurov, and V. Barsegov, *J. Chem. Phys.* **130**, 015102 (2009).
- [30] M. Rudemo, *Scand. J. Stat.* **9**, 65 (1982).
- [31] R. V. Hogg, J. W. McKean, and A. T. Craig, *Introduction to Mathematical Statistics*, 6th ed. (Pearson Prentice Hall, Upper Saddle River, NJ, 2005).
- [32] R. J. Hyndman, D. M. Bashtannyk, and G. K. Grunwald, *J. Comput. Graph. Stat.* **5**, 315 (1996).
- [33] D. M. Bashtannyk and R. J. Hyndman, *Comp. Stat. Data Anal.* **36**, 279 (2001).
- [34] J. Fan, *J. Am. Statist. Assoc.* **87**, 998 (1992).
- [35] D. Ruppert and M. P. Wand, *Ann. Stat.* **22**, 1346 (1994).
- [36] J. Fan, Q. Yao, and H. Tong, *Biometrika* **83**, 189 (1996).
- [37] F. Böttcher, J. Peinke, D. Kleinhans, R. Friedrich, P. G. Lind, and M. Haase, *Phys. Rev. Lett.* **97**, 090603 (2006).
- [38] D. Kleinhans, R. Friedrich, M. Wächter, and J. Peinke, *Phys. Rev. E* **76**, 041109 (2007).
- [39] B. Lehle, *Phys. Rev. E* **83**, 021113 (2011).