

Title	Kinetic analysis of reverse transcriptase activity of bacterial family A DNA polymerases.
Author(s)	Yasukawa, Kiyoshi; Konishi, Atsushi; Shinomura, Mayu; Nagaoka, Eriko; Fujiwara, Shinsuke
Citation	Biochemical and biophysical research communications (2012), 427(3): 654-658
Issue Date	2012-10-26
URL	http://hdl.handle.net/2433/161657
Right	© 2012 Elsevier Inc.
Type	Journal Article
Textversion	author

1 Biochem. Biophys. Res. Commun.

2

3 **Kinetic analysis of reverse transcriptase activity of bacterial family A DNA**
4 **polymerases**

5

6 Kiyoshi Yasukawa ^{a,*}, Atsushi Konishi ^a, Mayu Shinomura ^a, Eriko Nagaoka ^b,

7 Shinsuke Fujiwara ^{b,c}

8

9 ^a*Division of Food Science and Biotechnology, Graduate School of Agriculture, Kyoto*
10 *University, Sakyo-ku, Kyoto 606-8502, Japan*

11 ^b*Department of Bioscience, Graduate School of Science and Technology,*
12 *Kwansei-Gakuin University, 2-1 Gakuen, Sanda, Hyogo 669-1337, Japan*

13 ^c*Research Center for Environmental Bioscience, Graduate School of Science and*
14 *Technology, Kwansei-Gakuin University, 2-1 Gakuen, Sanda, Hyogo 669-1337, Japan*

15

16 * Corresponding author. Fax: +81-75-753-6265.

17 E-mail address: yasukawa@kais.kyoto-u.ac.jp (K. Yasukawa)

18

19 *Abbreviations:* K4pol, family A DNA polymerase from *Thermotoga petrophila* strain
20 K4; M1pol, family A DNA polymerase from *Thermus thermophilus* strain M1; MMLV,
21 Moloney murine leukemia virus; RT, reverse transcriptase; SDS-PAGE, sodium dodecyl
22 sulfate-polyacrylamide gel electrophoresis.

23

24

1 Some bacterial thermostable, wild-type or genetically engineered family A DNA
2 polymerases have reverse transcriptase activity. However, difference in reverse
3 transcriptase activities of family A DNA polymerases and retroviral reverse
4 transcriptases (RTs) is unclear. In this study, comparative kinetic analysis was
5 performed for the reverse transcriptase activities of the wild-type enzyme of family A
6 DNA polymerase (M1pol_{WT}) from *Thermus thermophilus* M1 and the variant enzyme of
7 family A DNA polymerase (K4pol_{L329A}), in which the mutation of Leu329→Ala is
8 undertaken, from *Thermotoga petrophila* K4. In the incorporation of dTTP into
9 poly(rA)-p(dT)₄₅, the reaction rates of K4pol_{L329A} and M1pol_{WT} exhibited a saturated
10 profile of the Michaelis-Menten kinetics for dTTP concentrations but a substrate
11 inhibition profile for poly(rA)-p(dT)₄₅ concentrations. In contrast, the reaction rates of
12 Moloney murine leukemia virus (MMLV) RT exhibited saturated profiles for both dTTP
13 and poly(rA)-p(dT)₄₅ concentrations. This suggests that high concentrations of
14 DNA-primed RNA template decrease the efficiency of cDNA synthesis with bacterial
15 family A DNA polymerases.

16

17 *Keywords:* family A DNA polymerase; Moloney murine leukemia virus; reverse
18 transcriptase; template-primer; *Thermotoga petrophila*; *Thermus thermophilus*.

1 **1. Introduction**

2
3 In cDNA synthesis, an elevated reaction temperature is desirable because it reduces
4 RNA secondary structure and nonspecific binding of the primer. Reverse transcriptases
5 (RTs) from Moloney murine leukemia virus (MMLV) and Avian myeloblastosis virus
6 (AMV) have been used in cDNA synthesis. MMLV RT and AMV RT are not
7 thermostable. To realize the cDNA synthesis reaction at high temperature, their thermal
8 stabilities were improved by eliminating the RNase H activity [1], random mutation [2],
9 and site-directed mutagenesis [3,4]. cDNA synthesis with thermostable RTs are carried
10 out at around 50°C.

11 Some bacterial or archaeal, wild-type or genetically engineered DNA polymerases
12 have RNA-dependent DNA polymerase (reverse transcriptase) activity [5–8]. One of the
13 advantages of such DNA polymerases over retroviral RTs are that the DNA polymerases
14 are stable even at around 90°C, enabling direct RT-PCR in a single tube [6–8]. Figure
15 1A shows domain structure of DNA polymerases. The proofreading 3'-5' exonuclease
16 domain confers high fidelity. Bacterial family A DNA polymerases, such as from
17 *Thermus thermophilus* and *Thermus aquaticus*, do not have the 3'-5' exonuclease
18 domain and exhibit reverse transcriptase activity. Bacterial family A DNA polymerases,
19 such as from *Thermotoga petrophila* and *Thermotoga maritime* have the 3'-5'
20 exonuclease domain and do not exhibit reverse transcriptase activity. Archaeal family B
21 DNA polymerases, such as *Thermococcus kodakarensis* and *Pyrococcus furiosus*, have
22 the 3'-5' exonuclease domain and exhibit reverse transcriptase activity. However, family
23 B DNA polymerases are not suitable in cDNA synthesis because the reaction stops
24 when a uracil-containing template is incorporated [9].

1 Figure 1B shows the modeled structure of Family A DNA polymerase (K4pol)
2 from *Thermotoga petrophila* K4. Nine residues (Tyr326, Leu329, Gln384, Lys387,
3 Phe388, Met408, Asn422, Tyr438, and Phe451) are predicted to be involved in
4 DNA/RNA distinction. Figure 1C and D show that the steric hindrance with the
5 2'-hydroxyl group of ribose in K4pol might be removed by the mutation of
6 Leu329→Ala. Previous site-directed mutagenesis study revealed that six variants
7 (Y326A, L329A, Q384A, F388A, M408A, and Y438A) exhibited the reverse
8 transcriptase activity [10]. However, the threshold PCR cycle numbers needed to reach
9 the constant fluorescent intensity in the direct RT-PCR with these variants were higher
10 by 5–10 cycles than that in the conventional RT-PCR using MMLV RT in cDNA
11 synthesis and family B DNA polymerase from hyperthermophilic archaeon
12 *Thermococcus kodakaraensis* in the subsequent PCR [10]. This indicates that cDNA
13 synthesis with these variants is less efficient than that with MMLV RT. In this study, to
14 address this issue, we made kinetic analysis of the reverse transcriptase activities of the
15 wild-type enzyme of family A DNA polymerase (M1pol_{WT}) from *T. thermophilus* M1
16 and the variant enzyme of K4pol (K4pol_{L329A}) in which the mutation of Leu329→Ala is
17 undertaken. The results have shown that high concentrations of DNA-primed RNA
18 template decrease their reverse transcriptase activities.

19

20 **2. Materials and methods**

21

22 *2.1. Expression and purification of recombinant form of K4pol_{L329A}*

23

24 K4pol_{L329A} was prepared as described previously [10]. Briefly, *E. coli* strain

1 BL21(DE3) codon plus [*F⁻ ompT hsdS(r_B⁻ m_B⁻) dcm⁺ Tet^r galλ (DE3) endA Hte [argU*
2 *proL Cam^r]*] (Agilent Technologies) was transformed with the pET-21a plasmid
3 harboring the K4pol_{L329A} gene. The K4pol_{L329A} was expressed in the soluble fraction of
4 the transformants. After the heat treatment of the soluble fraction at 85°C for 30 min,
5 the enzyme was purified. The enzyme concentration was determined by the method of
6 Bradford [11] using Protein Assay CBB Solution (Nacalai Tesque, Kyoto, Japan) with
7 bovine serum albumin (Nacalai Tesque) as standard.

8

9 2.2. Expression and purification of recombinant form of MMLV RT

10

11 Recombinant MMLV RT was prepared as described previously [3]. Briefly, *E. coli*
12 strain BL21(DE3) was transformed with the pET-22b plasmid harboring the gene
13 encoding the C-terminally His₆-tagged MMLV RT. The MMLV RT was expressed in the
14 soluble fraction of the transformants, from which active enzyme was purified.

15

16 2.3. Expression and purification of recombinant form of M1pol_{WT}

17

18 *T. thermophilus* strain M1 was isolated from a hot spring (73°C) at Kagoshima
19 (Japan) and identified as *T. thermophilus* based on 16S rDNA sequence. Amino acid
20 sequence of DNA polymerase and nucleotide sequence of 16S rDNA from M1 show the
21 highest identity to those of *T. thermophilus* HB8 (99% and 98%, respectively). DNA
22 polymerase gene (*M1pol*) of M1 was amplified by PCR using oligonucleotide primers
23 (5'-AAAAAAACATATGGAGGCGATGCTTCCGCT-3' and
24 5'-AAGAATTCTAACCCCTTGGCGGAAAGCCAGT-3') and M1 chromosomal DNA

1 as a template. PCR was performed using 1 unit of KOD Plus polymerase (Toyobo,
2 Osaka, Japan) under following condition (3 min at 94°C, followed by 30 cycle of 15 s at
3 94°C, 30 s at 50°C, 3 min at 68°C). The amplified DNA was digested with restriction
4 enzymes *NdeI* and *EcoRI*, and the obtained DNA was inserted into the pET-28a digested
5 with the same enzymes. The resultant plasmid was designed as pET-M1pol. BL21(DE3)
6 codon plus was transformed with pET-M1pol. The transformants were aerobically
7 grown at 30°C in 300 ml of LB medium (1 % tryptone, 0.5 % yeast extract, 1 % NaCl;
8 adjusted to pH 7.3 with NaOH) containing 20 µg/ml kanamycin and 30 µg/ml
9 chloramphenicol to OD_{660} of 0.4. IPTG was then added to the final concentration of 1
10 mM. The cells were further incubated at 30°C for 4 h and were harvested by
11 centrifugation (8,000 × g, 10 min). The cells were resuspended in 7 ml of buffer A (20
12 mM Tris-HCl, 500 mM NaCl, 5 mM imidazole, 0.1% Triton X-100, pH 7.9) and
13 disrupted by sonication. The lysate was then separated by centrifugation (8,000 × g, 10
14 min). The obtained supernatant was heated at 75°C for 30 min and centrifuged again
15 (8,000 × g, 10 min). The supernatant was applied to a Ni²⁺-TED Sepharose column
16 equilibrated with buffer A. Next, the column was washed several times with the same
17 buffer. The N-terminally His₆-tagged *T. thermophilus* M1 DNA polymerase (M1pol_{WT})
18 was eluted with buffer B (20 mM Tris-HCl, 500 mM NaCl, 100 mM imidazole, 0.1%
19 Triton X-100, pH 7.9). The active fractions were pooled and dialyzed against a buffer
20 containing 20 mM Tris-HCl, 500 mM NaCl, pH 7.9.

21

22 2.4. Nucleotide sequence accession number

23

24 The nucleotide sequences of DNA polymerase and 16S rDNA from *T.*

1 *thermophilus* M1 reported in this study have been submitted to the DDBJ nucleotide
2 sequence database under the accession numbers AB744210 and AB744654,
3 respectively.

4

5 2.5. DNA-dependent DNA polymerase assay

6

7 DNA-dependent DNA polymerase activity to incorporate dNTP into gapped
8 M13mp2 DNA was examined as described previously [12,13]. Briefly, the 6,789-bp
9 fragment was obtained by the digestion of the 7,196-bp double-strand circular M13mp2
10 DNA with *PvuII* and the precipitation by 6% w/v polyethylene glycol-8000, 550 mM
11 NaCl. The gapped DNA was prepared by the annealing of the 7,196-base single-strand
12 circular M13mp2 DNA (140 μ g in 10 mM Tris-HCl, pH 8.0) with the heat-denatured
13 6,789-bp fragment (150 μ g in 10 mM Tris-HCl, pH 8.0). The reaction (15 μ L) was
14 carried out with 670 nM K4pol_{L329A} and 57 μ g of gapped DNA in 50 mM bicine-KOH,
15 1 mM Mn(CH₃COO)₂, 115 mM CH₃COOK, 8% glycerol at pH 8.2 (buffer B) at 70°C.
16 The reaction products (13 μ L) were applied to 0.8% agarose gel, followed by staining
17 with ethidium bromide (1 μ g/ml).

18

19 2.6. Reverse transcriptase assay

20

21 Reverse transcriptase activity to incorporate dTTP into poly(rA)-p(dT)₄₅ was
22 examined, based on the previous method using poly(rA)-p(dT)₁₅ [3,4]. Briefly, 2.0 mM
23 poly(rA)-p(dT)₄₅ (T/P) (the concentration is expressed as that of p(dT)₄₅) was prepared

1 by mixing 300 μ L of 2.5 mM p(dT)₄₅ (Sigma-Aldrich Japan, Ishikari, Japan) and 75 μ L
2 of 100 mM poly(rA) (the concentration is expressed as that of rA) (GE Healthcare,
3 Buckinghamshire, UK) followed by the incubation at 70°C for 15 min and at room
4 temperature for 30 min. [³H]dTTP (1.85 Bq/pmol) was prepared by mixing 860 μ L of
5 water, 100 μ L of [³H]dTTP (1.52 TBq/mmol) (GE Healthcare), and 40 μ L of 100 mM
6 dTTP (GE Healthcare). The reaction was carried out with various concentrations of
7 dTTP and T/P in buffer B at 50°C for K4pol_{L329A} and M1pol_{WT}, or in 25 mM Tris-HCl,
8 50 mM KCl, 2 mM dithiothreitol, 5 mM MgCl₂, pH 8.3 at 37°C for MMLV RT.

9

10 **3. Results**

11

12 *3.1. DNA- and RNA-dependent DNA polymerase activities of K4pol_{L329A}*

13

14 Recombinant K4pol_{L329A}, MMLV RT, and M1pol_{WT} were expressed in *E. coli* and
15 purified from the soluble fractions of the cells. Figure 2A shows the result of
16 SDS-PAGE analysis of purified enzyme preparations. The preparations of K4pol_{L329A},
17 MMLV RT, and M1pol_{WT} yielded a single band with a molecular mass of 101, 75, and
18 90 kDa, respectively. Figure 2B shows the result of DNA-dependent DNA polymerase
19 assay. In this assay, the 407-base gap of the 7,196-bp double-strand circular M13mp2
20 DNA is filled by the reaction with DNA polymerase. The reaction product with
21 K4pol_{L329A} exhibited the band corresponding to the gap-filled DNA, but did not exhibit
22 the one corresponding to the gapped DNA. Figure 2C shows the result of reverse
23 transcriptase assay. The amounts of [³H]dTTP incorporated increased with increasing
24 time in the reaction with K4pol_{L329A}, M1pol_{WT}, and MMLV RT. But they did not

1 increase with increasing time in the reaction with the wild-type K4pol (K4pol_{WT}). This
2 indicates that the mutation of Leu329→Ala does not abolish the DNA-dependent DNA
3 polymerase activity and creates reverse transcriptase activity in K4pol.

4 5 *3.2. Inhibitory effect of T/P toward the reverse transcriptase activities of K4pol_{L329A} and* 6 *M1pol_{WT}*

7
8 Figure 3A shows the reaction rates of K4pol_{L329A}, M1pol_{WT}, and MMLV RT for
9 reverse transcriptase reaction at various dTTP concentrations. A saturated profile of the
10 Michaelis-Menten kinetics were obtained, and the K_m and k_{cat} values were determined to
11 be $280 \pm 80 \mu\text{M}$ and $1.7 \pm 0.3 \text{ s}^{-1}$, respectively, for K4pol_{L329A}, $230 \pm 70 \mu\text{M}$ and $1.0 \pm$
12 0.2 s^{-1} , respectively, for M1pol_{WT}, and $250 \pm 60 \mu\text{M}$ and $35 \pm 6 \text{ s}^{-1}$, respectively, for
13 MMLV RT. This indicates that the k_{cat} values of K4pol_{L329A} and M1pol_{WT} at 50°C were
14 5 and 3 %, respectively, of that of MMLV RT at 37°C while their K_m values were
15 similar.

16 Figure 3B shows the reaction rates of K4pol_{L329A}, M1pol_{WT}, and MMLV RT for
17 reverse transcriptase reaction at various T/P concentrations. A substrate inhibition
18 profile was obtained for K4pol_{L329A} and M1pol_{WT}: the reaction rates reached the
19 maximum at 10 μM T/P, decreased with increasing T/P concentration (10–100 μM), and
20 reached 50% of the maximum at 100 μM T/P. On the other hand, a saturated profile of
21 the Michaelis-Menten kinetics was obtained for MMLV RT: the K_m and k_{cat} values were
22 determined to be $5.6 \pm 0.8 \mu\text{M}$ and $29 \pm 6 \text{ s}^{-1}$, respectively.

23 24 *3.3. Temperature dependence of the reverse transcriptase activity of K4pol_{L329A}*

1

2 Considering that RNA is stable up to around 65°C, the optimal reaction
3 temperature for cDNA synthesis reaction with thermostable DNA polymerases is
4 thought to be around 65°C. However, poly(rA)-p(dT)₄₅ dissociates at 65°C. To estimate
5 the reverse transcriptase activity of K4pol_{L329A} at 65°C, the reaction rates of K4pol_{L329A}
6 to incorporate dTTP into T/P were measured at various temperatures (Fig. 4). They
7 increased with increasing temperature from 37 to 52°C. Inset shows Arrhenius plot on
8 the assumption that the observed reaction rates correspond to V_{\max} . Linear relationship
9 was held between the natural logarithm of V_{\max} and $1/T$. In this plot, $\ln(V_{\max})$ value of
10 6.4 (the V_{\max} value of 600 nM s⁻¹) was estimated at 65°C, and the k_{cat} value at 65°C was
11 calculated to be 12 s⁻¹. Considering that the k_{cat} values of MMLV RT obtained in this
12 study were 35 and 29 s⁻¹, the reverse transcriptase activity of K4pol_{L329A} at around 65°C
13 might be comparable to that of MMLV RT at 37°C.

14

15 **4. Discussion**

16

17 The mechanism of the template-primer-mediated inhibition of reverse transcriptase
18 activity is unclear. Because K4pol_{L329A} has the 3'-5' exonuclease domain but M1pol_{WT}
19 does not (Fig. 1), it is thought that the 3'-5' exonuclease domain is not critical for the
20 inhibition. Bacterial family A DNA polymerases and retroviral RTs share a common
21 structure: both groups comprise the fingers, palm, and thumb domains. Crystal
22 structures of *E. coli* DNA polymerase [14], *T. aquaticus* DNA polymerase [15], human
23 immunodeficiency virus type 1 (HIV-1) RT [16], and MMLV RT [17] revealed that the
24 palm domains are similar but the fingers and thumb domains are different between the

1 two groups [18]: particular amino acid residues in α -helix of the fingers domain are
2 involved in nucleotide binding in family A DNA polymerases while those in β -sheet of
3 the fingers domain are involved in RTs; and the sites in the thumb domain which
4 interact with the minor groove of the primer-template are different in the two groups.
5 There might be a possibility that the binding of template-primer with particular sites in
6 the fingers or thumb domain of family A DNA polymerase is involved in the inhibition.

7 The template-primer-mediated inhibition of reverse transcriptase activity suggests
8 that high concentrations of DNA-primed RNA template decrease the efficiency of
9 cDNA synthesis with bacterial family A DNA polymerases. We speculate that in
10 RT-PCR, this inhibition affects the quantification, but does not much affect the detection
11 of a target RNA. In clinical diagnosis, various nucleic acid tests are used for the
12 detection of a target RNA from pathogens. Sauter and Marx generated *T. aquaticus*
13 DNA polymerase variant, L322M/L459M/S515R/I638F/S739G/E773G/L789F, with
14 increased reverse transcriptase activity by random mutation [19]. They showed that
15 0.1–5 ng of a target RNA can be quantified by the direct real-time RT-PCR with this
16 variant enzyme in a single tube format [19]. This does not contradict our result because
17 the concentration of T/P is as high as 100 μ M (about 3 μ g/ μ L) in this study. To achieve
18 the efficient synthesis of cDNA using bacterial family A DNA polymerases with reverse
19 transcriptase activity, optimization is required for the concentration of T/P.

20 How are the template-primer-mediated inhibition of reverse transcriptase activities
21 of K4pol and M1pol removed? If particular region involved in the T/P-mediated
22 inhibition is identified, site-directed mutagenesis might be effective. Mayanagi et al.
23 reported the structure of the ternary complex of family B DNA polymerase from
24 hyperthermophilic archaeon *Pyrococcus furiosus*, proliferating cell nuclear antigen

1 (PCNA), and DNA [20]. Shimada et al. reported that cold inducible RNA helicase from
2 hyperthermophilic archaeon *Thermococcus kodakaraensis* is critical for its adaptation to
3 cold temperature [21]. There is a possibility that molecules that interact with nucleic
4 acids, such as sliding clump, a ring-shaped protein that slides on DNA, and RNA
5 helicase, an enzyme that unwinds single-strand paired RNA, might be effective to
6 remove the inhibition. Removal of the inhibition will open up the wide application of
7 bacterial family A DNA polymerase in quantitative RT-PCR in a single tube format.

8

9 **Acknowledgments**

10

11 This study was supported in part (K.Y.) by Grants-in-Aid for Scientific Research
12 (nos. 19580104 and 21580110) from the Japan Society for the Promotion of Science
13 and Daiwa Securities Health Foundation.

14

15 **References**

16

- 17 [1] G.F. Gerard, R.J. Potter, M.D. Smith, K. Rosenthal, G. Dhariwal, J. Lee, D.K.
18 Chatterjee, The role of template-primer in protection of reverse transcriptase from
19 thermal inactivation, *Nucleic Acids Res.*, 30 (2002) 3118–3129.
- 20 [2] B. Arezi, H. Hogrefe, Novel mutations in Moloney murine leukemia virus reverse
21 transcriptase increase thermostability through tighter binding to template-primer,
22 *Nucleic Acids Res.*, 37 (2009) 473–481.
- 23 [3] K. Yasukawa, M. Mizuno, A. Konishi, K. Inouye, Increase in thermal stability of
24 Moloney murine leukaemia virus reverse transcriptase by site-directed

- 1 mutagenesis, *J. Biotechnol.*, 150 (2010) 299–306.
- 2 [4] A. Konishi, K. Yasukawa, K. Inouye, Improving the thermal stability of avian
3 myeloblastosis virus reverse transcriptase α -subunit by site-directed mutagenesis,
4 *Biotechnol. Lett.*, 34 (2012) 1209–1215.
- 5 [5] T.W. Myers, D.H. Gelfand, Reverse transcription and DNA amplification by a
6 *Thermus thermophilus* DNA polymerase, *Biochemistry*, 30 (1991) 7661–7666.
- 7 [6] H. Shandilya, K. Griffiths, E.K. Flynn, M. Astatke, P.J. Shih, J.E. Lee, G.F.
8 Gerard, M.D. Gibbs, P.L. Bergquist, Thermophilic bacterial DNA polymerases
9 with reverse-transcriptase activity, *Extremophiles*, 8 (2004) 243–251.
- 10 [7] R. Kranaster, M. Drum, N. Engel, M. Weidmann, F.T. Hufert, A. Marx,
11 *Biotechnol. J.*, 5 (2010) 224–231.
- 12 [8] N.J. Schönbrunner, E.H. Fiss, O. Budker, S. Stoffel, C.L. Sigua, D.H. Gelfand,
13 T.W. Myers, Chimeric thermostable DNA polymerases with reverse transcriptase
14 and attenuated 3'–5' exonuclease activity, *Biochemistry*, 45 (2006) 12786–12795.
- 15 [9] S.J. Firbank, J. Wardle, P. Heslop, R.K. Lewis, B.A. Connolly, Uracil recognition
16 in archaeal DNA polymerases captured by X-ray crystallography, *J. Mol. Biol.*,
17 381 (2008) 529–538.
- 18 [10] S. Sano, Y. Yamada, T. Shinkawa, S. Kato, T. Okada, H. Higashibata, S. Fujiwara,
19 Mutations to create thermostable reverse transcriptase with bacterial family A
20 DNA polymerase from *Thermotoga petrophila*, *J. Biosci. Bioeng.*, 113 (2012)
21 315–321.
- 22 [11] M.M. Bradford, A rapid and sensitive method for the quantitation of microgram
23 quantities of protein utilizing the principle of protein-dye binding, *Anal.*
24 *Biochem.*, 72, 248–254 (1976).

- 1 [12] K. Bebenek, T.A. Kunkel, Analyzing fidelity of DNA polymerases, *Methods*
2 *Enzymol.*, 262 (1995) 217–232.
- 3 [13] V. Barrioluengo, M. Álvarez, D. Barbieri, L. Menéndez-Arias, Thermostable
4 HIV-1 group O reverse transcriptase variants with the same fidelity as murine
5 leukaemia virus reverse transcriptase, *Biochem. J.*, 436 (2011) 599–607.
- 6 [14] L.S. Beese, V. Derbyshire, T.A. Steitz, Structure of DNA polymerase I Klenow
7 fragment bound to duplex DNA, *Science*, 260 (1993) 352–355.
- 8 [15] S.H. Eom, J. Wang, T.A. Steitz, Structure of *Taq* polymerase with DNA at the
9 polymerase active site, *Nature*, 382 (1996) 278–281.
- 10 [16] L.A. Kohlstaedt, J. Wang, J.M. Friedman, P.A. Rice, T.A. Steitz, Crystal structure
11 at 3.5 Å resolution of HIV-1 reverse transcriptase complexed with an inhibitor,
12 *Science*, 256 (1992) 1783–1790.
- 13 [17] D. Das, M.M. Georgiadis, The crystal structure of the monomeric reverse
14 transcriptase from Moloney murine leukemia virus, *Structure*, 12 (2004)
15 819–829.
- 16 [18] T.A. Steitz, DNA polymerases: Structural diversity and common mechanism, *J.*
17 *Biol. Chem.*, 274 (1999) 17395–17398.
- 18 [19] K.B. Sauter, A. Marx, Evolving thermostable reverse transcriptase activity in a
19 DNA polymerase scaffold, *Angew. Chem. Int. Ed.*, 45 (2006) 7633–7635.
- 20 [20] K. Mayanagi, S. Kiyonari, H. Nishida, M. Saito, D. Kohda, Y. Ishino, T. Shirai,
21 K. Morikawa, Architecture of the DNA polymerase B-proliferating cell nuclear
22 antigen (PCNA)-DNA ternary complex, *Proc. Natl. Acad. Sci. USA*, 108 (2011)
23 1845–1849.
- 24 [21] Y. Shimada, W. Fukuda, Y. Akada, M. Ishida, J. Nakayama, T. Imanaka, S.

1 Fujiwara, Property of cold inducible DEAD-box RNA helicase in
2 hyperthermophilic archaea, *Biochem. Biophys. Res. Commun.*, 389 (2009)
3 622–627.
4

1 **Figure legends**

2

3 **Fig. 1.** Creation of reverse transcriptase activity in K4pol by the mutation of
4 Leu329→Ala. (A) Domain structure of bacterial DNA polymerases. (B-D) Modeled
5 structure of K4pol-*template* RNA complex. The structure is constructed by Swiss model
6 using *E. coli* Klenow fragment (1kfs) as a template. Carbon, nitrogen, oxygen, and
7 phosphorus atoms of RNA are shown in green, blue, red, and orange, respectively. (B)
8 Whole structure of K4pol_{WT}-*template* RNA complex. The amino acid residues predicted
9 to be located around an RNA *template* (Tyr326, Leu329, Gln384, Lys387, Phe388,
10 Met408, Asn422, Tyr438, and Phe451) are shown in green. (C) Close-up view of
11 Leu329 of K4pol_{WT}-*template* RNA complex. (D) Close-up view of Ala329 of
12 K4pol_{L329A}-*template* RNA complex. Electron densities of the side chain of Leu329 (C)
13 and Ala329 (D) are shown in red.

14

15 **Fig. 2.** DNA- and RNA-dependent DNA polymerase activities of K4pol_{L329A}. (A)
16 SDS-PAGE under reducing condition. Lane M, molecular-mass marker. (B) Agarose gel
17 electrophoresis of reaction products in the DNA-dependent DNA polymerase assay.
18 Lane M, molecular-mass marker. The DNA bands a, b, c, and d correspond to the
19 gap-filled DNA, the gapped DNA (the annealing product of the 6,789-base single-strand
20 DNA fragment and the 7,196-base single-strand circular DNA), the 6,789-bp
21 double-strand DNA fragment, and the 7,196-base single-strand circular DNA,
22 respectively. The reaction was carried out with 670 nM K4pol_{L329A} at 50°C for 5 min.
23 (C) Incorporation of radioactivity of reaction products in the reverse transcriptase assay.
24 The reaction was carried out with K4pol_{WT} (50 nM at 50°C), K4pol_{L329A} (50 nM at

1 50°C), M1pol_{WT} (50 nM at 50°C) or MMLV RT (2 nM at 37°C). The initial
2 concentrations of dTTP and T/P were 200 and 10 μM, respectively.

3

4 **Fig. 3.** Substrate dependence of the reaction rates (v_o) in the reverse transcriptase
5 reaction. The reaction was carried out with K4pol_{L329A} (50 nM at 50°C), M1pol_{WT} (50
6 nM at 50°C) or MMLV RT (2 nM at 37°C). Solid line represents the best fit of the
7 Michaelis-Menten equation. (A) Dependence of v_o on the dTTP concentration. The
8 initial T/P concentration was 10 μM. (B) Dependence of v_o on the T/P concentration.
9 The initial concentration of dTTP was 200 μM.

10

11 **Fig. 4.** Temperature dependence of the reaction rate (v_o) of K4pol_{L329A} in the reverse
12 transcriptase reaction. The reaction was carried out with 50 nM K4pol_{L329A}. The initial
13 concentrations of dTTP and T/P were 200 and 25 μM, respectively. Inset shows
14 Arrhenius plot.

15







A**B****C****D**

Fig. 1, Yasukawa *et al.*


A




B




C

Fig. 2, Yasukawa *et al.*

A



B

Fig. 3, Yasukawa *et al.*




Fig. 4, Yasukawa *et al.*