

Title	Symplectic capacity and short periodic billiard trajectory
Author(s)	Irie, Kei
Citation	Mathematische Zeitschrift (2012), 272(3-4): 1291-1320
Issue Date	2012-12
URL	http://hdl.handle.net/2433/162900
Right	The final publication is available at www.springerlink.com
Type	Journal Article
Textversion	author

SYMPLECTIC CAPACITY AND SHORT PERIODIC BILLIARD TRAJECTORY

KEI IRIE

ABSTRACT. We prove that a bounded domain Ω in \mathbb{R}^n with smooth boundary has a periodic billiard trajectory with at most $n + 1$ bounce times and of length less than $C_n r(\Omega)$, where C_n is a positive constant which depends only on n , and $r(\Omega)$ is the supremum of radius of balls in Ω . This result improves the result by C.Viterbo, which asserts that Ω has a periodic billiard trajectory of length less than $C'_n \text{vol}(\Omega)^{1/n}$. To prove this result, we study symplectic capacity of Liouville domains, which is defined via symplectic homology.

1. INTRODUCTION

Let Ω be a bounded domain in \mathbb{R}^n with smooth boundary. A *periodic billiard trajectory* on Ω is a continuous map $\gamma : \mathbb{R}/\tau\mathbb{Z} \rightarrow \bar{\Omega}$ such that there exists a finite set $\mathcal{B} \subset \mathbb{R}/\tau\mathbb{Z}$ and satisfies the following conditions:

- γ is smooth on $(\mathbb{R}/\tau\mathbb{Z}) \setminus \mathcal{B}$ and satisfies $\ddot{\gamma} = 0$.
- For each $t_0 \in \mathcal{B}$, $\gamma(t_0) \in \partial\Omega$, the left and right derivatives $\dot{\gamma}(t_0^\pm) := \lim_{t \rightarrow t_0^\pm} \dot{\gamma}(t)$ exist,

and satisfy the law of reflection (ν denotes the outward normal vector on $\partial\Omega$):

$$\begin{aligned} \langle \dot{\gamma}(t_0^+), \nu(\gamma(t_0)) \rangle &= -\langle \dot{\gamma}(t_0^-), \nu(\gamma(t_0)) \rangle \neq 0, \\ \dot{\gamma}(t_0^+) - \langle \dot{\gamma}(t_0^+), \nu(\gamma(t_0)) \rangle \cdot \nu(\gamma(t_0)) &= \dot{\gamma}(t_0^-) - \langle \dot{\gamma}(t_0^-), \nu(\gamma(t_0)) \rangle \cdot \nu(\gamma(t_0)). \end{aligned}$$

Elements of \mathcal{B} are called *bounce times*.

Before stating the main theorem, we introduce some notations. For $x \in \mathbb{R}^n$ and $r \geq 0$, $B(x, r) := \{y \in \mathbb{R}^n \mid |x - y| \leq r\}$. For $\Omega \subset \mathbb{R}^n$,

$$r(\Omega) := \sup\{r \geq 0 \mid \text{there exists } x \in \mathbb{R}^n \text{ such that } B(x, r) \subset \Omega\}.$$

The main theorem of this paper is the following:

Theorem 1.1. *Let Ω be a bounded domain in \mathbb{R}^n with smooth boundary. Then, there exists a periodic billiard trajectory γ on Ω with at most $n + 1$ bounce times and which satisfies the following length estimate:*

$$|\gamma| \leq C_n r(\Omega),$$

where C_n is a positive constant which depends only on n .

Date: February 6, 2012.

2010 Mathematics Subject Classification. Primary:34C25, Secondary:53D40.

Remark 1.2. The existence of a periodic billiard trajectory with at most $n + 1$ bounce times is due to [2]. They also consider arbitrary metrics on \mathbb{R}^n .

In [8], it is proved (theorem 4.1) that there exists a periodic billiard trajectory γ on Ω (with the flat metric) which satisfies a length estimate $|\gamma| \leq C'_n \text{vol}(\Omega)^{1/n}$, where C'_n is a positive constant which depends only on n . Notice that this result follows from theorem 1.1 and an obvious inequality $r(\Omega) \leq \omega_n^{-1/n} \text{vol}(\Omega)^{1/n}$, where ω_n denotes the volume of the n -dimensional unit ball.

In [1], it is proved (theorem 1.2, the case of a constant potential) that there exists a periodic billiard trajectory γ on Ω (with the flat metric) with at most $n + 1$ bounce times and which satisfies a length estimate $|\gamma| \leq C'' \text{diam}\Omega$, where C'' is a constant which does not depend on n , and

$$\text{diam}\Omega := \inf\{|v| \mid (v + \Omega) \cap \Omega = \emptyset\}.$$

Notice that our main theorem also implies this result for each fixed n , though we can not prove the independence of C'' on n by this argument.

Finally we remark that one can easily construct $(\Omega_k)_k$, a sequence of bounded domains in \mathbb{R}^n , such that $\lim_{k \rightarrow \infty} \text{vol}(\Omega_k) = \lim_{k \rightarrow \infty} \text{diam}(\Omega_k) = \infty$ and $r(\Omega_k) \leq 1$ for any k .

To prove theorem 1.1, we use *symplectic capacity* defined via symplectic homology, which was introduced in [7]. In the present paper, symplectic capacity is defined for Liouville domains (compact exact symplectic manifolds with convex boundaries), and it is denoted by cap_S . The definition is given at the beginning of section 3.

Using symplectic capacity cap_S , we introduce the notion of capacity for Riemannian manifolds (without boundaries), which is denoted by cap_R . Roughly speaking, it is defined by $\text{cap}_R(N) := \text{cap}_S(DT^*N)$, where $DT^*N := \{(q, p) \in T^*N \mid |p| \leq 1\}$. But when N is non-compact, the right hand side does not make sense since DT^*N is not a Liouville domain (since it is not compact). Hence we have to approximate DT^*N by compact domains. See definition 4.3 for the precise definition.

We prove that cap_R satisfies following properties:

- (A) Let Ω be a non-empty open set in \mathbb{R}^n . Then, $\text{cap}_R(\Omega) \leq C_n r(\Omega)$, where Ω is equipped with the flat Riemannian metric on \mathbb{R}^n .
- (B) If Ω is a bounded domain in \mathbb{R}^n with smooth boundary, there exists a periodic billiard trajectory on Ω with at most $n + 1$ bounce times and of length equals to $\text{cap}_R(\Omega)$.

Our main theorem 1.1 follows at once from (A) and (B).

We explain the structure of this paper. In section 2, we recall the notion of symplectic homology. We use the version introduced in [7].

In section 3, we define cap_S , and prove its properties. The most important result in this section is theorem 3.6, which asserts that when $\pi : Y \rightarrow X$ is a covering map between Liouville domains, then $\text{cap}_S(Y) \leq \text{cap}_S(X)$. Though its proof is not very difficult, it seems to the author that this result contains a novel idea.

In section 4, we define cap_R , and prove its properties. The main result in this section is theorem 4.13, which includes the property (A). Theorem 3.6 is used to prove that $\mathbb{R}^n \setminus \mathbb{Z}^n$ with the flat metric has a finite capacity (theorem 4.12). Theorem 4.13 is proved by theorem 4.12 and elementary geometric arguments.

In section 5, we prove the property (B) (theorem 5.1). The arguments in this section heavily rely on the techniques developed in the recent paper [1].

In the appendix, we prove theorem 2.12, which asserts that truncated symplectic homology of a Liouville domain (X, λ) depends only on $d\lambda$. It seems to the author that theorem 2.12 is well-known to experts. But we give a proof of the result since the author is unable to find its proof in the literature.

2. SYMPLECTIC HOMOLOGY

2.1. Liouville domains. First, we recall the notion of *Liouville domains*. A Liouville domain is a pair (X, λ) where X is a compact manifold with boundary and λ is a 1-form on X , with the following conditions:

- (1) $(X, d\lambda)$ is a symplectic manifold.
- (2) $Z \in \mathcal{X}(X)$ defined by $i_Z d\lambda = \lambda$ points strictly outwards on ∂X .

(2) implies that $(\partial X, \lambda)$ is a contact manifold. Let R be the Reeb vector field on $(\partial X, \lambda)$ (recall that R is characterized by $i_R d\lambda = 0, \lambda(R) = 1$).

In the rest of this paper, (X, λ) stands for a Liouville domain, and n stands for $\dim X/2$, unless otherwise stated.

$\mathcal{P}(\partial X, \lambda)$ denotes the set of periodic Reeb orbits on $(\partial X, \lambda)$, and $\mathcal{P}_0(\partial X, \lambda)$ denotes the set of elements of $\mathcal{P}(\partial X, \lambda)$ which is contractible in X . For each $x \in \mathcal{P}(\partial X, \lambda)$, its period is denoted by $\tau(x)$, and

$$\tau(\partial X, \lambda) := \{\tau(x) \mid x \in \mathcal{P}(\partial X, \lambda)\}.$$

It is well-known that $\tau(\partial X, \lambda)$ is a closed null set in $[0, \infty)$. Define $\delta(\partial X, \lambda) := \min \tau(\partial X, \lambda)$. It is clear that $\delta(\partial X, \lambda) > 0$.

There exists an unique (up to homotopy) almost complex structure on TX , which is compatible with $d\lambda$. In the rest of this paper, we only treat the case $c_1(TX) = 0$.

Let (X, λ) be a Liouville domain. We define $\Phi: \partial X \times (0, 1] \rightarrow X$ by

$$\Phi(z, 1) = z, \quad \partial_r \Phi(z, r) = r^{-1} Z(\Phi(z, r)).$$

It is easy to check that $\Phi^* \lambda = r\pi^* \lambda$, where $\pi: \partial X \times (0, 1] \rightarrow \partial X$ is the projection. Define \hat{X} and $\hat{\lambda} \in \Omega^1(\hat{X})$ by

$$\hat{X} := X \cup_{\Phi} \partial X \times (0, \infty), \quad \hat{\lambda} := \begin{cases} \lambda & (\text{on } X) \\ r\pi^* \lambda & (\text{on } \partial X \times (0, \infty)) \end{cases}.$$

We call $(\hat{X}, \hat{\lambda})$ the *completion* of (X, λ) .

By definition, there exists a natural embedding $I : \partial X \times (0, \infty) \rightarrow \hat{X}$. We often identify $\partial X \times (0, \infty)$ with its image via I . For $r \in (0, \infty)$, $X(r)$ denotes the bounded domain in \hat{X} with boundary $\partial X \times \{r\}$, i.e.

$$X(r) := \begin{cases} X \cup \partial X \times [1, r] & (r \geq 1) \\ X \setminus \partial X \times (r, 1] & (r < 1) \end{cases}.$$

Definition 2.1. Two Liouville domains (X, λ) , (X', λ') are called *equivalent* if and only if there exists a diffeomorphism $\varphi : X' \rightarrow X$ such that $\lambda' = \varphi^* \lambda$. (X, λ) , (X', λ') are called *isotopic* if there exists a smooth family of Liouville domains $(X, \lambda_t)_{0 \leq t \leq 1}$, such that $\lambda_0 = \lambda$ and (X, λ_1) is equivalent to (X', λ') .

2.2. Periodic orbits of Hamiltonian flows. For $H \in C^\infty(\hat{X})$, we define its *Hamiltonian vector field* X_H by $i_{X_H} d\hat{\lambda} = -dH$. For $H = (H_t)_{t \in \mathbb{R}/\tau\mathbb{Z}}$, a family of Hamiltonians on \hat{X} parametrized by $\mathbb{R}/\tau\mathbb{Z}$, let us denote by $\mathcal{P}_\tau(H)$ the set of $x : \mathbb{R}/\tau\mathbb{Z} \rightarrow \hat{X}$ which is contractible and satisfies $\partial_t x = X_{H_t}(x)$. $\mathcal{P}_1(H)$ is often abbreviated by $\mathcal{P}(H)$.

For $x \in \mathcal{P}_\tau(H)$, we define its *Conley-Zehnder index*. For later purposes, it is necessary to define the Conley-Zehnder index for degenerate periodic orbits. Hence we have to define the index for degenerate symplectic paths. We use the definition given in [5].

First we introduce some notations. Take a coordinate $(q_1, p_1, \dots, q_n, p_n)$ on \mathbb{R}^{2n} , and define

$$\begin{aligned} \lambda_n &:= \frac{1}{2} \sum_{1 \leq j \leq n} p_j \wedge dq_j - q_j \wedge dp_j, \\ \omega_n &:= d\lambda_n, \\ \text{Sp}(2n) &:= \{V \in \text{GL}(2n, \mathbb{R}) \mid V^* \omega_n = \omega_n\}. \end{aligned}$$

In the present paper, $\text{GL}(2n, \mathbb{R})$ acts on \mathbb{R}^{2n} from right, i.e. we denote the action of $\text{GL}(2n, \mathbb{R})$ on \mathbb{R}^{2n} by

$$(x_1, \dots, x_{2n}) \cdot (V_{ij})_{1 \leq i, j \leq 2n} := \left(\sum_{1 \leq j \leq 2n} x_j V_{1j}, \dots, \sum_{1 \leq j \leq 2n} x_j V_{2n, j} \right).$$

For $\tau > 0$, let us denote

$$\mathcal{P}_\tau(2n) := \{\gamma \in C^0([0, \tau], \text{Sp}(2n)) \mid \gamma(0) = 1_{2n}\}.$$

We define the index $i : \mathcal{P}_\tau(2n) \rightarrow \mathbb{Z}$ by several axioms. To spell out the axioms, we introduce more notations.

- For $\gamma_0, \gamma_1 \in \mathcal{P}_\tau(2n)$, $\gamma_0 \sim \gamma_1$ if and only if there exists $\delta : [0, 1] \times [0, \tau] \rightarrow \text{Sp}(2n)$ such that $\delta(i, \cdot) = \gamma_i(\cdot)$ ($i = 0, 1$), $\delta(s, 0) = 1_{2n}$, and $\text{rk}(\delta(s, 1) - 1_{2n})$ is constant on s .
- For $\gamma_i \in \mathcal{P}_\tau(2n_i)$ ($i = 0, 1$), define $\gamma_0 \diamond \gamma_1 \in \mathcal{P}_\tau(2n_0 + 2n_1)$ by $\gamma_0 \diamond \gamma_1(t) := \begin{pmatrix} \gamma_0(t) & 0 \\ 0 & \gamma_1(t) \end{pmatrix}$.

- For $\gamma_0, \gamma_1 \in C^0([0, \tau], \text{Sp}(2n))$ such that $\gamma_0(\tau) = \gamma_1(0)$, define $\gamma_1 * \gamma_0 \in C^0([0, \tau], \text{Sp}(2n))$ by $\gamma_1 * \gamma_0(t) = \begin{cases} \gamma_0(2t) & (t \leq \tau/2) \\ \gamma_1(2t - \tau) & (t \geq \tau/2) \end{cases}$.
- For $\tau > 0$ and $\theta \in \mathbb{R}$, define $\varphi_{\tau, \theta} \in \mathcal{P}_\tau(2)$ by $\varphi_{\tau, \theta}(t) := \begin{pmatrix} \cos(t\theta/\tau) & -\sin(t\theta/\tau) \\ \sin(t\theta/\tau) & \cos(t\theta/\tau) \end{pmatrix}$.

The index $i : \mathcal{P}_\tau(2n) \rightarrow \mathbb{Z}$ is defined by the following axioms:

Theorem 2.2 ([5], theorem 6.2.7). *For $\tau > 0$, there exists an unique map $i : \bigcup_{n \geq 1} \mathcal{P}_\tau(2n) \rightarrow \mathbb{Z}$ which satisfies the following five axioms:*

- (1) For $\gamma_0, \gamma_1 \in \mathcal{P}_\tau(2n)$, $\gamma_0 \sim \gamma_1 \implies i(\gamma_0) = i(\gamma_1)$.
- (2) For $\gamma_i \in \mathcal{P}_\tau(2n_i)$ ($i = 0, 1$), $i(\gamma_0 \diamond \gamma_1) = i(\gamma_0) + i(\gamma_1)$.
- (3) For any $\gamma \in \mathcal{P}_\tau(2)$ satisfying $\gamma(\tau) = \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}$ ($a = 0, \pm 1$), there exists $\theta_0 > 0$ such that $i([\gamma(\tau)\varphi_{\tau, -\theta}] * \gamma) = i(\gamma)$ for any $\theta \in (0, \theta_0]$.
- (4) For any $\gamma \in \mathcal{P}_\tau(2)$ satisfying $\gamma(\tau) = \begin{pmatrix} 1 & a \\ 0 & 1 \end{pmatrix}$ ($a = \pm 1$), there exists $\theta_0 > 0$ such that $i([\gamma(\tau)\varphi_{\tau, \theta}] * \gamma) = i(\gamma) + 1$ for any $\theta \in (0, \theta_0]$.
- (5) Define $\gamma_0 \in \mathcal{P}_\tau(2)$ by $\gamma_0(t) := \begin{pmatrix} 1 + t/\tau & 0 \\ 0 & (1 + t/\tau)^{-1} \end{pmatrix}$. Then, $i(\gamma_0) = 0$.

In [5], several equivalent definitions are given (definition 5.4.2, definition 6.1.10). In particular, definition 6.1.10 in [5] implies the following useful lemma:

Lemma 2.3. *Let us define the set of non-degenerate symplectic paths by*

$$\mathcal{P}_\tau^*(2n) := \{\gamma \in \mathcal{P}_\tau(2n) \mid \text{rk}(\gamma(\tau) - 1_{2n}) = 2n\}.$$

Then, for any $\gamma \in \mathcal{P}_\tau(2n)$, $i(\gamma) = \sup_{U \in N(\gamma)} \inf\{i(\beta) \mid \beta \in U \cap \mathcal{P}_\tau^(2n)\}$, where $N(\gamma)$ denotes the set of all open neighborhoods of γ in $\mathcal{P}_\tau(2n)$.*

The following lemma follows at once from the above lemma.

Lemma 2.4. *Assume that a sequence $(\gamma_k)_k$ in $\mathcal{P}_\tau(2n)$ converges to γ in $\mathcal{P}_\tau(2n)$. Then, $i(\gamma) \leq \liminf_{k \rightarrow \infty} i(\gamma_k)$.*

Next we define the Conley-Zehnder index $\mu_{\text{CZ}}(x)$ for $x \in \mathcal{P}_\tau(H)$. Let $D^2 := \{z \in \mathbb{C} \mid |z| \leq 1\}$, and take arbitrary $\bar{x} : D^2 \rightarrow \hat{X}$ such that $\bar{x}(e^{2\pi i \theta}) = x(\tau\theta)$ (such \bar{x} exists since x is contractible). Since D^2 is contractible, $\bar{x}^* T\hat{X}$ is a trivial symplectic vector bundle. Take the following trivialization of symplectic vector bundle:

$$F : (\mathbb{R}^{2n}, \omega_n) \times D^2 \rightarrow \bar{x}^* T\hat{X}; \quad (v, z) \mapsto (F_z(v), z).$$

Define $\gamma : \mathbb{R}/\tau\mathbb{Z} \rightarrow \text{Sp}(2n)$ by

$$\gamma(t) := (F_{e^{2\pi i t/\tau}})^{-1} \circ \Phi_t \circ F_1.$$

where $(\Phi_t)_t$ is the Poincaré map generated by $(X_{H_t})_t$. Finally, we define $\mu_{CZ}(x)$ by

$$\mu_{CZ}(x) := i(\gamma).$$

Since $c_1(TX) = 0$, the above definition is independent of choices of \bar{x} . An element $x \in \mathcal{P}_\tau(H)$ is called *non-degenerate* if and only if $\gamma \in \mathcal{P}_\tau^*(2n)$.

2.3. Floer homology on Liouville domains. For $r_0 \geq 1$, let $\mathcal{H}(X, \lambda : r_0)$ be the set of $H = (H_t)_{t \in \mathbb{R}/\mathbb{Z}}$, a family of Hamiltonians on \hat{X} parametrized by \mathbb{R}/\mathbb{Z} , with the following property:

There exist $a > 0$, $b \in \mathbb{R}$ such that $H_t(z, r) = ar + b$ for any $(z, r) \in \partial X \times [r_0, \infty)$ and $t \in \mathbb{R}/\mathbb{Z}$. (we denote a, b by a_H, b_H .)

We denote $\mathcal{H}(X, \lambda) := \bigcup_{r_0 \geq 1} \mathcal{H}(X, \lambda : r_0)$.

$H \in \mathcal{H}(X, \lambda)$ is called *admissible* if all elements of $\mathcal{P}(H)$ are non-degenerate, and $a_H \notin \tau(\partial X, \lambda)$. $\mathcal{H}_{\text{ad}}(X, \lambda)$ denotes the set of all admissible $H \in \mathcal{H}(X, \lambda)$. Note that when H is admissible, then $\sharp \mathcal{P}(H) < \infty$.

For $H \in \mathcal{H}_{\text{ad}}(X, \lambda)$, we define its Floer homology $\text{HF}_*(H)$. For each $k \in \mathbb{Z}$, let $\mathcal{P}_k(H)$ denote the set of $x \in \mathcal{P}(H)$ with $\mu_{CZ}(x) = k$, and let $C_k(H)$ denote the free \mathbb{Z}_2 -module over $\mathcal{P}_k(H)$.

To define the Floer homology, we need to equip \hat{X} with almost complex structures. For $r_0 \geq 1$, let $\mathcal{J}(X, \lambda : r_0)$ be the set of $J = (J_t)_{t \in \mathbb{R}/\mathbb{Z}}$, a family of almost complex structures on \hat{X} parametrized by \mathbb{R}/\mathbb{Z} , such that following properties hold for any $t \in \mathbb{R}/\mathbb{Z}$ (R and ξ denote the Reeb vector field and the contact distribution on $(\partial X, \lambda)$):

- J_t is compatible with $d\hat{\lambda}$.
- $J_t(\partial_r(z, r)) = r^{-1}R(z, r)$ for $(z, r) \in \partial X \times [r_0, \infty)$.
- There exists j_t , an almost complex structure on ξ such that $J_t|_{\xi(z, r)} = j_t$ for $(z, r) \in \partial X \times [r_0, \infty)$.

We denote $\mathcal{J}(X, \lambda) := \bigcup_{r_0 \geq 1} \mathcal{J}(X, \lambda : r_0)$.

Remark 2.5. Recall that an almost complex structure J on \hat{X} is compatible with $d\hat{\lambda}$ if and only if the bilinear form on $T\hat{X}$

$$\langle v, w \rangle_J := d\hat{\lambda}(v, Jw) \quad (v, w \in T\hat{X})$$

is a Riemannian metric. Let us denote $\langle v, v \rangle_J^{1/2}$ by $|v|_J$.

Let $H \in \mathcal{H}_{\text{ad}}(X, \lambda)$, $J \in \mathcal{J}(X, \lambda)$. For $x_-, x_+ \in \mathcal{P}(H)$, we consider the Floer equation for $u : \mathbb{R} \times \mathbb{R}/\mathbb{Z} \rightarrow \hat{X}$, namely:

$$\partial_s u - J_t(\partial_t u - X_{H_t}(u)) = 0, \quad u(s) \rightarrow x_\pm (s \rightarrow \pm\infty).$$

In the second formula, $u(s)$ denotes the map $\mathbb{R}/\mathbb{Z} \rightarrow \hat{X}; t \mapsto u(s, t)$. Let us denote the moduli space of solutions of the above Floer equations by $\hat{\mathcal{M}}(x_-, x_+ : H, J)$. $\hat{\mathcal{M}}(x_-, x_+ : H, J)$ admits a natural \mathbb{R} action:

$$s_0 \cdot u(s, t) := u(s - s_0, t).$$

$\mathcal{M}(x_-, x_+ : H, J)$ denotes the quotient of $\hat{\mathcal{M}}(x_-, x_+ : H, J)$ by the above \mathbb{R} action.

For generic J , $\mathcal{M}(x_-, x_+ : H, J)$ is a smooth manifold with dimension $\mu_{\text{CZ}}(x_-) - \mu_{\text{CZ}}(x_+) - 1$. For such J , we define the differential $\partial_{H,J} : C_k(H) \rightarrow C_{k-1}(H)$ by

$$\partial_{H,J}[x_-] := \sum_{x_+ \in \mathcal{P}_{k-1}(H)} \#_* \mathcal{M}(x_-, x_+ : H, J) \cdot [x_+] \quad (x_- \in \mathcal{P}_k(H)).$$

Then, $(C_*(H), \partial_{H,J})$ becomes a chain complex. It follows from the following C^0 bound for Floer trajectories (this is a special case of lemma 2.7, which is stated later):

Lemma 2.6. *There exists a compact set $B \subset \hat{X}$ such that for any $x_-, x_+ \in \mathcal{P}(H)$ and $u \in \hat{\mathcal{M}}(x_-, x_+ : H, J)$, $u(\mathbb{R} \times \mathbb{R}/\mathbb{Z}) \subset B$.*

It can be shown that the homology group of the complex $(C_*(H), \partial_{H,J})$ is independent of choices of J , and we denote it by $\text{HF}_*(H)$, or $\text{HF}_*(H : (X, \lambda))$, when we need to specify the Liouville domain.

Let $H_-, H_+ \in \mathcal{H}_{\text{ad}}(X, \lambda)$ and assume that $a_{H_-} \leq a_{H_+}$. Then, there exists a canonical morphism $\text{HF}_*(H_-) \rightarrow \text{HF}_*(H_+)$. This is constructed as follows: take $r_0 \geq 1$ and $(H_s)_{s \in \mathbb{R}}$, a family of elements in $\mathcal{H}(X, \lambda : r_0)$ and $(J_s)_{s \in \mathbb{R}}$, a family of elements in $\mathcal{J}(X, \lambda : r_0)$ which satisfy the following conditions:

- There exists $s_0 > 0$ such that $H_s = \begin{cases} H_- & (s \leq -s_0) \\ H_+ & (s \geq s_0) \end{cases}$, $J_s = \begin{cases} J_{-s_0} & (s \leq -s_0) \\ J_{s_0} & (s \geq s_0) \end{cases}$.
- $\partial_s a_{H_s} \geq 0$.

For $x_- \in \mathcal{P}(H_-)$ and $x_+ \in \mathcal{P}(H_+)$, consider the Floer equation for $u : \mathbb{R} \times \mathbb{R}/\mathbb{Z} \rightarrow \hat{X}$:

$$\partial_s u - J_{s,t}(\partial_t u - X_{H_{s,t}}(u)) = 0, \quad u(s) \rightarrow x_{\pm} (s \rightarrow \pm\infty),$$

where $H_{s,t} := (H_s)_t$, $J_{s,t} := (J_s)_t$.

We denote the moduli space of solutions of the above Floer equation by $\hat{\mathcal{M}}(x_-, x_+ : (H_s, J_s)_s)$. For generic $(J_s)_s$, $\hat{\mathcal{M}}(x_-, x_+ : (H_s, J_s)_s)$ is a smooth manifold of dimension $\mu_{\text{CZ}}(x_-) - \mu_{\text{CZ}}(x_+)$. Taking such $(J_s)_s$, we define a morphism $\varphi : C_k(H_-) \rightarrow C_k(H_+)$ by

$$\varphi[x_-] = \sum_{x_+ \in \mathcal{P}_k(H_+)} \#_* \hat{\mathcal{M}}(x_-, x_+ : (H_s, J_s)_s) \cdot [x_+] \quad (x_- \in \mathcal{P}_k(H_-)).$$

Then, φ is a chain map from $(C_*(H_-), \partial_{H_-, J_-})$ to $(C_*(H_+), \partial_{H_+, J_+})$. It follows from the following C^0 bound for Floer trajectories (it follows from lemma 1.5 in [6]):

Lemma 2.7. *There exists a compact set $B \subset \hat{X}$ such that for any $x_- \in \mathcal{P}(H_-)$, $x_+ \in \mathcal{P}(H_+)$ and $u \in \hat{\mathcal{M}}(x_-, x_+ : (H_s, J_s)_s)$, $u(\mathbb{R} \times \mathbb{R}/\mathbb{Z}) \subset B$.*

Therefore, φ defines a morphism $\varphi_*: \mathrm{HF}_*(H_-) \rightarrow \mathrm{HF}_*(H_+)$. This morphism does not depend on choice of $(H_s, J_s)_s$.

To sum up, we have constructed the canonical morphism $\mathrm{HF}_*(H_-) \rightarrow \mathrm{HF}_*(H_+)$ for $H_-, H_+ \in \mathcal{H}_{\mathrm{ad}}(X, \lambda)$ such that $a_{H_-} \leq a_{H_+}$. This morphism is called *monotone morphism*.

We also study truncated version of the Floer homology. For any $x: \mathbb{R}/\mathbb{Z} \rightarrow \hat{X}$, let

$$\mathcal{A}_H(x) := \int_{\mathbb{R}/\mathbb{Z}} x^* \hat{\lambda} - H(x(t)) dt.$$

For any interval $I \subset [-\infty, \infty]$, let $C_k^I(H)$ be the free \mathbb{Z}_2 -module generated over

$$\{x \in \mathcal{P}_k(H) \mid \mathcal{A}_H(x) \in I\}.$$

For $x_-, x_+ \in \mathcal{P}(H)$ and $u \in \hat{\mathcal{M}}(x_-, x_+ : H, J)$, by straightforward calculations we get

$$-\partial_s(\mathcal{A}_H(u(s))) = \int_{\mathbb{R}/\mathbb{Z}} |\partial_s u(s, t)|_{J_t}^2 dt.$$

In particular, if $\mathcal{A}_H(x) < \mathcal{A}_H(y)$, then $\hat{\mathcal{M}}(x, y : H, J) = \emptyset$. Hence for any interval $I \subset [-\infty, \infty]$, $(C_*^I(H), \partial_{H, J})$ is a chain complex. Then, we denote $H_*(C_*^I(H), \partial_{H, J})$ (which does not depend on J) by $\mathrm{HF}_*^I(H)$.

For $-\infty \leq a < b < c \leq \infty$, there exists a short exact sequence

$$0 \rightarrow C_*^{[a, b]}(H) \rightarrow C_*^{[a, c]}(H) \rightarrow C_*^{[b, c]}(H) \rightarrow 0.$$

Hence we get a long exact sequence

$$(1) \quad \cdots \rightarrow \mathrm{HF}_*^{[a, b]}(H) \rightarrow \mathrm{HF}_*^{[a, c]}(H) \rightarrow \mathrm{HF}_*^{[b, c]}(H) \rightarrow \mathrm{HF}_{*-1}^{[a, b]}(H) \rightarrow \cdots$$

2.4. Symplectic homology. Let (X, λ) be a Liouville domain. In this subsection, we define *symplectic homology* $\mathrm{SH}_*^I(X, \lambda)$ for any interval $I \subset \mathbb{R}$.

First, we define $\mathcal{H}^{\mathrm{rest}}(X, \lambda) \subset \mathcal{H}(X, \lambda)$ by

$$\mathcal{H}^{\mathrm{rest}}(X, \lambda) := \{H \in \mathcal{H}(X, \lambda : 1) \mid H_t|_X < 0 \text{ for any } t \in \mathbb{R}/\mathbb{Z}\}.$$

For $H_-, H_+ \in \mathcal{H}^{\mathrm{rest}}(X, \lambda)$, we denote $H_- \leq H_+$ if and only if $(H_-)_t \leq (H_+)_t$ for any $t \in \mathbb{R}/\mathbb{Z}$.

Let $H_-, H_+ \in \mathcal{H}_{\mathrm{ad}}^{\mathrm{rest}}(X, \lambda) := \mathcal{H}_{\mathrm{ad}}(X, \lambda) \cap \mathcal{H}^{\mathrm{rest}}(X, \lambda)$. When $H_- \leq H_+$, we can construct a morphism $\mathrm{HF}_*^I(H_-) \rightarrow \mathrm{HF}_*^I(H_+)$ for any interval $I \subset \mathbb{R}$. This is constructed as follows. First, take $(H_s)_{s \in \mathbb{R}}$, a family of elements of $\mathcal{H}^{\mathrm{rest}}(X, \lambda)$ and $(J_s)_{s \in \mathbb{R}}$, a family of elements of $\mathcal{J}(X, \lambda : 1)$ which satisfy the following properties:

- There exists $s_0 > 0$ such that $H_s = \begin{cases} H_- & (s \leq -s_0) \\ H_+ & (s \geq s_0) \end{cases}$, $J_s = \begin{cases} J_{-s_0} & (s \leq -s_0) \\ J_{s_0} & (s \geq s_0) \end{cases}$.
- $\partial_s H_{s, t}(x) \geq 0$ for any $(s, t) \in \mathbb{R} \times \mathbb{R}/\mathbb{Z}$ and $x \in \hat{X}$.

For $x_- \in \mathcal{P}(H_-)$, $x_+ \in \mathcal{P}(H_+)$ and $u \in \hat{\mathcal{M}}(x_-, x_+ : (H_s, J_s)_s)$,

$$-\partial_s(\mathcal{A}_{H_s}(u(s))) = \int_{\mathbb{R}/\mathbb{Z}} |\partial_s u|_{J_{s,t}}^2 + \partial_s H_{s,t}(u) dt \geq 0.$$

Hence if $\mathcal{A}_{H_-}(x_-) < \mathcal{A}_{H_+}(x_+)$, then $\hat{\mathcal{M}}(x_-, x_+ : (H_s, J_s)_s) = \emptyset$.

Therefore the morphism $\varphi : (C_*^I(H_-), \partial_{H_-, J_-}) \rightarrow (C_*^I(H_+), \partial_{H_+, J_+})$ defined by

$$\varphi[x_-] = \sum_{x_+ \in \mathcal{P}_k(H_+)} \# \hat{\mathcal{M}}(x_-, x_+ : (H_s, J_s)_s) \cdot [x_+] \quad (x_- \in \mathcal{P}_k(H_-))$$

is a chain map. Hence we get a morphism $\mathrm{HF}_*^I(H_-) \rightarrow \mathrm{HF}_*^I(H_+)$. This morphism does not depend on choices of $(H_s, J_s)_s$. Then, we define $\mathrm{SH}_*^I(X, \lambda)$ by

$$\mathrm{SH}_*^I(X, \lambda) := \varinjlim_{H \in \mathcal{H}_{\mathrm{ad}}^{\mathrm{rest}}(X, \lambda)} \mathrm{HF}_*^I(H).$$

For $-\infty \leq a < b < c \leq \infty$, by taking limit of (1), we get a long exact sequence

$$(2) \quad \cdots \rightarrow \mathrm{SH}_*^{[a,b]}(X, \lambda) \rightarrow \mathrm{SH}_*^{[a,c]}(X, \lambda) \rightarrow \mathrm{SH}_*^{[b,c]}(X, \lambda) \rightarrow \mathrm{SH}_{*-1}^{[a,b]}(X, \lambda) \rightarrow \cdots$$

For $a \in (-\infty, \infty]$, $\mathrm{SH}_*^{(-\infty, a)}(X, \lambda)$ is often denoted by $\mathrm{SH}_*^{<a}(X, \lambda)$. $\mathrm{SH}_*^{<\infty}(X, \lambda)$ is often abbreviated by $\mathrm{SH}_*(X, \lambda)$. The following lemma will be useful in later:

Lemma 2.8. *For any $H \in \mathcal{H}_{\mathrm{ad}}(X, \lambda)$, there exists a canonical isomorphism $\mathrm{SH}_*^{<a_H}(X, \lambda) \rightarrow \mathrm{HF}_*(H)$. When $H_-, H_+ \in \mathcal{H}_{\mathrm{ad}}(X, \lambda)$ satisfy $a_{H_-} \leq a_{H_+}$, the following diagram commutes:*

$$\begin{array}{ccc} \mathrm{SH}_*^{<a_{H_-}}(X, \lambda) & \xrightarrow{\cong} & \mathrm{HF}_*(H_-) \\ \downarrow & & \downarrow \\ \mathrm{SH}_*^{<a_{H_+}}(X, \lambda) & \xrightarrow{\cong} & \mathrm{HF}_*(H_+) \end{array}$$

Proof. It is not hard to check that the following natural morphisms are all isomorphic:

$$\begin{aligned} \varinjlim_{\substack{G \in \mathcal{H}_{\mathrm{ad}}^{\mathrm{rest}} \\ a_G \leq a_H}} \mathrm{HF}_*^{<a_H}(G) &\rightarrow \varinjlim_{G \in \mathcal{H}_{\mathrm{ad}}^{\mathrm{rest}}} \mathrm{HF}_*^{<a_H}(G) = \mathrm{SH}_*^{<a_H}(X, \lambda), \\ \varinjlim_{\substack{G \in \mathcal{H}_{\mathrm{ad}}^{\mathrm{rest}} \\ a_G \leq a_H}} \mathrm{HF}_*^{<a_H}(G) &\rightarrow \varinjlim_{\substack{G \in \mathcal{H}_{\mathrm{ad}}^{\mathrm{rest}} \\ a_G \leq a_H}} \mathrm{HF}_*(G), \\ \varinjlim_{\substack{G \in \mathcal{H}_{\mathrm{ad}}^{\mathrm{rest}} \\ a_G \leq a_H}} \mathrm{HF}_*(G) &\rightarrow \varinjlim_{\substack{G \in \mathcal{H}_{\mathrm{ad}} \\ a_G \leq a_H}} \mathrm{HF}_*(G), \\ \varinjlim_{\substack{G \in \mathcal{H}_{\mathrm{ad}} \\ a_G \leq a_H}} \mathrm{HF}_*(G) &\rightarrow \mathrm{HF}_*(H). \end{aligned}$$

By composing the above isomorphisms and their inverses, we get an isomorphism $\mathrm{SH}_*^{<a_H}(X, \lambda) \rightarrow \mathrm{HF}_*(H)$. This proves the first assertion. The second assertion follows from the above construction. \square

We recall three well-known results on symplectic homology. All these results were established in [7]. The first result is the following:

Theorem 2.9. *For any $0 < \delta \leq \delta(\partial X, \lambda)$, there exists a canonical isomorphism $\mathrm{SH}_*^{\leq \delta}(X, \lambda) \rightarrow H_{*+n}(X, \partial X)$.*

The second result is the following:

Theorem 2.10. *If (X, λ) and (Y, λ') are isotopic as Liouville domains, then $\mathrm{SH}_*(X, \lambda) \cong \mathrm{SH}_*(Y, \lambda')$.*

As a corollary of the above theorem, we can conclude that $\mathrm{SH}_*(X, \lambda)$ depends only on $d\lambda$. Assume that $(X, \lambda), (X, \lambda')$ are Liouville domains, and $d\lambda = d\lambda'$. Then, $(X, t\lambda + (1-t)\lambda')_{0 \leq t \leq 1}$ is a family of Liouville domains, and theorem 2.10 implies that $\mathrm{SH}_*(X, \lambda) \cong \mathrm{SH}_*(X, \lambda')$. Hence we often denote $\mathrm{SH}_*(X, \lambda)$ by $\mathrm{SH}_*(X, d\lambda)$.

The third result is the following:

Theorem 2.11. *For positive integer n and $r > 0$, let*

$$B^{2n}(r) := \{(q, p) \in \mathbb{R}^{2n} \mid |q|^2 + |p|^2 \leq r^2\}.$$

Then, $(B^{2n}(r), \lambda_n)$ is a Liouville domain, and $\mathrm{SH}_(B^{2n}(r), \lambda_n) = 0$.*

For proofs, see proposition 1.4 in [7] for theorem 2.9, theorem 1.7 in [7] for theorem 2.10, and section 4, example 1 in [7] for theorem 2.11.

Note that theorem 2.10 does not hold for truncated symplectic homology. However, the following result holds:

Theorem 2.12. *Let $(X, \lambda), (X, \lambda')$ be Liouville domains, and assume that $d\lambda = d\lambda'$. Then, for any $a \in (0, \infty]$, there exists a canonical isomorphism $\psi^{<a} : \mathrm{SH}_*^{<a}(X, \lambda) \rightarrow \mathrm{SH}_*^{<a}(X, \lambda')$. Moreover, for any $0 < a \leq b \leq \infty$,*

$$\begin{array}{ccc} \mathrm{SH}_*^{<a}(X, \lambda) & \xrightarrow{\psi^{<a}} & \mathrm{SH}_*^{<a}(X, \lambda') \\ \downarrow & & \downarrow \\ \mathrm{SH}_*^{<b}(X, \lambda) & \xrightarrow{\psi^{<b}} & \mathrm{SH}_*^{<b}(X, \lambda') \end{array}$$

commutes.

Theorem 2.12 is proved in the appendix.

3. SYMPLECTIC CAPACITY VIA SYMPLECTIC HOMOLOGY

Definition 3.1. Let (X, λ) be a Liouville domain. $\mathrm{cap}_S(X, \lambda)$ is defined by

$$\mathrm{cap}_S(X, \lambda) := \inf \{a \in (0, \infty] \mid \mathrm{SH}_n^{\leq \delta(\partial X, \lambda)}(X, \lambda) \rightarrow \mathrm{SH}_n^{<a}(X, \lambda) \text{ vanishes}\}.$$

Remark 3.2. The above capacity is introduced by C.Viterbo in [7], section 5.3.

Lemma 3.3. *Let (X, λ) be a Liouville domain.*

- (1) $\text{cap}_S(X, a\lambda) = a \cdot \text{cap}_S(X, \lambda)$ for any $a \in (0, \infty)$.
- (2) $\text{SH}_*(X, \lambda) = 0 \implies \text{cap}_S(X, \lambda) < \infty$.
- (3) $\text{cap}_S(X, \lambda)$ depends only on $d\lambda$.

Proof. (1) and (2) are immediate from the definition. (3) follows from theorem 2.12. \square

The goal of this section is to prove the following three properties of cap_S .

Theorem 3.4. *Let (X, λ) be a Liouville domain, and X_{in} be a submanifold of X of codimension 0. If (X_{in}, λ) is a Liouville domain, then $\text{cap}_S(X_{\text{in}}, \lambda) \leq \text{cap}_S(X, \lambda)$.*

Theorem 3.5. *Let (X, λ) be a Liouville domain. If $\text{cap}_S(X, \lambda) < \infty$, there exists $x \in \mathcal{P}_0(\partial X, \lambda)$ such that $\tau(x) = \text{cap}_S(X, \lambda)$, $\mu_{\text{CZ}}(x) \leq n + 1$.*

Theorem 3.6. *Let $\pi: Y \rightarrow X$ be a covering map such that $\deg \pi < \infty$. If (X, λ) is a Liouville domain, then $(Y, \pi^*\lambda)$ is a Liouville domain, and $\text{cap}_S(Y, \pi^*\lambda) \leq \text{cap}_S(X, \lambda)$.*

Remark 3.7. Conley-Zehnder index for elements in $\mathcal{P}_0(\partial X, \lambda)$, which appears on the statement of theorem 3.5 have not been defined. It is defined at the beginning of section 3.3.

It seems to the author that various variants of results similar to theorems 3.4, 3.5 are known or expected to hold by experts. We give its proof below for the sake of completeness since the author is unable to find their proofs in the literature. On the other hand, theorem 3.6 is new, though its proof is not very difficult. Theorem 3.6 plays a crucial role in the proof of theorem 4.13, which is the main result in section 4.

3.1. Proof of theorem 3.4. First we prove the following lemma.

Lemma 3.8. *Let (X, λ) and X_{in} be as in theorem 3.4, and $\varepsilon \in (0, 1)$. Let $H_-, H_+ \in \mathcal{H}_{\text{ad}}(X, \lambda)$, $(H_s)_{s \in \mathbb{R}}$ be a family of elements of $\mathcal{H}(X, \lambda)$, $(J_s)_{s \in \mathbb{R}}$ be a family of elements of $\mathcal{J}(X, \lambda)$. Assume that they satisfy the following conditions:*

- (i) *There exists $s_0 > 0$ such that $H_s = \begin{cases} H_- & (s \leq -s_0) \\ H_+ & (s \geq s_0) \end{cases}$.*
- (ii) *$\partial_s H_{s,t}(x) \geq 0$ for any $(s, t) \in \mathbb{R} \times \mathbb{R}/\mathbb{Z}$ and $x \in \hat{X}$.*
- (iii) *There exists $a \in C^\infty(\mathbb{R})$ such that $H_{s,t}(z, r) = a(s)(r - \varepsilon)$ for $(z, r) \in \partial X_{\text{in}} \times [\varepsilon^{2/3}, \varepsilon^{1/3}]$.*
- (iv) *$dr \circ J_{s,t} = -\lambda$ on $\partial X_{\text{in}} \times [\varepsilon^{2/3}, \varepsilon^{1/3}]$.*

Assume $x_- \in \mathcal{P}(H_-)$ and $x_+ \in \mathcal{P}(H_+)$ satisfy $x_-(\mathbb{R}/\mathbb{Z}), x_+(\mathbb{R}/\mathbb{Z}) \subset X_{\text{in}}(\varepsilon^{1/3})$. Then, for any $u \in \hat{\mathcal{M}}(x_-, x_+ : (H_s, J_s)_s)$, $u(\mathbb{R} \times \mathbb{R}/\mathbb{Z}) \subset X_{\text{in}}(\varepsilon^{1/3})$.

Proof. First notice that $x_-(\mathbb{R}/\mathbb{Z}), x_+(\mathbb{R}/\mathbb{Z})$ are contained in $X_{\text{in}}(\varepsilon^{2/3})$, since $(H_\pm)_t(z, r) = a(\pm s_0)(r - \varepsilon)$ for $(z, r) \in \partial X_{\text{in}} \times [\varepsilon^{2/3}, \varepsilon^{1/3}]$ and $a(\pm s_0) \notin \tau(\partial X_{\text{in}}, \lambda)$ (this follows from (i), (iii) and $H_\pm \in \mathcal{H}_{\text{ad}}(X, \lambda)$). We will prove that for any $u \in \hat{\mathcal{M}}(x_-, x_+ : (H_s, J_s)_s)$, $u(\mathbb{R} \times \mathbb{R}/\mathbb{Z}) \subset X_{\text{in}}(\varepsilon^{1/3})$. If this is not true, for any $r_0 \in (\varepsilon^{2/3}, \varepsilon^{1/3})$, $D_{r_0} := \mathbb{R} \times \mathbb{R}/\mathbb{Z} \setminus u^{-1}(\text{int} X_{\text{in}}(r_0))$ is non-empty. Note that D_{r_0} is compact since $x_\pm(\mathbb{R}/\mathbb{Z}) \subset X_{\text{in}}(\varepsilon^{2/3})$. For

generic r_0 , u is transverse to $\partial X_{\text{in}} \times \{r_0\}$, hence we may assume that D_{r_0} is a compact surface with boundary.

It is easy to verify that $\partial_s u$ is not constantly 0 on D_{r_0} . Hence

$$\int_{D_{r_0}} |\partial_s u|_{J_{s,t}}^2 ds dt > 0.$$

Since u satisfies the Floer equation $\partial_s u - J_{s,t}(\partial_t u - X_{H_{s,t}}(u)) = 0$,

$$\begin{aligned} \int_{D_{r_0}} |\partial_s u|_{J_{s,t}}^2 + \partial_s H_{s,t}(u) ds dt &= \int_{D_{r_0}} d\hat{\lambda}(\partial_t u, \partial_s u) + dH_{s,t}(\partial_s u) + \partial_s H_{s,t}(u) ds dt \\ &= \int_{\partial D_{r_0}} -u^* \lambda + H_{s,t}(u) dt. \end{aligned}$$

Since $u(\partial D_{r_0}) \subset \partial X_{\text{in}} \times \{r_0\}$, we get by (iii)

$$(s, t) \in \partial D_{r_0} \implies H_{s,t}(u(s, t)) = a(s)(r_0 - \varepsilon), \quad \lambda(X_{H_{s,t}}(u(s, t))) = a(s)r_0.$$

Therefore

$$\int_{\partial D_{r_0}} -u^* \lambda + H_{s,t}(u) dt = \int_{\partial D_{r_0}} \lambda(X_{H_{s,t}} \otimes dt - du) - \varepsilon \int_{\partial D_{r_0}} a(s) dt.$$

On the other hand, the Floer equation is equivalent to

$$J_{s,t} \circ (X_{H_{s,t}} \otimes dt - du) = -(X_{H_{s,t}} \otimes dt - du) \circ j,$$

where j is the complex structure on $\mathbb{R} \times \mathbb{R}/\mathbb{Z}$, defined by $j(\partial_s) = \partial_t$. Therefore by (iv),

$$\int_{\partial D_{r_0}} \lambda(X_{H_{s,t}} \otimes dt - du) = \int_{\partial D_{r_0}} \lambda(J_{s,t} \circ (X_{H_{s,t}} \otimes dt - du) \circ j) = \int_{\partial D_{r_0}} dr(X_{H_{s,t}} \otimes dt - du) \circ j.$$

$dr(X_{H_{s,t}}) = 0$ on $\partial X_{\text{in}} \times \{r_0\}$. Moreover, if V is a vector tangent to ∂D_{r_0} , and positive with respect to the boundary orientation, then jV points inwards, hence $dr(du(jV)) \geq 0$. Therefore,

$$\int_{\partial D_{r_0}} \lambda(X_{H_{s,t}} \otimes dt - du) \leq 0.$$

Finally,

$$\int_{D_{r_0}} |\partial_s u|_{J_{s,t}}^2 + \partial_s H_{s,t}(u) ds dt \leq -\varepsilon \int_{\partial D_{r_0}} a(s) dt = -\varepsilon \int_{D_{r_0}} \partial_s a(s) ds dt.$$

Since $\partial_s H_{s,t} \geq 0$ and $\partial_s a \geq 0$ by (ii), this implies

$$\int_{D_{r_0}} |\partial_s u|_{J_{s,t}}^2 ds dt \leq 0.$$

This is a contradiction. □

We prove theorem 3.4.

Proof. We prove that, if a satisfies $a > \text{cap}_S(X, \lambda)$ and $a \notin \tau(\partial X, \lambda) \cup \tau(\partial X_{\text{in}}, \lambda)$, then $a > \text{cap}_S(X_{\text{in}}, \lambda)$. This implies $\text{cap}_S(X, \lambda) \geq \text{cap}_S(X_{\text{in}}, \lambda)$, since $\tau(\partial X, \lambda)$ and $\tau(\partial X_{\text{in}}, \lambda)$ are null sets. In the rest of this proof, we assume that X and X_{in} are connected. General case follows at once from this particular case.

Take $\varepsilon > 0$ so that $[a(1-\varepsilon), a]$ is disjoint from $\tau(\partial X, \lambda)$ and $\tau(\partial X_{\text{in}}, \lambda)$. For any $c > 0$, define $H_c : \hat{X}_{\text{in}} \rightarrow \mathbb{R}$ and $K_c : \hat{X} \rightarrow \mathbb{R}$ as follows:

$$H_c(x) = \begin{cases} 0 & (x \in X_{\text{in}}(\varepsilon)) \\ c(r - \varepsilon) & (x = (z, r) \in \partial X_{\text{in}} \times [\varepsilon, \infty)) \end{cases},$$

$$K_c(x) = \begin{cases} H_c(x) & (x \in X_{\text{in}}) \\ c(1 - \varepsilon) & (x \in X \setminus X_{\text{in}}) \\ c(r - \varepsilon) & (x = (z, r) \in \partial X \times [1, \infty)). \end{cases}$$

Take $\delta > 0$ so small that $\delta < \min\{\delta(\partial X, \lambda), \delta(\partial X_{\text{in}}, \lambda)\}$. Then, perturbing K_a, K_δ and H_a, H_δ respectively, we can take $K'_a, K'_\delta \in \mathcal{H}_{\text{ad}}(X, \lambda)$ and $H'_a, H'_\delta \in \mathcal{H}_{\text{ad}}(X_{\text{in}}, \lambda)$ which satisfy the following properties:

- (i) For $c \in \{\delta, a\}$, the following holds:
 - (a) $(H'_c)_t = H_c$ on $\partial X_{\text{in}} \times [\varepsilon^{2/3}, \infty)$ for any $t \in \mathbb{R}/\mathbb{Z}$.
 - (b) $(K'_c)_t = K_c$ on $\partial X \times [2, \infty)$ for any $t \in \mathbb{R}/\mathbb{Z}$.
 - (c) $(K'_c)_t = (H'_c)_t$ on $X_{\text{in}}(\varepsilon^{1/3})$ for any $t \in \mathbb{R}/\mathbb{Z}$.
 - (d) For $x \in \mathcal{P}(K'_c)$, $\mathcal{A}_{K'_c}(x) > 0$ if and only if $x(\mathbb{R}/\mathbb{Z}) \subset X_{\text{in}}(\varepsilon^{1/3})$.
- (ii) $(K'_\delta)_t \leq (K'_a)_t$ and $(H'_\delta)_t \leq (H'_a)_t$ for any $t \in \mathbb{R}/\mathbb{Z}$.
- (iii) H'_δ and K'_δ are time independent. i.e. There exist $h \in C^\infty(\hat{X}_{\text{in}})$ and $k \in C^\infty(\hat{X})$ such that $(H'_\delta)_t = h$, $(K'_\delta)_t = k$. Moreover, $\mathcal{P}(H'_\delta) = \text{Crit}(h)$, $\mathcal{P}(K'_\delta) = \text{Crit}(k)$ and if $p \in \text{Crit}(k)$ satisfies $\text{ind } p = 0$, then $p \in X_{\text{in}}$.

Let $c \in \{\delta, a\}$. Then, by (i)-(c) and (i)-(d), if $x \in \mathcal{P}(K'_c)$ satisfies $\mathcal{A}_{K'_c}(x) > 0$, x can be identified with a solution of $\partial_t x = X_{(H'_c)_t}(x)$. We define $\psi_c : C_*^{>0}(K'_c) \rightarrow C_*(H'_c)$ by

$$\psi_c[x] = \begin{cases} [x] & (x \text{ is contractible in } X_{\text{in}}) \\ 0 & (\text{otherwise}) \end{cases}.$$

'if' part of (i)-(d) implies that ψ_c is an epimorphism.

Let $J = (J_t)_{t \in \mathbb{R}/\mathbb{Z}}$ be a family of almost complex structures on X_{in} , such that each J_t is compatible with $d\lambda$ and satisfies $dr \circ J_t = -\lambda$ on $X_{\text{in}} \times [\varepsilon^{2/3}, \varepsilon^{1/3}]$.

By (i)-(a) and (i)-(c), $(K'_c)_t(z, r) = c(r - \varepsilon)$ for $(z, r) \in X_{\text{in}} \times [\varepsilon^{2/3}, \varepsilon^{1/3}]$. Therefore, by lemma 3.8, if we extend J to $J^X \in \mathcal{J}(X, \lambda)$ and $J^{X_{\text{in}}} \in \mathcal{J}(X_{\text{in}}, \lambda)$, ψ_c defines a chain map from $(C_*^{>0}(K'_c), \partial_{K'_c, J^X})$ to $(C_*(H'_c), \partial_{H'_c, J^{X_{\text{in}}}})$.

It induces a morphism

$$\text{HF}_*(K'_c) \rightarrow \text{HF}_*^{>0}(K'_c) \rightarrow \text{HF}_*(H'_c).$$

We will denote this morphism also by ψ_c .

It follows from lemma 3.8 and (ii) that

$$(3) \quad \begin{array}{ccc} \mathrm{HF}_*(K'_\delta) & \xrightarrow{\psi_\delta} & \mathrm{HF}_*(H'_\delta) \\ \downarrow & & \downarrow \\ \mathrm{HF}_*(K'_a) & \xrightarrow{\psi_a} & \mathrm{HF}_*(H'_a) \end{array}$$

commutes, where vertical morphisms are monotone morphisms.

We complete the proof. We have to show that if $\mathrm{SH}_n^{<\delta}(X, \lambda) \rightarrow \mathrm{SH}_n^{<a}(X, \lambda)$ vanishes, then $\mathrm{SH}_n^{<\delta}(X_{\mathrm{in}}, \lambda) \rightarrow \mathrm{SH}_n^{<a}(X_{\mathrm{in}}, \lambda)$ vanishes.

By (i)-(b), $a_{K'_c} = a_{H'_c} = c$ for $c \in \{\delta, a\}$. Hence by lemma 2.8, it is enough to prove that if $\mathrm{HF}_n(K'_\delta) \rightarrow \mathrm{HF}_n(K'_a)$ vanishes, then $\mathrm{HF}_n(H'_\delta) \rightarrow \mathrm{HF}_n(H'_a)$ vanishes.

By (iii), $C_k(H'_\delta) = C_k(K'_\delta) = 0$ for $k \geq n+1$, and $C_n(K'_\delta)$ is identified with $C_n(H'_\delta)$. Hence $\psi_\delta : \mathrm{HF}_n(K'_\delta) \rightarrow \mathrm{HF}_n(H'_\delta)$ is injective, therefore isomorphic (recall that we have assumed X and X_{in} to be connected). Then, (3) implies that if $\mathrm{HF}_n(K'_\delta) \rightarrow \mathrm{HF}_n(K'_a)$ vanishes, then $\mathrm{HF}_n(H'_\delta) \rightarrow \mathrm{HF}_n(H'_a)$ vanishes. \square

3.2. Proof of theorem 3.5. First we define the Conley-Zehnder index for elements of $\mathcal{P}_0(\partial X, \lambda)$. We assume that $n \geq 2$. Let $x \in \mathcal{P}_0(\partial X, \lambda)$. Then, there exists $\bar{x} : D^2 \rightarrow X$ such that $\bar{x}(e^{2\pi i\theta}) = x(\tau\theta)$. Take a trivialization of \bar{x}^*TX as symplectic vector bundle,

$$F : (\mathbb{R}^{2n}, \omega_n) \times D^2 \rightarrow \bar{x}^*TX; \quad (v, z) \mapsto (F_z(v), z),$$

such that for any $\theta \in \mathbb{R}/\mathbb{Z}$, the following holds:

$$(4) \quad \begin{cases} F_{e^{2\pi i\theta}}(0, \dots, 0, 0, 1) = \partial_r(x(\tau\theta)), \\ F_{e^{2\pi i\theta}}(0, \dots, 0, 1, 0) = R(x(\tau\theta)), \\ F_{e^{2\pi i\theta}}(\mathbb{R}^{2n-2} \times (0, 0)) = \xi(x(\tau\theta)). \end{cases}$$

Note that such trivialization exists only if $n \geq 2$.

Define a symplectic path $\gamma \in \mathcal{P}_\tau(2n-2)$ by

$$\gamma(t) := (F_{e^{2\pi it/\tau}}|_{\mathbb{R}^{2n-2} \times (0,0)})^{-1} \circ \Phi_t|_\xi \circ F_1|_{\mathbb{R}^{2n-2} \times (0,0)},$$

where $(\Phi_t)_t$ is the Poincaré map of the flow generated by R on ∂X . Then, define

$$\mu_{\mathrm{CZ}}(x) := i(\gamma).$$

x is called *nondegenerate* if and only if $\gamma \in \mathcal{P}_\tau^*(2n-2)$. The following lemma will be useful in later (note that it also implies that the above definition is consistent, i.e. it does not depend on choices of \bar{x}).

Lemma 3.9. *Let $H \in C^\infty(\hat{X})$ such that $\partial X = H^{-1}(0)$ and $\partial_r H > 0$ on ∂X . Then, there exists 1 : 1 correspondence between elements of $\mathcal{P}_0(\partial X, \lambda)$ and periodic orbits of X_H on ∂X , which are contractible in X . For $x \in \mathcal{P}_0(\partial X, \lambda)$, denote the corresponding periodic orbit of X_H by x_H . When $n \geq 2$,*

$$\mu_{\mathrm{CZ}}(x) = \sup_H \mu_{\mathrm{CZ}}(x_H),$$

where H runs over all Hamiltonians satisfying the conditions as above.

Proof. The first assertion is obvious. We prove the second assertion. Let $x \in \mathcal{P}_0(\partial X, \lambda)$, and x_H be the corresponding periodic orbit of X_H with period τ . Take $\bar{x} : D^2 \rightarrow X$ such that $\bar{x}(e^{2\pi i\theta}) = x_H(\tau\theta)$, and take a trivialization of \bar{x}^*TX as symplectic vector bundle $F : D^2 \times (\mathbb{R}^{2n}, \omega_n) \rightarrow \bar{x}^*TX$, which satisfies (4).

Define $\Gamma \in \mathcal{P}_\tau(2n)$ by

$$\Gamma(t) := (F_{e^{2\pi it/\tau}})^{-1} \circ \Phi_t \circ F_1$$

where $(\Phi_t)_t$ is the Poincaré map of the flow generated by X_H . Then, $\Gamma(t)$ can be written in the form

$$\Gamma(t) = \begin{pmatrix} \gamma(t) & 0 & 0 \\ 0 & 1 & 0 \\ 0 & a(t) & 1 \end{pmatrix}.$$

Denote the symplectic path $t \mapsto \begin{pmatrix} 1 & 0 \\ a(t) & 1 \end{pmatrix}$ by α . Then, by theorem 2.2-(2), $i(\Gamma) = i(\alpha) + i(\gamma)$. By definition, $i(\gamma) = \mu_{\text{CZ}}(x)$. On the other hand, it is easy to verify that

$$i(\alpha) = \begin{cases} -1 & (a(1) \leq 0) \\ 0 & (a(1) > 0) \end{cases}.$$

This proves the second assertion. \square

By lemma 3.9, it is possible to define the Conley-Zehnder index for $x \in \mathcal{P}_0(\partial X, \lambda)$ in another way, i.e.

$$\mu_{\text{CZ}}(x) := \sup_H \mu_{\text{CZ}}(x_H),$$

where H runs over all elements in $C^\infty(\hat{X})$ such that $\partial X = H^{-1}(0)$ and $\partial_r H > 0$ on ∂X . Note that this definition makes sense even when $n = 1$.

Corollary 3.10. *Let $H \in C^\infty(\hat{X})$ such that $\partial X = H^{-1}(0)$ and $\partial_r H > 0$ on ∂X . Assume that there exist $0 < r_0 < 1$ and $h : [r_0, \infty) \rightarrow \mathbb{R}$ such that $H(z, r) = h(r)$ and $\partial_r^2 h(1) > 0$. Then, for any $x \in \mathcal{P}_0(\partial X, \lambda)$, $\mu_{\text{CZ}}(x_H) = \mu_{\text{CZ}}(x)$.*

Proof. First consider the case $n \geq 2$. We use notations in the proof of lemma 3.9. Then, if H satisfies the condition as the above statement, $a(1) > 0$. Hence $\mu_{\text{CZ}}(x_H) = \mu_{\text{CZ}}(x)$. The case $n = 1$ is proved by similar arguments. \square

In the rest of this subsection, we prove theorem 3.5. First we consider cases in which all elements of $\mathcal{P}_0(\partial X, \lambda)$ are non-degenerate.

Lemma 3.11. *Let (X, λ) be as in theorem 3.5. Assume that all elements in $\mathcal{P}_0(\partial X, \lambda)$ are non-degenerate. Then, there exists $x \in \mathcal{P}_0(\partial X, \lambda)$ such that $\tau(x) = \text{cap}_S(X, \lambda)$ and $\mu_{\text{CZ}}(x) \in \{n, n+1\}$.*

Proof. We claim that for any $\varepsilon > 0$, there exists $x_\varepsilon \in \mathcal{P}_0(\partial X, \lambda)$ such that $|\text{cap}_S(X, \lambda) - \tau(x_\varepsilon)| < \varepsilon$ and $\mu_{\text{CZ}}(x_\varepsilon) \in \{n, n+1\}$. Since all elements in $\mathcal{P}_0(\partial X, \lambda)$ are non-degenerate, $\tau(\partial X, \lambda) \cap (0, T)$ is a finite set for any $T > 0$. Therefore, for sufficiently small $\varepsilon > 0$, $\tau(x_\varepsilon) = \text{cap}_S(X, \lambda)$.

We prove the above claim. It is enough to show the claim for sufficiently small $\varepsilon > 0$. In particular, we may assume that $\varepsilon/2 < \text{cap}_S(X, \lambda)$. The proof consists of 3 steps.

Step 1. First, take $(G^i)_i$, a sequence of time-independent Hamiltonians on \hat{X} which satisfies the following properties:

- $(G^i)_i$ is a cofinal sequence in $\mathcal{H}^{\text{rest}}(X, \lambda)$, i.e. for any $G \in \mathcal{H}^{\text{rest}}(X, \lambda)$, $G_t \leq G^i$ for any $t \in \mathbb{R}/\mathbb{Z}$ when i is sufficiently large.
- $G^i|_{X(1/2)}$ is sufficiently small in C^2 norm.
- There exists $g^i : [1/2, \infty) \rightarrow \mathbb{R}$ such that $G^i(z, r) = g^i(r)$ on $\partial X \times [1/2, \infty)$ and $\partial_r^2 g^i > 0$ on $(1/2, 1)$.

Then, $\mathcal{P}(G^i)$ consists of constant maps to $\text{Crit}(G^i)$ and S^1 -family of degenerate periodic orbits. There exists a 1:1 correspondence between S^1 -family of periodic orbits and elements of $\mathcal{P}_0(\partial X, \lambda)$ with periods less than a_{G^i} . Let $x \in \mathcal{P}_0(\partial X, \lambda)$ such that $\tau(x) < a_{G^i}$, and let γ_x be an element of a S^1 -family of periodic orbits which corresponds to x . Then, it follows from corollary 3.10 and $\partial_r^2 g^i > 0$ on $(1/2, 1)$ that $\mu_{\text{CZ}}(\gamma_x) = \mu_{\text{CZ}}(x)$. Moreover, by replacing G^i if necessary, we may assume that $|\mathcal{A}_{G^i}(\gamma_x) - \tau(x)| < \varepsilon/2$.

Step 2. Perturbing each $(G^i)_i$, we can construct $(H^i)_i$, a sequence in $\mathcal{H}_{\text{ad}}^{\text{rest}}(X, \lambda)$ with the following properties:

- (i) $(H^i)_i$ is a cofinal sequence in $\mathcal{H}_{\text{ad}}^{\text{rest}}(X, \lambda)$. i.e. for any $H \in \mathcal{H}_{\text{ad}}^{\text{rest}}(X, \lambda)$, $H_t \leq H^i_t$ for any $t \in \mathbb{R}/\mathbb{Z}$ for sufficiently large i .
- (ii) $H^i|_{X(1/2)}$ is time-independent, i.e. there exists $h^i \in C^\infty(X(1/2))$ such that $H^i_t|_{X(1/2)} = h^i$ for any $t \in \mathbb{R}/\mathbb{Z}$.
- (iii) For each $x \in \mathcal{P}_0(\partial X, \lambda)$ such that $\tau(x) < a_{H^i}$, there exists $x^\pm \in \mathcal{P}(H^i)$ such that $\mu_{\text{CZ}}(x^\pm) = \mu_{\text{CZ}}(x) + (1 \pm 1)/2$, $|\mathcal{A}_{H^i}(x^\pm) - \tau(x)| < \varepsilon/2$.
- (iv) $\mathcal{P}(H^i)$ consists of constant maps to $\text{Crit}(h^i)$ and $\{x^\pm \mid x \in \mathcal{P}_0(\partial X, \lambda), \tau(x) < a_{H^i}\}$.

Precise arguments on perturbations are carried out as in [4], proposition 2.2.

Step 3. Abbreviate $\text{cap}_S(X, \lambda)$ by c . By definition of cap_S , $\text{SH}_n^{<c-\varepsilon/2}(X, \lambda) \rightarrow \text{SH}_n^{<c+\varepsilon/2}(X, \lambda)$ is not injective. Then $\text{SH}_{n+1}^{[c-\varepsilon/2, c+\varepsilon/2)}(X, \lambda) \neq 0$, for the long exact sequence

$$\dots \rightarrow \text{SH}_{n+1}^{[c-\varepsilon/2, c+\varepsilon/2)}(X, \lambda) \rightarrow \text{SH}_n^{<c-\varepsilon/2}(X, \lambda) \rightarrow \text{SH}_n^{<c+\varepsilon/2}(X, \lambda) \rightarrow \dots$$

Therefore, by (i), $\text{HF}_{n+1}^{[c-\varepsilon/2, c+\varepsilon/2)}(H^i) \neq 0$ for sufficiently large i . This implies that there exists $x_i \in \mathcal{P}(H^i)$ such that $\mathcal{A}_{H^i}(x_i) \in [c - \varepsilon/2, c + \varepsilon/2)$ and $\mu_{\text{CZ}}(x_i) = n + 1$. Since $(H^i)_i$ is cofinal in $\mathcal{H}_{\text{rest}}(X, \lambda)$, we may assume that $\inf h^i > \varepsilon/2 - c$. Hence x_i is not a constant map to $\text{Crit}(h^i)$, and by (iv), there exists $x \in \mathcal{P}_0(\partial X, \lambda)$ such that $x_i = x^+$ or $x_i = x^-$. By (iii), $\tau(x) \in [c - \varepsilon, c + \varepsilon)$ and $\mu_{\text{CZ}}(x) \in \{n, n + 1\}$. Hence we have proved the claim. \square

We prove theorem 3.5.

Proof. Let $(\hat{X}, \hat{\lambda})$ be the completion of (X, λ) . For any positive smooth function f on ∂X , let Σ_f be the hypersurface in \hat{X} defined by $\{(z, f(z)) \mid z \in \partial X\}$, and D_f be the bounded domain in \hat{X} with boundary Σ_f . Then $(D_f, \hat{\lambda})$ is a Liouville domain.

If $|\log f|_{C^0(\partial X)} \leq c$, $X(e^{-c}) \subset D_f \subset X(e^c)$. Hence by theorem 3.4,

$$e^{-c} \leq \frac{\text{cap}_S(D_f, \hat{\lambda})}{\text{cap}_S(X, \lambda)} \leq e^c.$$

In particular, if $|\log f|_{C^0(\partial X)}$ is sufficiently small, then $\text{cap}_S(D_f, \hat{\lambda})$ is sufficiently close to $\text{cap}_S(X, \lambda)$.

Let $(f_m)_m$ be a sequence of $C^\infty(\partial X)$, such that all periodic Reeb orbits on $(\Sigma_{f_m}, \hat{\lambda})$ are non-degenerate, and $|\log f_m|_{C^2(\partial X)} \rightarrow 0$ as $m \rightarrow \infty$. By lemma 3.11, for each integer m there exists $x_m \in \mathcal{P}_0(\Sigma_{f_m}, \hat{\lambda})$ such that $\tau(x_m) = \text{cap}_S(D_{f_m}, \hat{\lambda})$ and $\mu_{\text{CZ}}(x_m) \in \{n, n+1\}$. Since $|\log f_m|_{C^2(\partial X)} \rightarrow 0$, $f_m \lambda$ converges to λ in C^2 . Hence, setting R_m to be the Reeb vector field on $(\partial X, f_m \lambda)$, R_m converges to R in C^1 . On the other hand, $\tau(x_m)$ converges to $\text{cap}_S(X, \lambda) > 0$. Hence, up to a subsequence, $(x_m)_m$ converges to $x_\infty \in \mathcal{P}_0(\partial X, \lambda)$ such that $\tau(x_\infty) = \text{cap}_S(X, \lambda)$. Moreover,

$$\mu_{\text{CZ}}(x_\infty) \leq \liminf_{m \rightarrow \infty} \mu_{\text{CZ}}(x_m) \leq n+1,$$

where the first inequality follows from lemma 2.4. \square

Theorem 3.5, together with lemma 3.9 implies the following corollary:

Corollary 3.12. *Let (X, λ) be a Liouville domain, and $\text{cap}_S(X, \lambda) < \infty$. Then, for any $H \in C^\infty(X)$ such that $\partial X = H^{-1}(0)$ and $\partial_r H > 0$ on ∂X , there exists $x : \mathbb{R}/\tau\mathbb{Z} \rightarrow \partial X$ such that $\partial_t x = X_H(x)$, $\int_{\mathbb{R}/\tau\mathbb{Z}} x^* \lambda = \text{cap}_S(X, \lambda)$ and $\mu_{\text{CZ}}(x) \leq n+1$.*

3.3. Proof of theorem 3.6. We prove the first assertion. Since $\deg \pi < \infty$, Y is compact. Define $Z \in \mathcal{X}(X)$ by $i_Z d\lambda = \lambda$. Then, $i_{\pi^* Z} d\pi^* \lambda = \pi^* \lambda$, and $\pi^* Z$ points outwards on ∂Y . Hence $(Y, \pi^* \lambda)$ is a Liouville domain. Now we prove the second assertion: $\text{cap}_S(Y, \pi^* \lambda) \leq \text{cap}_S(X, \lambda)$. Define $\hat{\pi} : \hat{Y} \rightarrow \hat{X}$ by

$$\hat{\pi}(y) = \begin{cases} \pi(y) & (y \in Y) \\ (\pi(z), r) & (y = (z, r) \in \partial Y \times [1, \infty)). \end{cases}$$

Then, $\hat{\pi} : \hat{Y} \rightarrow \hat{X}$ is a covering map and $\deg \hat{\pi} = \deg \pi$.

For $H \in \mathcal{H}(X, \lambda)$, denote $H \circ \hat{\pi}$ by \bar{H} . Since $\mathcal{P}(H)$ and $\mathcal{P}(\bar{H})$ consist of contractible solutions, $\mathcal{P}(\bar{H}) \rightarrow \mathcal{P}(H) : y \mapsto \hat{\pi} \circ y$ is $\deg(\pi) : 1$. We denote this map also by $\hat{\pi}$.

Denote the Poincaré map generated by X_H (resp. $X_{\bar{H}}$) by $(\Phi_t^H)_t$ (resp. $(\Phi_t^{\bar{H}})_t$). Clearly, $d\hat{\pi} \circ d\Phi_1^{\bar{H}} = d\Phi_1^H \circ d\hat{\pi}$. Hence $y \in \mathcal{P}(\bar{H})$ is non-degenerate if and only if $\hat{\pi}(y) \in \mathcal{P}(H)$ is non-degenerate. Moreover, since $\tau(\partial Y, \pi^* \lambda) \subset \tau(\partial X, \lambda)$ and $a_{\bar{H}} = a_H$, if $a_H \notin \tau(\partial X, \lambda)$ then $a_{\bar{H}} \notin \tau(\partial Y, \pi^* \lambda)$. Therefore, if $H \in \mathcal{H}_{\text{ad}}(X, \lambda)$ then $\bar{H} \in \mathcal{H}_{\text{ad}}(Y, \pi^* \lambda)$.

Let $H \in \mathcal{H}_{\text{ad}}(X, \lambda)$ and $J = (J_t)_{t \in \mathbb{R}/\mathbb{Z}} \in \mathcal{J}(X, \lambda)$. Recall that J is said to satisfy the *transversality condition* with respect to H if and only if for any $x, x' \in \mathcal{P}(H)$ and

$$u \in \hat{\mathcal{M}}(x, x' : H, J),$$

$$D_u : L^{1,p}(u^*T\hat{X}) \rightarrow L^p(u^*T\hat{X}); \quad \xi \mapsto \nabla_s \xi - J_t \nabla_t \xi_t - (\nabla_\xi J_t \cdot \partial_t u + \nabla_\xi(\nabla H_t))$$

is onto (p is an arbitrary real number satisfying $p > 2$). Let $\mathcal{J}_H(X, \lambda)$ be the set of elements of $\mathcal{J}(X, \lambda)$ which satisfy the transversality condition with respect to H .

Define $\bar{J} \in \mathcal{J}(Y, \pi^* \lambda)$ by $\bar{J}_t := \hat{\pi}^* J_t$. Then, for any $x \in \mathcal{P}(H)$ and $y \in \mathcal{P}(\bar{H})$, the following map is bijective:

$$\bigsqcup_{y' \in \hat{\pi}^{-1}(x)} \hat{\mathcal{M}}(y', y : \bar{H}, \bar{J}) \rightarrow \hat{\mathcal{M}}(x, \hat{\pi}(y) : H, J); \quad u \mapsto \hat{\pi} \circ u.$$

Clearly, $d\hat{\pi} \circ D_u = D_{\pi \circ u} \circ d\hat{\pi}$ for any u . Hence, $J \in \mathcal{J}_H(X, \lambda)$ if and only if $\bar{J} \in \mathcal{J}_{\bar{H}}(Y, \pi^* \lambda)$.

Let $H \in \mathcal{H}_{\text{ad}}(X, \lambda)$ and $J \in \mathcal{J}_H(X, \lambda)$. Then, we claim that

$$\psi_H : (C_*(H), \partial_{H,J}) \rightarrow (C_*(\bar{H}), \partial_{\bar{H},\bar{J}}); \quad [x] \mapsto \sum_{y \in \hat{\pi}^{-1}(x)} [y]$$

is a chain map. Let k be an integer and $x \in \mathcal{P}_k(H)$. Then, by definition

$$\begin{aligned} \psi_H(\partial_{H,J}[x]) &= \psi_H \left(\sum_{x' \in \mathcal{P}_{k-1}(H)} \# \mathcal{M}(x, x' : H, J) \cdot [x'] \right) \\ &= \sum_{y' \in \mathcal{P}_{k-1}(\bar{H})} \# \mathcal{M}(x, \hat{\pi}(y') : H, J) \cdot [y'], \\ \partial_{\bar{H},\bar{J}}(\psi_H[x]) &= \sum_{y \in \hat{\pi}^{-1}(x)} \partial_{\bar{H},\bar{J}}[y] = \sum_{y' \in \mathcal{P}_{k-1}(\bar{H})} \left(\sum_{y \in \hat{\pi}^{-1}(x)} \# \mathcal{M}(y, y' : \bar{H}, \bar{J}) \right) \cdot [y']. \end{aligned}$$

Hence it is enough to prove that

$$\# \mathcal{M}(x, \hat{\pi}(y') : H, J) = \sum_{y \in \hat{\pi}^{-1}(x)} \# \mathcal{M}(y, y' : \bar{H}, \bar{J})$$

for any $x \in \mathcal{P}_k(H)$, $y' \in \mathcal{P}_{k-1}(\bar{H})$. But it is clear since

$$\bigsqcup_{y \in \hat{\pi}^{-1}(x)} \mathcal{M}(y, y' : \bar{H}, \bar{J}) \rightarrow \mathcal{M}(x, \hat{\pi}(y') : H, J); \quad [u] \mapsto [\hat{\pi} \circ u]$$

is a bijection. Therefore we have proved that ψ_H is a chain map. Hence we can define a morphism

$$\text{HF}_*(H : X, \lambda) \rightarrow \text{HF}_*(\bar{H} : Y, \pi^* \lambda).$$

We denote this morphism also by ψ_H . Let $H, H' \in \mathcal{H}_{\text{ad}}(X, \lambda)$. If $a_H \leq a_{H'}$,

$$(5) \quad \begin{array}{ccc} \text{HF}_*(H) & \xrightarrow{\psi_H} & \text{HF}_*(\bar{H}) \\ \downarrow & & \downarrow \\ \text{HF}_*(H') & \xrightarrow{\psi_{H'}} & \text{HF}_*(\bar{H}') \end{array}$$

commutes, where vertical morphisms are monotone morphisms.

To prove $\text{cap}_S(Y, \pi^*\lambda) \leq \text{cap}_S(X, \lambda)$, it is enough to show that if $a \notin \tau(\partial X, \lambda)$ satisfies $a > \text{cap}_S(X, \lambda)$, then $a > \text{cap}_S(Y, \pi^*\lambda)$. $a > \text{cap}_S(X, \lambda)$ implies that $\text{SH}_n^{<\delta}(X, \lambda) \rightarrow \text{SH}_n^{<a}(X, \lambda)$ vanishes for any $0 < \delta < \delta(\partial X, \lambda)$. Take $H_\pm \in \mathcal{H}_{\text{ad}}(X, \lambda)$ such that $a_{H_-} = \delta$, $a_{H_+} = a$. Then, by lemma 2.8, $\text{HF}_n(H_-) \rightarrow \text{HF}_n(H_+)$ vanishes.

In the rest of this proof, we assume that X and Y are connected (general case follows at once from this case). Take H_- so that it satisfies following conditions:

- H_- is time independent.
- $H_-(z, r) = \delta r + \text{const}$ for $(z, r) \in \partial X \times [1, \infty)$.
- $H_-|_X$ is sufficiently small in C^2 .

Then, $\mathcal{P}(H_-)$ and $\mathcal{P}(\bar{H}_-)$ consist only of constant maps to $\text{Crit}(H_-)$, $\text{Crit}(\bar{H}_-)$. In particular, $C_k(\bar{H}_-) = C_k(H_-) = 0$ for $k \geq n + 1$. Hence $\psi_{H_-} : \text{HF}_n(H_-) \rightarrow \text{HF}_n(\bar{H}_-)$ is injective, therefore isomorphism (since X and Y are connected).

Hence, the commutative diagram (5) implies that $\text{HF}_n(\bar{H}_-) \rightarrow \text{HF}_n(\bar{H}_+)$ vanishes. Again by lemma 2.8, $\text{SH}_n^{<\delta}(Y, \pi^*\lambda) \rightarrow \text{SH}_n^{<a}(Y, \pi^*\lambda)$ vanishes. Hence $a > \text{cap}_S(Y, \pi^*\lambda)$. This completes the proof of theorem 3.6. \square

4. CAPACITY OF RIEMANNIAN MANIFOLDS

In this section, we introduce the notion of capacity for Riemannian manifolds without boundaries, which is denoted by cap_R . The main result in this section is theorem 4.13, which includes property (A) which we have stated in the introduction. In 4.1, we give the definition of cap_R , and prove its basic properties. In particular, proposition 4.6, which is an easy consequence of theorem 3.6, is important. In 4.2, we prove that when N is a compact connected Riemannian manifold with non-empty boundary, then $\text{cap}_R(\text{int}N) < \infty$ (theorem 4.7). In 4.3, first we prove that $\mathbb{R}^n \setminus \mathbb{Z}^n$ with the flat metric has a finite capacity (theorem 4.12). This is proved by combining proposition 4.6 and theorem 4.7. Theorem 4.13 is obtained by theorem 4.12 and elementary geometric arguments.

4.1. The definition and basic properties. First we introduce some notations. Let N be a n -dimensional Riemannian manifold. Let us denote the natural projection $T^*N \rightarrow N$; $(q, p) \mapsto q$ by π_N . We define $\lambda_N \in \Omega^1(T^*N)$ by

$$\lambda_N(v) := p(d\pi_N(v)) \quad (q \in N, p \in T_q^*N, v \in T_{(q,p)}(T^*N)).$$

Then, $\omega_N := d\lambda_N$ is a symplectic form on T^*N . Define $\nu_N \in \mathcal{X}(T^*N)$ by $i_{\nu_N}\omega_N = \lambda_N$.

For $V \in C^\infty(N)$, define $H_V \in C^\infty(T^*N)$ by

$$H_V(q, p) = V(q) + |p|^2/2,$$

and denote $\{H_V \leq 0\} \subset T^*N$ by D_V .

$\mathcal{V}(N)$ denotes the set of $V \in C^\infty(N)$ such that 0 is a regular value of V , and $\{V \leq 0\} \subset N$ is compact.

For $\xi \in \mathcal{X}(N)$, define $F_\xi \in C^\infty(T^*N)$ and $\tilde{\xi} \in \mathcal{X}(T^*N)$ by $F_\xi(q, p) := p(\xi_q)$ and $\tilde{\xi} := X_{F_\xi}$. Then, $L_{\tilde{\xi}}\omega_N = 0$ and $\tilde{\xi}_{(q,0)} = \xi_q$.

Lemma 4.1. *Let N be a Riemannian manifold without boundary. Then, for any $V \in \mathcal{V}(N)$, there exists $\lambda \in \Omega^1(T^*N)$ such that $d\lambda = \omega_N$ and (D_V, λ) is a Liouville domain. If $V, V' \in \mathcal{V}(N)$ satisfy $V > V'$, then $\text{cap}_S(D_V, \omega_N) \leq \text{cap}_S(D_{V'}, \omega_N)$.*

Remark 4.2. For any Liouville domain (X, λ) , $\text{cap}(X, \lambda)$ depends only on $d\lambda$ (lemma 3.3, (3)). Hence formulas $\text{cap}_S(D_V, \omega_N)$ and $\text{cap}_S(D_{V'}, \omega_N)$ make sense.

Proof. We prove the first assertion. Take $\xi \in \mathcal{X}(N)$ such that $dV(\xi) > 0$ on $\{V = 0\}$. For $\delta > 0$, define $Z_\delta \in \mathcal{X}(T^*N)$ by $Z_\delta := \nu_N + \delta\tilde{\xi}$. Then, $L_{Z_\delta}\omega_N = \omega_N$ for any δ . Moreover, $dH_V(Z_\delta) > 0$ on $H_V^{-1}(0)$ for sufficiently small $\delta > 0$. Hence $\lambda_\delta := i_{Z_\delta}\omega_N$ satisfies $d\lambda_\delta = \omega_N$ and (D_V, λ_δ) is a Liouville domain for sufficiently small $\delta > 0$.

We prove the second assertion. If $V > V'$, $\{V = 0\} \cap \{V' = 0\} = \emptyset$. Hence there exists $\xi \in \mathcal{X}(N)$ such that $dV(\xi) > 0$ on $\{V = 0\}$ and $dV'(\xi) > 0$ on $\{V' = 0\}$. Then, for sufficiently small $\delta > 0$, (D_V, λ_δ) and $(D_{V'}, \lambda_\delta)$ are both Liouville domains. On the other hand, $D_V \subset D_{V'}$. Hence by theorem 3.4, $\text{cap}_S(D_V, \omega_N) \leq \text{cap}_S(D_{V'}, \omega_N)$. \square

We define the notion of capacity for Riemannian manifolds without boundary.

Definition 4.3. Let N be a Riemannian manifold without boundary. Then, capacity of N is defined by

$$\text{cap}_R(N) := \sup\{\text{cap}_S(D_V, \omega_N) \mid V \in \mathcal{V}(N), V > -1/2\}.$$

Remark 4.4. As is clear from the above definition, when N is a compact Riemannian manifold without boundary $\text{cap}_R(N) = \text{cap}_S(DT^*N, \omega_N)$, where $DT^*N := \{(q, p) \in T^*N \mid |p| \leq 1\}$.

In the following, we sometimes denote N by (N, g) , where g is the Riemannian metric on N . We also sometimes denote H_V and D_V by $H_{V, g}$, $D_{V, g}$.

Lemma 4.5. *Let (N, g) be a Riemannian manifold without boundary.*

- (1) *For any open set $\Omega \subset N$, $\text{cap}_R(\Omega, g) \leq \text{cap}_R(N, g)$.*
- (2) *$\text{cap}_R(N, g) = \sup\{\text{cap}_R(\Omega, g) \mid \Omega \subset N \text{ is a open set such that } \bar{\Omega} \text{ is compact}\}$.*
- (3) *Let a be a positive number. Then $\text{cap}_R(N, ag) = a \cdot \text{cap}_R(N, g)$, where ag is defined by $(ag)(v) := a \cdot g(v)$ ($v \in TN$).*
- (4) *Let g' be a Riemannian metric on N , and assume that $g \leq g'$ (which means that $g(v) \leq g'(v)$ for any $v \in TN$). Then $\text{cap}_R(N, g) \leq \text{cap}_R(N, g')$.*

Proof. (1) and (2) are clear from the definition. (3) follows from $D_{V, ag} = \{(q, ap) \mid (q, p) \in D_{V, g}\}$. (4) follows from $g \leq g' \implies D_{V, g} \subset D_{V, g'}$. \square

Proposition 4.6. *Let (N, g) be a Riemannian manifold without boundary, and $\pi : M \rightarrow N$ be a covering map such that $\deg \pi < \infty$. Then, $\text{cap}_R(M, \pi^*g) \leq \text{cap}_R(N, g)$.*

Proof. For any $V \in \mathcal{V}(M)$ such that $V > -1/2$, there exists $W \in \mathcal{V}(N)$ such that $W > -1/2$ and $V > W \circ \pi$. Hence

$$\text{cap}_S(D_V, \omega_M) \leq \text{cap}_S(D_{W \circ \pi}, \omega_M) \leq \text{cap}_S(D_W, \omega_N) \leq \text{cap}_R(N, g).$$

The first inequality follows from lemma 4.1, the second inequality follows from theorem 3.6, and the last inequality is clear from the definition of cap_R . Therefore $\text{cap}_R(M, \pi^*g) \leq \text{cap}_R(N, g)$. \square

4.2. Capacity of interiors of compact Riemannian manifolds with boundaries.

The goal of this subsection is to prove the following theorem:

Theorem 4.7. *Let N be a compact connected Riemannian manifold with non-empty boundary. Then, $\text{cap}_R(\text{int}N) < \infty$.*

At first, notice the following consequence of theorem 2.10:

Lemma 4.8. *Let N be a Riemannian manifold without boundary, and $V \in \mathcal{V}(N)$. Then, $\text{SH}_*(D_V, \omega_N)$ depends only on diffeomorphism type of $\{V \leq 0\}$.*

We prove the following lemma:

Lemma 4.9. *Let N be a Riemannian manifold without boundary, and $V \in \mathcal{V}(N)$. Assume that $\{V \leq 0\}$ is connected and $\{V = 0\} \neq \emptyset$. Then, $\text{SH}_*(D_V, \omega_N) = 0$. In particular, $\text{cap}_S(D_V, \omega_N) < \infty$.*

Proof. First note that the second assertion follows from the first assertion by lemma 3.3-(2). We prove the first assertion. By lemma 4.8, for any $W \in \mathcal{V}(N)$ such that $\{W \leq 0\} = \{V \leq 0\}$, $\text{SH}_*(D_V, \omega_N) \cong \text{SH}_*(D_W, \omega_N)$. Since $\{V \leq 0\}$ is a compact connected manifold with non-empty boundary, we can take W so that it is a Morse function and $\{P_1, \dots, P_m\} := \text{Crit}(W) \cap \{W \leq 0\}$ satisfies the following:

- $\min W = W(P_1) < W(P_2) < \dots < W(P_m) < 0$.
- $\text{ind}P_j \leq n - 1$ for all $1 \leq j \leq m$.
- $\text{ind}P_j = 0$ if and only if $j = 1$.

To complete the proof, we extend the definition of symplectic homology. Let (X, λ) be a Liouville domain and H be a Hamiltonian on its completion. Then, let $\text{HF}_*^{\text{all}}(H)$ be the homology of $(C_*^{\text{all}}(H), \partial)$, where $C_*^{\text{all}}(H)$ is a \mathbb{Z}_2 -graded free \mathbb{Z}_2 module generated by all (not only contractible) periodic orbits of X_H . We define $\text{SH}_*^{\text{all}}(X, \lambda) := \varinjlim_H \text{HF}_*^{\text{all}}(H)$.

Obviously, $\text{SH}_*^{\text{all}}(X, \lambda) = 0 \implies \text{SH}_*(X, \lambda) = 0$.

In the following, we prove that $\text{SH}_*^{\text{all}}(D_W, \omega_N) = 0$. For $a \in \mathbb{R}$, abbreviate $\{H_W \leq a\} \subset T^*N$ by $D_{\leq a}$. Then, for any $a \in (W(P_1), W(P_2))$,

$$\text{SH}_*(D_{\leq a}, \omega_N) \cong \text{SH}_*(B^{2n}(1), \omega_n) = 0.$$

The first isomorphism follows from lemma 4.8, and the second equality follows from theorem 2.11. Since $D_{\leq a}$ is simply connected, $\text{SH}_*^{\text{all}}(D_{\leq a}, \omega_N) = 0$.

Hence it is enough to show that for any $j \in \{2, \dots, m\}$, $a \in (W(P_{j-1}), W(P_j))$ and $b \in (W(P_j), W(P_{j+1}))$, the isomorphism $\text{SH}_*^{\text{all}}(D_{\leq a}, \omega_N) \cong \text{SH}_*^{\text{all}}(D_{\leq b}, \omega_N)$ holds. By lemma 4.8, it is enough to show that there exists $\varepsilon > 0$ such that $\text{SH}_*^{\text{all}}(D_{\leq W(P_j) - \varepsilon}, \omega_N) \cong \text{SH}_*^{\text{all}}(D_{\leq W(P_j) + \varepsilon}, \omega_N)$.

Let $j \in \{2, \dots, m\}$ and set $k := \text{ind}P_j$. Take a local coordinate (q_1, \dots, q_n) around P_j such that P_j corresponds to $(0, \dots, 0)$ and

$$W(q_1, \dots, q_n) = W(P_j) - \sum_{1 \leq i \leq k} q_i^2 + \sum_{k+1 \leq i \leq n} q_i^2.$$

Take $\varepsilon > 0$ sufficiently small, and let

$$\Sigma_\varepsilon := \{(q_1, \dots, q_k, 0, \dots, 0) \in N \mid q_1^2 + \dots + q_k^2 = \varepsilon\}.$$

For $\delta > 0$, let $Z_\delta := \nu_N + \delta \widetilde{\nabla W}$, and $\lambda_\delta := i_{Z_\delta} \omega_N$. Then, $(D_{\leq W(P_j) \pm \varepsilon}, \lambda_\delta)$ are Liouville domains for sufficiently small $\delta > 0$, and Σ_ε is an isotropic submanifold of $(\partial D_{\leq W(P_j) - \varepsilon}, \lambda_\delta)$. Moreover, $(D_{\leq W(P_j) + \varepsilon}, \lambda_\delta)$ is isotopic as Liouville domain (see definition 2.1) to the Liouville domain obtained by attaching k -handle to $(D_{\leq W(P_j) - \varepsilon}, \lambda_\delta)$ along Σ_ε in the sense of [9]. Hence by theorem 1.11 (1) in [3],

$$\text{SH}_*^{\text{all}}(D_{\leq W(P_j) - \varepsilon}, \omega_N) \cong \text{SH}_*^{\text{all}}(D_{\leq W(P_j) + \varepsilon}, \omega_N).$$

This completes the proof. \square

Remark 4.10. The above proof shows that (D_W, ω_N) carries a structure of so called "subcritical Weinstein domain".

Finally, we prove theorem 4.7.

Proof. For any Riemannian metrics g and g' on N , there exists $a > 0$ such that $g \leq ag'$ since N is compact. Then, $\text{cap}_R(\text{int}N, g) \leq a \cdot \text{cap}_R(\text{int}N, g')$. Therefore it is enough to show that there exists a Riemannian metric g on N such that $\text{cap}_R(\text{int}N, g) < \infty$.

Take a Riemannian manifold (N', g') without boundary, and an embedding $i : N \hookrightarrow N'$. We show that $\text{cap}_R(\text{int}N, i^*g') < \infty$.

Since N is a compact connected manifold with non-empty boundary, there exists $V \in \mathcal{V}(N')$ such that $V \circ i < -1/2$ and $\{V \leq 0\}$ is connected, $\{V = 0\} \neq \emptyset$. For any $W \in \mathcal{V}(\text{int}N)$ such that $W > -1/2$, $\text{cap}_S(D_W, \omega_N) \leq \text{cap}_S(D_V, \omega_{N'})$. Hence $\text{cap}_R(\text{int}N) \leq \text{cap}_S(D_V, \omega_{N'})$. On the other hand, $\text{cap}_S(D_V, \omega_{N'}) < \infty$ by lemma 4.9. This completes the proof. \square

The following corollary of theorem 4.7 plays an important role in the next subsection.

Corollary 4.11. *Let N be a compact connected Riemannian manifold (possibly with boundary), and $x \in \text{int}N$. Then, $\text{cap}_R(\text{int}N \setminus \{x\}) < \infty$.*

Proof. Let $n := \dim N$. When $n = 1$, the assertion is easily confirmed. Hence in the following, we consider the case $n \geq 2$. It is enough to show that there exists a Riemannian metric g on N such that $\text{cap}_R(\text{int}N \setminus \{x\}, g) < \infty$.

Let U be a coordinate neighborhood containing x , and (q_1, \dots, q_n) be a local coordinate on U , such that x corresponds to $(0, \dots, 0)$. We may assume that $B(x, 1) \subset U$ and $dg = \sum_{1 \leq j \leq n} dq_j^2$ on U .

Set $S := \partial B(x, 1) \subset U$, and $g_S := g|_S$. Take arbitrary smooth function $\mu : [0, 1] \rightarrow \mathbb{R}_{>0}$ such that $\mu \equiv 1/2$ on $[0, 1/3]$ and $\mu(r) = r$ on $[2/3, 1]$, and $\mu(r) \geq r$. Consider a cylinder $C = S \times [0, 1]$ equipped with a metric h defined by

$$|v + a\partial_r(z, r)|_h := (|v(z)|_{g_S}^2 \mu(r)^2 + a^2)^{1/2} \quad (v \in TS, a \in \mathbb{R}).$$

Then,

$$I : (S \times (2/3, 1], h) \rightarrow (B(x, 1) \setminus B(x, 2/3), g); \quad (z, r) \mapsto zr$$

is an isometry.

Let (\tilde{N}, \tilde{g}) be a Riemannian manifold which is obtained by pasting $(N \setminus B(x, 2/3), g)$ with (C, h) via I . Then, \tilde{N} is a compact manifold with non-empty boundary (since $S \times \{0\} \subset \partial\tilde{N}$), and connected (since N is connected and $n \geq 2$). Hence theorem 4.7 implies that $\text{cap}_R(\text{int}\tilde{N}, \tilde{g}) < \infty$.

Define a diffeomorphism $J : \text{int}\tilde{N} \rightarrow \text{int}N \setminus \{x\}$ by

$$J(y) := \begin{cases} y & (y \in N \setminus B(x, 2/3)) \\ rz & (y = (z, r) \in C = S \times (0, 1]) \end{cases}.$$

Then, since $\mu(r) \geq r$, $J^*g \leq \tilde{g}$. Hence

$$\text{cap}_R(\text{int}N \setminus \{x\}, g) = \text{cap}_R(\text{int}\tilde{N}, J^*g) \leq \text{cap}_R(\text{int}\tilde{N}, \tilde{g}) < \infty.$$

□

4.3. Capacity of domains in \mathbb{R}^n .

Theorem 4.12. *Let g_n denote the flat metric on \mathbb{R}^n . Then, $\text{cap}_R(\mathbb{R}^n \setminus \mathbb{Z}^n, g_n) < \infty$.*

Proof. When $n = 1$, $\text{cap}_R(\mathbb{R} \setminus \mathbb{Z}, g_1) = \text{cap}_R((0, 1), g_1) < \infty$. In the following, we assume that $n \geq 2$.

By lemma 4.5-(2), it is enough to show that, for any bounded open set Ω in $\mathbb{R}^n \setminus \mathbb{Z}^n$, $\text{cap}_R(\Omega)$ is bounded from above by some constant which depends only on n .

Let Ω be a bounded open set in $\mathbb{R}^n \setminus \mathbb{Z}^n$. Then, for sufficiently large integer l , $\Omega \subset (-l, l)^n$. Hence Ω can be considered as an open set in $(\mathbb{R}^n \setminus \mathbb{Z}^n)/2l\mathbb{Z}^n$.

Consider the natural covering map of degree $(2l)^n$: $(\mathbb{R}^n \setminus \mathbb{Z}^n)/2l\mathbb{Z}^n \rightarrow (\mathbb{R}^n \setminus \mathbb{Z}^n)/\mathbb{Z}^n$. Then

$$\text{cap}_R(\Omega) \leq \text{cap}_R((\mathbb{R}^n \setminus \mathbb{Z}^n)/2l\mathbb{Z}^n) \leq \text{cap}_R((\mathbb{R}^n \setminus \mathbb{Z}^n)/\mathbb{Z}^n).$$

The first inequality follows from lemma 4.5-(1), and the second inequality follows from proposition 4.6. On the other hand, $(\mathbb{R}^n \setminus \mathbb{Z}^n)/\mathbb{Z}^n$ is $\mathbb{R}^n/\mathbb{Z}^n$ minus a point. Hence, by corollary 4.11, $\text{cap}_R((\mathbb{R}^n \setminus \mathbb{Z}^n)/\mathbb{Z}^n) < \infty$. This completes the proof. □

Theorem 4.13. *For each integer n , there exists $c_0(n), c_1(n) > 0$ such that for any non-empty open set Ω in \mathbb{R}^n ,*

$$c_0(n) \leq \frac{\text{cap}_R(\Omega, g_n)}{r(\Omega)} \leq c_1(n).$$

Proof. If $r < r(\Omega)$, then there exists $x \in \mathbb{R}^n$ such that $B(x, r) \subset \Omega$. Hence, by lemma 4.5-(1) and (3),

$$\text{cap}_R(\Omega) \geq \text{cap}_R(B(x, r)) = r \cdot \text{cap}_R(B(x, 1)).$$

Hence $\text{cap}_R(B^n(1)) \leq \frac{\text{cap}_R(\Omega)}{r(\Omega)}$ for any $\Omega \neq \emptyset$.

Next we bound $\frac{\text{cap}_R(\Omega)}{r(\Omega)}$ from above. Take an arbitrary positive number r so that $r > r(\Omega)$. Then, for any $x \in \mathbb{R}^n$, $B(x, r) \setminus \Omega \neq \emptyset$. For any $j = (j_1, \dots, j_n) \in \mathbb{Z}^n$, take an arbitrary point p_j on $B(4rj, r) \setminus \Omega$, where $4rj = (4rj_1, \dots, 4rj_n)$. Then, $\text{cap}_R(\Omega) \leq \text{cap}_R(\mathbb{R}^n \setminus \{p_j\}_{j \in \mathbb{Z}^n})$.

Take sufficiently large $\alpha_n > 0$ so that for any $x \in B^n(1)$, there exists a diffeomorphism φ on $B^n(2)$ with compact support such that $\varphi(x) = (0, \dots, 0)$, and $g_n \leq \alpha_n \cdot \varphi^* g_n$. Then, since $B(4ri, 2r) \cap B(4rj, 2r) = \emptyset$ when $i \neq j$, there exists a diffeomorphism $\psi : \mathbb{R}^n \setminus \{p_j\}_{j \in \mathbb{Z}^n} \rightarrow \mathbb{R}^n \setminus 4r\mathbb{Z}^n$ such that $g_n \leq \alpha_n \cdot \psi^* g_n$. Then,

$$\begin{aligned} \text{cap}_R(\mathbb{R}^n \setminus \{p_j\}_{j \in \mathbb{Z}^n}, g_n) &\leq \alpha_n \cdot \text{cap}_R(\mathbb{R}^n \setminus \{p_j\}_{j \in \mathbb{Z}^n}, \psi^* g_n) \\ &= \alpha_n \cdot \text{cap}_R(\mathbb{R}^n \setminus 4r\mathbb{Z}^n, g_n) = 4\alpha_n r \cdot \text{cap}_R(\mathbb{R}^n \setminus \mathbb{Z}^n, g_n). \end{aligned}$$

To sum up,

$$r > r(\Omega) \implies \text{cap}_R(\Omega) \leq 4\alpha_n r \cdot \text{cap}_R(\mathbb{R}^n \setminus \mathbb{Z}^n).$$

Hence $\frac{\text{cap}_R(\Omega)}{r(\Omega)} \leq 4\alpha_n \cdot \text{cap}_R(\mathbb{R}^n \setminus \mathbb{Z}^n) < \infty$. □

5. SHORT PERIODIC BILLIARD TRAJECTORY

The goal of this section is to prove the following theorem.

Theorem 5.1. *Let Ω be a bounded domain in \mathbb{R}^n with smooth boundary. Then, there exists a periodic billiard trajectory on Ω with at most $n + 1$ bounce times and length equal to $\text{cap}_R(\Omega)$.*

Theorem 5.1 is exactly the same as property (B) of cap_R which we have introduced in the introduction. Hence, as we have explained in the introduction, it completes the proof of our main theorem 1.1.

We start to prove theorem 5.1. The proof heavily relies on the arguments in [1]. First we recall the settings in [1]. Fix $d_0 \in (0, 1/2)$ so small that $\text{dist}_{\partial\Omega} : q \mapsto \min\{|q - q'| \mid q' \in \partial\Omega\}$ is smooth on $\{\text{dist}_{\partial\Omega} \leq 2d_0\}$. Let $k : [0, \infty) \rightarrow [0, 2d_0]$ be a smooth function such that $0 \leq k' \leq 1$, $k(x) = x$ if $x \leq d_0$ and k is constant on $[2d_0, \infty)$. Then, we define a function $h \in C^\infty(\bar{\Omega})$ by $h(q) := k(\text{dist}_{\partial\Omega}(q))$, and define $U \in C^\infty(\Omega)$ by $U(q) := h^{-2}(q)$. Then, U is a positive function on Ω which grows like $(\text{dist}_{\partial\Omega})^{-2}$ near $\partial\Omega$ and is constant on the region $\{\text{dist}_{\partial\Omega} \geq 2d_0\}$.

For each $\varepsilon > 0$, consider the modified Lagrangian

$$L_\varepsilon : T\Omega \rightarrow \mathbb{R}; (q, v) \mapsto |v|^2/2 - \varepsilon U(q).$$

For each energy value $E \in \mathbb{R}$ the free-time action functional $\mathcal{L}_\varepsilon^E : L^{1,2}(\mathbb{R}/\mathbb{Z}, \Omega) \times \mathbb{R}_{>0} \rightarrow \mathbb{R}$ is given by

$$\mathcal{L}_\varepsilon^E(\Gamma, \tau) := \tau \int_0^1 \left[L_\varepsilon(\Gamma(t), \tau^{-1} \partial_t \Gamma(t)) + E \right] dt.$$

For $(\Gamma, \tau) \in L^{1,2}(\mathbb{R}/\mathbb{Z}, \Omega) \times \mathbb{R}_{>0}$, let γ be the corresponding τ -periodic curve, i.e. $\gamma : \mathbb{R}/\tau\mathbb{Z} \rightarrow \Omega; t \mapsto \Gamma(t/\tau)$. Then, straightforward calculations show that (Γ, τ) is a critical point of $\mathcal{L}_\varepsilon^E$ if and only if γ satisfies

$$\partial_t^2 \gamma + \nabla(\varepsilon U)(\gamma) = 0$$

with energy

$$E_\varepsilon(\gamma) := |\partial_t \gamma|^2/2 + \varepsilon U(\gamma) = E.$$

When (Γ, τ) is a critical point of $\mathcal{L}_\varepsilon^E$, $\mu_{\text{Morse}}(\Gamma : \mathcal{L}_\varepsilon^E|_{L^{1,2}(\mathbb{R}/\mathbb{Z}) \times \{\tau\}})$ denotes the number of negative eigenvalues of the Hessian of $\mathcal{L}_\varepsilon^E|_{L^{1,2}(\mathbb{R}/\mathbb{Z}) \times \{\tau\}}$ at (Γ, τ) .

For $\varepsilon > 0$, define $H_\varepsilon : T^*\Omega \rightarrow \mathbb{R}$ by $H_\varepsilon(q, p) := \varepsilon U(q) + |p|^2/2$, and $D_\varepsilon := \{H_\varepsilon \leq 1/2\} \subset T^*\Omega$.

Lemma 5.2. *For any $\varepsilon > 0$, there exists $(\Gamma_\varepsilon, \tau_\varepsilon)$, a critical point of $\mathcal{L}_\varepsilon^{1/2}$, which satisfies the following properties:*

- (1) $\int_{\mathbb{R}/\tau_\varepsilon\mathbb{Z}} |\partial_t \gamma_\varepsilon|^2 dt = \text{cap}_S(D_\varepsilon, \omega_\Omega)$.
- (2) $\mu_{\text{Morse}}(\Gamma_\varepsilon; \mathcal{L}_\varepsilon^{1/2}|_{L^{1,2}(\mathbb{R}/\mathbb{Z}) \times \{\tau_\varepsilon\}}) \leq n + 1$.

Proof. Take arbitrary $\lambda \in \Omega^1(T^*\Omega)$ such that $d\lambda = \omega_\Omega$ and (D_ε, λ) is a Liouville domain. By corollary 3.12, there exists $x_\varepsilon : \mathbb{R}/\tau_\varepsilon\mathbb{Z} \rightarrow H_\varepsilon^{-1}(1/2)$, which is a periodic orbit of X_{H_ε} and satisfies

$$\int_{\mathbb{R}/\tau_\varepsilon\mathbb{Z}} x_\varepsilon^* \lambda = \text{cap}_S(D_\varepsilon, \omega_\Omega), \quad \mu_{\text{CZ}}(x_\varepsilon) \leq n + 1.$$

Define $\gamma_\varepsilon : \mathbb{R}/\tau_\varepsilon\mathbb{Z} \rightarrow \Omega$ and $\Gamma_\varepsilon : \mathbb{R}/\mathbb{Z} \rightarrow \Omega$ by $\gamma_\varepsilon := \pi_\Omega \circ x_\varepsilon$, $\Gamma_\varepsilon(t) := \gamma_\varepsilon(\tau_\varepsilon t)$. Then, it is obvious that $(\Gamma_\varepsilon, \tau_\varepsilon)$ is a critical point of $\mathcal{L}_\varepsilon^{1/2}$. Moreover,

$$\int_{\mathbb{R}/\tau_\varepsilon\mathbb{Z}} |\partial_t \gamma_\varepsilon|^2 dt = \int_{\mathbb{R}/\tau_\varepsilon\mathbb{Z}} x_\varepsilon^* \lambda_\Omega = \int_{\mathbb{R}/\tau_\varepsilon\mathbb{Z}} x_\varepsilon^* \lambda = \text{cap}_S(D_\varepsilon, \omega_\Omega).$$

In the second equality, we use that x_ε is contractible in $T^*\Omega$ and $d\lambda = d\lambda_\Omega$.

Finally,

$$\mu_{\text{Morse}}(\Gamma_\varepsilon; \mathcal{L}_\varepsilon^{1/2}|_{L^{1,2}(\mathbb{R}/\mathbb{Z}) \times \{\tau_\varepsilon\}}) = \mu_{\text{CZ}}(x_\varepsilon).$$

This identity follows from theorem 7.3.1 in [5]. (2) follows immediately from this identity and $\mu_{\text{CZ}}(x_\varepsilon) \leq n + 1$. \square

Lemma 5.3. *For each $\varepsilon > 0$, take $(\Gamma_\varepsilon, \tau_\varepsilon)$, a critical point of $\mathcal{L}_\varepsilon^{1/2}$ which satisfies properties in lemma 5.2. Then,*

$$0 < \liminf_{\varepsilon \rightarrow 0} \tau_\varepsilon \leq \limsup_{\varepsilon \rightarrow 0} \tau_\varepsilon < \infty.$$

Proof. First we show that $\liminf_{\varepsilon \rightarrow 0} \tau_\varepsilon > 0$. Assume that $\liminf_{\varepsilon \rightarrow 0} \tau_\varepsilon = 0$, i.e. there exists a sequence $(\varepsilon_k)_k$ such that $\varepsilon_k, \tau_{\varepsilon_k} \rightarrow 0$ as $k \rightarrow \infty$. Then, there exists a sequence of integers $(N_k)_k$ such that $1 < \tau_{\varepsilon_k} N_k < 2$. Set $\Theta_k \in L^{1,2}(\mathbb{R}/\mathbb{Z}, \Omega)$ by $\Theta_k(t) := \Gamma_{\varepsilon_k}(N_k t)$. Then, $(\Theta_k, \tau_{\varepsilon_k} N_k)$ is a critical point of $\mathcal{L}_{\varepsilon_k}^{1/2}$. By proposition 2.1 in [1], a certain subsequence of $(\Theta_k)_k$ converges in $L^{1,2}(\mathbb{R}/\mathbb{Z}, \bar{\Omega})$. On the other hand, by lemma 5.2-(1),

$$|\partial_t \Theta_k|_{L^2(\mathbb{R}/\mathbb{Z})}^2 = N_k^2 \int_{\mathbb{R}/\mathbb{Z}} |\partial_t \Gamma_{\varepsilon_k}(t)|^2 dt = N_k^2 \tau_{\varepsilon_k} \text{cap}_S(D_{\varepsilon_k}, \omega_\Omega).$$

Since $\text{cap}_S(D_{\varepsilon_k}, \omega_\Omega) \rightarrow \text{cap}_R(\Omega)$ as $k \rightarrow \infty$ and $N_k \tau_{\varepsilon_k} > 1$, the last term goes to ∞ as $k \rightarrow \infty$, contradicting that a certain subsequence of $(\Theta_k)_k$ converges in $L^{1,2}(\mathbb{R}/\mathbb{Z}, \bar{\Omega})$.

Next we show that $\limsup_{\varepsilon \rightarrow 0} \tau_\varepsilon < \infty$. For each $\varepsilon > 0$, define $x_\varepsilon : \mathbb{R}/\tau_\varepsilon \mathbb{Z} \rightarrow T^*\Omega$ by $x_\varepsilon = (\gamma_\varepsilon, \partial_t \gamma_\varepsilon)$. Then, x_ε is an integral curve of X_{H_ε} on $H_\varepsilon^{-1}(1/2)$. On the other hand, by proposition 3.2 in [1], when $\varepsilon > 0$ is sufficiently small, there exists $\lambda_\varepsilon \in \Omega^1(T^*\Omega)$ such that $d\lambda_\varepsilon = \omega_\Omega$ and the following inequality holds on $H_\varepsilon^{-1}(1/2)$:

$$\lambda_\varepsilon(X_{H_\varepsilon}) \geq \frac{(1/2 - 0)^3}{2[(1/2 - 0)^2 + 48(1/2 - 0)^2]} = \frac{1}{196}.$$

Notice that $(D_\varepsilon, \lambda_\varepsilon)$ is a Liouville domain, since setting $Z_\varepsilon \in \mathcal{X}(T^*\Omega)$ by $i_{Z_\varepsilon} \omega_\Omega = \lambda_\varepsilon$, then $dH_\varepsilon(Z_\varepsilon) = \omega_\Omega(Z_\varepsilon, X_{H_\varepsilon}) = \lambda_\varepsilon(X_{H_\varepsilon}) > 0$ on $H_\varepsilon^{-1}(1/2)$.

Since

$$\int_{\mathbb{R}/\tau_\varepsilon \mathbb{Z}} \lambda_\varepsilon(X_{H_\varepsilon}(x_\varepsilon(t))) dt = \int_{\mathbb{R}/\tau_\varepsilon \mathbb{Z}} x_\varepsilon^* \lambda_\varepsilon = \text{cap}_S(D_\varepsilon, \omega_\Omega),$$

$\tau_\varepsilon \leq 196 \cdot \text{cap}_S(D_\varepsilon, \omega_\Omega) \leq 196 \cdot \text{cap}_R(\Omega)$ for sufficiently small $\varepsilon > 0$. This completes the proof. \square

Finally, we prove theorem 5.1.

Proof. For each $\varepsilon > 0$, take $(\Gamma_\varepsilon, \tau_\varepsilon)$, a critical point of $\mathcal{L}_\varepsilon^{1/2}$ which satisfies properties in lemma 5.2. Then by lemma 5.3, we can apply proposition 2.1 in [1] to the sequence $(\Gamma_\varepsilon, \tau_\varepsilon)_{\varepsilon > 0}$. i.e. a certain subsequence of $(\Gamma_\varepsilon, \tau_\varepsilon)_{\varepsilon > 0}$ converges to (Γ, τ) in $L^{1,2}(\mathbb{R}/\mathbb{Z}, \bar{\Omega}) \times \mathbb{R}_{>0}$, and $\gamma : \mathbb{R}/\tau \mathbb{Z} \rightarrow \bar{\Omega}; t \mapsto \Gamma(t/\tau)$ is a periodic billiard trajectory on Ω , such that $E(\gamma) = \lim_{\varepsilon \rightarrow 0} E_\varepsilon(\gamma_\varepsilon) = 1/2$.

Since $\mu_{\text{Morse}}(\Gamma_\varepsilon; \mathcal{L}_\varepsilon^{1/2}|_{L^{1,2}(\mathbb{R}/\mathbb{Z}) \times \{\tau_\varepsilon\}}) \leq n+1$ for each $\varepsilon > 0$, proposition 2.2 in [1] implies that Γ has at most $n+1$ bounce times.

Finally we prove that $\tau = \text{cap}_R(\Omega)$. For each $\varepsilon > 0$, by lemma 5.2-(1),

$$\text{cap}_S(D_\varepsilon, \omega_\Omega) = \int_{\mathbb{R}/\tau_\varepsilon \mathbb{Z}} |\partial_t \gamma_\varepsilon|^2 dt = \tau_\varepsilon^{-1} \int_{\mathbb{R}/\mathbb{Z}} |\partial_t \Gamma_\varepsilon|^2 dt.$$

Since $\lim_{\varepsilon \rightarrow 0} \tau_\varepsilon = \tau$ and $\lim_{\varepsilon \rightarrow 0} \Gamma_\varepsilon \rightarrow \Gamma$ in $L^{1,2}(\mathbb{R}/\mathbb{Z}, \bar{\Omega})$, by taking limit of the above identity we get

$$\text{cap}_R(\Omega) = \lim_{\varepsilon \rightarrow 0} \text{cap}_S(D_\varepsilon, \omega_\Omega) = \tau^{-1} \int_{\mathbb{R}/\mathbb{Z}} |\partial_t \Gamma|^2 dt.$$

On the other hand, since $E(\gamma) = 1/2$, $|\partial_t \gamma| = 1$ for almost every $t \in \mathbb{R}/\tau\mathbb{Z}$. Hence $|\partial_t \Gamma| = \tau$ for almost every $t \in \mathbb{R}/\mathbb{Z}$. Therefore $\text{cap}_R(\Omega) = \tau^{-1} \cdot \tau^2 = \tau$. \square

Acknowledgements. The author would like to appreciate his advisor professor Kenji Fukaya for reading the manuscript and giving precious comments, and also the anonymous referee for many suggestions. The author is supported by Grant-in-Aid for JSPS fellows.

APPENDIX: PROOF OF THEOREM 2.12

Assume that (X, λ) , (X, λ') are Liouville domains such that $d\lambda = d\lambda'$.

Take arbitrary smooth function $\rho: \mathbb{R} \rightarrow [0, 1]$ such that for sufficiently large $s_0 > 0$ $\rho(s) = \begin{cases} 0 & (s \leq -s_0) \\ 1 & (s \geq s_0) \end{cases}$. Let $\lambda_s := (1 - \rho(s))\lambda + \rho(s)\lambda'$. Then, $(X, \lambda_s)_{s \in \mathbb{R}}$ is a smooth family of Liouville domains. Denote the completion of (X, λ_s) by $(\hat{X}_s, \hat{\lambda}_s)$.

Our aim is to define a morphism from the Floer chain complex on (X, λ) to the Floer chain complex on (X, λ') . To define a morphism, we study the Floer equation on a fiber bundle over \mathbb{R} , which is constructed as follows. First, consider trivial bundles

$$E_X : X \times \mathbb{R} \rightarrow \mathbb{R}, \quad E_{\partial X} : (\partial X \times (0, \infty)) \times \mathbb{R} \rightarrow \mathbb{R}.$$

Define an embedding

$$I : (\partial X \times (0, 1]) \times \mathbb{R} \rightarrow X \times \mathbb{R}; \quad ((z, r), s) \mapsto (I_s(z, r), s)$$

by (Z_s denotes the vector field on X characterized by $i_{Z_s} d\lambda_s = \lambda_s$):

$$\begin{aligned} I_s(z, 1) &= z & (z \in \partial X), \\ \partial_r I_s(z, r) &= r^{-1} Z_s(I_s(z, r)) & (z \in \partial X, r \in (0, 1]). \end{aligned}$$

Let $E := E_X \cup_I E_{\partial X}$. E is a fiber bundle over \mathbb{R} , and each fiber E_s is identified with \hat{X}_s . Note that there exist natural bundle maps over \mathbb{R} :

$$\begin{aligned} j_1 &: E_X \rightarrow E, \\ j_2 &: (\partial X \times [1, \infty)) \times \mathbb{R} \rightarrow E. \end{aligned}$$

To study the Floer equation on E , we equip E with a connection ∇ , and denote the horizontal lift of ∂_s to E by W . We take ∇ so that W satisfies

- $j_1^*(W) = (0, \partial_s)$,
- $j_2^*(W) = (0, \partial_s)$ outside $(\partial X \times [1, 2]) \times [-s_0, s_0]$.

Let $H_- \in \mathcal{H}_{\text{ad}}^{\text{rest}}(X, \lambda)$, $H_+ \in \mathcal{H}_{\text{ad}}^{\text{rest}}(X, \lambda')$ and $(H_s)_{s \in \mathbb{R}}$ be a family of Hamiltonians with the following properties:

- $H_s \in \mathcal{H}^{\text{rest}}(X, \lambda_s)$ for any $s \in \mathbb{R}$.
- $\partial_s H_{s,t}(x) \geq 0$ for any $(s, t) \in \mathbb{R} \times \mathbb{R}/\mathbb{Z}$ and $x \in X$.
- There exists $s_1 \geq s_0$ such that $H_s = \begin{cases} H_- & (s \leq -s_1) \\ H_+ & (s \geq s_1) \end{cases}$.

- $\partial_s a_{H_s} \geq 0$.

Let $(J_s)_s$ be a family of (time-dependent) almost complex structures on $E_s = \hat{X}_s$, such that $J_s \in \mathcal{J}(X, \lambda_s : 1)$ for any s , and

$$J_s = \begin{cases} J_{-s_0} & (s \leq -s_0) \\ J_{s_0} & (s \geq s_0) \end{cases}.$$

We denote $J_{\pm s_0}$ by J_{\pm} .

Then, for $x_- \in \mathcal{P}(H_-)$, $x_+ \in \mathcal{P}(H_+)$, we study the following Floer equation for $u : \mathbb{R} \times \mathbb{R}/\mathbb{Z} \rightarrow E$:

- $u(s, t) \in E_s$,
- $\nabla_s u - J_{s,t}(\partial_t u - X_{H_{s,t}} \circ u) = 0$,
- $u(s) \rightarrow x_{\pm}$ as $s \rightarrow \pm\infty$.

We denote the moduli space of solutions of the above Floer equation by $\hat{\mathcal{M}}(x, y : (H_s, J_s)_s)$.

In the following, we abbreviate a_{H_s} by $a(s)$. The key step in the proof of theorem 2.12 is to prove the following lemma:

Lemma 5.4. *There exist $c_0, c_1 > 0$, which depend only on $(J_s)_s$ and ρ , such that: if a satisfies $\partial_s a \geq c_0 a + c_1$ on $[-s_0, s_0]$, then for any $x_- \in \mathcal{P}(H_-)$, $x_+ \in \mathcal{P}(H_+)$ and $u \in \hat{\mathcal{M}}(x_-, x_+ : (H_s, J_s)_s)$, $u(\mathbb{R} \times \mathbb{R}/\mathbb{Z}) \subset j_1(X \times \mathbb{R})$.*

Proof. Step 1. First note that for any $x \in \mathcal{P}(H_-) \cup \mathcal{P}(H_+)$, $x(\mathbb{R}/\mathbb{Z}) \subset X$. This is because $H_{\pm} \in \mathcal{H}_{\text{ad}}^{\text{rest}}(X, \lambda)$. Our aim is to show that $u(\mathbb{R} \times \mathbb{R}/\mathbb{Z})$ is contained in $j_1(X \times \mathbb{R})$. If this is not true, for some $r_0 > 1$

$$D_{r_0} := u^{-1}\left(j_2(\partial X \times [r_0, \infty)) \times \mathbb{R}\right)$$

is a non-empty surface with boundary. Note that D_{r_0} must be compact, since both $x_-(\mathbb{R}/\mathbb{Z})$ and $x_+(\mathbb{R}/\mathbb{Z})$ are contained in X . Define $v : D_{r_0} \rightarrow \partial X \times [1, \infty)$ by

$$u(s, t) = j_2(v(s, t), s),$$

and define $z : D_{r_0} \rightarrow \partial X$ and $r : D_{r_0} \rightarrow [1, \infty)$ by $v(s, t) = (z(s, t), r(s, t))$.

We will prove that there exist $c_0, c_1 > 0$, which depend only on $(J_s)_s$ and ρ , such that if a satisfies $\partial_s a \geq c_0 a + c_1$ on $[-s_0, s_0]$, then $\Delta r \geq 0$. Since $r \equiv r_0$ on ∂D_{r_0} , this implies $r \leq r_0$ on D_{r_0} . On the other hand, by definition $r > r_0$ on $\text{int} D_{r_0}$, hence we get a contradiction.

Step 2. We calculate $\Delta r(s, t)$ for $(s, t) \in D_{r_0}$. Recall that u satisfies the Floer equation

$$\nabla_s u - J_{s,t}(\partial_t u - X_{H_{s,t}}(u)) = 0.$$

Since $H_{s,t}(z, r) = a(s)r + b(s)$ on $\partial X \times [1, \infty)$,

$$J_{s,t}X_{H_{s,t}} = -\nabla_{s,t}H_{s,t} = -ar\partial_r.$$

$\nabla_{s,t}H_{s,t}$ denotes the gradient of $H_{s,t}$ with respect to $\langle \cdot, \cdot \rangle_{J_{s,t}}$.

Moreover, by definition of W ,

$$\partial_s u = \nabla_s u + W(u).$$

Hence, the Floer equation can be written as:

$$(6) \quad \partial_s u - W(u) - J_{s,t} \partial_t u - ar \partial_r = 0.$$

It is convenient to define $\hat{\lambda} \in \Omega^1(E)$ by

- $\hat{\lambda}|_{E_s} = \hat{\lambda}_s$ for any $s \in \mathbb{R}$.
- $\hat{\lambda}(W) \equiv 0$.

Then, by applying dr and $\hat{\lambda}$ to (6) respectively, we get

$$\begin{aligned} \partial_s r + \hat{\lambda}(\partial_t u) - ar - dr(W(u)) &= 0, \\ \hat{\lambda}(\partial_s u) - \partial_t r &= 0. \end{aligned}$$

Then,

$$\begin{aligned} \Delta r &= \partial_t(\hat{\lambda}(\partial_s u)) - \partial_s(\hat{\lambda}(\partial_t u)) + \partial_s(ar) + \partial_s(dr(W(u))) \\ &= d\hat{\lambda}(\partial_t u, \partial_s u) + \partial_s(ar) + \partial_s(dr(W(u))) \\ &= |\nabla_s u|_{J_{s,t}}^2 + d\hat{\lambda}(\partial_t u, W(u)) + \partial_s a \cdot r + a \cdot dr(W(u)) + \partial_s(dr(W(u))). \end{aligned}$$

In the following, we abbreviate $|\cdot|_{J_{s,t}}$ by $|\cdot|_{s,t}$.

Step 3. We prove that $\Delta r(s, t) \geq 0$ when $s \notin [-s_0, s_0]$. Assume that $s \notin [-s_0, s_0]$. Then, since $j_2^* W = (0, \partial_s)$ on $(\partial X \times [1, \infty)) \times (\mathbb{R} \setminus [-s_0, s_0])$,

$$dr(W(u(s, t))) = 0.$$

Moreover, since $\partial_s \lambda_s = 0$ for $s \notin [-s_0, s_0]$,

$$i_{W(u(s, t))} d\hat{\lambda} = 0.$$

Hence

$$\Delta r(s, t) = |\nabla_s u(s, t)|_{s,t}^2 + \partial_s a(s) \cdot r(s, t) \geq 0.$$

Step 4. Next we prove the following: there exist $c_2, c_3 > 0$ which depend only on $(J_s)_s$ and ρ , such that if a satisfies $\partial_s a \geq c_2 a + c_3$ on $[-s_0, s_0]$, then $r(s, t) \leq 2$ for any $(s, t) \in \mathbb{R} \times \mathbb{R}/\mathbb{Z}$.

Assume that $r(s, t) > 2$ for some $(s, t) \in \mathbb{R} \times \mathbb{R}/\mathbb{Z}$. Then, since $j_2^* W = (0, \partial_s)$ on $(\partial X \times [2, \infty)) \times \mathbb{R}$,

$$dr(W(u(s, t))) = 0.$$

Hence

$$\Delta r(s, t) = |\nabla_s u|_{s,t}^2 + d\hat{\lambda}(\partial_t u, W(u)) + \partial_s a \cdot r.$$

Since $j_2^* \hat{\lambda} = r \lambda_s$,

$$i_{\partial_s}(j_2^* d\hat{\lambda}) = i_{\partial_s}(dr \wedge \lambda_s + r(d\lambda_s + ds \wedge \partial_s \lambda_s)) = r \partial_s \lambda_s.$$

Hence

$$-d\hat{\lambda}(\partial_t u, W(u)) = i_{\partial_s}(j_2^* d\hat{\lambda})(\partial_t v) = r \partial_s \lambda_s(\partial_t v) = r \partial_s \lambda_s(\partial_t z).$$

Therefore, there exists $c_4 > 0$ which depends only on $(J_s)_s$ and ρ such that

$$|d\hat{\lambda}(\partial_t u, W(u))| = r \cdot |\partial_s \lambda_s(\partial_t z)| \leq c_4 r^{1/2} |\partial_t u|_{s,t}.$$

Moreover, since u satisfies the Floer equation $\nabla_s u - J_{s,t} \partial_t u - ar \partial_r = 0$,

$$|\partial_t u|_{s,t} \leq |\nabla_s u|_{s,t} + |ar \partial_r|_{s,t} = |\nabla_s u|_{s,t} + ar^{1/2}.$$

Therefore

$$\begin{aligned} \Delta r &\geq |\nabla_s u|_{s,t}^2 + \partial_s a \cdot r - c_4 \cdot r^{1/2} (|\nabla_s u|_{s,t} + ar^{1/2}) \\ &\geq |\nabla_s u|_{s,t}^2 + \partial_s a \cdot r - (|\nabla_s u|_{s,t}^2 + c_4^2 r) / 2 - c_4 ar \\ &\geq (\partial_s a - c_4 a - c_4^2 / 2) r. \end{aligned}$$

Hence setting $c_2 := c_4$, $c_3 := c_4^2 / 2$, the following holds:

Assume $\partial_s a \geq c_2 a + c_3$ on $[-s_0, s_0]$. Then, $\Delta r(s, t) \geq 0$ if $s \in [-s_0, s_0]$ and $r(s, t) > 2$.

On the other hand, by Step 3, $\Delta r(s, t) \geq 0$ if $s \notin [-s_0, s_0]$. Hence if $\partial_s a \geq c_2 a + c_3$ on $[-s_0, s_0]$, $\Delta r(s, t) \geq 0$ on $\{r > 2\} \subset \mathbb{R} \times \mathbb{R}/\mathbb{Z}$. This implies that $\{r > 2\} = \emptyset$, by same arguments as step 1.

Step 5. Finally we prove that there exist $c_0, c_1 > 0$, which depend only on $(J_s)_s$ and ρ , such that if a satisfies $\partial_s a \geq c_0 a + c_1$ on $[-s_0, s_0]$, then $\Delta r \geq 0$.

It is convenient to equip E with a Riemannian metric $\langle \cdot, \cdot \rangle_t$ such that

- $\langle \cdot, \cdot \rangle_t|_{E_s}$ is equal to $\langle \cdot, \cdot \rangle_{s,t}$.
- For any $q \in E_s$, $T_q E_s$ and W_q is orthogonal with respect to $\langle \cdot, \cdot \rangle_t$.
- For any $q \in E$, $|W_q|_t = 1$.

By step 4, $r(s, t) \leq 2$ for any $(s, t) \in \mathbb{R} \times \mathbb{R}/\mathbb{Z}$. Therefore there exist $c_5, c_6, c_7 > 0$ such that

$$\begin{aligned} |d\hat{\lambda}(\partial_t u, W(u))| &\leq c_5 |\partial_t u|_t, \\ |dr(W(u))| &\leq c_6, \\ |\partial_s(dr(W(u)))| &\leq c_7 |\partial_s u|_t \end{aligned}$$

on $[-s_0, s_0] \times \mathbb{R}/\mathbb{Z}$. On the other hand,

$$\begin{aligned} |\partial_t u|_t &\leq |\nabla_s u|_t + |ar \partial_r|_t = |\nabla_s u|_t + ar^{1/2} \leq |\nabla_s u|_t + a\sqrt{2}, \\ |\partial_s u|_t &\leq |W(u)|_t + |\nabla_s u|_t = 1 + |\nabla_s u|_t. \end{aligned}$$

Hence,

$$\begin{aligned} \Delta r &\geq |\nabla_s u|_t^2 + \partial_s a \cdot r - c_5 (|\nabla_s u|_t + a\sqrt{2}) - c_6 a - c_7 (1 + |\nabla_s u|_t) \\ &= |\nabla_s u|_t^2 + \partial_s a \cdot r - (\sqrt{2}c_5 + c_6)a - (c_5 + c_7)|\nabla_s u|_t - c_7. \end{aligned}$$

Therefore, setting $c_0 := \sqrt{2}c_5 + c_6$, $c_1 := c_7 + (c_5 + c_7)^2 / 2$, $\Delta r \geq \partial_s a - c_0 a - c_1$ (we use $\partial_s a \geq 0$ and $r \geq 1$). Hence setting c_0 and c_1 as above, the following holds:

If $\partial_s a \geq c_0 a + c_1$ on $[-s_0, s_0]$, then $\Delta r \geq 0$ on $[-s_0, s_0] \times \mathbb{R}/\mathbb{Z}$.

On the other hand, $\Delta r(s, t) \geq 0$ when $s \notin [-s_0, s_0]$, as we have proved in step 3. This completes the proof of step 5. \square

Corollary 5.5. *Let $c_0, c_1 > 0$ satisfy the condition in lemma 5.4, and assume that $\partial_s a \geq c_0 a + c_1$ on $[-s_0, s_0]$. Then, for any $x_- \in \mathcal{P}(H_-)$, $x_+ \in \mathcal{P}(H_+)$ and $u \in \hat{\mathcal{M}}(x_-, x_+ : (H_s, J_s)_s)$,*

$$\partial_s \mathcal{A}_{H_s}(u(s) : X, \lambda_s) \leq 0.$$

In particular, if $\mathcal{A}_{H_+}(x_+) > \mathcal{A}_{H_-}(x_-)$, then $\hat{\mathcal{M}}(x_-, x_+ : (H_s, J_s)_s) = \emptyset$.

Proof. By lemma 5.4, $u(\mathbb{R} \times \mathbb{R}/\mathbb{Z}) \subset j_1(X \times \mathbb{R})$. Since $j_1^* W = (0, \partial_s)$,

$$\partial_s \mathcal{A}_{H_s}(u(s) : X, \lambda_s) = \int_{\mathbb{R}/\mathbb{Z}} u(s)^* \partial_s \lambda_s - \int_{\mathbb{R}/\mathbb{Z}} \{|\nabla_s u(s, t)|^2 + \partial_s H_{s,t}(u(s, t))\} dt.$$

Since $u(s) : \mathbb{R}/\mathbb{Z} \rightarrow X$ is contractible, one can extend $u(s)$ to $\overline{u(s)} : D^2 \rightarrow X$ so that $\overline{u(s)}(e^{2\pi i \theta}) = u(s)(\theta)$. Then,

$$\int_{\mathbb{R}/\mathbb{Z}} u(s)^* \partial_s \lambda_s = \int_{D^2} \overline{u(s)}^* \partial_s (d\lambda^s) = 0.$$

Hence

$$\partial_s \mathcal{A}_{H_s}(u(s) : X, \lambda_s) = - \int_{\mathbb{R}/\mathbb{Z}} \{|\nabla_s u(s, t)|^2 + \partial_s H_{s,t}(u(s, t))\} dt \leq 0,$$

where the last inequality follows from $\partial_s H_{s,t} \geq 0$ on X . \square

Finally we prove theorem 2.12. Define a morphism $\psi : C_*(H_-, \partial_{H_-, J_-}) \rightarrow C_*(H_+, \partial_{H_+, J_+})$ by

$$\psi([x_-]) = \sum_{x_+ \in \mathcal{P}_k(H_+)} \# \hat{\mathcal{M}}(x_-, x_+ : (H_s, J_s)_s) \cdot [x_+] \quad (x_- \in \mathcal{P}_k(H_-)).$$

Then, by lemma 5.4, ψ is a chain map. Moreover, by corollary 5.5, ψ defines a chain map $\psi^{<a} : C_*^{<a}(H_-, \partial_{H_-, J_-}) \rightarrow C_*^{<a}(H_+, \partial_{H_+, J_+})$ for any $a \in (0, \infty]$. This defines a morphism $\text{HF}_*^{<a}(H_-) \rightarrow \text{HF}_*^{<a}(H_+)$, and we denote this morphism also by $\psi^{<a}$. It is clear from the construction that

$$\begin{array}{ccc} \text{HF}_*^{<a}(H_- : X, \lambda) & \xrightarrow{\psi^{<a}} & \text{HF}_*^{<a}(H_+ : X, \lambda') \\ \downarrow & & \downarrow \\ \text{HF}_*^{<b}(H_- : X, \lambda) & \xrightarrow{\psi^{<b}} & \text{HF}_*^{<b}(H_+ : X, \lambda') \end{array}$$

commutes for any $a \leq b$.

By taking a limit, we get a morphism $\text{SH}_*^{<a}(X, \lambda) \rightarrow \text{SH}_*^{<a}(X, \lambda')$ (still denoted by $\psi^{<a}$), and the following diagram commutes:

$$\begin{array}{ccc} \text{SH}_*^{<a}(X, \lambda) & \xrightarrow{\psi^{<a}} & \text{SH}_*^{<a}(X, \lambda') \\ \downarrow & & \downarrow \\ \text{SH}_*^{<b}(X, \lambda) & \xrightarrow{\psi^{<b}} & \text{SH}_*^{<b}(X, \lambda') \end{array}$$

It is easy to check that $\psi^{<a}$ does not depend on choices of ρ and $(J^s)_s$, $\psi^{<a} : \text{SH}^{<a}(X, \lambda) \rightarrow \text{SH}^{<a}(X, \lambda)$ is the identity, and the following diagram commutes:

$$\begin{array}{ccc}
 \text{SH}_*^{<a}(X, \lambda) & \xrightarrow{\psi^{<a}} & \text{SH}_*^{<a}(X, \lambda') \\
 & \searrow \psi^{<a} & \swarrow \psi^{<a} \\
 & \text{SH}_*^{<a}(X, \lambda'') &
 \end{array}$$

Then, it follows that $\psi^{<a}$ is isomorphic. Hence this completes the proof of theorem 2.12.

REFERENCES

- [1] Albers, P., Mazzucchelli, M.: Periodic bounce orbits of prescribed energy, Int. Math. Res. Notices, doi:10.1093/imrn/rnq193 (2010)
- [2] Benci, V., Giannoni, F.: Periodic bounce trajectories with a low number of bounce points, Ann. Inst. Henri Poincaré, Anal. Non Linéaire, 6, No.1, 73–93 (1989)
- [3] Cieliebak, K.: Handle attaching in symplectic homology and the Chord Conjecture, J. Eur. Math. Soc, 4, 115–142 (2002)
- [4] Cieliebak, K., Floer, A., Hofer, H., Wysocki, K.: Applications of symplectic homology II: Stability of the action spectrum, Math. Z, 223, 27–45 (1996)
- [5] Long, Y.: Index theory for symplectic paths with applications, Progr. Math, vol.207, Birkhäuser, Basel (2002)
- [6] Oancea, A.: A survey of Floer homology for manifolds with contact type boundary or symplectic homology, Ensaios Math, 7, 51–91 (2004)
- [7] Viterbo, C.: Functors and computations in Floer homology I, Geom. Funct. Anal, 9, 985–1033 (1999)
- [8] Viterbo, C.: Metric and isoperimetric problems in symplectic geometry, J. Amer. Math. Soc, 13, No.2, 411–431 (2000)
- [9] Weinstein, A.: Contact surgery and symplectic handlebodies, Hokk. Math. J, 20, 241–251 (1991)

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN

E-mail address: iriek@math.kyoto-u.ac.jp