

Title	A CASE OF ECONOMICAL DAMAGE DONE BY NEBALIA
Author(s)	Nishimura, Saburo; Hamabe, Mototsugu
Citation	PUBLICATIONS OF THE SETO MARINE BIOLOGICAL LABORATORY (1964), 12(2): 173-175
Issue Date	1964-10-15
URL	http://hdl.handle.net/2433/175356
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

A CASE OF ECONOMICAL DAMAGE DONE BY *NEBALIA*¹⁾

SABURO NISHIMURA

Seto Marine Biological Laboratory, Sirahama

and

MOTOTSUGU HAMABE

Kasumi Branch of the Japan Sea Regional Fisheries
Research Laboratory, Kasumi, Hyogo-Ken

With 2 Text-figures

It is generally known that typical nebalians, i.e. members of the genera *Nebalia* and *Nebaliella*, are filter-feeders upon detritus; they live on the muddy bottom and feed, it is supposed, on small organic particles suspended just above the mud or take coarser particles in by filtrating them rapidly before they settle as the animals burrow or kick up the mud (ROWETT 1943). Recently the authors happened to learn an interesting case of feeding habit of this peculiar crustacean, which has been, so far as the present authors are concerned, never recorded in the Japanese waters and seems to suggest that the animal may affect to some extent the natural economy in the sea under special circumstances.

In the early summer of 1957, the junior author was engaged in gathering specimens of the sharp-nosed shark, *Scoliodon walbeehmi* (BLEEKER), caught by long line in the waters around the Oki Islands in the southern Japan Sea for the purpose of investigating their stomach contents; and then his attention was called on the frequent occurrences of not a few individuals more or less heavily damaged among the fished sharks. These damaged sharks showed a common symptom that the muscles had variously been scraped off and lost so that the skin had been detached for the most part of the body from the underlying muscle layers, leaving spacious hollows between the skin and the skeleton. And in these hollows were found numerous little crustaceans attaching to or encroaching, it was supposed, into muscles. The frequency of occurrence and the degree of damage done by this little animal varied with circumstances. When the long lines were kept sunk for two or three days longer on


1) Contributions from the Seto Marine Biological Laboratory, No. 418.

account of rough weather, then more damaged sharks were met with and the degree of damage was more intensive. In an extreme instance, all musculature and other soft parts of the shark body were almost completely robbed off and there were left only skin and skeleton.


The offensive behavior of this crustacean upon the fishes caught by fishing gears was first reported by the junior author in a short article in Japanese with the title "On the uso" (HAMABE 1958). *Uso* is the local name for this little crustacean talked among the fishermen of the Oki Islands. In this article appeared in a publication of limited circulation, the junior author presented a sketch of the crustacean, which is here reproduced in Text-fig. 1, but erroneously referred the animal to a species of amphipods. The senior author, examining some specimens sent to him from the junior author, confirmed that the crustacean was a species of the genus *Nebalia* LEACH, 1814.

連 絡 ニ ュ ー ス 第 85 号 (4)

妙な魚体があることに気付いた。二時化のため、二三日揚縄が遅れると、時間の長短によりいろいろであるが、内部の肉質を掠奪され、骨と肉が遊離して、甚だしいのは、五月の鯉のほりのようになつて上つてくるものがあることである。体内には漁夫の俗稱する「ワソ」という小甲殻類が無数に寄生している。蚊の蛹のようなこの「ワソ」を動物園監で調べてみると、どうも筋足動物、甲殻類、海脚目で、「リシアナタクタス科」のものは死魚を好み、網にかかった魚体に附着したまま引上げられること稀ならず」とある点からみて、多分リシアナタクタス科の一種であろうと考えられる。全長一五・一〇ミリメートルの小虫で、簡単にスケッチすると図



死体を採集する直接的漁業「底曳」がないため、浦郷では、間接的に他の大型の最終捕食魚の胃からスルメイカを抽出するという方法をとるはかないのであるが、イカや魚の死体が、冷塩水漬のような具合に、深海底に、ワソをはじめ、ものもの動物から食害されることなく転がっていき、問題はないであろうが、自然はしかく簡単に禁断の果実を味わわしてくれぬ。(浦郷支所 浜部)



Text-fig. 1. The sketch of "uso" given by the junior author in his 1958 paper, together with part of the explanation.

Text-fig. 2. *Nebalia* sp. taken from an affected sharp-nosed shark. x2.5

Dissecting the nebalians (Text-fig. 2) collected from a damaged sharp-nosed shark, the senior author noted that, though some individuals had ingested bottom deposits containing many siliceous particles imbedded in debris matrix, others had the intestine packed with a matter referable to masticated animal tissues together with a rich oily substance. It seems that certain nebalians had actually been feeding on the muscles and other soft parts, such as liver,

of the shark*, while the specimens containing only bottom deposits in the intestine but no trace of animal tissues might be the individuals arrived at that shark just before it was taken out of the water.

A part of the specimens of *Nebalia* were sent to Dr. Masao IWASA of the Seikei University, Tokyo, for specific identification.

The fact that nebalians swarm to rotten animal substance such as dead crabs or shrimps on the sea bottom was already noticed by ROBINSON (1906; cited by KOMAI 1917), but yet there has been no report mentioning that the animals attack and devour the fish caught by fishing gears as certain species of necrophagous isopods and amphipods do. The latter crustaceans have been well known among Japanese coastal fishermen as loathsome pests to their catch. It may be surprising to see that in certain localities at least nebalians are distributed so densely as to swarm enormously to and devour such large-sized dead animals as sharks till these are finally disintegrated in a rather short time. In these localities, the role of nebalians as scavengers in the sea must be re-evaluated.

Before closing this short note, the authors express their sincere gratitude to Dr. Takasi TOKIOKA of the Seto Marine Biological Laboratory who kindly read the manuscript.

REFERENCES

- HAMABE, M. 1958. [On the *uso*]. Nihonkai-ku Suisan Shiken Kenkyū Renraku Nyūsu, No. 85, pp. 3-4. (In Japanese.)
KOMAI, T. 1917. [Nebaliacea]. Dobutsugaku Zasshi, Vol. 29, pp. 297-298. (In Japanese.)
ROWETT, H. G. Q. 1943. The gut of Nebaliacea. Discovery Repts., Vol. 23, pp. 3-17.

* The sharp-nosed shark contains an exceptionally large amount of oil in the liver, and this shark was once fished in a great scale around the Oki Islands for the purpose of gathering liver oil and extracting vitamin A from it.