

Title	A New Species of the Commensal Crab Genus <i>Aphanodactylus</i> (Crustacea : Brachyura : Pinnotheridae) from the Yaeyama Islands, Southern Japan
Author(s)	Konishi, Kooichi; Noda, Hirokuni
Citation	PUBLICATIONS OF THE SETO MARINE BIOLOGICAL LABORATORY (1999), 38(5-6): 223-229
Issue Date	1999-09-30
URL	http://hdl.handle.net/2433/176286
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

**A New Species of the Commensal Crab Genus *Aphanodactylus*
(Crustacea: Brachyura: Pinnotheridae) from the Yaeyama
Islands, Southern Japan**

KOICHI KONISHI¹⁾ & HIROKUNI NODA²⁾

¹⁾National Research Institute of Aquaculture, Nansei, Mie, 516-0193 Japan

²⁾Department of Biology, Tokyo Women's Medical University, Shinjuku,
Tokyo, 162-8666 Japan

Abstract A new commensal crab, *Aphanodactylus loimiae*, is described from the Yaeyama Islands, Okinawa Prefecture, Japan. The present species is the third species of this rare pinnotherid genus. It differs from the two previously known species in carapace shape, number of segments of antennal flagellum, and number of merus teeth on walking legs. The single specimen was found in the tube of a sessile polychaete, *Loimia ingens* Grube. A dichotomous key and a diagrammatic key to the subfamilies of the Pinnotheridae and the genera of the Asthenognathinae in Japan are given.

Key words: Crustacea, Pinnotheridae, *Aphanodactylus*, Polychaeta, *Loimia*, key, new species, taxonomy

Since the description of *Pinnotheres pholadis* in the "Fauna Japonica" by de Haan (1835), the pinnotherid crab fauna of Japan has come to include 31 species in 13 genera (Sakai, 1976; Miyake, 1983; Konishi, 1996). Most of the work, however, has been focused on the subfamily Pinnotherinae, with relatively little attention directed toward other subfamilies, e.g., the Asthenognathinae. At present, the Asthenognathinae consists of seven genera in the world, two of which, *Asthenognathus* and *Tritodynamia*, have been known from Japan. The genus *Aphanodactylus* was proposed by Tesch (1918), with the description of a new species, *A. sibogae*, from Sumbawa Island, Indonesia. Another species, *A. edmondsoni* was described by Rathbun (1932) from the Hawaiian Islands. No other records of *Aphanodactylus* crabs have been published. A single specimen of a small pinnotherid crab, representing the genus *Aphanodactylus*, but differing morphologically from the two previously known species, was collected from a polychaete tube near the Yaeyama Marine Park Research Station in the Yaeyama Islands, southwest of Okinawa, Japan.

This paper gives a description of this new species of *Aphanodactylus* and a diagrammatic key to the subfamilies of the Pinnotheridae and the genera of the Asthenognathinae in Japan. The specimen has been deposited in the National Science Museum, Tokyo, under the accession number NSMT-Cr 12484.

***Aphanodactylus loimiae* sp. nov.**

[Japanese name: Koyubi-pinno]

(Figs. 1, 2)

Material examined: Holotype (NSMT-Cr 12484) ovigerous female with carapace length (CL) 7.9 mm and width (CW) 13.9 mm, from a tube of the terebellid polychaete *Loimia ingens* Grube, 1878, collected by F. Iwase, June 28, 1988. Type locality: ca. 3 m depth off Kuroshima Island, the Yaeyama Islands, Japan, 24°14'N, 123°59'E.

Etymology: The specific epithet was formed from the generic name of the host terebellid




Fig. 1. *Aphnodactylus loimiae* sp. nov., holotype ovigerous female (NSMT-Cr 12484). a: whole animal in dorsal view, setae of appendages only shown on right side, b: antenna, c: maxilliped 3, d: left cheliped, e: merus of walking leg 3, f: walking leg 4 with detail of seta. Scale bar = 1 mm.




Fig. 2. *Aphanodactylus loimiae* sp. nov., holotype ovigerous female (NSMT-Cr 12484). Photographs in dorsal (a), ventral (b), and frontal view (c). Scale bar = 10 mm.

polychaete *Loimia*.

Description: Carapace (Figs. 1a, 2a) slightly subquadrate, transversely elongated, about 1.76 in CW:CL ratio. Anterior border of carapace almost straight, but slightly deflexed in middle portion. Surface of carapace smooth. Eye-stalks small, oval in frontal view. Antennule folded into a groove. Antenna (Fig. 1b) minute, peduncle four-segmented, and flagellum consisting of seven segments with a long seta and two small setae on distal segment. Maxilliped 3 (Fig. 1c) oriented longitudinally (Fig. 2c), ischium distinctly longer than merus, and palp segments jointed end to end; inner side of merus and ischium fringed with long, plumose setae; exopod with two-segmented flagellum with long distal setae. Cheliped (Fig. 1d) smooth, but carrying a number of setae on its inner margin and on edges of movable and immovable fingers; length of palm about 1.5 times that of fingers. Meri of walking legs 1-3 almost equal in length and with 4-6 teeth on posterior border (Fig. 1e), while those of walking legs 4 with a well developed tooth and two vestigial ones (Fig. 1f). Dactyli of walking legs remarkably tiny and almost invisible, as suggested by the generic name (Gk. $\alpha\phi\alpha\nu\eta\varsigma$ = invisible + dactylus): shorter than 1/3 of propodus. Each leg with rows of plumose setae along the edges. Abdomen (Fig. 2b) consisting of seven segments. Body colour ivory-white in living specimen.

Remarks

The CW:CL ratio of the present specimen (=1.76) is different from those of *A. sibogae* (=1.88) and *A. edmondsoni* (=1.69). The present species differs from *A. sibogae* in lacking two pairs of surface pits on the carapace and in the large number of segments of the antennal flagellum: 2-3 and 7 segments in *A. sibogae* and *A. loimiae*, respectively. Another congener, *A. edmondsoni*, is distinguished by the outline of carapace and the posterior teeth on its walking legs 1-3: a large triangular spine instead of 5-6 spines in the present species. Edmondson (1962) stated that the walking leg 4 of *A. edmondsoni* was armed with

Table 1. Diagnostic characters of three species of *Aphanodactylus*

Characters	<i>A. loimiae</i> This study	<i>A. sibogae</i> Tesch (1918)	<i>A. edmondsoni</i> Rathbum (1932)
Carapace:			
width (mm)	13.9	11.25	16.2
length (mm)	7.9	6	9.6
width/length	1.76	1.88	1.69
anterior border	straight	straight	slightly convex
pits on dorsal surface	absent	present	absent
Antenna:			
flagellar segments	7	2-3	10
Walking legs:			
merus teeth (1st-3rd)	4-6	some teeth	1
merus teeth (4th)	1+2v	?	2
Locality:	Kuroshima Is. [Japan]	Sumbawa Is. [Indonesia]	Oahu Is. [Hawaii]
Host:	<i>Loimia ingens</i>	<i>Loimia</i> sp.*	<i>Loimia medusa</i>

v: vestigial

*Possibly *L. ingens*, judging from Caullery (1944)

two small teeth on lower border of the merus, while there are prominent tooth and two vestigial ones in *A. loimiae*. The diagnostic characters of the three species of *Aphanodactylus* are summarized in Table 1.

Other than the present genus, close similarity has also been noted to a species of the pinnothereinid genus *Pinnixa*. Serène (1964) commented on the similarities found in *Aphanodactylus* and *Pinnixa brevipes* H. Milne Edwards, 1853 and stated, "Cependant *brevipes* a les pattes ambulatoires beaucoup plus longues comparativement à celles de *A. sibogae* et c'est une espèce différente qui justifie une nouvelle description avec figure d'après le type." Detailed information on the type specimen of *P. brevipes* is required before any conclusions can be drawn.

The present crab was found in a tube of the sessile polychaete *Loimia* just as *A. sibogae* was. According to Edmondson (1946), *A. edmondsoni* lives in the shelly tube of a large terebellid worm, and Schmitt *et al.* (1973) suggested "*Loimia medusa* (Savigny)" for this worm which is commonly found around Hawaii. No other host animals have been known for *Aphanodactylus* (Table 1). It is possible that the large tubes of *Loimia* provide a good refuge for commensal crabs with a smooth and transversely elongated carapace. In fact, the pinnotherinid species *Sakaina japonica* Serène, 1964, which has similar external morphology to species of *Aphanodactylus*, lives within the tubes of *L. medusa* (Yamaguchi *et al.*, 1976; Konishi, 1981). Caullery (1944) noted that commensal anomuran crabs of the genus *Polyonyx* were commonly found in the tubes of *L. ingens*.

Aphanodactylus loimiae is the third record and species of its genus in the world. The subfamily Asthenognathinae of Japan now consists of three genera. Up-dated keys to the pinnotherid subfamilies and the asthenognathinid genera of Japan are given below, and Fig. 3 reproduces the same information diagrammatically.

Key to the subfamilies of the Pinnotheridae in Japan.

- F1a. Orbits transverse in the usual position, invisible in dorsal view..... F2
 F1b. Orbits and eye-stalks parallel longitudinally, wholly visible in dorsal view
 Subfamily Xenophthalminae
 F2a. Ischium and merus of maxilliped 3 separated..... F3
 F2b. Ischium and merus of maxilliped 3 fused to a single piece
 Subfamily Pinnotherinae
 F3a. Ischium of maxilliped 3 smaller than merus, palp often larger than ischium-merus
 Subfamily Pinnotherelinae
 F3b. Ischium of maxilliped 3 larger than merus, palp smaller than ischium-merus
 Subfamily Asthenognathinae

Key to the genera of the Asthenognathinae in Japan

- G1a. Merus of maxilliped 3 slightly shorter than ischium, palp segments jointed end to end
 G2
 G1b. Merus of maxilliped 3 as long as ischium, dactylus of palp jointed to inner border
 of propod *Tritodynamia*
 G2a. Dactylus of walking leg 4 not considerably reduced, longer than 1/2 of propodus;
 carapace roughly hexagonal *Asthenognathus*
 G2b. Dactylus of walking leg 4 considerably reduced in size, shorter than 1/3 of propodus;
 carapace subquadrate *Aphanodactylus*




Fig 3. Diagrammatic key to pinnotherid subfamilies and asthenognathinid genera of Japan. Abbreviations on the various branch segments correspond to the headers of the key in the text. Diagnostic parts are indicated by hatching. The figures of *Xenophthalmus*, *Tritodynamia*, and *Asihenognathus* are redrawn from Sakai, 1976.

Acknowledgements

We express our sincere gratitude to Mr. F. Iwase of Stella-Chemifa Corporation for collecting the material and Dr. H. Uchida of the Sabiura Marine Park Research Station for identifying the polychaete worm. Dr. H. Takahashi Prof. Emer. of Hokkaido University kindly provided us a literature. Our cordial thanks also due to Dr. M.J. Grygier of the Lake Biwa Museum for critical reading of the manuscript and Dr. M. Takeda of the National Science Museum for his encouragement during the study.

References

- Caullery, M. 1944. Polychètes sédentaires de l'expédition du Siboga. Ariciidae, Spionidae, Chaetopteridae, Chlorhaemidae, Opheliidae, Oweniidae, Sabellariidae, Sternaspidae, Amphipotentidae, Ampharetidae, Terebellidae. Siboga-Expeditie, 24i, 204 pp.
- Edmondson, C.H. 1946. Reef and shore fauna of Hawaii. Occasional Papers of the Bernice Pauahi Bishop Museum, Special Publication 22: 1-381.
- Edmondson, C.H. 1962. Hawaiian Crustacea: Goneplacidae, Pinnotheridae, Cymopoliidae, Ocypodidae and Gecarcinidae. Occasional Papers of the Bernice Pauahi Bishop Museum 23: 1-23.
- De Haan, W. 1835. Crustacea. In: F. von Siebold, Fauna Japonica. 244 pp. 55 pls.
- Miyake, S. 1983. Japanese crustacean decapods and stomatopods in color. Vol. II. Brachyura. Hoikusha, Osaka. 261 pp. 56 pls. [In Japanese]
- Konishi, K. 1981. A description of laboratory-reared larvae of the pinnotherid crab *Sakaina japonica* Serène (Decapoda, Brachyura). Journal of Faculty of Science, Hokkaido University, Series VI, Zoology 22: 165-176.
- Konishi, K. 1996. Some topics on pea crabs of Japan. Cancer, 5: 15-21. [In Japanese]
- Rathbun, M.J. 1932. A new pinnotherid crab from the Hawaiian Islands. Journal of the Washington Academy of Sciences 22: 181-182.
- Sakai, T. 1976. Crabs of Japan and adjacent seas. xxix+773 pp. (English part), 461 pp. (Japanese part), 16 pp. +251 pls. Kodansha, Tokyo.
- Schmitt, W.L., McCain, J.C., and Davidson, E.S. 1973. Crustaceorum catalogus. Pars 3. Decapoda I. Brachyura I. Fam. Pinnotheridae. Dr. W.Junk B.V., 160 pp.
- Serène, R. 1964. Papers from Dr. Th. Mortensen's Pacific Expedition 1914-1916 80. Goneplacidae et Pinnotheridae. Videnskabelige Meddelelser fra Dansk Naturhistorisk Forening København 126: 181-282.
- Tesch, J.J. 1918. The Decapoda Brachyura of the Siboga Expedition. II. Goneplacidae and Pinnotheridae. Siboga-Expeditie, 39c, 147 pp., pls. 7-18.
- Yamaguchi, T., Takeda, M., and Tokudome, K. 1976. A list of crabs collected in the vicinity of the Aitsu Marine Biological Station and a preliminary report on the cheliped asymmetry of the crabs. Calanus, 5: 31-46. [In Japanese]