

Title	Revision of Pelagic Ostracods of the Subfamily Archiconchoecinae
Author(s)	Chavtur, V. G.; Stovbun, G. G.
Citation	PUBLICATIONS OF THE SETO MARINE BIOLOGICAL LABORATORY (2003), 39(4-6): 139-219
Issue Date	2003-03-31
URL	http://hdl.handle.net/2433/176312
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

Rivision of Pelagic Ostracods of the Subfamily Archiconchoecinae

V. G. CHAVTUR and G. G. STOVBUN

Laboratory of Planktonology, Institute of Marine Biology,
Far East Branch, Russian Academy of Science,
Vladivostok 690041, Russia

Abstract Pelagic ostracods of the subfamily Archiconchoecinae have been revised. Their single genus *Archiconchoecia* Müller, 1894 has been divided into seven genera and two subgenera: *Archiconchoecia* (*Archiconchoecia*) Müller, 1894, *Archiconchoecia* (*Archiconchoecidia*) subgen. nov., *Archiconchoecilla* gen. nov., *Archiconchoecissa* gen. nov., *Archiconchoecemma* gen. nov., *Archiconchoecinna* gen. nov., *Archiconchoecetta* gen. nov. and *Archiconchoecerra* gen. nov. Six new genera, a new subgenus and seven new species have been described in detail. A redescription was made for the additional known taxa, *Archiconchoecilla maculata* (Chavtur, 1977) and *Archiconchoecemma orientalis* (Chavtur, 1987). A detailed synonymy and data on distribution are given for all species, and for some species morphological comments also have been added. Keys were elaborated for all taxa. The distribution of ostracods of the subfamily Archiconchoecinae has been discussed.

Key words: Pelagic ostracods, Halocyprididae, Archiconchoecinae, Taxonomy

Introduction

Ostracods of the subfamily Archiconchoecinae are being studied for a little more than a century. The first species of this group was found in the surface waters of the Naples Bay, Mediterranean Sea, in late 19th century. Müller (1894) included it into a new genus *Archiconchoecia* and described as *A. striata*. In the early 20th century he described two more species belonging to this genus, *A. ventricosa* (0°-2°N and 3°E-6°W, in a vertical tow from around 3000 m) and *A. cuneata* (19°S and 20°W, from 800-0 m), from a deep-water zone in the Atlantic Ocean (Müller, 1906, 1908). In the same years, from the area of the British Isles, a new bathypelagic species, *A. cucullata*, was recorded by Brady (1902). The following communication on ostracods of this genus did not appear in the literature for 60 years, until Poulsen (1969) published a monograph, and gave this group of ostracods the rank of subfamily (Archiconchoecinae). Later, Deevey (1978 c) described a part of specimens described as *A. ventricosa* (22-28°S and 166-177°E, layer 50-1500 m) as a new species *A. poulseni*.

In the period of 1970-80's, active research were done in Archiconchoecinae. For example, Chavtur (1977a, 1987) described novel species (*A. maculata* and *A. orientalis*) for areas of the Kurile-Kamchatka and Japan Trenches. Six new species (*A. bimucronata*, *A. bifurcata*, *A. gastroides*, *A. falcata*, *A. pilosa*, *A. longiseta*) of this subfamily in various forms were found by Deevey (1978c) in deep-water samples from the Sargasso Sea at 32°N, 64°W (1000-2000 m), and two from the Caribbean Sea, i.e. *A. fabiformis* (11°N and 66°W, in a tow from 1200-0 m) and *A. bispicula* (17°N and 85°W, in a tow from 2000-0 m). She also described another three species from the South Pacific (Deevey, 1978a) and the Pacific Sector of the Southern Ocean (Deevey, 1982); i.e. *A. versicula* (39°S, 107°W; between 1000-2000 m), *A. similis* (65°-71°S, 160°-180°E; between 1015-1829 m) and *A. bidens* (71°S, 179°E; between 1400-1600 m).

Recently, Angel (1983) found *A. sp. nov. 1* and *A. sp. nov. 2* in the northeastern Atlantic at depths 1900-2100 and 2100-2300 m, respectively. Yet, the description and figures of these species were not given, and 20 species are thus known in this subfamily currently.

Material and Methods

In this work, collections of Russian scientific expeditions to the North-West Pacific (R/V "Vityaz" in 1954, 1966, 1969; SRTM - 662 in 1959; SRTR "Orlyk" in 1969 and R/V "Cavalerovo" in 1980), North-East Pacific (R/V "Vityaz" in 1969), South-West Pacific (R/V "Vityaz" in 1957), North Atlantic (R/V "Lomonosov" in 1959) and Central Arctic (Drifting Station "North Pole - 2" in 1951) were used. Ten species of the subfamily Archiconchoecinae were found, of which seven were new to science. The collection data (coordinates and depth) of each species are given in the section "Systematics" of the description of species.

A detailed study of our collection of ostracods and critical analysis of the literature sources have revealed high morphological heterogeneity of this group and we assigned six genera, one subgenus, and 32 species (Table 1).

Main taxonomic characters used are as follows: shape and sculpturing of the shell; location of the asymmetrical glands on the valves; structure and shape of the frontal organ, and shape and sizes of the claspers on the 2nd antenna; shape of the teeth on the mandible basal endite; shape, structure and sizes of copulatory appendage; number of setae on the dorsal side of the 1st antenna, on the 2nd segment of the endopodite of the 2nd antenna ("c"- "e" setae in male), and on the exopodite of the 6th limb; number of claws (claw-like setae) on the 3rd and 4th segments of the exopodite of the 6th limb and on the furca.

Followings were considered secondary important characters: sizes of the shell, limbs and its segments, setae and claws; armament of the setae; number of the setae on the epipodial appendages of the 5th and 6th limbs; correlation between length and height of the copulatory appendage.

The form of the teeth on the mandibular basal endite enabled us to divide the entire ostracod assembly into two main groups (they may correspond to a rank of tribe). In Group I these teeth were triangular, while in Group II they were serrated (saw-like). In the former, the right asymmetrical gland was at the topmost part of the posterior valve margin, whereas in the latter, in its middle (seldomly a little higher) or bottom part. Group I was very diverse, and can be divided into several subgroups. One (*Archiconchoecissa* gen. nov.) was definitely marked by a strong rostrum, developed posterodorsal spine (on right valve) and a distinct (as tile) sculpture and also by having two dorsal setae on the 1st antenna, setae "c"- "e" with very thick base on the 2nd antenna (in male), and 2-3 setae on the 2nd segment of the exopodite of the 6th limb. Part of the remaining species could be easily united into another subgroup (*Archiconchoecia*) by the form of the shell (valves were rounded with dorsomedial indentation) and by the presence of a very thick (in the middle part) clasper on the 2nd antenna. The third subgroup (*Archiconchoecilla* gen. nov.) was isolated mainly on the basis of the following characters: the posterior margin of the shell was asymmetrical rounded (without spines) on the right and left valves; protopodite was longer than exopodite on the 2nd antenna; exopodite of the mandible was represented by one long seta, and its endopodite had usual dorsal seta. The rest species of Group I were mainly characterized by a short oval shell, the presence of 2-3 long setae on the mandibular exopodite and very short and fragile slender 2nd (from the dorsal margin) seta on the 3rd segment of its endopodite (*Archiconchoecemma* gen. nov.).

Group II were also morphologically not uniform. The species of this group have both distinctly and slightly separated serrated (saw-like) teeth on the basal endite of the mandible. The first assembly of the group with distinctly separated teeth, i.e. *Archiconchoecinna* gen. nov. was additionally distinguished principally by the following characters: posterior margin about 1/3-1/4 of the way down was straight, the rest of margin was round and slopes downward to the anteroventral side; terminal segment of the 5th and 6th limbs beared a very short middle claw; dorsal margin on the 3rd segment of the exopodite of the 6th limb had a claw or a claw-like seta instead of usual seta.

For all established genera and subgenera, we prepared descriptions in a uniform pattern, and elaborated keys. The description of new species was given meticulous descriptions and illustrations. For the known species, the information on the synonymy and distribution was updated and for some

Table 1. Systematic position of ostracods in subfamily Archiconchoecinae

Literary data	Proposed system
<i>Archiconchoecia</i> Müller, 1894	<i>Archiconchoecia</i> (<i>Archiconchoecia</i>) Müller, 1894
1. <i>A. striata</i> Müller, 1894 - type-species	1. <i>A. (A.) striata</i> Müller, 1894 - type-species
	2. <i>A. (A.)</i> aff. <i>striata</i> 1 (<i>A. striata</i> on George, 1979)
	3. <i>A. (A.)</i> aff. <i>striata</i> 2 (<i>A. striata</i> on Angel, 1981)
	4. <i>A. (A.) instriata</i> sp. nov.
	5. <i>A. (A.) propinqua</i> sp. nov.
	<i>Archiconchoecia</i> (<i>Archiconchoecidia</i>) subgen. nov.
	6. <i>A. (A.) apertesulcata</i> sp. nov.
	<i>Archiconchoecissa</i> gen. nov.
	7. <i>A. pljusnini</i> sp. nov. - type-species
2. <i>A.</i> aff. <i>cucullata</i> (on Angel, 1993)	8. <i>A.</i> aff. <i>pljusnini</i> (<i>A.</i> aff. <i>cucullata</i> on Angel, 1993)
3. <i>A. cucullata</i> (Brady, 1902)	9. <i>A. cucullata</i> (Brady, 1902)
	10. <i>A.</i> aff. <i>cucullata</i> (<i>A. cucullata</i> on Angel, 1983)
	11. <i>A.</i> aff. <i>cucullata</i> 2 (<i>A. cucullata</i> small form on Angel, 1993)
	<i>Archiconchoecilla</i> gen. nov.
4. <i>A. maculata</i> Chavtur, 1977	12. <i>A. maculata</i> (Chavtur, 1977) - type-species
5. <i>A. versicula</i> Deevey, 1978	13. <i>A. versicula</i> (Deevey, 1978)
	<i>Archiconchoecemma</i> gen. nov.
6. <i>A. simula</i> Deevey, 1982	14. <i>A. simula</i> (Deevey, 1982)
7. <i>A. orientalis</i> Chavtur, 1987	15. <i>A. orientalis</i> (Chavtur, 1987)
	<i>Archiconchoecinna</i> gen. nov.
	16. <i>A. ecuneata</i> sp. nov. - type-species
8. <i>A. cuneata</i> (Müller, 1908)	17. <i>A. arctica</i> sp. nov. (<i>A. cuneata</i> on Chavtur, 1978)
	18. <i>A. cuneata</i> (Müller, 1908)
	19. <i>A.</i> aff. <i>cuneata</i> 1 (<i>A. cuneata</i> on Deevey, 1968)
	20. <i>A.</i> aff. <i>cuneata</i> 2 (<i>A. cuneata</i> on Deevey, 1978)
	<i>Archiconchoecetta</i> gen. nov.
9. <i>A. ventricosa</i> Muller, 1906	21. <i>A. ventricosa</i> (Müller, 1906) type-species
10. <i>A. poulsenii</i> Deevey, 1978	22. <i>A. poulsenii</i> (Deevey, 1978)
	23. <i>A. inventricosa</i> sp. nov.
11. <i>A. bidens</i> Deevey, 1982	24. <i>A. bidens</i> (Deevey, 1982)
12. <i>A. fabiformis</i> Deevey, 1978	25. <i>A. fabiformis</i> (Deevey, 1978)
13. <i>A. bimucronata</i> Deevey, 1978	26. <i>A. bimucronata</i> (Deevey, 1978)
14. <i>A. bispicula</i> Deevey, 1978	27. <i>A. bispicula</i> (Deevey, 1978)
15. <i>A. bifurcata</i> Deevey, 1978	28. <i>A. bifurcata</i> (Deevey, 1978)
16. <i>A. gastrodes</i> Deevey, 1978	29. <i>A. gastrodes</i> (Deevey, 1978)
17. <i>A. falcata</i> Deevey, 1978	30. <i>A. falcata</i> (Deevey, 1978)
18. <i>A. pilosa</i> Deevey, 1978	31. <i>A. pilosa</i> (Deevey, 1978)
	<i>Archiconchoecerra</i> gen. nov.
19. <i>A. longiseta</i> Deevey, 1978	32. <i>A. longiseta</i> (Deevey, 1978) - type-species
20. <i>Archiconchoecia</i> sp. nov. 1 (on Angel, 1983)*	33. <i>Archiconchoecia</i> sp. nov. 1 (on Angel, 1983)*
21. <i>Archiconchoecia</i> sp. nov. 2 (on Angel, 1983)*	34. <i>Archiconchoecia</i> sp. nov. 2 (on Angel, 1983)*

*without description and figures

species comments on the morphology are given. Species previously described by the present author (Chavtur, 1977a, 1987), were redescribed on a unified pattern and its new illustrations were given.

Keys were provided for all species of each genus and subgenus.

Among the species of another assembly with slightly separated teeth on the mandible, a special place was occupied by one species due to an unusual structure of the shell and some limbs (*Archiconchoecerra* gen. nov.), namely, there was well-defined lens-like structure near the posteroventral corner of each valve; one from 6 filaments placed on the distal segment of the 1st antenna was about twice as long as others; the 3rd segment of the 6th limb did not have a dorsal seta. The rest species had a specific "ventricosa"-shape (anterior margin was extended beyond the rostrum), a low setting of the right asymmetrical gland (below the midheight of the posterior margin), and very thin claspers on the 2nd antenna in male (*Archiconchoecetta* gen. nov.).

All species, specified in Table 1 as "affinis", are actually undescribed. But since the literature does not specify the nomenclature of the specimens researched, we did not venture to isolate new taxa due to the lack of guarantees for their conservation.

The scientific text and illustrations to this work belong to V. G. Chavtur. G. G. Stovbun helped preparation of the manuscript. This work was supported by the Russian Foundation for Basic Research.

Systematics

Order HALOCYPRIDA Dana, 1853

Suborder HALOCYPRIDINA Dana, 1853

Superfamily HALOCYPRIDOIDEA Dana, 1853

Family HALOCYPRIDIDAE Dana, 1853

Subfamily ARCHICONCHOECINAE Poulsen, 1969

The subfamily Archiconchoecinae contains 7 genera, 2 subgenera and 34 species (Table 1).

Key to Subgenera and Genera of Subfamily Archiconchoecinae

1. Basal endite of mandible has toothrow of serrated teeth 2
 Basal endite of mandible has toothrow of triangular teeth 4
2. Right asymmetrical gland is located usual by less below 1/2 (rare about 1/2) shell height on the posterior margin, basal endite of mandible is armed with slightly separated teeth, exopodite of mandible is represented by usual seta 3
 Right asymmetrical gland is located about or above 1/2 the shell height on the posterior margin, basal endite of mandible is armed with distinctly separated teeth, exopodite of mandible is represented by very short spinous bristle *Archiconchoecinna* gen. nov.
3. One filament on the 1st antenna about twice as long as other 5, filaments on the endopodite of the 2nd antenna in female short and about 2 1/2 - 3 shorter than its exopodite and protopodite *Archiconchoecerra* gen. nov.
 All filaments on the 1st antenna is about equal in length, filaments on the endopodite of the 2nd antenna in female are somewhat longer than its exopodite and protopodite *Archiconchoecetta* gen. nov.
4. Rostrum is strong, prolonged, shell has posterodorsal spines, endopodite of the 2nd antenna in male bears "c" - "e" bristles with thick base *Archiconchoecissa* gen. nov.
 Rostrum is slight and not long, shell has no posterodorsal spines, endopodite of the 2nd antenna in male bears only "c" and "d" usual bristles 5
5. Dorsal margin of shell with distinct medial indentation 6
 Dorsal margin of shell without indentation 7
6. Shell less than 0,7 mm, rostrum small, rostra incisure absent, claspers on the 2nd antenna in male pointed at the tip *Archiconchoecia* (*Archiconchoecia*)

Shell larger than 0,8 mm, rostrum is developed, rostral incisure is present, claspers on the 2nd antenna in male thickened at the tip
 *Archiconchoecia* (*Archiconchoecidia*) nov. subgen.

7. Shell has asymmetrical posterior margins on the right and left valves, mandible exopodite is represented by one seta *Archiconchoecilla* gen. nov.
 Shell has symmetrical posterior margins on the right and left valves, mandible exopodite is represented by 2-3 setae *Archiconchoecemma* gen. nov.

Genus *Archiconchoecia* Müller, 1894

Archiconchoecia Müller, 1894: 225; 1906: 43-44 (part); 1912: 55 (part); Deevey, 1968: 22 (part).

Type-species: *Archiconchoecia striata* Müller, 1894.

Composition

This genus contains two subgenera: *Archiconchoecia* (*Archiconchoecia*) Müller, 1894 and *A.* (*Archiconchoecidia*) subgen. nov. First subgenus includes 5 species and new subgenus consists of one species.

Redescription

Shell. Males are ranged in length from 0.60-0.85 mm and females from 0.57-0.85 mm (Angel, 1981 - both sexes from 0.5-0.6 mm). Valves in male are rounded and slightly prolonged, and those in female are nearly rounded (in *A.* (*A.*) *apertesulcata* sp. nov. same as in male). Height of shell in male and female is 52-63% and 58-78% of the length respectively. Greatest height is at the posterior part. Rostrum is tiny or relatively developed, and rostral incisure is absent or barely noticeable. Dorsal margin is straight with distinct faint in *A.* (*A.*) *propinqua* medial indentation (except *A.* (*A.*) *apertesulcata* sp. nov.). Anterior and posterior margins are evenly rounded. Posterior margin without tubercles or with small knob-like tubercles (there asymmetrical glands in *A.* (*A.*) *apertesulcata* sp. nov.). Left and right asymmetrical glands are approximately at 70-80% and 60-70% of the way up the posterior margin respectively. Shell is with concentric striations (cross lines between concentric striations in *A.* (*A.*) *propinqua* sp. nov.). Rostrum is tiny or relatively developed.

Frontal organ. It is straight or slightly down-curved, short and narrow and extended barely beyond the down-curving distal segments of the 1st antenna. The organs in male and female are about 80-110% and 65-100% the length of the dorsal side of the 1st antenna respectively. Capitulum section is unseparated, rounded at the tip, and without hairs and spinules.

First antenna. The limb is 6-jointed, with one dorsal seta placed on the 2nd segment. This seta is long, slim and armed with short hairs (or spinules). The 5th segment bears 2, and the 6th - 4 long filaments subequal in length, which are widened or not towards the tip and pointed or nearly pointed at the tip. Dorsal margin of the limb is 55-70% (120% in *A.* (*A.*) aff. *striata* 1 according to George, 1979) of the filament length. Height of antenna (on the 2nd segment) is about 20% of its length. The 4th segment is bare (without hairs).

Second antenna. The exopodite is thick or thin (in *A.* (*A.*) *apertesulcata* sp. nov.) and shorter than the protopodite (barely longer in *A.* (*A.*) *apertesulcata* sp. nov.). The total length of the 2nd - 9th segments of the exopodite is approximately 30-50% of the length of the 1st segment. Basal segment of the endopodite has warty surface (bare surface in *A.* (*A.*) *apertesulcata* sp. nov.), and distal segment is without long hairs and is armed with "c" and "d" bristles in male. Right clasping organ is very thick in the middle. Right and left claspers are tapering towards the tip (with terminal swollen in *A.* (*A.*) *apertesulcata* sp. nov.). The 5 filaments of the distal segment of the endopodite are distally barely tapering towards the tip or have an equal thickness throughout; they are subequal in length or 2 proximal filaments are some what longer than other 3. All filaments are shorter than protopodite

(barely longer in male *A. (A.) apertesulcata* sp. nov.) and exopodite.

Mandible. The epipodite is without bristle. The exopodite is represented by one long or 2 relatively short setae. The 1st segment of the endopodite bears one dorsal and 4 ventral (one in female *A. (A.) propinqua* sp. nov.) setae; the 2nd segment is with 3 dorsal (1-3 claw-like) and 2 ventral (one in female *A. (A.) propinqua* sp. nov.) setae. The 3rd segment bears 5 (in *A. (A.) propinqua* sp. nov.) or 7 setae, of which one (only in *A. (A.) apertesulcata* sp. nov.) or 3 dorsal are claw-like. The 1st dorsal seta is shorter than the 2nd dorsal seta. The main terminal claw-seta of the mandible is longer than its endopodite (on the dorsal side) in female, and shorter, equal or longer in male. The basale is armed with 2 anterior setae and 2 lateral setae and one anterolateral seta near the articulation. Basal endite has tooththrow with 6 distinctly separated triangular teeth (additional one large lateral tooth in *A. (A.) aff. striata* 1) and 2 short posterior bristles.

Maxilla. The basal segment of the endopodite bears 6 anterior and 3-4 posterior (one may be placed laterally) setae. The distal segment has 2 claw-like setae and 3 usual setae.

Fifth limb. The epipodial appendage has 3 groups of 4+4(3)+4(5) plumose setae (16 total). The 1st segment of the exopodite bears 5 (7 in *A. (A.) apertesulcata* sp. nov.) ventral and 1-3 (one in *A. (A.) aff. striata* 1 and 3 in *A. (A.) apertesulcata* sp. nov.) dorsal setae, and the 2nd segment has 1-2 (one in *A. (A.) striata*) ventral and one dorsal setae. The middle claw-like seta on the distal segment is 50-65% the length of the exopodite (on the dorsal side) and subequal to the dorsal seta in length.

Sixth limb. The epipodial appendage is armed with 3 groups of 4(5)+4(5-6)+5(6) plumose setae (14-16 totally). The endopodite is with one long and one (or without it) short setae. The 1st segment of exopodite is without (only in *A. (A.) striata*) or with one dorsal and 2-3 (5 in *A. (A.) apertesulcata* sp. nov.) ventral setae; the 2nd segment has only one ventral (plus one dorsal seta in female *A. (A.) instriata* sp. nov.) and the 3rd segment with one dorsal and one ventral setae. The 4th segment bears 2 setae and middle claw-like seta. Latter claw-like seta is 65-70% in male and 55-70% in female, and distodorsal seta (absent in *A. (A.) striata*) of the 1st segment is 20-50% length of the exopodite (on the dorsal side). The height of the exopodite is about 30-40% of the length.

Seventh limb. The limb is short, thick (50-60% of the length) and 40-45 % of its longest seta in length.

Copulatory appendage. The greatest height is nearly at the middle, and is 45-50% of the length from tip to dorsomedial indentation. It is tapering towards the base (untapering in *A. (A.) striata*) and the tip from the middle. The pointed tip is upturned (blunt protuberance in place of point in *A. (A.) apertesulcata* sp. nov.).

Caudal furca. This limb has 6 pairs of claws and a single unpaired bristle (it is absent in *A. (A.) instriata* sp. nov.).

Distribution. Members of this genus have been collected mainly in the tropical-subtropical surface water (rare lower 500 m) of the World's Oceans in latitudinal range 36°N-42°S (subgen *A. Archiconcoecia*) and were caught in the Kurile-Kamchatka Trench (44-48°N) from 2500-4000 m (subgen *Archiconchoecidia*).

Key to Subgenera of Genus *Achiconchoecia* (Adult Male and Female)

1. Shell length is less than 0.7 mm, it is prolonged in female and rounded in male; claspers of 2nd antenna have long base and are distally narrowed; exopodite of is mandible represented by 2 short setae (one long seta in *A. (A.) aff. striata* 1) *A. (Archiconchoecia)* Müller, 1894
- Shell length over 0.7 mm, it is prolonged in female and in male; claspers of 2nd antenna have short base and are terminal swollen; exopodite of mandible is represented by one long seta *A. (Archiconchoecidia)* subgen. nov.

Subgenus *Archiconchoecia* (*Archiconchoecia*) Müller, 1894

Composition

This subgenus contains 5 species: *A. (A.) striata* Müller, 1894, *A. (A.) instriata* sp. nov., *A. (A.) propinqua* sp. nov., *A. (A.)* aff. *striata* 1 (*A. striata* sensu George, 1979) and *A. (A.)* aff. *striata* 2 (*A. striata* sensu Angel, 1981).

Description

Shell. Valves in male are prolonged and in of female rounded. Rostrum is tiny. Rostral incisure is absent (obscure in *A. (A.)* aff. *striata* 2). Posterior margin is without tubercles.

Second antenna. The exopodite is thick and shorter than the protopodite. Basal segment of the endopodite has warty surface. The right is clasper with prolonged base and distally narrowed. The left clasper is curved and about as long as the right one. The 5 filaments on the distal segment of the endopodite are shorter than the protopodite.

Mandible. The exopodite is represented by 2 somewhat short setae (one long seta in *A. (A.)* aff. *striata* 1). All dorsal setae on the 2nd segment and 2 dorsal setae on the 3rd segment are claw-like.

Fifth limb. The 1st segment of the exopodite bears 5 ventral and 1-2 dorsal setae.

Sixth limb. The 1st segment of the exopodite has 2-3 ventral setae.

Copulatory appendage. Pointed tip upturned.

**Key to Species of Subgenus *Archiconchoecia* (*Archiconchoecia*)
(Adult Female and Male)**

1. Exopodite of mandible is represented by 2 short setae 2
 Exopodite of mandible is represented by 1 long seta *A. (A.)* aff. *striata* 1
2. Each valve has one lens-like structure near ventral margin; dorsal side of the 3rd segment on exopodite of mandible is straight 3
 Valves lack lens-like structure; dorsal side of the 3rd segment on exopodite of mandible with distal protuberance *A. (A.) propinqua* sp. nov.
3. Rostrum narrow and prolonged; endopodite of 2nd antenna has short "a" and "b" bristles *A. (A.)* aff. *striata* 2
 Rostrum is wide and not prolonged; endopodite of 2nd antenna has long "a" and "b" bristles 4
4. Medioventral lens-like structure on each valve is small; middle part of right clasper of 2nd antenna is about twice as wide as the base; distodorsal seta of 1st segment of 5th limb is about 60% length of this limb (on the dorsal side); endopodite of 6th limb has one long and one short setae and 1st segment of exopodite is without dorsal seta *A. (A.) striata* Müller
 Medioventral lens-like structure one each valve is large; middle part of right clasper on 2nd antenna is about 2.5-3 times as breadth as the base; distodorsal seta of 1st segment of 5th limb is barely over 100% the length of this limb; endopodite of 6th limb has only one long seta and 1st segment of exopodite is with dorsal seta *A. (A.) instriata* sp. nov.

***Archiconchoecia* (*Archiconchoecia*) *instriata* Chavtur sp. nov.**

(Figs. 1-3 and 11, A)

? *Archiconchoecia striata*: Juday, 1906: 16, pl. 3, figs. 1-3; Tseng, 1970: 285, 286, 289, 295, table 1, figs. 5,10; Deevey, 1978a: 44, 46, table 1; 1983: table 1; Martens, 1979: 309-311, Abb. 4a-g; 1981: 68, 88, 89, 90; Hanai *et al.*, 1980: 51; Chen *et al.*; 1983: 91, fig. 9; Yin, 1991: 78; Chen and Lin, 1994a: 446, 447, 449; 1995: 41-42, fig. 47.

Material examined

Holotype N1100 - adult male, length 0.60 mm and height 0.34 mm, appendages are mounted on slide and valves remained in alcohol. In collection of the Museum of Institute of Marine Biology, Vladivostok, Russia (together with paratypes). Type-Locality - SRTR "Orlik", 36°10'N, 142°21'E, depth 100-0 m, 3 June 1967.

Paratypes. N1101 - adult male, length 0.62 mm, height 0.35 mm from same sample as of holotype; N1102 - adult female, length 0.58 mm, height 0.36 mm, SRTR "Orlik", 21°54'N, 138°18'E, depth 100-0 m, 21 May 1967; N1103 - adult female, length 0.57 mm, height 0.35 mm, SRTR "Orlik", approximately 33°N, 138°E, depth 100-0 m, date in 1967, appendages of paratypes on slides and valves in alcohol.

Additional Specimens - Adult male, length 0.60 mm, SRTR "Orlik", 21°54'N, 38°18'E, depth 100-0 m, 21 May 1967; adult female, length 0.62 mm, R/V "Cavalerovo" station 43, 33°15'N, 148°27.5'E, depth 100-0 m, 28 August 1980; juvenile, length 0.5 mm, "SRT - 662", station 21, 35°14'N, 152°07'E, depth 200-0 m, 16 September 1959; adult female and juvenile (in poor condition), R/V "Vityaz", station 3210, 34°02'N, 146°30'E, depth 230-108 m, 23 October 1954; adult female (in poor condition), R/V "Vityaz", station 3212, 35°46'N, 145°42'E, depth 600-0 m, 24 October 1954.

All materials including holotype and paratypes were collected by Juday's Plankton Net (S=0.1 m²).

Etymology

The specific name "*instriata*", from Latin "in" [=not] and "*striata*" [=lined], refers to closeness, but not identity to species *A. (A.) striata* Müller, 1894.

Description of adult male

Shell (Fig. 1, A-D). The length is 0.60-0.62 mm (from 0.62 to 0.68 mm according to Chen *et al.*, 1983 and Martens, 1979). Valves are slightly prolonged. Height of shell is 56% of the length (from 56 to 59% according to Martens, 1979 and Chen *et al.*, 1983). Greatest height is at the posterior part. Rostrum is tiny. Rostral incisure absent. Dorsal margin has slight and small medial indentation. Numerous small glands placed are along anterior, ventral and posterior margins. Posterior margin is evenly rounded. Each valve has one medial large clear lens-like structure near ventral margin. Left and right asymmetrical glands are located approximately at 70 and 60% of the way up the posterior margin respectively. The surface of the shell is sculptured with concentric striae.

Frontal organ (Fig. 1, A-G). It is straight, extended slightly beyond the down-curving distal segments of the 1st antenna, in nearly equal thickness, and is approximately 80% of the length of the 1st antenna (on the dorsal side). The stem is of equal thickness throughout. The capitulum section is unseparated, rounded at the tip, and without hairs or spinules.

First antenna (Fig. 1, E-H). The limb is 6-jointed. Dorsal seta of the 2nd segment is long, slim and armed with short coarse hairs. Long filaments on the 5th and 6th segments are distally unwidened, slightly pointed at the tip and subequal in length. Dorsal margin of the limb is 65-70% of the filament length. Height of antenna (on the 2nd segment) is 18-21% of the length. The 1st segment is wide, about 75% or equal in length with the 2nd segment, and has a few internal dark brown pigment spots.

Second antenna (Fig. 1, I-M). The exopodite is thick (thicker at the proximal part) and shorter than the protopodite (about 70% of its length). The total length of the 2nd-9th segments of the exopodite is about 40% of the 1st segment in length. Basal segment of the endopodite has warty surface are long bristles "a" and "b". The bristles "c" and "d" placed on its 2nd segment are thin and of usual type. The right clasper is very thick of the middle (about 2.5-3 times as wide as its base), with base prolonged and tapering towards the tip. The left clasper is developed, curved, without terminal thickening and about same as long as the right one, but considerably thinner than it. The 5 filaments on the endopodite are subequal in length and have an equal thickness throughout, rounded or barely pointed at the tip and are about half and two-thirds the length of the protopodite and exopodite

respectively.

Mandible (Figs. 2, A, B and 11, A). The epipodite is well developed and without bristle. The exopodite is represented by 2 relatively short setae. The 1st segment of the endopodite is with tuberos dorsal margin and bears one dorsal and 4 ventral (2 long and 2 medium length) setae, the 2nd segment bears 3 stout (claw-like) dorsal and 2 usual long ventral setae. The distal segment has straight dorsal margin and 7 setae (3 dorsal are claw-like), longest seta of which is about as long as the endopodite (on the dorsal side), and the 1st dorsal seta shorter than the 2nd seta. Basale is armed with 2 anterior and 2 lateral setae and one anterolateral seta and the anterior row of the hairs near the articulation. Basal endite has tooththrow with distinctly separated triangular teeth and 2 short posterior bristles. Coxal endite is armed with 3 closely set tooththrows. Numerous thin ventral spines are placed on the flat masticatory pad.

Maxilla. The basal segment of the endopodite bears 6 (5-6 plumose) anterior and 3 (one plumose) posterior setae. The distal segment is armed with 2 claw-setae and 3 usual setae.

Fifth limb (Fig. 2, C, D). The epipodite is torn. The 1st segment of the exopodite bears 5 (one plumose) ventral and 2 (one plumose) dorsal setae, one of the latter is about as long as the dorsal margin of the exopodite. The 2nd segment is with 2 ventral and one dorsal setae, and the latter seta is about 75% length of the exopodite (on the dorsal side). Dorsal side of the 1st segment and the ventral one of the 2nd segment are hairy. The dorsal claw-seta is borne on the distal segment is subequal to the middle claw-seta in length.

Sixth limb (Fig. 2, E, F). The epipodial appendage has 3 groups of 5+5(6)+5(6) plumose setae. The endopodite bears one long plumose seta. The 1st segment of the exopodite is with one dorsal and 2 ventral plumose setae. The 2nd segment bears one ventral and 3rd segment is with one long dorsal and one short ventral setae. Two usual setae and one middle claw-seta are placed on the 4th segment. Latter claw-seta is 65% and dorsal seta of the 1st segment is about 50% the length of the exopodite (on the dorsal side).

Seventh limb. This is deformed.

Copulatory appendage. (Figs. 2, G, H and 3, A). It is in poor condition. Penis is distally and proximally narrowed and tapers towards the base and tip from its middle. The height is greatest at the middle. The pointed tip upturned. Penis is with 2 or without (unnoted) muscles.

Caudal furca (Figs. 2, G, H and 3, H). This limb has 6 slim and long paired claws. Unpaired bristle is probably absent.

Description of adult female

Shell (Fig. 3, B-D). The length is 0.57-0.58 mm (0.58mm according to Chen and Lin, 1995 and 0.62-0.63mm in Martens, 1979). Height of shell is 60-62% of the length (65% according to Chen and Lin, 1995 and 70% in Martens, 1979). Medioventral lens-like structures are less than in male.

Frontal organ (Fig. 3, E). It is about 70 % of the 1st antenna in length (on the dorsal side).

First antenna (Fig. 3, E). Dorsal margin of the limb is 65% the length of the filaments. Height (of the 2nd segment) is 21% of the length. The 1st segment is as long as the 2nd segment and thinner than in male.

Second antenna (Fig. 3, F). The exopodite is thicker at the proximal part and is about 90% of the its length. The 2nd segment of the endopodite lacks "c"- "e" bristles. The 5 terminal filaments are subequal in length, barely pointed at tip, and are of equal thickness throughout.

Mandible (Fig. 3, G). Main terminal claw-seta is longer than the endopodite (on the dorsal side).

Fifth limb (Fig. 3, H). The endopodite bears one midlong plumose seta. The 2nd segment of the exopodite has one dorsal and one ventral setae.

Sixth and seventh limbs. They are torn.

Caudal furca. As in male.




Fig. 1. *Archiconchoecia (Archiconchoecia) instriata* sp. nov. (male: 1101 - A-C, E, I-L; 1100 - D, F-H, M, N). A and B - lateral view of left and right valves of shell, C and D - ventral view of shell, E and F - frontal organ and 1st antenna, G - distal part of frontal organ and 1st antenna, H - 1st antenna, I and J - right endopodite of 2nd antenna, K and L - left endopodite of 2nd antenna, M - right clasper of 2nd antenna, N - fragment of endopodite of the 2nd antenna.



Fig. 2. *Archiconchoecia (Archiconchoecia) instriata* sp. nov. (male: 1100). A - mandible, B - coxal segment of mandible, C and D - fifth limb, E and F - sixth limb, G and H - furca and penis.




Fig. 3. *Archiconchoecia (Archiconchoecia) instriata* sp. nov. (male: 1101 - A, female: 1102 - B, female: 1103 - F-H) A - furca and penis, B and C - lateral view of left and right valves of shell, D - ventral view of shell, E - frontal organ and 1st antenna, F - endopodite of 2nd antenna, G - endopodite of mandible, H - proximal part of sixth limb.

Distribution.

This species was caught in the north-western Pacific (21-35°N and 138-152°E) from 0-200 m (and one sample: in a vertical tow from 600-0 m). Apparently this species occurs in vicinity of San-Diego (Juday, 1906), south-eastern Pacific (30-36°S) between 0-500 m (Deevey, 1978c, 1983; Martens, 1979) South-China Sea and tropical surface and deep water of the north-western Pacific (Chen *et al.*, 1983; Chen and Lin, 1994a; 1995; Yin, 1991).

Comparison

This species is most close to *A. (A.) striata* and *A. (A.) propinqua* sp. nov. But it differs from the first species by larger medial lens-like structure on the valves (near of the middle its ventral margin), by longer distodorsal seta on the 1st segment and dorsal seta on the 2nd segment and middle claw-seta on the 3rd segment of the exopodite of the 5th limb, by presence of only one long seta on the endopodite of the 6th limb, by distally narrowed penis, by absence on the caudal furca of unpaired bristle. From the second species it differs in the following: asymmetrical glands are located lower on the posterior margin of valves; shell has concentric striae and middle lens-like structure on the surface; frontal organ is smaller and rounded at the tip; dorsal margin on the 3rd segment of the mandible endopodite is without distal ledge; on the 1st and 2nd segments are greater number of ventral setae in female, and main (middle) claw-seta on the 3rd segment of the mandible endopodite are larger; on the 1st segment of the maxilla are smaller number of setae; distodorsal seta and main middle claw-seta on the 1st and the 3rd segments of the 5th limb; are longer respectively distodorsal seta on the 1st segment of the exopodite of the 6th limb is longer; furca is without unpaired bristle.

Archiconchoecia (Archiconchoecia) propinqua Chavtur sp. nov.

(Figs. 4-6 and 11, B, C)

Material examined

Holotype. N1104 - adult male, length 0.64 mm and height 0.40 mm, appendages are mounted on slide and valves remained in alcohol. In collection of the Museum of Institute of Marine Biology, Vladivostok, Russia (together with paratype). Type-Locality - R/V "Cavalerovo", station 74, 33°02'N and 139°01'E, depth 100-0 m, 5 September 1980 (Plankton Juday's Net, S=0.1 m²).

Paratype. '1105 - adult female, length 0.65 and height 0.45 mm (appendages are mounted on slide and valves in alcohol). R/V "Cavalerovo", station 58, 34°29'N and 140°45'E, depth 100-0 m, 1 September 1980 (Plankton Juday's Net, S=0.1 m²).

Etymology

The species name "*propinqua*", from Latin [=closely related], refers to morphological relationship of *A. (A.) striata* Müller, 1894.

Description of adult male

Shell (Fig. 4, A-C). The is length 0.64 mm. Valves are slightly prolonged. The greatest height is at the posterior part and is about 60% of the length. Rostrum is tiny. Rostral incisure is absent. Dorsal margin has faint indication of medial indentation. Numerous small glands are placed along anterior, ventral and posterior margins. Upper half of rounded posterior margin is more convex than lower part. Medial lens-like structures are absent near ventral margin. Left and right asymmetrical glands are approximately 80 and 70% of the way up the posterior margin respectively. Surface of the shell is sculptured concentric striae and cross lines between them.

Frontal organ (Fig. 4, D). It is straight, extended slightly beyond the down-curving distal segments of the 1st antenna, and is about 110% the length of the dorsal side of the 1st antenna. The stem is of equal thickness throughout. Capitulum section is slightly separated, pointed at the tip, and without hairs and spinules.

First antenna (Fig. 4, D). The limb is 6-jointed. Second segment has long, slim dorsal seta, which is armed with short hairs. The 5th and 6th segments bear 6 long filaments equal in length, distally unwidened and pointed at the tip. Dorsal margin of the limb is 58% the length of the filaments. Height of antenna (on the 2nd segment) is 22% of the length. The 1st segment is slightly longer and higher than the 2nd segment, and has no dark pigment spots.

Second antenna (Fig. 4, E-G). The exopodite is thick (thicker at the middle) and shorter than the protopodite (about 80% of its length). The total length of the 2nd-9th segments of the exopodite is about 40% the length of the 1st segment. Basal segment of the endopodite has warty surface, are long bristles "a" and "b". Bristles "c" and "d", placed on its segment are somewhat thick proximally. The right clasper is about twice as thick as its base, and tapering towards the tip; its base is prolonged. The left clasper is developed and in curved, and without terminal thickening and is about as long as the right one. The filament "g" on the 2nd segment is longer and thicker than the other 4. These filaments are slightly tapering towards the tip, rounded at the tip and are about two-thirds the length of the protopodite and exopodite.

Mandible (Figs. 5, A and 11, B, C). The epipodite is well developed and without bristle. The exopodite is represented by 2 relatively short setae. The 1st segment of the endopodite has tuberos dorsal margin, and bears one dorsal and 4 ventral (2 long and 2 midlong) setae. The 2nd segment is with 3 stout (claw-like) dorsal and 2 usual long ventral setae. The 3rd segment has noticeable distal ledge on the dorsal margin and bears 5 setae, 3 dorsal of which are claw-like and 2 ventral setae are usual type. Basale is armed with 2 anterior and 2 lateral setae, one anterolateral seta, and the anterior row of the hairs near the articulation. Basal endite has tooththrow with 6 distinctly separated triangular teeth and 2 short posterior bristles. Coxal endite is armed with 3 closely set tooththrows. Numerous thin ventral spines are located on the flat masticatory pad.

Maxilla (Fig. 5, B). The basal segment of the endopodite has 6 anterior (all plumose), 3 (one plumose) posterior and one medial setae. The distal segment is armed with 2 claw-setae and 3 usual setae.

Fifth limb (Figs. 5, C and 6, H). The epipodial appendage has 3 groups of 4+3+5 plumose setae. The 1st segment of the exopodite bears 5 (one plumose) ventral and 2 (one plumose) dorsal setae, of which distodorsal seta is about 90% the length of the exopodite (on the dorsal side). The 2nd segment is with 2 ventral and one dorsal setae. Latter seta is about 75% and main (middle) terminal claw-like seta is 50% the length of the exopodite. The dorsal and middle terminal claw-setae are subequal in length. The ventral side of the 2nd segment is covered with hairs.

Sixth limb (Fig. 5, D). The epipodial appendage bears 3 groups of 4+5+6 plumose setae. Only one long plumose seta is placed on the endopodite. The 1st segment of the exopodite is with one dorsal and 2 ventral plumose setae. The 2nd segment bears one ventral and 3rd segment one long dorsal and one short ventral setae. The terminal segment is armed with two usual setae and middle one claw-seta. These claw-seta and dorsal seta are about 70% and 30% the length of the exopodite (on the dorsal side).

Seventh limb. This limb is torn.

Caudal furca (Fig. 4, H). This limb has 6 pairs of slim and long claws, and single unpaired bristle.

Copulatory appendage (Fig. 4, H). It is tapering towards the base and the tip from the middle. The greatest height is at the middle. The pointed tip is upturned. No muscles are noted on the penis.

Description of adult female

Shell (Fig. 6, A-C). The length is 0.65 mm. Shell is rounded and with strongly convex ventral margin. The height is about 70% of the length.

Frontal organ. It is in poor condition.

First antenna. As in male.

Second antenna (Fig. 6, D, E). The 1st segment of the exopodite is twice the total length of 2nd-




Fig. 4. *Archiconchoecia (Archiconchoecia) propinqua* sp. nov. (male: 1104) A and B - lateral view of left and right valves of shell, C - ventral view of shell, D - frontal organ and 1st antenna, E - 2nd antenna, F and G - right and left endopodites of 2nd antenna, H - furca and penis.



Fig. 5 *Archiconchoecia (Archiconchoecia) propinqua* sp. nov. (male: 1104) A - mandible, B - maxilla, C - fifth limb, D - sixth limb.




Fig. 6 *Archiconchoecia (Archiconchoecia) propinqua* sp. nov. (female: 1105 - A-G, male: 1104 - H) A and B - lateral view of left and right valves of shell, C - ventral view of right valve, D - 2nd antenna, E - endopodite of 2nd antenna, F - mandible, G and H - distal part of fifth limb.

9th segments. Endopodite without "a"-"e" bristles. Terminal "g" filament is slightly tapering towards the tip or has an equal thickness throughout, pointed at the tip and is about 3/4 the length of the protopodite and exopodite ("h"-"f" filaments are broken).

Mandible (Fig. 6, F). The 1st segment of the endopodite bears one dorsal and one ventral and 2nd segment has 3 dorsal and one ventral setae. Main terminal claw-seta is shorter than the endopodite (on the dorsal side).

Maxilla and fifth limb (Fig. 6, G, H). They are as in male.

Seventh limb. It is torn. The setation of the distal segment and the length of its main claw-seta are as in male.

Caudal furca. As in male.

Comparison

This new species is close to *A. (A.) instriata* sp. nov., but differs in many respects: asymmetrical glands are located above on the posterior margin of valves; shell with concentric striations and cross lines between these striations, and without middle lens-like structure on the surface; frontal organ is larger and pointed at the tip; dorsal margin on the 3rd segment of the mandible endopodite is with distal ledge; on the 1st and 2nd segments of the mandible in female are placed less ventral setae, and there is smaller main middle claw-seta on the 3rd segment there are more setae on the 1st segment of the maxilla; the distodorsal seta on the 1st segment, the main middle claw-seta on the 3rd segments of the 5th limb, and distodorsal seta on the 1st segment of the exopodite of the 6th limb are shorter; furca is with unpaired bristle.

Distribution

Collected only in subtropical surface water of the north-western Pacific (33-34°N and 139-140°E) from 100-0 m.

Archiconchoecia (Archiconchoecia) striata Müller, 1894

Archiconchoecia striata Müller, 1894: 225, Taf. 6, Fig. 31-46, Taf. 8, Fig. 34; 1906: 45, Taf. 7, Fig. 13-17; 1908: 63; 1912: 56; Schweiger, 1912: (cit. on Gooday, Angel, 1977); Gamulin, 1948: (cit. on Gooday, Angel, 1977); Hure, 1961: 11, 54, Sl. 6, 7, Tab. 2; Grice, Hart, 1962: 302; Puri, 1963: 2 (list); Leveau, 1965: 178; 1967: 67, 1969: 129, 130, 131, 136, 138, 140; Deevey, 1968: 23, fig. 4; 1970: 801; 1971: 231; 1974: 358; 1978b: 57, 59, tab. 1, 2; 1982: 469, 470; Deevey, Brooks, 1980: 59-61, tab. 2-4; Vucetic, 1977; (cit. on Gooday, Angel, 1977); Alcaraz, Manriques, Vasquez, 1975: 382, 383, 384, Fig. 1: 16, 17; Alcaraz, 1977: 5, 23; Angel, Fasham, 1975: 727, 736, fig. 5, tab. 2-4; Angel, 1979: 81-82; 1981: 556, 561 (part); Gooday, Angel, 1977: 140, 144, 146; Moraitou-Apostolopoulou, 1981: 175, 176, fig. 3; Gonzales, Breman, 1982: 204, 205, 206; Drapun, 1981: 75; 1983: 34; Benassi, Ferrari, Rossi, Sei, Angel and McKenzie, 1998.

Not *Archiconchoecia striata*: Chavtur, 1977a: 142, 157, 158, fig. 8, tab. 6; 1977b: 31, tab. 2 (list); 1977c: 19, tab. 2 (list); 1991: 47.

Not *Archiconchoecia striata*: George, 1979: 123-127, figs. 1-16; George, Nair, 1980: 31, 41, fig. 6; Angel, 1981: 556 (part), 561 (part), fig. 194 (6 A-H); Chavtur, 1992: tab. 2 (list).

Material examined

Adult female (in poor condition), R/V "Lomonosov", station 466, 22°19'N 61°02'W, in tow from 2600-0 m, 2 October 1959.

Distribution

This species is known from tropical - subtropical zone of the Mediterranean Sea (Müller, 1894; Schweiger, 1912; Gamulin, 1948; Hure, 1961; Grice and Hart, 1962; Puri, 1963; Leveau, 1965;

Vucetic, 1970; Alcaraz, 1977; Gooday and Angel, 1977; Moraitou-Apostolopoulou, 1981; Benassi et al., 1998) and Atlantic Ocean within latitudes 36°N-42°S (NE Atlantic - Müller, 1906; Alcaraz, et al., 1975; Alcaraz, 1977; Angel and Fasham, 1975; Angel, 1979; 1981. NW Atlantic - Deevey, 1968; 1970; 1971; 1978b; 1982; Deevey, Brooks, 1980; Angel, 1979; 1981; Gonzales and Breman, 1982. S. Atlantic - Müller, 1906; 1908; Deevey, 1974; Angel, 1981; Drapun, 1981; 1983).

Archiconchoecia striata was found at levels between the surface and 1000 m with mostly in the upper 500 m (Deevey, 1978b; Deevey and Brooks, 1980) and (as exclusion) 1 juvenile was recovered from 2700-2400 m (Müller, 1906). This species was captured also in tows from 3000-0 m. Besides that, it is known in the Indian Ocean (7-34°N and 27-32°S) between the surface and 400 m (Müller, 1906; 1908; Leveau, 1967; 1969).

Archiconchoecia (Archiconchoecia) aff. striata 1

Archiconchoecia striata: George, 1979: 123-127, figs. 1-16; George and Nair, 1980: 31, 41, fig. 6.

Remarks

This species distinctly differs from Müller's description in small rostrum on the shell, suture on the frontal organ, thin exopodite, short filaments, and strongly curved right clasper on the 2nd antenna, and only one seta on the mandible exopodite.

Distribution

Collected in the epipelagial of the Indian Ocean within latitudes 4-24°N. Probably, specimens noted by Müller (1906, 1908) and Levean (1967, 1969) from this ocean also belong to *A. (A.) aff. striata* 1.

Archiconchoecia (Archiconchoecia) aff. striata 2

Archiconchoecia striata: Angel, 1981: 556 (part), 561 (part), fig. 194 (6 A-H).

Remarks

This species is separated from Müller's description by prolonged rostrum on the shell, and very small size of "a" and "b" bristles on the 2nd antenna (Angel's publication contains only information for the shell and 1st and 2nd antenna).

Distribution

Atlantic Ocean (precise locality is unknown).

Archiconchoecia (Archiconchoecidia) Chavtur subgen. nov.

Composition

This subgenus contains only *A. (A.) apertesulcata* n. sp.

Description

Shell. Valves of both sexes are prolonged. Rostrum is relatively developed. Rostral incisure is barely noticeable. Posterior margin is with one small knob-like tubercle on each valve.

Second antenna. The exopodite is thin and longer than the protopodite. Basal segment of the endopodite has bare surface. The right clasper is with short base and with terminal thickening. The left clasper is straight and shorter than the right one. The 5 filaments placed on the distal segment of the endopodite in male are longer than protopodite.

Mandible. The exopodite is represented by one long seta. All dorsal setae borne on the 2nd segment and 2 dorsal setae on the 3rd segment of the exopodite are usual type.

Fifth limb. The 1st segment of the exopodite bears 7 ventral and 3 dorsal setae.

Sixth limb. The 1st segment of the exopodite is armed with 5 ventral setae.

Copulatory appendage. The blunt protuberance is placed at the tip.

***Archiconchoecia (Archiconchoecidia) apertesulcata* Chavtur sp. nov.**

(Figs. 7-10 and 11, D-G)

Archiconchoecia striata: Chavtur, 1977a: 142, 157, 158, fig.8, table 6; 1977b: 31, table 2 (list); 1977c: 19, table 2 (list); 1991: 47.

Archiconchoecia striata: Chavtur, 1992: table 2 (list).

Material examined

Holotype. N1106 - adult male, length 0.85 and height 0.34 mm, appendages are mounted on slide and valves remained in alcohol. In collection of the Museum of Institute of Marine Biology, Vladivostok, Russia (together with paratypes). Type-locality - R/V "Vityaz", station 5635, 44°25N, 149°10E, depth 4000-3000 m, 28 August 1966 (Plankton Bogorov-Rass's Net, S=1,0 m²).

Paratypes. 1107 - adult male (length 0.85 and height 0.50 mm), 1108 - adult female (length 0.82 and height 0.50 mm), 1109 - adult female (length 0.85 and height 0.50 mm) from the same sample as holotype. Appendages of all specimens are remained on slides and valves in alcohol.

Additional specimens - R/V "Vityaz": 3 adult males, length 0.85 mm, from the same sample as holotype; one adult female, length 0.85 mm, station and date as holotype, depth 3000-2500; 2 adult females, length 0.82-0.84 mm, station 5610, 48°48N, 153°18E, depth 3000-2500 m, 23 July 1966.

All material including holotype and paratypes were caught by Plankton Bogorov-Rass's Net (S=1,0 m²).

Etymology

The specific name "*apertesulcata*", from the Latin "*aperte*" [=distinctly] and "*sulcata*" [=grooved, lined], refers to sculpturing of the shell.

Description of adult male

Shell (Fig. 7, A, B). Length is 0.85 mm. Valves are slightly prolonged. Height of shell is 56-58% of the length. Greatest height is at the posterior part. Rostrum is wide and relatively developed, and there rostral incisure is barely noticeable. Dorsal margin with is distinct medial indentation. Numerous varisized glands are placed along anterior and ventral margins. Posterior margin is evenly rounded and with a small knob-like tubercle on each valve (there placed asymmetrical gland). Left and right asymmetrical glands on the valve are approximately at 70 and 60% of the way up the posterior margin respectively. Shell is with concentric striations.

Frontal organ (Fig. 7, C, D). It slopes downward, extends beyond curved distal segments of the 1st antenna, and is nearly equal the length of the dorsal side of the 1st antenna. Capitulum section is unseparated, rounded at the tip, without hairs and spinules and thicker than stem.

First antenna (Fig. 7, D). The limb is 6 - jointed. Dorsal seta of the 2nd segment is long, slim and armed with spinules. The 5th and 6th segments are with long filaments of equal length, distally widened and pointed at the tip. Dorsal margin of the limb is about 60% of the filaments length. Height of antenna (on the 2nd segment) is approximately 20% of the length. The 1st segment is slightly thicker than the 2nd segment and almost 1/4 as long as it. There are few dark pigment spots on the 1st segment.

Second antenna (Fig. 7, E-G). The exopodite is thin, approximately equal in thickness along the shaft and is about 110% length of the protopodite. The total length of the 2nd-9th segments of the exopodite is about 30% length of the 1st segment. Basal segment of the endopodite has a surface, and bristles "a" and "b" are The bristles "c" and "b" on the 2nd segment are of usual type. The right




Fig. 7 *Archiconchoecia (Archiconchoecidia) apertesulcata* sp. nov. (male: 1107 - A, B, D; 1106 - C, E-I)
 A and B - lateral and ventral views of shell, C - frontal organ, D - frontal organ and 1st antenna, E -
 2nd antenna, F and G - left and right endopodites of 2nd antenna, H and I - mandible.




Fig. 8 *Archiconcoecia (Archiconchoecidia) apertesulcata* sp. nov. (Male: 1106 - 1-5; 1107 - 6) A - maxilla, B and C - fifth limb, D - sixth limb, E - seventh limb, F - penis.




Fig. 9 *Archiconchoecia (Archiconchoecidia) apertesulcata* sp. nov. (male: 1106- A, D, F, female 1109 - B, C, E, G-I) A- furca and penis, B and C - lateral and ventral views of shell, D - margin of right valve, E - anterior part of left valve, F and G - posterior part of shell, H and I - frontal organ and 1st antenna.




Fig.10 *Archiconchoecia (Archiconchoecidia) apertesulcata* sp. nov. (female: 1109 - A, B, D-F, 1108 -C)
 A - distal part of 1st antenna, B - 2nd antenna, C - endopodite of 2nd antenna, A - endopodite of mandible, E - sixth limb, F - furca.

clasper is thick in the middle (about 1.5 times thicker than its base), has a short base and tapers towards the tip. The left clasper is developed, straight, with terminal thickening, shorter and thinner than the right one. All the filaments borne on 2nd segment are equal in length, slightly widened in the middle part, rounded and have spinule at the tip, and are approximately as long as the exopodite and protopodite.

Mandible (Figs. 7, H, I and 11, D). The epipodite is well developed and lacks bristle. The exopodite is represented by one long seta. The 1st segment of the endopodite has slightly tuberosous dorsal margin and is armed with one dorsal and 4 ventral (one short and 3 long) setae, and the 2nd




Fig. 11 *Archiconchoecia (Archiconchoecidia) instriata* sp. nov. (male: 1100 - A), *A. (A.) propinqua* sp. nov. (female: 11105 - B and C), *A. (Archiconchoecidia) apertesulcata* (male: 1106 - D, female: 1108 - E, female: No. 1109 - F,G) A-G - tooththrows and masticatory pad of coxa of mandible.

segment is with 3 slight dorsal and 2 usual long ventral setae. The 3rd segment has straight dorsal margin and bears 7 setae, longest seta of which is claw-like and shorter than the endopodite (on the dorsal side); the 1st dorsal seta is longer than the 2nd seta. Basale is armed with 2 anterior and 2 lateral setae, one anterolateral seta, and the anterior row of the hairs are near the articulation. Basal entite has toothrow with 6 distinctly separated triangular teeth and 2 short proximal bristles. Coxal endite bears 3 closely set toothrows. Numerous thin ventral spines are borne on the flat masticatory pad.

Maxilla (Fig. 8, A). The basal segment (of the endopodite) bears 6 anterior (one plumose), 3 posterior (2 plumose), and one medial, setae. The distal segment is armed with 2 claw-like and 3 usual setae.

Fifth limb (Fig. 8, B, C). The epipodite has 3 groups of 4+4+4 plumose setae.

The 1st segment of the exopodite bears 7 ventral (2 plumose) and 3 dorsal (2 plumose) setae, of which the distodorsal seta is about 75% length of the exopodite (on the dorsal side). The 2nd segment is with 2 ventral and one dorsal setae, and the last seta is approximately 50%, and the middle claw-seta of the 3rd segment is about 60% the length of the exopodite. The dorsal and middle terminal claw-setae are subequal in length. No hairs are recorded the limb.

Sixth limb (Fig. 8, D). The epipodial appendage has 3 groups of the 5+4+6 plumose setae. One long plumose and one short nonplumose setae are borne on the endopodite. The 1st segment of the exopodite has one dorsal and 5 ventral (3 plumose) setae. The 2nd segment is armed with one ventral seta and the 3rd one with one long dorsal and one short ventral setae. Two usual setae and one middle claw-seta are placed on the 4th segment. Last claw-seta and dorsal seta of the 1st segment are about 70% and 40% length of the exopodite (on the dorsal side) respectively.

Seventh limb (Fig. 8, E). It is short and thick (65% of the length), and is about 40% length of its longest seta. The short seta is 75% length of the long one.

Copulatory appendage. (Figs. 8, F and 9, A). It is tapering towards the base and tip from the middle. The greatest height is at the middle. The rounded tip is with blunt tubercle. No muscles are noted on the penis.

Caudal furca (Fig. 9, A). This limb has 6 slim and long paired claws and single unpaired bristle.

Description of adult female.

Shell (Fig. 9, B-G). Length is 0.82-0.85 mm.

Frontal organ (Fig. 9, H, I). Capitulum section bears some terminal hairs (or spinules?).

First antenna (Figs. 9, H, I and 10, A). The 1st segment is about 1/3 as long as the 2nd segment.

Second antenna (Fig. 10, B, C). The endopodite is without "c"- "e" bristles. The terminal filaments are equal in thickness (along the entire length) and about 3/4 the length of the protopodite and exopodite.

Mandible (Figs. 10, D and 11, E-G). Main terminal claw-seta is longer than the endopodite (on the dorsal side).

Maxilla and fifth limb. As in male.

Sixth limb (Fig. 10, E). The epipodial appendage has 3 groups of the 5+4+5 plumose setae. Dorsal seta of the 1st segment is about 30% length of the exopodite (on the dorsal side).

Seventh limb and Caudal furca (Fig. 10, F). As in male.

Comparison

New species sharply differs from all the other species of this genus in having a rostral incisure and tubercles on the posterior margins of the shell. Its exopodite is thin and longer than protopodite on the 2nd antenna. Claspers on the 2nd antenna have a short base and terminal thickening, and its filaments are longer than protopodite in male. Mandible bears only one seta on the exopodite and usual (not claw-like) dorsal setae on the 2nd and 3rd (setae near dorsal margin) segments of the endopodite. This new species is also differs from other by the number of setae on the 5th and 6th

limbs and the presence of blunt protuberance at the tip.

Distribution

This species was caught only in the region of Kurile-Kamchatka Trench from 2500-4000 m.

Archiconchoecilla Chavtur gen. nov.

Type-species - *Archiconchoecia maculata* Chavtur, 1977.

Composition

This genus contains only 2 species: *A. maculata* (Chavtur, 1977) and *A. versicula* (Deevey, 1978).

Description

Shell. Male and female range from 0.8 to 1.4 mm in length. Valves are slightly prolonged. Height of shell in male and female is 52-56% and 52-55% of the length respectively, and approximately equal at the anterior and posterior parts. Rostrum is developed, and there is only a slight rostral incisure. Dorsal and ventral margins are straight or nearly straight. Anterior and posterior margins are rounded (posterior part of the right valve in *A. maculata* is unrounded). Posterior margins of the right and left valves are asymmetrical at the upper half: left margin is slightly extended beyond right margin. Left and right asymmetrical glands are about 80-90 and 60-70% of the way up the posterior margin respectively. These glands open on tiny tubercles. The surface of the shell is sculptured with wavy lines or with varying shape cells (reticulated).

Frontal organ. It is straight or slightly down-curved, short or middle length, and barely or considerably extended beyond the down-curving distal segments of the 1st antenna. Organs in male and female are approximately 90-110% and 80% of the length of the dorsal side of the 1st antenna respectively. Capitulum section is unseparated, rounded at the tip (slightly pointed in *A. versicula* on Angel, 1981).

First antenna. The limb is 6-jointed, with one dorsal seta placed on the 2nd segment. This seta long, slim and armed coarsely with short hairs or spinules. The 5th and 6th segments bear 6 long filaments in equal length, slightly widened towards the tip or unwidened and pointed or rounded at the tip. Dorsal margin of the limb is 58-60% of the filaments length. Height of antenna (on the 2nd segment) is less 20% (17% in all species) of the length. Hairs are placed dorsally on the 4th segment. The limb has numerous dark pigment spots.

Second antenna. The exopodite is thin, equal in length or longer than the protopodite. The total length of the 2nd-9th segments of the exopodite is approximately 35-45% length of the 1st segment. Basal segment of the endopodite bears long hairs (female *A. versicula* unknown) and without warty surface. Its 2nd segment is armed with long hairs in female and with bristles "c" and "d" in male. The right clasping organ is slightly thick. The right and left claspers distally are not tapering, with or without usual terminal thickening and rounded at the tip. The filaments of the 2nd segment have an equal thickness throughout, and pointed at the tip. One or two proximal filaments are slightly longer than the 3 others, protopodite and exopodite.

Mandible. The epipodite is without bristles. The exopodite is represented by one long seta. The 1st segment is armed with one dorsal and 4 ventral setae; the 2nd segment with 3 dorsal and 2 ventral setae; the 3rd segment has 7 setae, the largest of which is claw-like and 2 first dorsal setae are slight (the 1st shorter than 2nd dorsal seta). Main claw-seta borne on the 3rd segment is longer than the exopodite (on the dorsal side) in female, and shorter in male. Basale is armed with 2-3 anterior, 2 lateral and 1 anterolateral setae and anterior row of the long hairs near the articulation. Basal endite has toothrow with 6 distinctly separated triangular teeth (in addition to one lateral tooth in *A. maculata*) and 2 short posterior bristles.

Maxilla. The basal segment of the endopodite bears 6 anterior and 4 posterior setae. The 2nd segment has 2 claw-setae and 3 usual setae.

Fifth limb. The epipodial appendage has 3 groups of 4+4+4 plumose setae. The 1st segment of the exopodite bears 6-7 ventral and 3 dorsal setae, and 2nd segment is with 1-2 ventral and one dorsal setae. The middle claw-seta of the 3rd segment is about 60% the length of the exopodite (on the dorsal side) and approximately as long as its dorsal seta.

Sixth limb. The epipodial appendage has 3 groups of 4(5)+4+5(6) plumose setae (13-15 totally) in male and 4(?) + 4 + 4 in female. One long plumose seta is borne on the endopodite. The 1st segment of exopodite is with 1-2 dorsal and 4-5 ventral setae, the 2nd segment is only with one ventral, and 3rd segment is with one dorsal and one ventral setae. The middle claw-seta of the 4th segment is longer than its 2 usual setae, and is 75-80% the length of exopodite in male and 75% in female. The distodorsal seta of the 1st segment is 35-80% the length of the exopodite (on the dorsal side). Height of the exopodite is approximately 25-35% of its length.

Seventh limb. It is thick and prolonged (50% of its length) in *A. maculata* and short (25%) in *A. versicula*. The limb is about 30% of the longest seta in length (this seta is broken in *A. versicula*).

Copulatory appendage. It is tapering towards the base and tip from the middle. The greatest height is at the middle and approximately 30-40% of the length (from tip to dorsomedial indentation). The tip is rounded.

Caudal furca. This limb has 6 or 8 pairs of claws and with or without unpaired bristle.

Comparison

New genus differs from all the other genera in the following respects: posterior margins on the right and left valves are asymmetrical at the upper half; the exopodite of the 2nd antenna is longer than the protopodite; the exopodite of the mandible is represented only by one long seta, and its endopodite is armed with usual dorsal setae.

This genera is close to *Archiconchoecinna* gen. nov., but separated (in addition to the above mentioned) by equal height in the anterior and posterior halves of the shell, triangular teeth on the basal endite of the mandible, less number of the posterior setae placed on the 1st segment of the endopodites of the maxilla and of the 6th limb, longer distodorsal seta on the 1st segment of the exopodite of the 5th limb, and longer middle-terminal seta on the 6th limb.

Distribution

Members of the genus *Archiconchoecilla* were collected in the North Pacific, North Atlantic and Southern Ocean (Pacific sector). The northernmost and southernmost latitude are 45°N and 44°S. The known depth range of the genus is 1000-9500 m.

Key to Species of Genus *Archiconchoecilla* (Adult Male and Female)

1. Shell length is over 1mm, sculpture is represented by varying shape cells, furca has 8 pairs of claws *A. maculata* (Chavtur)
- Shell length is less than 1mm, sculpture is represented by wavy lines, furca has 6 pairs of claws *A. versicula* (Deevey)

Archiconchoecilla maculata (Chavtur, 1977)

(Figs. 12-16)

Archyconchoecia maculata Chavtur, 1977a: 140-142, 155, 158, 160, figs. 3c-d, 4a-d; 1977b: table 2; 1977c: tabl. 2; 1991: 42.

Archiconchoecia maculata Chavtur, 1992: table 2 (list).

Material examined

Holotype. 1110 - (1526 - obsolete number) adult male, length 1.4 mm, appendages are mounted

on slide and valves preserved in alcohol. In collection of the Museum of Institute of Marine Biology, Vladivostok, Russia (together with paratypes). Type-locality- R/V "Vityaz", station 5262, 45°11N and 152°28E, depth 9500-7280 m, 24 August 1966 (Plankton Bogorov-Rass's Net, S=1.0 m²).

Paratypes. 1111 (1527 - obsolete number) -adult female (length 1.4 mm), 1112 - adult female (1.3 mm), 1113 - adult male (1.4mm), 1114 - adult male (1.4 mm) from same sample as of holotype.

Additional specimens - 24 adult females, 20 adult males and 13 juveniles from sample as holotype.

Redescription of adult male

Shell (Fig. 12, A, B). The length is 1.2-1.4 mm. Valves are slightly prolonged. Height of shell is 52-53% of the length, and is equal at the anterior and posterior parts. Rostrum is developed and wide; there is only slight rostral incisure. Dorsal and ventral margins are straight. Posterior margins of the right and left valves are asymmetrical at the upper half, left margin is extended slightly beyond right margin. Left and right asymmetrical glands are approximately at 90 and 70% of the way up the posterior margin respectively. These glands are open on tiny tubercles. The surface of the shell is sculptured with varying shape cells.

Frontal organ (Fig. 12, C). It is straight, short, thicker at the proximal part and barely extends beyond the down-curving distal segments of the 1st antenna and it is approximately 90-95% of the dorsal side in length. Capitulum section is not separated, rounded at the tip and without hairs and spinules.

First antenna (Fig. 12, C-E). The limb is 6-jointed. Second segment has long, slim seta and is armed with short hairs. The 5th and 6th segments bear 6 long filaments equal in length, which are slightly widened towards the tip or not and pointed at the tip. Dorsal margin of the limb is nearly 60% the length of the filaments. Height of antenna (on the 2nd segment) is about 17% of the length. The 1st segment is slightly higher than the 2nd segment and approximately is 70% of its length. Long hairs are placed dorsally on the 4th and 5th segments. The 1st and 2nd segments have numerous dark pigment spots.

Second antenna. (Figs. 12, F, G and 13, A, B). The exopodite is thin (more thin in the middle part) and longer than the protopodite. The total length of the 2nd-9th segments of the exopodite is about 35% of the 1st segment in length. Basal segment of the endopodite bears long hairs; the bristles "a" and "b" are very developed and The bristles "c" and "d" placed on the 2nd segment are stout and spine-like type (bristle "e" is absent). The right clasper is curved and has terminal thickening; the base and the middle parts of this clasper are approximately equal in thickness. The left clasper is nearly straight and shorter than right one. Claspers are rounded at the tip. The 5 filaments on the 2nd segment are approximately equal in thickness lengthwise and pointed at the tip. One or 2 proximal filaments are longer than other 3, the protopodite and exopodite.

Mandible (Fig. 13, C-F). The epipodite is developed and without bristle. The exopodite is represented by one long seta (2 only in one limb). The 1st segment of the endopodite has almost straight dorsal margin and is armed with one dorsal and 4 ventral long setae; the 2nd segment has 3 dorsal and 2 ventral setae. The 3rd segment bears 7 setae, of which main claw-setae is 77-81% of the endopodite in length (on the dorsal side) and 2 first dorsal setae are slight (the 1st one is shorter than the 2nd). Basale is armed with 2-3 anterior and 2 lateral setae and with one anterolateral seta and anterior row of the hair near the articulation. Basal endite has tooththrow with 6 distinctly separated terminal and one lateral triangular teeth and 2 short posterior bristles. Coxal endite is armed with 3 closely set tooththrows. Numerous thin ventral spines and lateral hairs are located on the flat masticatory pad.

Maxilla (Fig. 14, A-C). The basal segment of the endopodite bears 6 plumose anterior and 4 (of which 2 are plumose) posterior setae. The distal segment is armed with 2 claw-setae and 3 (only 4 in one limb of male) usual setae.

Fifth limb (Fig. 14, D, E). The epipodite is armed with 3 groups of 4+4+4 plumose setae. The




Fig. 12 *Archiconchoecilla maculata* (male: 1110 - A-D, G; 1113 - E; 1114 - F) A - lateral view of right valve of shell, B - posterodorsal part of shell, C - frontal organ and 1st antenna, D and E - 1st antenna, F and G - left and right 2nd antenna.




Fig. 13 *Archiconchoecilla maculata* (male: 1110 - A, B; 1114 - C, F; 1113 - D, E) A - right endopodite of 2nd antenna, B - left clasper of 2nd antenna, C and D - mandible, E - endopodite of mandible, F - tooththrows and masticatory pad of coxa of mandible.




Fig. 14 *Archiconchoecilla maculata* (male: 1114 - A, B; 1113 - C, E, F) A, B and C - maxilla, D and E - fifth limb, F - sixth limb.

1st segment of the exopodite bears 6 (one plumose) ventral and 3 (all plumose) dorsal setae, of which distodorsal seta is subequal (95-96%) to the dorsal margin of the exopodite in length. The 2nd segment is with 2 ventral and one dorsal setae; the last seta is about 50% and middle claw-seta of 3rd segment is 60% of the exopodite in length (on the dorsal side). The terminal dorsal and middle claw-setae are subequal in length. Dorsal and ventral sides of the 1st and 2nd segments bear long hairs.




Fig. 15 *Archiconchoecilla maculata* (male: 1113 - A, B, D; male: 1114 - C, D, female: 1112 - F) A - sixth limb, B and C - penis, D - distal part of penis, E - furca, F - 2nd antenna.




Fig. 16 *Archiconchoecilla maculata* (female: 1111 - A; 1112 - B-F) A - endopodite of 2nd antenna, B and C - mandible, F - basale endite of mandible, E - sixth limb, F - toothrows and masticatory pad of coxa of mandible.

Sixth limb (Fig. 15, A). The epipodite appendage has 3 groups of 4+4+5 plumose setae. One long plumose seta is placed on the endopodite. The 1st segment of the exopodite is with 2 (one plumose) dorsal and 4 (all plumose) ventral setae. The 2nd segment bears one ventral, and 3rd - one dorsal and one ventral setae. Two usual setae and one middle claw-seta are borne by the 4th segment. This claw seta and dorsodistal seta of the 1st segment are about 80% of the exopodite in length (on the dorsal side). Height of the exopodite is approximately 35% its length.

Seventh limb (Fig. 14, F). It is prolonged and thick (25% of the length) and is about 30% of the longest seta in length. The short seta is 75% the length of the long one.

Copulatory appendage (Fig. 15, B-D). It is tapering towards the base and tip from the middle. The greatest height is at the middle. It is rounded at the tip and has 3 slightly developed muscles.

Caudal furca (Fig. 15, E). It has 8 pairs of slim claws and without odd bristle.

Redescription of adult female

Shell. The length is 1.2-1.4 mm, as in male.

Frontal organ. It does not extend beyond the down-curving distal segments of the 1st antenna and is about 80% of its dorsal side in length.

First antenna. As in male.

Second antenna. (Figs. 15, F and 16, A). The total length of the 2nd-9th segments of the exopodite is about 45% of the 1st segment in length. The 1st and 2nd segments of the endopodite are covered with long hairs.

Mandible (Fig. 16, B-D, F). Basale is armed with 3 anterior, one anterolateral and 3 lateral setae. Main terminal claw - seta is 93-110% of the endopodite in length (on the dorsal side).

Maxilla and fifth limb. As in male.

Sixth limb (Fig. 16, E). The epipodial appendage has 3 groups of 4+4+4 plumose setae. Middle claw-seta of the 4th segment is approximately 75% and distodorsal seta is 50% of the exopodite in length (on the dorsal side). Height of the exopodite is about 30% of the length.

Seventh limb and Caudal furca. As in male.

Comparison

The shell of *A. maculata* differs from that of *A. versicula* in having a large size, sculpture with waving shape cells, more asymmetrical another shape posterior margins on the left and right valves. Besides *A. maculata* has smaller frontal organ, longer 1st segment on the 1st antenna, and bears dorsal hairs on the 5th segment of this limb. It has longer exopodite and right clasper on the 2nd antenna, greater number of setae and teeth are situated on the basal segment of the mandible, less ventral setae on the 1st and greater number their 2nd segments of the 5th limb, greater number and length of the dorsal setae on the 1st segment of the 6th limb, more prolonged 7th limb, more straighter and narrower penis, greater number of claws and no odd bristle on the furca.

Distribution

Kurile-Kamchatka Trench (45°11N-152°28E), depth 7280-9500m (Chavtur, 1977a).

Archiconchoecilla versicula (Deevey, 1978)

Archiconchoecia versicula Deevey, 1978a: 46-50, figs. 2, 3; 1978 c: 106, 111, table 1; 1983: table 1; Angel, 1981: 558, fig. 194 (7, A - H); 1983: 553, 554, fig. 2; Chen and Lin, 1995: 42-43, fig. 48.

Distribution

Species is known from Pacific sector of the South Ocean (37-44°S) at depth range of 1000 to 2000m (Deevey, 1978a; 1982; 1983). Also, *A. versicula* was found in the Caribbean Sea (Deevey, 1978b) and in the Northeastern Atlantic (42°N, 17°W) at depth of 3300-3900m.

Archiconchoecissa Chavtur gen. nov.

Archiconchoecia Müller, 1906: 43-44 (part); 1912: 55 (part); Granata, Caporiacco, 1949: 14; Deevey, 1968: 22 (part); Poulsen, 1969: 14-15 (part).

Type-species - *Archiconchoecissa pljusnini* Chavtur n. sp.

Composition

This genus contains only 5 species: *A. pljusnini* n. sp., *A. cucullata*, *A. aff. cucullata* 1, *A. aff. cucullata* 2, *A. aff. pljusnini*.

Description

Shell. Males range from 1.62 to 2.50 mm in length and females from 1.58 to 2.60 mm. Valves are prolonged. Height of shells in male in and female approximately 25-40% and 35-45% of the length respectively (without posterodorsal spines); it is subequal at the anterior and posterior parts or slightly greater at the posterior part of shell (considerably greater in female *A. cucullata* on Angel, 1981, fig. 194-4a). Rostrum is strongly developed, long and wide; rostral incisure is absent. Dorsal margin is straight, ventral margin also, slightly convex or barely concave. Anterior margin is nearly straight or slightly rounded. Posterior margin is rounded or slightly rounded. Right valve has long posterodorsal spine. Ventral and posterior margins are serrated. Left asymmetrical gland is approximately at 65-90% of the way up the posterior margin and placed on developed tubercle; right gland is respectively at 55-65% and located on small tubercle. There are distinct primary reticular and secondary tile-like (with denticles) sculptures.

Frontal organ. It is straight or barely down-curved, long, narrow, considerably extended beyond the down-curving distal segments of the 1st antenna. Organ in male and female are respectively about 100-135% and 100-130% the length of the dorsal side on the 1st antenna. Capitulum section is not separated (separated only in female *A. cucullata* on Angel, 1981, fig. 194-4c) and without hairs (with hairs only in male *A. cucullata* on Poulsen, 1969, fig. 4h).

First antenna. The limb is 5-jointed (6-jointed only in *A. cucullata* on Poulsen, 1969: 16, Fig. 36). The 2nd and 5th segments have each one long, slim seta and armed with hairs (seta of the 5th segment is shorter than 2nd one). Six long filaments are borne on the 5th and 6th segments equal in length, distally not dilating (or very slightly) and rounded or pointed at the tip. Dorsal margin of the limb is approximately 60-80% of the filaments in length. Height of antenna (on the 2nd segment) is about 15-20% of its length. All or some segments bear several hairs.

Second antenna. The exopodite is thin, shorter or about as long as the protopodite. The total length of the 2nd-9th segments of the exopodite is approximately 30-35% the length of the 1st segment. Basal segment of the endopodite is with or without hairs, and without warty surface. The bristles "c", "d" (in *A. aff. pljusnini*) and also "e" (in *A. pljusnini* and *A. cucullata* 1) of the 2nd segment in male are developed and have thick base or have only "c" and "d" undeveloped bristles (*A. aff. pljusnini*). The middle part of the right clasper is not thick. The right and left claspers are tapering towards the tip or not, with a slight terminal thickening. The 5 filaments of the 2nd segment of the endopodite are equal in length, distally somewhat widened and are slightly longer or about as long as the protopodite and exopodite.

Mandible. The epipodite is developed moderately, with or without bristle. The exopodite is represented by 1-2 (medium length) midlong setae. The 1st segment of the endopodite is armed with one dorsal and 4 ventral setae, the 2nd with 3 dorsal (one stout and 2 usual type) and 2 ventral setae. The 3rd segment bears 7 setae, of which the 1st and 3rd dorsal setae are claw-like and 2nd longer than the 1st dorsal seta. The main terminal claw-seta is shorter than the endopodite (on the dorsal side). Basale is armed with 2 anterior and 2 lateral setae, and 1 anterolateral seta, and anterior row of the long hairs near the articulation. Basal endite has toothrow with 6

distinctly separated terminal triangular teeth (plus one lateral in *A. cucullata* on Poulsen, 1969) and 2 short posterior bristles.

Maxilla. The basal segment of the endopodite bears 6 anterior and 5-6 posterior setae. The 2nd segment has 1-2 claw-setae and 4-5 usual setae.

Fifth limb. The epipodial appendage has 3 groups of 6+4(5)+4(5) plumose setae (unknown for *A. pljusnini* n. sp.). The 1st segment of the exopodite bears 7 ventral and 3 dorsal setae and the 2nd with 3 ventral and one dorsal setae. The middle claw-seta of the 3rd segment is about 60-80% the length of the exopodite (on the dorsal side) and about as long as its dorsal seta.

Sixth limb. The epipodite is armed with 3 groups of 5(6-7)+5(6)+5(6) plumose setae (16-18 total). One long plumose and one short nonplumose setae are placed on the endopodite. The 1st segment of the exopodite bears 1-2 dorsal and 3-6 ventral setae, the 2nd is with 3 ventral setae and the 3rd segment has one dorsal and one ventral setae. Two usual setae and one middle slim claw-seta are borne on the distal segment. The last claw-seta is about 60-70% the length of the exopodite (on the dorsal side). Height of the exopodite is 20-25% of the length.

Seventh limb. It is thin, prolonged (25-35% of the length). The limb is 20-45% of its longest seta in length.

Copulatory appendage. It is slightly tapering towards the tip and fairly curved at the proximal parts. The greatest height is at the proximal part and about 25-35% of the length (from tip to dorsomedial indentation). The tip is wide, blunt and with small protuberance or without it.

Caudal furca. This limb has 7-8 pairs of claws and single unpaired bristle.

Comparison

New genus sharply differs from all the other genera in having larger and more prolonged rostrum, developed right posterodorsal spine, serrated ventral and posterior margins, and distinct primary reticular and secondary tile-like (with denticles) sculptures on the shell; additional dorsal seta is borne on the 4th segment of the 1st antenna, and developed "e" bristle on the endopodite of the 2nd antenna in male (bristles "c"-"e" with very thick base); greater number of setae are placed on the 1st segment of the endopodite of the maxilla, and also plumose setae on the 1st segment (on ventral side) of the exopodite of the 5th limb and on the 2nd segment of the exopodite of the 6th limb.

This genus is somewhat close to *Archiconchoecilla* gen. nov., but has following differences (in addition to the above mentioned): larger shell; longer frontal organ pointed at the tip; shorter protopodite of the 2nd antenna and seta (setae) on the endopodite of the mandible; greater number of posterior setae on the 1st segment of the endopodite of the 5th limb and on the endopodite of the 6th and 7th limbs.

Distribution

Members of this genus were collected in deep-sea zone of the Atlantic, Indian and Southern Ocean and also West Pacific. The northernmost and southernmost latitudes are 60°N and 63°S. The known depth range of the genus is 250(300)-5000 m.

Key to Species of Genus *Archiconchoecissa* (without *A. aff. cucullata* 1)

1. Rostrum is down-curving and less than 15% of the shell in length, frontal organ is approximately as long as the first antenna (on the dorsal side)..... 2
 Rostrum is up-curving and over 20% of the shell in length, frontal organ is considerably longer than the 1st antenna 3
2. Capitulum of frontal organ is fused with stem, terminal setae of 1st antenna is twice as long as limb, seta "e" on 2nd endopodite antenna of male is developed*A. pljusnini* n. sp.
 Capitulum of frontal organ is delimited from stem, terminal setae of 1st antenna is 1.5 length of limb, seta "e" on 2nd antenna of male is lacking*A. aff. pljusnini*

3. Size is more than 2.3mm, setae "c", "d" and "e" on 2nd endopodite antenna are developed *A. cucullata*
 Size is less than 2.1mm, only "c" and "d" minute setae are on 2nd endopodite antenna *A. aff. cucullata*

***Archiconchoecissa pljusnini* Chavtur sp. nov.**

(Figs. 17-20, 21, A, B)

Archiconchoecia cucullata: Chavtur, 1977a: 142, 158, fig. 8, table 6.

Archiconchoecia cucullata: Chavtur, 1977b: table 2; 1977c: table 2.

Archiconchoecia cucullata: Chavtur, 1991: 37.

Archiconchoecia sp. nov. Chavtur, 1992: table 2.

Material examined

Holotype. N1115 - adult male, length is 2.30 mm, appendages are mounted on slide and valves preserved in alcohol. In collection of the Museum of Institute of Marine Biology, Vladivostok, Russia together (with paratypes). Type-locality. R/V "Vityaz", station 7260, 5°38'0-5°33'5N and 130°48'3-48'2E, depth 5000-4079 m, 18 March 1975.

Paratypes. 1116 - adult female (length 2.20 mm) from same sample as holotype; 1117 - adult female (length 2.25 mm) station and date as of holotype, depth 4000-3495 m.

Additional specimens - R/V "Vityaz": juvenile, length 1.50 mm, station and date as holotype, depth 4000-3495 m; adult female, length 2.30 mm, station 7229, 5°21'7-5°00'4N, 124°21'7-124°04'4E, depth 3500-2994 m, 24 February 1975; juvenile, length 1.7 mm, station 5628, 43°54' N, 149°57' E, depth 4100-3090 m, 31 August 1966.

All material including holotype and paratypes were caught by Plankton Bogorov-Rass's Net (S=1.0 m²).

Etymology

The species is named after Dr. Vladimir N. Pljusnin.

Description of adult male

Shell. (Fig. 17, A-C). Length 2.3 mm (with rostrum and posterodorsal spine). Valves are prolonged. Height of shell about 40% of the length (without posterodorsal spines), and nearly equal at the anterior and posterior parts. Rostrum is strongly developed, long, wide, down-curving and about 15% of the shell in length (without posterodorsal spines). Rostral incisure is absent. Dorsal margin is straight and ventral margin is slightly concave. Posterior and anterior margins are slightly rounded. Posterior margins of the left and right valves are asymmetrical: right valve is with long and pointed posterodorsal spine and left without it. Ventral and posterior margins are serrated. Left asymmetrical gland is located approximately at 90% of the way up the posterior margin and placed on developed tubercle, right gland is at 60% and on small tubercle respectively. There are distinct primary reticular and secondary tile-like (with denticles) sculpturing.

Frontal organ (Fig. 17, D, E). It is straight, long, narrow considerably extended beyond the curved distal segments of the 1st antenna, and is about as long as the dorsal side of the 1st antenna. Capitulum section is down-curved distally, unseparated, pointed at the tip, slightly thicker than stem and without hairs and spinules.

First antenna. (Fig. 17, D, E). The limb is 5-jointed, and with one dorsal seta on each of the 2nd and 6th segments. These setae are long, slim and armed with hairs. Long filaments are borne on the 5th segment equal in length, distally widened and pointed at the tip. Dorsal margin of the limb is approximately 80% of the filaments in length. Height of antenna (on the 2nd segment) is about 15-20% of the length. Some hairs are placed laterally on the 2nd, 3rd and 5th segments. The first

segment has several dark pigment spots.

Second antenna (Fig. 17, F-J). The exopodite is thin, barely shorter than the protopodite. The total length of the 2nd-9th of the exopodite is 30-35% of the 1st segment in length. Basal segment of the endopodite is armed with tiny hairs and without warty surface. Its bristles "a" and "b" are long, thick in the base and with hairs. The bristles "c", "d" and "e" placed on the 2nd segment are very thick in the base. The right clasper is curved, with short base and terminal thickening; the base is slightly thinner than the middle part. The left is clasper curved, without terminal thickening and shorter than the right one. The 5 filaments placed on the 2nd segment are about equal in length, distally somewhat widened, slightly pointed at the tip, barely shorter than the protopodite, and about as long as the exopodite.

Mandible (Fig. 18, A-D). The epipodite is developed moderately and without bristles. The exopodite is represented by 2 midlong setae. The 1st segment of the endopodite is with untuberos dorsal margin and armed with one dorsal and 4 long ventral setae, the 2nd segment has 3 dorsal (one stout and 2 usual type) and 2 ventral setae. The distal segment bears 7 setae, of which the 1st and 3rd dorsal setae are claw-like and 2nd longer than the 1st dorsal seta. The main terminal claw-seta is approximately 75% of the endopodite in length (on the dorsal side). Basale is armed with 2 anterior and 2 lateral setae, and one anterolateral seta, and anterior row of the long hairs near the articulation. Basal endite has tooththrow with 6 distinctly separated triangular teeth and 2 short posterior proximal bristles. Coxale endite carries out 3 closely set tooththrows. Numerous spines (or short filaments) and hairs are placed on the flat masticatory pad.

Maxilla (Fig. 18, E, F). The basal segment has 6 (all plumose) anterior and 5 (4 plumose) posterior setae. The distal segment is armed with 2 claw-setae and 4 usual setae.

Fifth limb (Fig. 19, A). The epipodite is torn. The 1st segment of the exopodite bears 7 (4 plumose) ventral and 3 (2 plumose) dorsal setae, of which distodorsal seta is approximately as long as the exopodite (on the dorsal side). The 2nd segment has 3 ventral and one dorsal setae, and the last seta is about 40% and middle claw-seta of the 3rd segment is 60% of the exopodite in length. Terminal dorsal and middle claw-setae are subequal in length. Hairs are placed on the ventral side of the 1st and 2nd segments of the exopodite.

Sixth limb (Fig. 19, B, C). The epipodial appendage has 3 groups of 5+5+6 plumose setae. One long and one short plumose setae are placed on the endopodite. The 1st segment of the exopodite bears 2 dorsal and 3 ventral plumose setae. The 2nd segment is with 3 ventral and the 3rd has one dorsal and one ventral setae. Two usual setae and one middle slim claw-seta are borne on the 4th segment. This claw-seta is 65-70% of the exopodite in length (on the dorsal side). Height of the exopodite is 20-25% of the length.

Seventh limb. It is thin and prolonged (35% of the length). The longest seta is about twice the length of the short seta and 5 times as long as the limb.

Copulatory appendage (Fig. 18, H). It is tapering towards the tip and obviously curved at the proximal part. The greatest height is at the base and is about 25% of the length (from tip to dorsomedial indentation). The tip is wide, blunt and bears a small protuberance. Penis is with a few muscles.

Caudal furca. (Fig. 18, G). This limb has 7 pairs of slim claws and single unpaired bristle.

Description of adult female. Shell (Fig. 20, A, B). Length is 2.2-2.3 mm (with rostrum and posterodorsal spines). Left and right asymmetrical glands are at 85-90 and 60-65% of the way up the posterior margin respectively.

Frontal organ (Fig. 19, D, E) and First antenna. As in male.

Second antenna (Fig. 20, C). Bristles "a" and "b" placed on the 1st segment are long, usual in the base and covered with short hairs. The 2nd segment is without "c"- "e" bristles.

Mandible (Figs. 19, F and 21, A, B). Main terminal claw-seta is about 80% of the endopodite in length (on the dorsal side).




Fig. 17 *Archiconchoecissa pljusnini* sp. nov. (male: 1115) A - lateral view of left valve of shell, B - ventral view of shell, C - sculpture of shell, D and E - frontal organ and 1st antenna, F and G - left endopodite of 2nd antenna, H - right endopodite of 2nd antenna, I - bristles "c", "d" and "e" of endopodite of 2nd antenna, J - 2nd antenna.




Fig. 18 *Archiconchoecissa pljusnini* sp. nov. (male: 1115) A - mandible, B and C - endopodite of mandible, D - basal endite of mandible, E - maxilla, F - seta of basale segment of maxilla, G - furca, H - penis.




Fig. 19 *Archiconchoecissa pljusnini* sp. nov. (male: 1116 - A-C, female: 1116 - D-F, female: 1117 - G) A - fifth maxilla, B and C - sixth limb, D and E - distal part of frontal organ, F - endopodite of mandible, G - seventh limb.




Fig. 20 *Archiconchoecissa pljusnini* sp. nov. (female: 1116 - A, B, G, H; 1117 - C-F) A - lateral view of left valve of shell, B - posterodorsal margin of the shell, C - endopodite of 2nd antenna, D - maxilla, E and F - distal and proximal parts of fifth limb, G and H - sixth limb.

Maxilla (Fig. 20, D). The 1st segment of the endopodite has 6 (one plumose) anterior and 5 (2 plumose) posterior setae.

Fifth limb (Fig. 20, E, F). As in male.

Sixth limb (Fig. 20, G, H). The 1st segment of the exopodite bears one dorsal and 3 ventral plumose setae. Middle terminal claw-seta is about 60% of the exopodite in length (on the dorsal side).

Seventh limb (Fig. 19, G) and Caudal furca. As in male.

Comparison

The new species differs from *A. cucullata* in following respects: rostrum is shorter and down-curving; frontal organ is also shorter; exopodite of the mandible is with 2 setae; 1st segment of the sixth limb bears lesser number of setae; furca has only 7 pairs of the claws and single unpaired bristle.

Distribution

This species was collected in the West Pacific between latitudes of 43°N to 5°S (Kurile - Kamchatka Trench, the Banda and Sulawesi Seas) at the depth range 3000-5000 m.

Archiconchoecissa aff. *plusnini*

Archiconchoecia aff. *cucullata*: Angel, 1993: 74, Fig. 24.

Remarks

This is new abyssopelagic species requires formal description (Angel, 1993). It is closely related to *A. plusnini*, but differs in some respects (in bracket for *A. plusnini*): the capitulum of the frontal organ in both sexes is delimited from the stem by suture (fused), terminal setae of the 1st antenna in both sexes are about one and half (twice as long as) the length of the limb, the "e"-bristle placed on the 2nd endopodite antenna in male is lacking (it is developed and have very thick base).

Distribution.

It occurs at 20°-49°N in deep water in the Atlantic (Angel, 1993).




Fig. 21 *Archiconchoecissa plusnini* sp. nov. (female: 1117) A and B - tooththrows and masticatory pad of coxa of mandible.

Archiconchoecissa cucullata (Brady, 1902)

Conchoecissa cucullata Brady, 1902: 191, 199, pl. 24, figs. 1-8; 1903: 338, 339; Cleve, 1905: 131.

Archiconchoecia cucullata: Müller, 1906: 44, Taf. 7, Fig. 7-12, 18; 1908: 63; 1912: 56; Fowler, 1909: 256, 279, 296; Skogsberg, 1931: 4; Granata, Caporiacco, 1949: 18; Leveau, 1967: 67; Deevey, 1968: 23-25, fig. 5; 1971: 231; 1974: 355, 358; 1978a: 45, 46; 1978b: 54, 59; 1982: 134, 136; 1983: 411; Deevey, Brooks, 1980: 59, 61-62, tab 2, 3; Hillman, 1969: 30; Poulsen, 1969: 15-21, figs. 3-5; Angel, 1969a: 519, 540; 1969b: 78, 79; 1979: 79-81, fig. 71; 1981: 555, fig. 194 (4A-H); 1983: 552, 554; 1984: 321; 1993: 70, Figs. 22; Angel and Fasham, 1975: 718, 719, 720, 721, 725, 728, 730, 738; Angel *et al.*, 1982: 300, 307; 1993: 70, Fig. 22; Fasham *et al.*, 1974: 98; Alcaraz, 1977: 3, 6 10; Hanai *et al.*, 1980: 53; Drapun, 1981: 75; 1983: 34; 1988: 49; Chen *et al.*, 1983: 92-93, fig. 10; Ellis, 1985: 929, 942; Yin, 1991: 77; Chen and Lin, 1995: 41, fig. 46.

Not *Archiconchoecia cucullata*: Chavtur, 1977a: 142, 158, fig. 8, tab. 6.

Not *Archiconchoecia cucullata*: Chavtur, 1977b: table. 2; 1977c: table. 2.

Not *Archiconchoecia cucullata*: Chavtur, 1991: 37.

Material examined - R/V "Lomonosov", 1959: adult female (length 2.2 mm) and juvenile (1.5 mm), station 466, 22°19' N and 61°02' W, depth 2600-0 m, 2 October: adult female (1.8 mm), station 505, 31°23' N and 16°48' W, depth 1618-0 m, 31 October; adult female (1.8 mm), station 508, 34°53' N 14°52' W, depth 1620-0 m, 2 November (collected by Plankton Conic Net, S=1.0 m²).

Distribution

This species inhabits Atlantic Ocean in latitudes range of 60°N to its south boundary (NE Atlantic - Brady, 1902; 1903; Müller, 1906; 1908; 1912; Folwler, 1909; Skogsberg, 1931; Granata, Caporiacco, 1949; Poulsen, 1969; Angel, 1969a; 1969b; 1979; 1981; 1983; 1984; Angel, Fashim, 1975; Angel *et al.*, 1982; Fasham *et al.*, 1974; Alcaraz, 1977; Ellis, 1985. NW Atlantic - Skogsberg, 1931; Granata, Caporiacco, 1949; Deevey, 1968; 1971; 1978b; 1982; Deevey, Brooks, 1980; Poulsen, 1969; Angel, 1979; 1981. S Atlantic - Müller, 1908; Deevey, 1974; Angel, 1981; Drapun, 1981; 1983; 1988), Indian Ocean between 7°N and its south boundary (Cleve, 1905; Müller, 1906; 1908; Leveau, 1967; Poulsen, 1969; Hanai *et al.*, 1980), Pacific Ocean between Paraselsky Islands (South China Sea) and 5°S (NW Pacific-Poulsen, 1969; Hanai *et al.*, 1980; Chen *et al.*, 1983; Chen and Lin, 1995. S Pacific-Poulsen, 1969) and South Ocean from its north boundary to 63°S (Atlantic sector - Müller, 1906; Hillman, 1969; Poulsen, 1969; Deevey, 1974. Indian sector - Müller, 1908; Hillman, 1969; Poulsen, 1969; Deevey, 1982. Pacific sector - Hillman, 1969; Deevey, 1978a; 1982, 1983). *A. cucullata* is known in the depth range of 250-300 to 3000-4000 m, but usually bellow 500 m. Most numerous were caught in the level 500-1000 m (Angel, 1979) and 2000-2500 m (Poulsen, 1969).

Archiconchoecissa aff. *cucullata* 1

Archiconchoecia aff. *cucullata* Angel, 1983: 553 (tab.1).

Remark

Angel's publication contains no morphological information for this species.

Distribution

Collected only in the region 42°N-17°W from levels 3500-3700 and 3700-3900 m.

Archiconchoecissa aff. *cucullata* 2

Archiconchoecissa aff. *cucullata*: Angel, 1993: 72 (part), Fig. 23

Remarks

This is closely related to *A. cucullata*, but differs in following: the carapace is noticeably shorter, the endopodite of the 2nd antenna is bare and armed with (in male) minute "c" and "d" bristles ("e"-bristle is lacking).

Distribution

This is more abundant than *A. cucullata* and sometimes sub-dominant at deep mesopelagic depths at lower latitudes (Angel, 1993: location is not reported).

Archiconchoecemma Chavtur gen. nov.

Type-species - *Archiconchoecia simula* Deevey, 1982

Composition

This genus contains only 2 species: *A. simula* (Deevey, 1982) and *A. orientalis* (Chavtur, 1987).

Description

Shell. Males range from 1.3 to 1.4 mm in length, and female from 1.4 to 1.44 mm. Valves are slightly prolonged. Height of shell in male and female is approximately 55 and 60% of the length respectively. Greatest height is at the posterior part. Rostrum is slightly developed, and rostral incisure is absent. Dorsal and ventral margins are straight or nearly straight. Anterior and posterior margins are evenly rounded and symmetrical. Posterior margin is without tubercles and points. Left and right asymmetrical glands are about 85-90 and 65-70% of the way up the posterior margin respectively. No sculpturing is noted on the shell or there is slight indication of sculpturing, as lines approximately paralleling the dorsal margin and near anterior and posterior margins.

Frontal organ. It is slightly down-curved, short, narrow and barely extended beyond the down-curved distal segments of the 1st antenna. Organ in male and female is approximately 75-95% and 85-90% the length of the dorsal side of the 1st antenna, respectively. Capitulum section is down-curved distally, unseparated, pointed at the tip, and with tiny hairs at its posterior part.

First antenna. The limb is 6-jointed, with one dorsal seta placed on the 2nd segment. The seta is short, slim and armed with hairs. The 5th and 6th segments are with 6 long filaments unequal in length, distally not widened and rounded or pointed at the tip. Dorsal margin of the limb is 75-80% of the filaments in length. Height of antenna (on the 2nd segment) is 15-20% of the length. Hairs on the 4th segment are absent.

Second antenna (male of *A. orientalis* unknown). The exopodite is thin and nearly as long as the protopodite. The total length of the 2nd-9th segments of the exopodite is approximately 30-35% of the 1st segment in length. Basal segment of endopodite is armed with spinules around the bases of the bristles. The 2nd segment is covered with long hairs in female and has short bristles "c" and "d" in male. The right clasper is hook-like, sharply curved and slightly widened in the middle. The left clasper is quite small and thumb-like. Claspers do not taper distally. The 5 filaments placed on the 2nd segment are distally somewhat widened towards the tip and are unequal in length (2 proximal filaments are longer than the other 3).

Mandible. The epipodite is developed and with short bristle on the thick base (unknown for *A. simula*). The exopodite is represented by 2 long setae (3 in female of *A. simula*). The 1st segment of the endopodite has one dorsal and 4 ventral setae and the 2nd segment has 3 dorsal and 2 ventral setae. The 3rd segment bears 7 setae, of which 1st and 3rd (3rd and 4th in *A. orientalis*) dorsal setae are claw-like, 2nd seta is very slight and considerably shorter than the 1st dorsal seta, and main claw-seta is longer than the endopodite (on the dorsal side) in female and shorter in male. Basale is armed with 3-4 anterior and 2 lateral setae and 1 anterolateral seta near the articulation. Basal endite has toothrow with 6 distinctly separated triangular teeth (plus one lateral tooth in *A. orientalis*).

Maxilla. The basal segment of the endopodite bears 6 anterior and 4-6 posterior setae. The distal segment is armed with 2 claw-setae and 3 usual setae.

Fifth limb. The epipodite has 3 groups of 4+4+4(5) plumose setae. The 1st segment of the exopodite bears 7 ventral and 3 dorsal setae and 2nd segment has 2 ventral and one dorsal setae. The middle claw-seta of the 3rd segment is about 65-70% of the exopodite in length (on the dorsal side) is slightly longer than its dorsal seta.

Sixth limb. The epipodial appendage has groups of 5+5+5(6) plumose setae. One long plumose seta is placed on the endopodite. The 1st segment of the exopodite bears 2 dorsal and 5 ventral setae, the 2nd segment has only one ventral and 3rd - one dorsal and one ventral setae. Two usual setae and one middle slim claw-seta are located on the distal segment. This claw-seta is approximately 80% of the exopodite in length (on the dorsal side). Height of the exopodite is 30-45% of the length.

Seventh limb. It is very thin (torn in *A. orientalis*), prolonged (about 10% of the length). The limb is 30% the length of the longest seta.

Copulatory appendage (known only in *A. simula*). It is tapering towards the base and the tip from the middle, and is rounded at the tip. The greatest height is at the middle and approximately 40% of the length (from tip to dorsomedial indentation).

Caudal furca. This limb has 7-8 pairs of claws and single odd bristle.

Comparison

New genus differs from all the other genera by greater number of long setae placed on the exopodite of the mandible, and very short slight seta on the 3rd segment of its endopodite.

This genus is close to *Archiconchoecia* Müller, 1894 in shape, sculpture, and location of the asymmetrical glands of the shell. But it is separated (in addition to the above said) by some larger and more prolonged shell without dorsomedial indentation. Besides, its frontal organ is pointed at the tip, protopodite and exopodite of the 2nd antenna are equal in length, right clasper is of usual type (without thickening in the middle), the 1st segment on the exopodite of the 6th limb and furca are armed with more of setae and claws, respectively.

Distribution

Members of the new genus have been taken in a deep-sea zone of the Southern Ocean, North Pacific and North Atlantic. The northernmost and southernmost latitudes are 37°N-72°S. The known depth range of the genus is 1015-4000 m.

Key to Species of Genus *Archiconchoecemma* (Adult male and female)

1. Shell has linear sculpture, 1st segment of endopodite of maxilla bears 6 posterior setae, furca has 7 pairs of claws *A. orientalis* (Chavtur)
Shell without noted sculpture, 1st segment of endopodite of maxilla bears 4 posterior setae, furca has 8 pairs of claws *A. similia* (Deevey)

***Archiconchoecemma orientalis* (Chavtur, 1987)**

(Figs. 22-24)

Archiconchoecia orientalis Chavtur, 1987: 943-945, fig.1, 1-8; 3, 1-3; 1991: 43.

Archiconchoecia orientalis Chavtur, 1992: table 2 (list).

Material examined

Holotype. N1118 - (N18815 - obsolete number) - adult female, length 1.4 mm, appendages are mounted on slide and valves conserved in alcohol. In collection of the Museum of Institute of Marine Biology, Vladivostok, Russia. Type-locality - R/V "Vityaz", station 6151, 37°38'6-38'48N, 143°51'

5-52°6E, depth 2500-2000 m, 29 June 1969 (Plankton Bogorov-Rass' s Net, S=0.5 m²).

Redescription of adult female.

Shell (Fig. 22, A, B). Length 1.4 mm. Valves are slightly prolonged. Height of shell approximately is 60% of the length. Highest at the posterior part. Rostrum very small, and rostral incisure absent. Dorsal and ventral margins are nearly straight and barely concave at the medial part. Anterior and posterior margins are evenly rounded. Left and right asymmetrical glands are about 85-90% and 65-70%, of the way up the posterior margin, respectively. Shell has slight indication of sculpturing, as lines, approximately paralleling the dorsal margin and near anterior and posterior margins.

Frontal organ (Fig. 22, C). It is curved, short, subequal thickness throughout, barely extending beyond the down-curving distal segments of the 1st antenna and about 75% of the 1st antenna in length (of the dorsal side). Capitulum section is down-curved, unseparated, pointed at the tip and has tiny hairs at its posterior part.

First antenna (Fig. 22, A). The limb is 6-jointed. Dorsal seta of the 2nd segment is short, slim and armed with hairs. Long filaments on the 5th and 6th segments are unequal in length: 2 most distal filaments being somewhat shorter than others. One of proximal filaments is slightly widened in distal and pointed at the tip, and the other 5 filaments are unwidened and rounded respectively. Dorsal margin of the limb is about 70% of the filaments in length. Height of antenna (on the 2nd segment) is 20% less (17%) than the length. The 1st and 2nd segments are covered with lateral hairs. The 1st segment has some dark pigment spots.

Second antenna (Fig. 22, D). The exopodite is thin and about as long as the protopodite. The total length of the 2nd-9th segments is about 30% of the 1st segment in length. Basal segment of the endopodite is armed with long hairs, its bristles "a" and "b" are short. The 2nd segment has long hairs and without "c"-"e" bristles. The 5 filaments are slightly widened distally and unequal in length (2 proximal filaments are longer than the other 3).

Mandible (Figs. 22, E-G, 23, A, E, F and 24, A). The epipodite is developed and has one short bristle placed on the thick base. The exopodite is represented by 2 long setae are armed with long proximal and short distal hairs. The 1st segment of the endopodite has untuberous dorsal margin and covered with one dorsal and 4 ventral setae; the 2nd segment has 3 dorsal (1st dorsal is claw-like) and 2 ventral setae. The 3rd segment bears 7 distal setae, of which 1st, 3rd and 4th setae from the dorsal side are claw-like and 2nd seta is about 1/3 of the 1st seta in length. Basale is armed with 3-4 anterior and 2 lateral setae and one anterolateral seta near the articulation. Basal endite has toothrow of 6 distinctly separated terminal and one lateral triangular teeth and 2 short posterior bristles. Coxal endite is armed with 3 closely set tooththrows and with tiny hairs. The flat masticatory pad is covered with numerous varisized filaments.

Maxilla (Fig. 23, B). The basal segment of the endopodite bears 6 plumose anterior and 6 (2 plumose) posterior setae. The 2nd segment has 2 claw-setae and 3 usual setae.

Fifth limb (Fig. 23, C). The epipodite has 3 groups of 4+4+5 plumose setae. The 1st segment of the exopodite bears 7 ventral and 3 dorsal setae, of which distodorsal seta is about 90% the length of the exopodite (on the dorsal side). The 2nd segment has 2 ventral and one dorsal setae, the last seta is 55% and middle claw-seta of the 3rd segment is about 70% of the exopodite in length, respectively. This claw-seta is barely longer than the dorsal seta its segment. Long hairs are placed only on the ventral surface of the 1st segment.

Sixth limb (Fig. 23, D). The epipodial appendage is covered with 15 plumose setae. Only one long plumose seta is located on the endopodite. The 1st segment of the exopodite has 5 ventral and 2 dorsal setae, the 2nd segment has one ventral and the 3rd has one ventral and one dorsal setae. Two usual setae and one midlong claw-seta are placed on the distal segment. This claw-seta is about 90% and distodorsal seta of the 1st segment is approximately 40% the length of the exopodite (on the dorsal




Fig. 22 *Archiconchoecemma orientalis* (female: 18815) A, B - lateral view of right and left of shell, C - frontal organ and 1st antenna, D - endopodite of 2nd antenna, E - mandible, F and G - tooththrows and masticatory pad of coxa of mandible.




Fig. 23 *Archiconchoecemma orientalis* (female: 1118) A - basal endite of mandible, B - maxilla, C - fifth limb, D - sixth limb, E and F - epipodial bristle of mandible.

side). Height of the exopodite is 30-35% of the length.

Seventh limb. It is lost.

Caudal furca. This limb has 7 pairs of claws and single unpaired bristle.

Male. It is unknown.

Comparison

A. orientalis nov. sp. differs from another species of this genus - *A. simula* in having a sculpture on the shell, greater number of posterior setae on the 1st segment of the mandible endopodite, longer distodorsal seta on the 1st segment and longer dorsal seta on the 2nd segment of the 5th limb, thinner of the 6th limb and less number of claws on the furca.

Distribution

New species was captured in the North Pacific (37°38' 6-43' 0N and 143°51' 5-52' 6E) between 2000 and 2500 m.

Archiconchoecemma simula (Deevey, 1982)

Archiconchoecia simula Deevey, 1982: 137-140, figs. 6-12, tab. 1, 2; 1983: tab.1.

Archiconchoecia n. sp. Angel, 1981: 556, 561, fig. 194 (5A-H).

Distribution

Species was described from the South Ocean. It is known from the Pacific sector between 65-72°S and 160-180°E in depth range 1015-1829 m (Deevey, 1982) and from the Atlantic sector in the region 62-69°N and 75-85°W in depth 2196-2562 m (Deevey, 1983). Besides that, this species have been taken in the North Atlantic (10°30N-20°0W and 20°0N-21°0W) between 1500-2000 and 2500-4000 m (Angel, 1981; Deevey, 1982).

Archiconchoecinna Chavtur gen. nov.

Archiconchoecia Müller, 1912: 55 (part); Deevey, 1968: 22 (part).

Type species - *Archiconchoecinna ecuneata* Chavtur, new species




Fig. 24 *Archiconchoecemma orientalis* (female: 1118) A - tooththrows and masticatory pad of coxa of mandible.

Composition

This genus contains 5 species: *A. ecuneata* n. sp., *A. cuneata* (Müller, 1908), *A. arctica* n. sp., *A. aff. cuneata* 1 (*A. cuneata* in Deevey, 1968) and *A. aff. cuneata* 2 (*A. cuneata* in Deevey, 1978).

Description

Shell. Males range from 0.72 to 1.3 mm in length, and females from 0.82 to 1.25 mm. Valves are slightly (considerably in *A. cuneata* - Müller, 1908) prolonged. Height of shells in male and female are 58-60% (52% in *A. arctica* n. sp.) and 50-60% of the length respectively. Greatest height of shell in male is at the anterior part and in female is at the posterior or subequal at both parts. Rostrum is developed and wide, and rostral incisure is not deep (deepened in *A. cuneata* - Müller, 1908). Dorsal margin is straight (slightly slopes downward at the posterior part in male *A. aff. cuneata* 2). Ventral and anterior margins are rounded. Posterior margin about 1/3-1/4 of the way down is straight, the rest of margin is slightly rounded and slopes downward to anteroventral side. Posterior margin is without tubercles and points. Left asymmetrical gland is located near posterodorsal corner (at corner in *A. cuneata* - Müller, 1908), and right gland is at 1/2 or above of the shell height on the posterior margin. The shell has not visible sculpturing.

Frontal organ. It is slightly down-curved, short, barely extended beyond the down-curving distal segments of the 1st antenna in length. Organ in male and female are 60-70% and 80-90% the length of the 1st antenna on the dorsal side respectively. Capitulum section is not separated, is rounded or nearly rounded at the tip and without hairs and spinules.

First antenna. The limb is 6-jointed, with one dorsal seta placed on the 2nd segment. This seta is long, stout and coarsely spinous. Long 6 filaments placed on the 5th and 6th segments are equal in length, widened and pointed at the tip (distally not widened and rounded only in *A. cuneata* - Müller, 1908, Taf. 8, Fig. 4). Dorsal margin of the limb in the male and female are approximately 60-70% and 70-90% the length of filaments respectively. Height of antenna (on the 2nd segment) is about 20% (17-22%) of the length. Hairs on the limb are missing (5th segment is covered with hairs in *A. aff. cuneata* 2 - Deevey, 1978c: 112).

Second antenna. The exopodite is shorter or about as long as the protopodite. The total length of the 2nd-9th segments of the exopodite is approximately 50-70% of the 1st segment in length (35% in *A. arctica* n. sp.). Basal segment is without warty surface and with or without hairs. The 2nd segment is not covered with hairs, in male there are bristles "c" and "d" (without bristles in *A. arctica* n. sp.). The right clasper is narrow in the middle of the way (not narrow in *A. arctica* n. sp.). The right and left claspers are not distally tapering and have terminal thickening. The 5 filaments placed on the 2nd segment are distally widened or equal in thickness throughout, equal or unequal in length, pointed (rounded in *A. arctica* n. sp.) at the tip and longer than the exopodite and about as long as the protopodite.

Mandible. The epipodite has no bristle. The exopodite is represented by very short spinous bristle (in *A. ecuneata* n. sp.) or without it. The 1st segment of the endopodite bears one dorsal and 4 (2 in *A. arctica* n. sp.) ventral setae, the 2nd segment has 3 dorsal and 2 ventral setae. The 3rd segment has 7 setae, of which the largest is claw-like and 1st dorsal seta shorter than the 2nd dorsal seta (both setae are slight). Main terminal claw-seta is shorter than the endopodite (on the dorsal side). Basale is armed with 2 anterior and 2 lateral setae, one anterolateral seta and anterior long hairs near the articulation. Basal endite has toothrow of 6 distinctly separated serrated teeth (with or without lateral tooth), and 2 short posterior setae.

Maxilla. The basal segment of the endopodite bears 6 (5 on one limb of male *A. ecuneata* n. sp.) anterior and 5 posterior setae. The distal segment is armed with 2 claw-setae and 3 usual setae.

Fifth limb. The quantity of setae placed on the epipodite is unknown. The 1st segment of the exopodite has 5-7 ventral and 2-3 dorsal, and the 2nd segment has 2 ventral and one (absent in *A. arctica* n. sp.) dorsal setae. The middle claw-seta of the 3rd segment is about 30-40% (60% in *A. arctica* n. sp.) of the exopodite in length (on the dorsal side) and shorter or subequal in length to the

dorsal claw-setae.

Sixth limb. The epipodial appendage has 3 groups of 5+5+6 plumose setae (known only for *A. ecuneata* n. sp.). The endopodite bears 2 (1-2 plumose) setae. The 1st segment of the exopodite bears 0-2 dorsal and 5 ventral setae, the 2nd segment has only one ventral and the 3rd segment has one dorsal (claw-like) and one ventral setae. The 4th segment has dorsal and middle claw-setae and ventral usual seta. The middle claw-seta is 13-23% (43% in *A. arctica* n. sp.), and distodorsal seta (absent in *A. arctica* n. sp.) of the 1st segment is 10-25% in male and 15-35% in female of the exopodite in length (on the dorsal side). Height of the exopodite is 30-40 in male and 35-40% in female of the length.

Seventh limb. It is unknown.

Copulatory appendage. It is tapering towards the tip. The greatest height is at the distal half or only at the base and is about 40-50% of the length (from tip to dorsomedial indentation). The tip is pointed and upward (rounded and not upward in *A. arctica* n. sp.).

Caudal furca. This limb has 8 (7 in *A. arctica* n. sp.) pairs of claws. Furca is with (without in *A. arctica* n. sp.) single unpaired bristle.

Remarks. *A. arctica* nov. sp. sharply separates from the description of *Archiconchoecinna* gen. nov. This species differs in many respects from other known species: "c" and "d" setae are lacking and the left clasper is short and thick on the endopodite of the 2nd antenna in male; the endopodite of mandible is only with 2 ventral setae placed on the 1st segment and without dorsal seta on the 2nd segment; longest terminal claw-setae of the 6th limb are longer, penis is not upward and is rounded at the tip. However, *A. arctica* nov. sp. is related to *Archiconchoecinna* in having shell "cuneata" - shaped of shell and typical location of the asymmetric glands, the exopodite of the mandible has no seta, serrated teeth of the basale are distinctly separated; the 6th limb is armed with one on the 1st segment and 2 dorsal claw-setae on the 4th segments.

Comparison

New genus differs from all the other in following: exopodite of the mandibular has not seta, teeth on the basal endite of its limb are distinctly separated and serrated middle claw on the distal segment of the 5th and 6th limbs very short; and dorsal claw (or claw-like seta) on the 3rd segment of the exopodite of the 6th limb is also very short.

This genus is close to *Archiconchoecilla* gen. nov., but distinct (in addition to the above side) on the following respects: the shell is higher in the anterior part; the basal endite of the mandible is armed with serrated teeth; greater number of the setae are placed on the 1st segment (posterior side) of the endopodite of the maxilla and on the endopodite of the 6th limb; distodorsal seta on the 1st segment of the 5th limb and middle seta on the distal segment of the 6th limb are shorter.

Shell shape of *Archiconchoecinna* gen. nov. is close to *Archiconchoecilla* gen. nov., but differs from it by having symmetrical left and right valves, invisible sculpturing, and shorter (to the length of the protopodite) exopodite on the 2nd antenna.

Distribution

Members of this genus were collected in a deep-sea zone of the tropical-subtropical Atlantic, boreal Pacific and central Arctic. The known depth range of the genus is 100 (200)-3800 m.

Key to Species of Genus *Archiconchoecinna* (Adult male and female)

1. Terminal filaments of the 1st antenna are widened distally 2
 Terminal filaments of the 1st antenna are not widened distally 4
2. Terminal claw-setae of the 6th limb are less than 25% of the exopodite in length (on the dorsalside), furca has 8 pairs of claws and single unpaired bristle 3
 Terminal claw-setae of the 6th limb exceed 40% of the exopodite in length, furca has 7pairs of

- claws and lacks unpaired bristle *A. arctica* n. sp.
3. Shell length is over 1mm, right asymmetrical gland is more half way up the posterior margin, exopodite of mandible has minute bristle and basal endite has lateral tooth, 1st segment of the exopodite of 6th limb bears one lateral seta *A. ecuneata* n. sp.
Shell length is less than 1mm, right asymmetrical gland is approximately half way up the posterior margin, exopodite of mandible is without bristle and basal endite without lateral tooth, 1st segment of exopodite of 6th limb bears 2 lateral setae *A. aff. cuneata* 2.
4. Left asymmetrical gland is located at the posterodorsal corner, bristle "b" of 2nd antenna is shorter than its 1st segment and is about 20% length of terminal filaments *A. cuneata* (Müller).
Left asymmetrical gland is located near posterodorsal corner, bristle "b" of 2nd antenna are longer than its 1st segment and is over 50% length of terminal filaments *A. aff. cuneata* 1

***Archiconchoecinna ecuneata* Chavtur sp. nov.**

(Figs. 25-28 and 31, A, B)

Archiconchoecia cuneata: Chavtur, 1992: table 2 (list).

Material examined

Holotype. N1119 - adult male, length 1.1 mm, appendages are mounted on slide and valves remained in alcohol. In collection of the museum of Institute of Marine Biology, Vladivostok, Russia (together with paratype). Type-locality - R/V "Vityaz", station 6111, lat, 56°16' 6-12' 0N, 138°20' 0-137°17' 8W, depth 1500-1000 m, 16 May 1966.

Paratype. N1120 - adult female, length 1.25 mm; appendages are mounted on slide and valves remained in alcohol, from sample as in holotype.

Additional specimens - R/V "Vityaz", 1969: adult male (length 1.3 mm), station 6118, 56°26' 9-28' 0N, 137°14' 9-136°56' 5W, depth 1000-500 m, 18 May; 2 adult males (1.15 and 1.15 mm), station 6139, 53°11' 5-21' 5N, 163°09' 0-162°56' 0W, depth 2000-1500 m, 12 June; juvenile (1.05 mm), station 6144, 51°42' 0N, 167°55' 0W, depth 500-300 m, 17 June.

All material (including holotype and paratype) were caught by Plankton Bogorov-Rass's Net (S=0.5 m²).

Etymology

The specific name "*ecuneata*", from Latin "e" [=no] and "*cuneata*" [=narrowed], refers to closeness, but not identity to species *A. cuneata* (Müller, 1908).

Description of adult male

Shell (Fig. 25, A). Length in 1.1-1.3 mm. Valves are slightly prolonged. Height of shell is about 60% of the length. Greatest height is at the anterior part. Rostrum developed and wide, and rostral incisure is not deepened. Dorsal margin is straight and ventral and anterior margins rounded. Posterior margin about 1/4 of the way down is straight, the rest of margin is slightly rounded and slopes downward to anteroventral side. This margin is without tubercles and points. Left asymmetrical gland is located near posterodorsal corner and right gland is at 1/2 or above of the height on the posterior margin. Sculpturing on shell is invisible.

Frontal organ (Fig. 25, B, C). It is slightly curved, short, barely beyond the down-curving distal segments of the 1st antenna and is approximately 60% of the 1st antenna in length (of the dorsal side). Capitulum section is somewhat thicker than stem, down-curved, rounded at the tip and without hairs and spinules.

First antenna (Fig. 25, B, C). The limb is 6-jointed. Dorsal seta of the 2nd segment is long, stout and coarsely spinous. The 5th and 6th segments bear 6 long filaments equal in length, distally widened and pointed at the tip. Dorsal margin of the limb is approximately 60% of the filaments length. Height of antenna (on the 2nd segment) is about 20% of its length. Hairs on the limb are absent. The 1st segment has some dark pigment spots.

Second antenna (Fig. 25, D-F). The exopodite is thick and shorter than the protopodite. The total length of the 2nd-9th segments of the exopodite is approximately 50% length of the 1st segment. Basal segment of the endopodite is without hairs and warty surface; the bristle "a" is thin, short and is 35-45% of the bristle "b" in length. The bristles "c" and "d" of the 2nd segment are thin and long. The right clasper is narrow, curved, with short base (its thickness is same as in the middle part) and with slight terminal thickening. The left clasper is similar to the right but shorter than it. Claspers are rounded at the tip. The 5 filaments borne on the 2nd segment are distally widened, unequal (?) in length, pointed at the tip and longer than the exopodite and protopodite.

Mandible (Figs. 26, A, B and 31, A, B). The epipodite is slightly (?) developed and without bristle. The exopodite is represented by very short spinous bristle. The 1st segment of the endopodite has straight dorsal margin and is armed with one dorsal and 4 ventral setae, the 2nd segment with 3 dorsal and 2 ventral setae. The 3rd segment bears 7 setae, of which largest seta is claw-like and about 70% of the endopodite in length (on the dorsal side), 2 first dorsal setae are slight (the 1st shorter than the 2nd seta). Basale is armed with 2 anterior, 2 lateral, 1 anterolateral setae, and anterior row long hairs near the articulation. Basal endite has toothrow with 6 distinctly separated terminal teeth (5 serrated and one proximal bare triangular) and lateral tooth (triangular) tooth and 2 short posterior bristles. Coxale endite is armed with 3 closely set toothrows and tiny hairs. The masticatory surface is especially protuberant and bears rather large pointed denticles as well as bristles and with numerous long hairs.

Maxilla (Fig. 26, C, D). The basal segment of the endopodite bears 5-6 (all plumose) anterior and 5 (one plumose) posterior setae; plumose setae are armed with long proximal and short distal hairs. The 2nd segment has 2 claw-setae and 3 (one long and 2 short) usual setae.

Fifth limb (Figs. 26, E and 27, A). The epipodite is in poor condition. The 1st segment of the exopodite bears 5 ventral and 3 (2 plumose) dorsal setae, of which distodorsal seta is about 50% of the exopodite in length (on the dorsal side). Second segment is with 2 ventral and one dorsal setae, last seta is 40-45% and middle claw-seta of the 3rd segment is 35-40% of the exopodite in length. The terminal dorsal claw-seta is longer than the middle one. Ventral surface of the 1st and 2nd segments is covered with hairs.

Sixth limb (Fig. 27, B, C). The epipodial appendage has 3 groups of 5+5+6 plumose setae. Two long plumose setae are placed on the endopodite. The 1st segment of the exopodite bears one very short (bare) dorsal and 5 (4 plumose) ventral setae. The 2nd segment is with one ventral and the 3rd with one dorsal spinous claw-seta and one ventral ordinary seta. The 4th segment has one dorsal and one middle short claw-setae and one ventral very short usual seta, of which dorsal claw-seta is largest. Middle claw-seta of its segment is about 20% and distodorsal seta of the 1st segment is 10-15% of the exopodite in length (on the dorsal side).

Height of the exopodite is about 40% of the length.

Seventh limb. It is lost.

Copulatory appendage (Fig. 25, G). Greatest height is at the distal half and is approximately 50% of the length (from tip to dorsomedial indentation). The tip is pointed and upturned. There is wedge-like muscle directed from the base to a pointed tip.

Caudal furca. This has 8 pairs of slim claws and single unpaired bristle.

Descriptions of adult female

Shell. It is in poor condition. Length is 1,25 mm.

Frontal organ (Fig. 27, D). It is longer than in male and is about 80% of 1st antenna in length (on the dorsal side).

First antenna. (Fig. 27, D). Limb is about 70% of the terminal filaments in length. Height of antenna (on the 2nd segment) is barely over 20% of the length.

Second antenna (Fig. 27, E, F). Terminal filaments are approximately equal and as long as the protopodite.




Fig. 25 *Archiconchoecinna ecuneata* sp. nov. (male: 1119) A - lateral view of shell, B and C - frontal organ 1st antenna, D - 2nd antenna, E and F - right and left endopodites of 2nd antenna, G - penis.




Fig. 26 *Archiconchoecinna ecuneata* sp. nov. (male: 1119) A - mandible B - basal endite of mandible, C and D - maxilla, E - proximal part of fifth limb.




Fig. 27 *Archiconchoecinna ecuneata* sp. nov. (male: 1119 - A-C, female: 1120 - D-G) A - fifth limb, B and C - sixth limb, D - frontal organ and 1st antenna, E - 2nd antenna, F - endopodite 2nd antenna, G - distal arming of 1st segment of endopodite maxilla.




Fig. 28 *Archiconchoecinna ecuneata* sp. nov. (female: 1120) A - mandible, B and C - maxilla, D - fifth limb, E - endopodite fifth limb, F - sixth limb.

Mandible (Fig. 28, A). Main terminal claw-seta is scarcely over 70% of the endopodite in length (on the dorsal side)

Maxilla (Figs. 27, G and 28, B, C). One seta of other 5 anterior setae placed on the basal segment is plumose.

Fifth limb (Fig. 28, D, E). The 1st segment of exopodite bears 7 ventral and 3 dorsal setae, of which distodorsal seta is 65% of the exopodite in length (on the dorsal side).

Sixth limb (Fig. 28, F). The endopodite is with one long common and one short plumose setae. Setae placed on the 1st exopodite are nonplumose. Claw-setae of the 4th segment are thicker and claw-seta of the 3rd is thicker and longer than in male. Distodorsal seta on the 1st segment is longer and about 35% of the exopodite in length (on the dorsal side). Height of the exopodite is 45-50% of the length.

Seventh limb. It is lost.

Caudal furca. As in male.

Comparison

The new species is closest to *A. aff. cuneata* 2, but differs in many respects: shell is longer, right asymmetrical gland is located more half of the way up the posterior margin; frontal organ is somewhat shorter; the 1st segment of the exopodite on the 2nd antenna is longer than its 2nd-9th segments, terminal filaments on the endopodite of the 2nd antenna are longer (to the length of its exopodite and protopodite); the exopodite of the mandible is represented by very short spinous bristle (it is lacking in *A. aff. cuneata* 2), the basal endite on the mandible is armed with additional lateral tooth (it is missing in *A. aff. cuneata* 2); plumose setae placed on the maxilla have longer hairs; the 1st segment of the exopodite on the 6th limb bears one (2 in *A. aff. cuneata* 2) dorsal seta, the 6th limb is thicker.

Distribution

New species was captured in the Gulf of Alaska and in the region of the Aleutian Trench at water depth of 300 to 2000 m.

Archiconchoecinna arctica Chavtur sp. nov.

(Figs. 29, 30 and 31, C, D)

Archiconchoecia cuneata: Chavtur, 1978: 1791-1793.

Archiconchoecia species nov. Chavtur, 1991: 94; 1992: table 2 (list).

Holotype. N1121 - adult male, length 1.05 mm, appendages and valves are mounted on slide. In collection of the Museum of Institute Marine Biology, Vladivostok, Russia. Type-locality - Drifting Station "North Pole - 2", 81°29N and 197°16E, depth 3800-995 m, 3 April 1951 Nansen's (Plankton Net).

Etymology

The specific name from the locality of the holotype.

Description of adult male

Shell (Fig. 29, A). Length is 1.05 mm. Valves are slightly prolonged. Height of shell is about 50% of the length. Greatest height is at the anterior part. Rostrum is developed and wide, and rostral incisure is not deepened. Dorsal margin is straight and ventral and anterior margins rounded. Posterior margin is approximately 1/4 of the way down straight, the rest of margin is slightly rounded and strongly slopes downward to anteroventral side; it is without tubercles and points. Left asymmetrical gland is located near posterodorsal corner, and right gland is located above the shell height on the posterior margin. Sculpture on the shell is not visible.

Frontal organ (Fig. 29, B, C). It is slightly curved, short, and slightly beyond the down-curving distal segments of the 1st antenna (% the length of the 1st antenna is unknown since the proximal part




Fig. 29 *Archiconchoecinna arctica* sp. nov. (male: 1121) A - lateral view of right valve of shell, B - distal part of frontal organ of 1st antenna, C - distal part of frontal organ, D - dorsal seta of 1st antenna, E - 2nd antenna, F and G - endopodite 2nd antenna, H - distal part of endopodite mandible, I - fifth limb.




Fig. 30 *Archiconchoecinna arctica* sp. nov. (male: 1121) A - mandible, B - basal endite of mandible, C and D - dorsal claw-seta of 3rd segment of exopodite of sixth limb, E - maxilla, F and G - sixth limb, H - penis.

of the 1st antenna is broken). Capitulum section is rounded at the tip and without hairs and spinules (proximal part of frontal organ is torn).

First antenna (Fig. 29, B, D). The limb is 6-jointed. It has one long, stout and coarsely spinous dorsal seta on the 2nd segment. Six long filaments placed on the 5th and 6th segments are equal in length, which are distally widened and pointed at the tip (% the length and height of the filaments from the length limb and the presence of pigment spots are unknown since 1st segment is broken). Hairs on the limb are absent.

Second antenna (Fig. 29, E-G). The exopodite is thick and subequal to protopodite in length. The total length of the 2nd-9th segments of the exopodite is approximately 35-40% of the 1st segment in length. Basal segment of the endopodite is without hairs and warty surface, and the bristles "a" and "b" are thin and short ("a" is about 60% of "b" in length). There are no bristles "a"- "c" on the 2nd segment. The right clasper is stout, relatively wide, curved, with short base (its thickness as in the middle part) and with slight terminal thickening. The left clasper is also stout, but considerably shorter and with large terminal thickening. Claspers are rounded at the tip. The 5 filaments of the 2nd segment are distally widened, approximately equal in length, pointed at the tip and longer than the exopodite and protopodite.

Mandible (Figs. 29, H; 6, A, B and 31, C, D). The epipodite is developed and without bristle. The




Fig. 31 *Archiconchoecinna ecuneata* sp. nov. (male: 1119 - A, female: 1120 - B), *A. arctica* n. sp. (male: 1121 - C-D) A-D - toothrows and masticatory pad of coxa of mandible.

exopodite is also without bristle. The 1st segment of the endopodite has straight dorsal margin and is armed with one dorsal and 2 (4 in Chavtur, 1978 is not correct) ventral setae, the 2nd segment is with 3 dorsal and 2 ventral setae. The 3rd segment bears 7 setae, of which largest seta is claw-like and about 80% of the endopodite in length (on the dorsal side) and 2 first dorsal setae are slight (the 1st is shorter than the 2nd seta). Basale is armed with 2 anterior and 2 lateral setae, one anterolateral seta and anterior row long hairs near the articulation. Basal endite has tooththrow with 6 distinctly separated serrated terminal and one triangular lateral teeth and 2 short posterior setae. Coxal endite is armed with 3 tooththrows. The masticatory pad is protuberant (?) and bears small rounded terminal denticle and a row of long hairs.

Maxilla (Fig. 30, E). The basal segment of the endopodite has 6 anterior and 5 posterior nonplumose setae and also a distal row of long spines near the articulation. The 2nd segment is armed with 2 claw-setae and 3 usual (short) setae.

Fifth limb. The epipodite is torn. The 1st segment of the exopodite has 6 ventral and 2 dorsal nonplumose setae, of which distodorsal seta is about 50% of the exopodite in length (on the dorsal side). The 2nd segment has only 2 ventral setae. The middle claw-seta of the 3rd segment is approximately 60% of the exopodite in length. The terminal dorsal claw-seta is about as long as the middle one. The limb has no hairs.

Sixth limb (Fig. 30, C, D, F, G). The epipodite is in poor condition. Two long plumose are placed on the endopodite. The 1st segment of the exopodite has 5 (all nonplumose) ventral only. The 2nd segment with one ventral seta and the 3rd has one minute pectinate claw-like and one usual ventral setae. The 4th segment is armed with one dorsal and one middle long claw-setae and one usual very short ventral seta, of which dorsal claw-seta is the largest, and middle claw-seta is 40-45% of the exopodite in length (on the dorsal side). Height of the exopodite is 35% of the length.

Seventh limb. It is lost.

Copulatory appendage (Fig. 30, H). Greatest height is at the base and about 35% of the length (from tip to dorsomedial indentation). Penis is tapering from the base towards the tip. The tip is rounded and upturned. No muscles are noted on the penis.

Caudal furca. This limb has 7 pairs of claws and no unpaired bristle.

Female. Unknown.

Comparison

New species is separated from all other members of the genus (see the "Comparison" for the description of this genus). It is somewhat close to *A. ecuneata* nov. sp., but differs by longer 1st segment of exopodite on the 2nd antenna, lesser number of ventral setae are placed on the 1st segment of the mandible endopodite, absence of the dorsal seta is on the 2nd segment of the exopodite of the 5th limb and on the 1st segment of the exopodite of the 6th limb, longer terminal claw-setae are on the 5th and 6th limbs, lesser number of the claws and absence of the unpaired bristle on the furca, and also it differs by the shape of the penis.

Distribution

New species was caught in the central Arctic only in a tow from 935 to 3800 m.

Archiconchoecinna cuneata (Müller, 1908)

Archiconchoecia cuneata Müller, 1908: 63-64, Taf. 8, Fig. 1-4; 1912: 55.

Archiconchoecia cuneata: Angel, Fasham, 1975: 736; Angel, 1979: 79-80; Chen and Lin, 1994a: 448, 450.

Not *Archiconchoecia cuneata*: Chavtur, 1978: 1791-1793, fig. 2, 1-11.

Distribution

Species was described from South Atlantic (19°3S-20°0W) in a tow from 800 to 0 m. Probably it inhabits the North Atlantic between 40-18°N in the depth range 200-1250 m (Angel, 1979; Angel, Fasham, 1975) and North-West Pacific near Philippines in the deep zone (also a small amount of this species appeared in the 0-150 m layer of this area, resulting from the upwelling effects (Chen and Lin, 1994a).

***Archiconchoecinna* aff. *cuneata* 1**

Archiconchoecia cuneata: Deevey, 1968: 26, fig. 6; 1971: 231; Deevey, Brooks, 1980: 45, 60 (part).

Remarks

This species differs from *A. cuneata* by a larger height of shell, shorter rostrum, and smaller depth of rostral incisure, low position of the left asymmetrical gland; longer "b" bristle and shorter filaments placed on the epipodite of the 2nd antenna.

Distribution

The species was taken in the Sargasso Sea from 400-100 and 500-0 m.

***Archiconchoecinna* aff. *cuneata* 2**

Archiconchoecia cuneata: Deevey, 1978c: 107, 111, 112-115, figs. 3, 4.

Remarks

This species differs from *A. cuneata* in the following: larger shell height, shorter rostrum, and smaller depth of rostral incisure, low location of the left asymmetrical gland; filaments with distal widened on the 1st antenna, and longer "b" bristle on the 2nd antenna.

It differs from *A. aff. cuneata* 1 in having a distally widened filaments of the 1st antenna, and longer filaments on the endopodite of the 2nd antenna.

Distribution

Species has been taken only in the Sargasso Sea from 1000 to 500 m.

***Archiconchoecetta* Chavtur gen. nov.**

Archiconchoecia Müller, 1906: 43-44 (part); Poulsen, 1969: 14-15 (part).

Type-species - *Archiconchoecia ventricosa* Müller, 1906.

Composition

This genus consists of 11 species: *A. ventricosa* (Müller, 1906), *A. poulsenii* (Deevey, 1978), *A. fabiformis* (Deevey, 1978), *A. bimucronata* (Deevey, 1978), *A. bispicula* (Deevey, 1978), *A. bifurcata* (Deevey, 1978), *A. gastroides* (Deevey, 1978), *A. falcata* (Deevey, 1978), *A. pilosa* (Deevey, 1978), *A. bidens* (Deevey, 1982), *A. inventricosa* n. sp.

Description

Shell. Males range from 0.8 to 1.7 mm in length, female from 0.9 to 1.9 mm. Valves are slightly or fairly prolonged. Height of shell in male and female are 54-61% and 47-58% of the length, respectively. Greatest height of male is at the anterior part, and in the female it is approximately equal at the anterior and posterior parts (highest at anterior part only in female of *A. inventricosa* n. sp). Rostrum is not developed and narrow, and rostral incisure is deepened. Dorsal margin of male is straight and slopes downward to posterior end and in female it is barely swollen or straight (for *A. inventricosa* n. sp. same as in male). Anterior margin is straight and extends beyond rostrum.

Posterior margin is evenly rounded or slopes downward at the upper part, without tubercles and points. Left asymmetrical gland is located at posterodorsal corner, right gland of male is about 30-40% and female 40% of the way up the posterior margin (no indication of asymmetrical glands for *A. bidens*). Slight indication of sculpturing is, as lines parallel to the ventral margin (at an angle of 45° to this margin in *A. poulsenii*) or sculpturing invisible on the shell (*A. bifurcata*, *A. gastrodes*, *A. falcata*).

Frontal organ. It is straight or slightly down-curved, slightly or considerably extended beyond the down-curving distal segments of the 1st antenna (organ lost in *A. gastrodes*). Organ in male and female is 85-100% and 90-110% of the 1st antenna in length (of the dorsal side) respectively. Capitulum section is not separated or separated, with 1-2 thin needle-like prolongations or rounded at the tip, and with or without hairs.

First antenna. The limb is 6-jointed, with one long, stout and coarsely spinous dorsal seta on the 2nd segment. Six long filaments on the 5th and 6th segments are equal in length, which are distally widened or not and pointed or rounded (only in *A. ventricosa* - Muller, 1906) at the tip. Dorsal margin of the limb is about 50-80% of the filament in length. Height of antenna (on the 2nd segment) is about 20% (17-22%) of the length. Hairs on the limb are missing.

Second antenna. The exopodite is thin or relatively thick and shorter than the protopodite. The total length of the 2nd-9th segments of the exopodite is approximately 35-50% of the 1st segment in length. Basal segment is without warty surface. The 2nd segment is without hairs in female (with hairs in *A. fabiformis*) and bears usual bristles "c" and "d" in male. The right clasper is narrow in the middle. The right and left claspers are not distally tapering, with or without terminal thickening and rounded at the tip. The 5 filaments placed on the 2nd segment are approximately equal in thickness throughout (distally widened in *A. bidens*), equal or unequal in length, pointed at the tip (rounded in *A. ventricosa* - Müller, 1906, Taf. 7, Fig. 5), and longer than the exopodite and protopodite.

Mandible. The epipodite is without bristle. The exopodite is represented by one short or midlong seta. The 1st segment of the endopodite has one dorsal and 4 (3-4 in *A. inventricosa* n.sp.) ventral setae, the 2nd segment with 3 dorsal and 2 ventral setae. The 3rd segment bears 7 setae, of which largest seta is claw-like and 2 first dorsal setae are slight (they probably are claw-like in *A. pilosa* and *A. falcata*), the 1st dorsal seta is longer than the 2nd dorsal seta. Main terminal claw-seta is shorter, equal or longer than the endopodite in female (on the dorsal side) and shorter in male. Basale is armed with 2 lateral setae, 2 anterior (4 in *A. fabiformis*), one anterolateral setae and anterior row long hairs near the articulation. Basal endite has tooththrow with 6 (7 in *A. poulsenii*) slightly separated serrated teeth and 2 short posterior bristles.

Maxilla. The basal segment of the endopodite bears 5-6 anterior and 4-6 posterior setae. The distal segment has 3 (4 in *A. poulsenii*) usual setae and 2 claw-setae (posterior claw-seta shorter than the distal one in *A. ventricosa* and longer than in others species).

Fifth limb. The epipodite has 3 groups of 4+4+4(5) plumose setae. The 1st segment of the exopodite bears 5-8 ventral and 2-3 dorsal and 2nd segment is with 2 ventral and one dorsal setae. The middle claw-seta of the 3rd segment is 50-90% of the exopodite in length (on the dorsal side) and longer than its dorsal seta.

Sixth limb. The epipodial appendage has 3 groups of 5+5+6 in male and 4(5)+4(5)+5(6) (13-16 totally) in female plumose setae. Two (only one in *A. fabiformis*) plumose setae are placed on the endopodite. The 1st segment of the exopodite is armed with 1-2 dorsal and 4-5 ventral setae, the 2nd segment has only one ventral seta and the 3rd has one dorsal and one ventral setae. The 4th segment has 2 usual setae and one middle claw-seta (it is lacking in juv. *A. falcata*). This claw-seta is about 50-110% (20% in *A. pilosa*) in male and 50-100% in female and distodorsal seta of the 1st segment is about 15-40% in male and 25-40% in female of the exopodite in length (on the dorsal side). Height of the exopodite is 15-40% of the length.

Seventh limb. It is varying in size and shape.

Copulatory appendages. It is tapering towards the base and the tip from the middle. Highest is at the middle and about 40-50% of the length (from tip to dorsomedial indentation). The tip is rounded

(small protuberance is placed at the tip in *A. pilosa*).

Caudal furca. This limb has 7-8 pairs of claws and single unpaired bristle.

Remarks. Morphology of *A. pilosa* considerably deviates from the description of this genus in the following: left clasper of the 2nd antenna is thick and short; ventral seta on the distal segment of 5th limb is longest (middle seta in others species); distal segment of 6th limb bears 2 short claws (one in others species). This structure is such as in genus *Archiconchoecina*. However some indications point out the relationship to the new genus: character "ventricosa"-shape and sculpturing of the shell, presence of seta on the exopodite and slight separation of teeth on the basal endite of the mandible, presence of one long dorsal and one medial setae on the distal segment of the 5th limb and usual dorsal seta on the 3rd segment of the endopodite of the 6th limb.

Comparison

New genus differs from other genera in having the anterior margin extended beyond rostrum, low location of right asymmetrical gland (usually less than 1/2 of the shell height on the posterior margin), very thin claspers on 2nd antenna and slightly separated teeth on the basale endite of the mandible.

It is more close to genus *Archiconchoecia*, but differs (in addition to the above characters) by short rostrum, developed seta on the mandibular exopodite and usual (not claw-like) dorsal seta placed on the 3rd segment of the exopodite of 6th limb.

Distribution

Members of this genus have been collected in of all oceans (except the Arctic ocean) and the Sargasso and Caribbean Seas. The northernmost and southernmost latitudes are 51°N and 72°S. Most species were caught from 500 (1000) to 2000 m. Only *A. bifurcata* was taken from 500 (1000) - 3700 m and *A. inventricosa* nov. sp. - from between 3000 and 5500 m.

Key to Species of Genus *Archiconchoecetta*

- 1. Furca has 7 pairs of claws 2
- Furca has 8 pairs of claws 5
- 2. Shell length is less than 1.1 mm, 1st segment of maxilla bears 4 posterior setae 3
- Shell length is over 1.4 mm, 1st segment of maxilla bears 5-6 posterior setae 4
- 3. Frontal organ has 2 terminal needle-like extensions; 1st segment of exopodite of fifth limb bears 2 dorsal and 6 ventral setae, its terminal claw-seta is approximately 90% of exopodite in length (on the dorsal side)..... *A. bispicula* (Deevey)
- Frontal organ is with one terminal needle-like prolongations; 1st segment of exopodite of fifth limb bears 3 dorsal and 6 ventral setae, its terminal claw-seta is approximately 70% of exopodite in length *A. ventricosa* (Müller)
- 4. Shell in female is highest at the anterior part, basal endite of mandible is armed with a row of 6 teeth, 1st segment of its endopodite bears 6 anterior and 6 posterior setae; posterior claw-seta is longer than the anterior on the 2nd segment of maxilla *A. inventricosa* new species.
- Height of shell in female is equal at anterior and posterior parts, basale endite of mandible is armed with a row of 7 teeth, 1st segment of its endopodite bears 5 anterior and 5 posterior setae; posterior claw-seta is shorter than anterior one on the 2nd segment of maxilla *A. poulsenii* (Deevey)
- 5. Ventral seta placed on the distal segment of fifth limb is longest, 6th limb is with 2 short claws *A. pilosa* (Deevey)
- Medial seta placed on the distal segment of fifth limb is longest, 6th limb has one long claw 6
- 6. 1st segment of endopodite of maxilla has 9-10 setae, 1st segment of exopodite of 5th

- limb has 5 ventral setae 7
 1st segment of endopodite of maxilla has 11-12 setae, 1st segment of exopodite of 5th
 limb has 6-8 ventral setae 8
7. Frontal organ has 2 terminal needle-like prolongations; basale of mandible has 2
 anterior setae, 1st segment of endopodite of maxilla bears 9 setae *A. bimucronata* (Deevey)
 Frontal organ lacks terminal prolongations; basale of mandible has 4 anterior setae, 1st
 segment of endopodite of maxilla bears 10 setae *A. fabiformis* (Deevey)
8. Shell length is less than 1.4 mm, terminal filaments of 1st antenna are not distally
 widened *A. falcata* (Deevey)
 Shell length is over 1.5 mm; terminal filaments of 1st antenna are distally widened 9
9. 1st segment of endopodite of maxilla has 11 (5 plumose) setae, 1st segment of exopodite
 of 5th limb has 6 (4 plumose) ventral setae *A. bidens* (Deevey)
 1st segment of endopodite of maxilla has 12 (all nonplumose) setae, 1st segment of
 exopodite of 5th limb has 7-8 (one plumose) setae 10
10. Distal surface of endopodite of maxilla is armed with some spinules; only female - shell
 height is over 50% of length, main terminal claw-seta of mandible is shorter than its
 endopodite (on the dorsal side), exopodite height of 6th limb is approximately 40% of
 length *A. bifurcata* (Deevey)
 Distal surface of endopodite of maxilla is without spinules; only female-shell height is
 less 50% of length, main terminal claw-seta of mandible is longer than its endopodite,
 exopodite height of 6th limb is less 20% of length *A. gastrodes* (Deevey)

***Archiconchoecetta inventricosa* Chavtur sp. nov.**

(Figs. 32-34)

Archiconchoecia ventricosa: Chavtur, 1977a: 142, 158, fig.8, table 6, 1977b: table 2; 1977c: table 2.

Archiconchoecia aff. *ventricosa* Chavtur, 1991: 49.

Archiconchoecia sp. nov. Chavtur, 1992: table 2 (list).

Material examined

Holotype. N1122 - adult female, length 1.9 mm, appendages are mounted on slide and preserved valves in alcohol. In collection of the Museum of Institute of Marine Biology, Vladivostok, Russia (together with paratype). Type-locality - R/V "Vityaz", station 6144, 51°42N, 167°55W, depth 5500-3300 m, 17 April (Plankton Bogorov-Rass' s Net, S = 0.5 m²).

Paratype. 1123 - adult female, length 1.5 mm, R/V "Vityaz", station 5621, 45°06N, 155°55E, depth 4000-3000 m, 17 August, Plankton Bogorov-Rass' s Net, S = 1.0 m² (appendages are mounted on slide and valves remained in alcohol).

Etymology

The species name "*inventricosa*", from the Latin "*in*" [=no] and "*ventricosa*" [big - bellied], refers to closeness, but not identity with species *A. ventricosa* (Müller, 1906).

Description of adult female

Shell (Fig. 32, A). Length is in 1.5-1.9 mm. Valves are slightly prolonged. Height of shell is 58-60% of the length. Greatest height is at the anterior part. Rostrum is not developed and narrow, and rostral incisure deepened. Dorsal margin is almost straight and slopes downward to posterior end. Ventral margin is straight or barely concave. Anterior margin is rounded and considerably extended beyond rostrum. Posterior margin is evenly rounded. Left asymmetrical gland is located at posterodorsal corner, right gland is approximately at 30% of the way up the posterior margin. Slight indication of sculpturing, is as wavy lines paralleling the ventral margin.

Frontal organ. (Fig. 32, B). It is almost straight, narrow, considerably extended beyond the

down-curving distal segments of the 1st antenna and is approximately 90-95% of the 1st antenna in length (on the dorsal side). Capitulum section is unseparated, with 2 thin and small needle-like prolongations at the tip and without hairs.

First antenna. (Fig. 32, B-D). The limb is 6-jointed. Dorsal seta of the 2nd segment is long, stout and coarsely spinous (proxidorsally and distoventrally) and with short hairs (distodorsally and proximoventrally). Six long filaments on the 5th and 6th segment are in equal length, which are distally widened and pointed at the tip. Dorsal margin of the limb is about 60% of the filament in length. Height of antenna (on the 2nd segment) is nearly 20% of the length. Hairs and pigment spots on the limb are missing.

Second antenna (Figs. 32, F and 33, A). The exopodite is relatively thick and shorter than the protopodite. The total length of the 2nd-9th segments of the exopodite is approximately 40% of the 1st segment in length. Basal segment is without hairs and warty surface, the bristles "a" and "b" are thin and armed with long hairs, bristle "a" is about twice as long as the "b" bristle. Bristles "c"- "e" on the 2nd segment are lacking. The 5 filaments placed on this segment have an equal length and thickness throughout, pointed at the tip, and longer than the exopodite and protopodite.

Mandible (Figs. 33, B, C and 34, G, H). The epipodite is slightly developed and without bristle. The exopodite is represented by one midlong seta. The 1st segment has straight dorsal margin and bears one dorsal and 3-4 ventral setae, the 2nd segment has 3 dorsal (usual type) and 2 ventral setae. The 3rd segment is armed with 7 setae, of which the largest is claw-like and about as long as the endopodite (on the dorsal side), and 2 first dorsal setae are slight (the 1st shorter than the 2nd seta). Basale bears 2 lateral setae, 2 anterior, one anterolateral setae and anterior row long hairs near the articulation. Basal endite has toothrow with 6 slightly separated serrated teeth and 2 short posterior bristles. Coxale endite is armed with 3 toothrows, 4 rounded denticles and some tiny hairs. The masticatory pad is flat and bears 2 very wide rounded denticles, terminal row of short filaments and numerous long hairs.

Maxilla (Fig. 33, D, F). The basal segment bears 6 unplumose anterior and 6 (one plumose) posterior setae. The distal segment has 2 claw-setae and 3 (one lateral is short) usual setae.

Fifth limb (Fig. 34, A). The epipodite has 3 groups of 4+4+5 (one short) plumose setae. The 1st segment of the exopodite has 7 ventral and 3 dorsal setae, of which distodorsal seta is approximately 65% of the exopodite in length (on the dorsal side). The 2nd segment has 2 ventral and one dorsal setae, last seta is about 30% and middle claw-seta of the 3rd segment is 55% of the exopodite in length. Two other terminal setae are slight and short.

Sixth limb (Fig. 34, B-E). The epipodial appendage is armed with 3 groups of 5+4+6 (one short) plumose setae. Two short plumose setae are borne on the endopodite. The 1st segment of the exopodite has 2 dorsal and 5 ventral setae, its distodorsal seta about half the length of the exopodite (on the dorsal side). The 2nd segment is with one ventral and the 3rd segment with one dorsal and one ventral setae. The 4th segment has 2 thin setae and middle claw-seta, the last is longest them and about 50% of the exopodite is length. Height of the exopodite is approximately 30% of the its length.

Seventh limb (Fig. 3, F). It is medium in length and thickness (25% of the length) and prolonged. The longest seta is about twice as long as the short seta and 3 times as long as the limb.

Furca (Fig. 34, G). This limb has 7 pairs of claws and single unpaired bristle.

Comparison

The new species differs from other species of this genus in having a greatest height of the shell at the anterior half. It is closely related to *A. ventricosa* and *A. bispicula*. The new species differs from the first species in the following: the shell is larger and less prolonged; frontal organ is with 2 thin and small needle-like prolongations at the tip; filaments of the 1st antenna are widened distally; the 1st segment of the maxilla exopodite has more posterior setae; and the 1st segment on the exopodite of the 5th limb has more ventral setae. New species differs from *A. bispicula* in the following: the shell is larger; terminal claw-seta on the mandible is longer; and in 5th and 6th limbs this claw-seta is shorter,




Fig. 32 *Archiconchoecetta inventricosa* sp. nov. (female: 1123 - A; 1122 - B-F) A - lateral view of left valve of shell, B - frontal organ and 1st antenna, C - distal part of 1st antenna D - dorsal seta of 1st antenna, E - 2nd antenna, F - endopodite 2nd antenna.




Fig. 33 *Archiconchoecetta inventricosa* sp. nov. (female: 1122 - A, C-F; 1123 - F) A - endopodite of 2nd antenna, B - mandible, C - basale endite of mandible, D and E - maxilla, F - distal part of endopodite of maxilla.




Fig. 34 *Archiconchoecetta inventricosa* sp. nov. (female: 1122 - A-E; 1123 - F-I) A - fifth limb, B and C - sixth limb, D and E - distolateral seta of 1st segment of exopodite sixth limb and proximal part of this segment, F - seventh limb, G - furcal lamella, H and I - tooththrows and masticatory pad of coxa of mandible.

and the rest as is the case with *A. ventricosa*.

Distribution

New species has been taken in the Kurile - Kamchatka and Aleutian Trenches between 3000 and 5500 m.

Archiconchoecetta ventricosa (Müller, 1906)

Archiconchoecia ventricosa Müller, 1906: 45-46, Taf. 7, Fig. 1-6; 1908: 63; 1912: 55; Alcaraz, Manriquez and Vasquez, 1975: 381, 382; Alcaraz, 1977: 9; Angel and Fasham, 1975: 716-718, 720, tab. 2-4; Angel, 1977: 243; 1979: 120, 121; Deevey, 1978a: 63-67, Figs. 4, 5; Deevey and Brooks, 1980: 63-65, Fig. 10a-c; McKenzie, Duccasse, Dufour and Peypouquet, 1979: 2; Drapun, 1981:75; Angel, 1993: 80, Fig.27.

? *Archiconchoecia ventricosa*: Angel, 1981: 556, 562, Fig. 194-7 (A-H).

Not *Archiconchoecia ventricosa*: Poulsen, 1969: 21-24, figs. 5-7.

Not *Archiconchoecia ventricosa*: Chavtur, 1977a: 142, 158, fig. 8, tab. 6; 1977b: tab.2; 1977c: table. 2.

Remarks

Probably, specimens noted by Angel (1981) are not *A. ventricosa*, since the frontal organ is without terminal long spine (description and illustrations of many appendages are missing in this publication).

Distribution

Species inhabits the Atlantic Ocean in the latitudes range 44°N-24°S (NE Atlantic - Müller, 1906; 1912; Alcaraz *et al.*, 1975; Alcaraz, 1977; Angel and Fasham, 1975; Angel, 1977; 1979; McKenzie *et al.*, 1979. NW Atlantic - Deevey, 1978a; Deevey and Brooks, 1980; Angel, 1979. S Atlantic - Müller, 1908; Drapun, 1981). Also it was taken in the Indian Ocean from 7°N - 85°W (Müller, 1906). It is known from levels in the depth range 2000-500 m and in tows from 3500 m to the surface. Angel (1993) reported that *A. ventricosa* is widespread (but never common) and occurs in the Indian and Atlantic Oceans between 0°-49°N.

Archiconchoecetta poulseni (Deevey, 1978)

Archiconchoecia ventricosa: Poulsen, 1969: 21-24, figs. 5-7.

Archiconchoecia poulseni Deevey, 1978c: 106, 107.

Remarks

Deevey (1978c) singled out new species since "...Poulsen's female differs from Muller's specimens in size, the shape of frontal organ, and in the number of other ways, it is here renamed as *A. poulseni* n. sp." In addition, *A. poulseni* is separated from *A. ventricosa* in the following: shell is twice as long as that in *A. ventricosa*; frontal organ is with 2 (one in *A. ventricosa*) long terminal spines; basale endite is armed with 7 (6 in *A. ventricosa*) teeth, posterior seta is longer than anterior seta on the 2nd segment of the maxilla endite (shorter in *A. ventricosa*). Besides, these species differ by sizes and setation of all the limbs.

Distribution

Collected in the South Pacific (22°43S-166°06E and 28°18S- 177°01E) from levels 1500-50 m (3000-100 mw) and 500-50 m (1000-100 mw).

***Archiconchoecetta bidens* (Deevey, 1982)**

Archiconchoecia bidens Deevey, 1982: 140-143, figs. 13-15.

Distribution

Species has been taken only in the Southern Ocean (71°57'S-179°58'E) from 1600-1400 m.

***Archiconchoecetta fabiformis* (Deevey, 1978)**

Archiconchoecia fabiformis Deevey, 1978a: 59-63, Figs. 2, 3.

Distribution

Species was caught in the Sargasso Sea (11°45'N-66°56'W) in a tow from 1200-0 m.

***Archiconchoecetta bimucronata* (Deevey, 1978)**

Archiconchoecia bimucronata Deevey, 1978c: 128-131, Figs. 13-14; 1983: 411; Deevey, Brooks, 1980: 65, Fig. 10e, f.

Distribution

Known from the North Atlantic (32°0'N - 64°30'W) between 1500-1000 m (Deevey, 1978b; Deevey, Brooks, 1980) and from the South Pacific (42°28'S-long) in level 2000-1000 m (Deevey, 1983).

***Archiconchoecetta bispicula* (Deevey, 1978)**

Archiconchoecia bispicula Deevey, 1978c: 123-125, Fig. 10; Deevey, Brooks, 1980: 63, 65, Fig. 10g,h.; Ellis, 1985: 928; Angel, 1993: 68, Fig. 21.

Distribution

Species was described from the Caribbean Sea (17°18'N - 85°27'W, in a vertical tow, probably from around 2000 m). Later, it has been taken in the North Atlantic from region 32°10'N - 64°30'W between 1000 and 500 m (Deevey and Books, 1980) and 49°N - 13°W in 900 to 800 m (Ellis, 1985). Angel (1993) reported that *A. bispicula* known from 17°-53°N in the North Atlantic (a rare benthopelagic / deep mesopelagic species usually taken at 800-1500 m).

***Archiconchoecetta bifurcata* (Deevey, 1978)**

Archiconchoecia bifurcata Deevey, 1978c: 131-134, Figs. 15,16; 1983: 411; Deevey and Brooks, 1980: 62-63, 65, Fig. i, j.

Distribution

Species was caught in the Sargasso Sea (32°10'N-64°30'W) from levels 1000-500 and 1500-1000 m (Deevey, 1978c; Deevey, Brooks, 1980) and in the Pacific sector of the Southern Ocean (62°20'S-? long) from 3700-3500 m (Deevey, 1983).

***Archiconchoecetta gastrodes* (Deevey, 1978)**

Archiconchoecia gastrodes Deevey, 1978c: 134-137, Figs. 17-18; Deevey, Brooks, 1980: 65, Fig. 10d; Angel, 1983: 552, 1993: 76, Fig. 25; Ellis, 1985: 929.

Distribution

Collected in the Sargasso Sea from region 32°10'N-64°30'W in the level 1500-1000 m (Deevey, 1978c; Deevey, Brooks, 1980) and in the North Atlantic from 42°N-17°W (Angel, 1983) and 44°N-13°W (Ellis, 1985) between 2700-1500 and 1650-800 m respectively.

Archiconchoecetta falcata (Deevey, 1978)

Archiconchoecia falcata Deevey, 1978c: 125-128, Figs. 11, 12; 1983: 411; Deevey, Brooks, 1980: 43, 65; Chen and Lin, 1994b: 416, table 2.

Distribution

This species was caught in the Sargasso Sea (32°10N-64°30W) from 2000-1000 m (Deevey, 1978c; Deevey and Brooks, 1980), in the Pacific sector of the Southern Ocean (56°S-? long) from 1830 m (Deevey, 1983), and in the central South China Sea in the depth range 1000-4000 m (Chen and Lin, 1994b).

Archiconchoecetta pilosa (Deevey, 1978)

Archiconchoecia pilosa Deevey, 1978c: 115-119, Figs. 5,6; 1983: 411; Deevey, Brooks, 1980: 63, tab. 2; Ellis, 1985: 929.

Distribution

Species was collected in the Sargasso Sea from the region 32°10N - 64°30W (Deevey, 1978c; Deevey, Brooks, 1980) and the North Atlantic from 49°N-13°W (Ellis, 1985) between 2000-1500 and 600-500 m, respectively. Also, it was taken in the South Pacific (37°N-? long) between 1000 and 500 m (Deevey, 1983). Angel (1993) reported that this is a rare bathypelagic species which known from 10°-49°N in the North Atlantic.

Archiconchoecerra Chavtur gen. nov.

Type-species - *Archiconchoecia longiseta* Deevey, 1978.

Composition

This genus contains only *A. longiseta* (Deevey, 1978).

Description

Shell. Females range from 1.27-1.3 mm in length, male unknown. Valves are prolonged. Greatest height of the shell is at the posterior part and around 60% of the length. Rostrum is developed and wide, and rostral incisure is not deepened. Dorsal margin is straight, ventral, anterior and posterior margins are rounded. Posterior margin is without tubercles and points. Clear lens-like structure is near posteroventral corner of each valve. Left asymmetrical gland is near posterodorsal corner, right gland is 1/3 to 1/2 of the way up the posterior margin. The shell has not visible sculpture.

Frontal organ. It is slightly down-curved and considerably extended beyond the down-curving distal segments of the 1st antenna and is approximately 90% of this antenna in length (on the dorsal side). Capitulum section is not separated or barely separated, with 1-2 tiny points at the tip and with tiny ventral hairs.

First antenna. The limb is 6-jointed, with one long, stout and coarsely spinous dorsal seta on the 2nd segment. The 5th and 6th segments bear 6 filaments unequal in length, one is about twice as long as the other 5. These filaments are not distally widened and are pointed at the tip. Dorsal margin of the limb is approximately equal in length to 5 short filaments. Height of antenna (on the 2nd segment) is about 20% of the length. Hairs are placed on the 2nd, 3rd and 4th segments.

Second antenna. The exopodite is relatively thick and shorter than the protopodite. The total length of the 2nd-9th segments of the exopodite is approximately 40-50% of the 1st segment in length. Basal segment of the endopodite is with long bristles "a" and "b" and hairs near its base, and without warty surface. The 2nd segment is armed with hairs and without bristles "c"-"e". The 5 filaments are very short, distally tapering, pointed at the tip and about 1/4-1/3 the length of the exopodite and protopodite.

Table 2. Vertical distribution of species, genera and subgenera of Archiconchoecinae.

Taxon	Depth in km											
	0-0.5	0.5-1	1-2	2-3	3-4	4-5	5-6	6-7	7-8	8-9	9-10	
<i>A.(A.) propinqua</i>	+											
<i>A.(A.) instriata</i>	+											
<i>A.(A.) aff. striata 1</i>	+											
<i>A.(A.) aff. striata 2</i>	+											1)
<i>A. cuneata</i>	+	+										
<i>A.(A.) striata</i>	+	+		?								2)
<i>A. fabiformis</i>	+	+	+									3)
<i>A. poulseni</i>	+	+	+									
<i>A. aff. cuneata 1</i>	+											
<i>A. cucullata</i>	+	+	+	+	+							
<i>A. ecuneata</i>	+	+	+									
<i>A. aff. cucullata 2</i>		+										
<i>A. aff. cuneata 2</i>		+										
<i>A. bispicula</i>		+	+									
<i>A. pilosa</i>		+	+									
<i>A. ventricosa</i>	?	+	+	?	?							4)
<i>A. gastrodes</i>		+	+	+								
<i>A. versicula</i>			+	+								
<i>A. falcata</i>			+									
<i>A. bimucronata</i>			+									
<i>A. arctica</i>			+	+	+							5)
<i>A. bidens</i>			+									
<i>A. longiseta</i>			+									
<i>A. simula</i>			+	+	+							
<i>A. sp. nov. 1</i>			+	+								
<i>A. orientalis</i>				+								
<i>A. sp. nov. 2</i>				+								
<i>A.(A.) apertesulcata</i>				+	+							
<i>A. pljusnini</i>					+	+						
<i>A. inventricosa</i>					+	+	+					
<i>A. bifurcata</i>					+							
<i>A. aff. cucullata 1</i>					+							
<i>A. aff. pljusnini</i>						+						
<i>A. maculata</i>									+	+	+	6)
Total	11	12	16	9-?11	8-?9	3	1		1	1	1	

1) correct locality is unknown

2) also 1 juv. from 2700-2400 m

3) only in a tow from 1200 m to surface

4) also in a tows from 3500 m to surface

5) only 1 male from 3800-935 m

6) only 1 sample from 9500-7280 m

Mandible. The epipodite is without bristles. The exopodite is represented by one short seta. The 1st segment of the endopodite has straight dorsal margin and with one dorsal and 4 ventral setae, the 2nd segment with 3 dorsal and 2 ventral setae. The 3rd segment bears 7 setae, of which largest seta is claw-like and about as long as the endopodite (on the dorsal side), and 2 first dorsal setae are slight (the 1st barely shorter than the 2nd seta). Basale is armed with 2 anterior and 2 lateral setae, and one anterolateral seta near the articulation. Basale endite has toothrow with 6 slightly separated serrated teeth and 2 short proximal bristles.

Maxilla. The basal segment bears 6 anterior and 4-5 posterior setae. The distal segment is armed with 2 claw-setae (posterior claw-seta long and anterior - very short) and 3 usual setae.

Fifth limb. The epipodite has 3 groups of 4+5+4 plumose setae. The 1st segment of the exopodite bears 5 ventral and 2 dorsal and the 2nd segment is with only 2 ventral setae. The middle claw-seta of the 3rd segment is about 60-70% of the exopodite in length and approximately as long as (slightly shorter) the dorsal claw-seta.

Sixth limb. The epipodial appendage is armed with 3 clusters of 5+5+6 plumose setae. Two long plumose setae are placed on the endopodite. The 1st segment of the exopodite has 2 dorsal and 4 ventral, the 2nd segment is with only one ventral and the 3rd segment with one dorsal and one ventral setae. The 4th segment bears 2 long and slim dorsal and middle claw-setae and one short usual posterior seta; this middle claw-seta is about 80% and distodorsal seta of the 1st segment is 40-50% the length of the exopodite (on the dorsal side). Height of the exopodite is 60-65% of the length.

Caudal furca. This limb has 8 pairs of claws and single unpaired bristle.

Comparison

New genus sharply differs from the all other genera in having a clear lens-like structure near posteroventral corner of the each valve; longest seta on the 1st antenna is approximately twice as long as the other its setae; very short filaments are borne on the 2nd antenna; longer posterior and shorter anterior setae are placed on the 3rd segment of the exopodite of the 6th limb.

This genus is close to *Archiconchoecetta* gen. nov., but separated (in addition to the above characters) mainly by the shape of shell, location of the right asymmetrical gland; shorter seta on the exopodite, and slightly separated teeth on the basal endite of the mandible; longest dorsal (no middle) seta is borne on the distal segment of the 5th limb.

Distribution

Known only in the Sargasso Sea from 1500-2000 m.

Distribution of ostracods of subfamily Archiconchoecinae

The members of the subfamily Archiconchoecinae occur worldwide from the North to the South Pole and from the surface to maximal depth. They are common for the tropical and subtropical zones of the World Ocean (Deevey, 1987c; Deevey, Brooks, 1980), and less frequent in the temperate waters of the Northern Hemisphere (Chavtur, 1977a; 1977b; 1992) and in the South Ocean (Deevey, 1978a; 1982; 1983), and as exception in the Arctic Ocean (Chavtur, 1992) (Fig. 35). In waters of the Arctic Ocean only one specimen of ostracods of this group was recorder (Chavtur, 1978; 1992). The Archiconchoecinae ostracods prefer the deepwater zone of the World Ocean; only several of the subgenus *Archiconchoecia* (*Archiconchoecia*) are typical inhabitants of the epipelagical zone. Here belong only *A. (A.) instriata* nov. sp., *A. (A.) propinqua* nov. sp. and probably *A. (A.) aff. striata* 1. To interzonal species (term by Vinogradov, 1968) belong *A. (A.) striata* and perhaps *Archiconchoecetta fabiformis* (the latter is known only in a vertical tow from 1200-0 m: Deevey, 1978c). As to the rest species they are typical representatives of the deepwater fauna, while *Archiconchoecilla maculata* occurs only in the hadal zone (deeper than 6000 m).

With increasing depth the number of species rises (Table 2) to form the maximum in the layer

1000-2000 m. However it should be expected that this maximum boundary will shift somewhat lower with time, since in the tropical-subtropical zone layer-by-layer catches were done only down to 2000 m depth, and total catches (from around 3000-4000 m) did not provide realistic information on vertical distribution of ostracods. Most genera were recoded between 3000-4000 m.

The species of Archiconchoecinae do not form mass accumulation. In catches they are usually represented by single specimens. The greatest abundance of 500 specimens/1000 m³ was found for *A. (A.) striata* from the Sargasso Sea in 1970 (Deevey, 1978c).

Ostracods of this subfamily are filter-feeding organisms and belong (depending on depth) either to herbivores or detritivores. The universal structure of the mandible as well as tiny sizes of these organisms enable them to command the greatest depth and lead in abundance among ostracods of other subfamilies in the hadal zone (Chavtur, 1977a; 1977b; 1992).

The number species of this group in the World Ocean is not high as compared to other pelagic ostracod groups. One should expect descriptions of new species mainly from tropical and subtropical regions. In the number of species only of the subfamilies Halocypridinae and temporarily Euconchoecinae, rank second to this group. The composition of the latter subfamily is very diverse at great depths (according to unpublished author's data).

Widely-distributed (literature data) species *Archiconchoecia striata* is actually a set of morphologically close species: *Archiconchoecia (Archiconchoecia) striata*, *A. (A.) instriata* nov. sp., *A. (A.) propinqua* nov. sp., *A. (A.) aff. striata* 1, *A. (A.) aff. striata* 2 and *Archiconchoecia (Archiconchoecidia) apertesulcata* nov. sp. With time, possibly this list will be augmented.

Acknowledgments

I wish to thank Dr. M. E. Vinogradov and Dr. J. A. Rudjakov of Institute of Oceanology, Russian Academy of Science, for the plankton materials upon which partially study is based and for the environmental data presented in the station data herein. Also I am greatly indebted to Dr. E. I. Shornikov of Institute of Marine Biology, Far East Branch, Russian Academy of Science, for his advice during the course of this study. This work was supported by the grant from the Russian Fundamental Fund of Investigation (RFFI).

References

- Alcaraz M., 1977. Cladoceros y ostracodas de los alrededores del Estrecho de Gibraltar en junio-julio de 1972. Resultados Expediciones científicas del Bugue Oceanográfico Cornide de Saavedra, 6, 41-63.
- Alcaraz M., Manrígués M., and Vazgues A., 1975. Ostracodas pelagicos del SW de Portugal: Estructura de las comunidades. Investigaciones Pesqueras, 39 (2), 379-396.
- Angel M. V. 1969a. Planktonic ostracods the Canary Island region; their depth distributions, journal migrations, and community organization. Journal of the Marine Biological Association of the United Kingdom, 49, 515-553.
- Angel M. V. 1969b. Repeated samples from a deep midwater planktonic ostracod community. Journal of Experimental Marine Biology and Ecology, 3, 76-89.
- Angel M. V. 1977. Studies on Atlantic halocyprid ostracods: vertical distribution of the species in the 1000 m in the vicinity of 44°N, 13°W. Journal of the Marine Biological Association of the United Kingdom, 57, 239-252.
- Angel M. V., 1979. Studies on Atlantic halocyprid ostracods: their vertical distributions and community in the central gyre region along latitude 30° N from off Africa to Bermuda. Progress of Oceanography, 8, 3-124.
- Angel M. V., 1981. Ostracoda. Atlas de zooplancton del Atlantico sudoccidental y metodos de trabajo con el zooplancton marino. Publication especial del INIDEP Mar del Plata, Argentina, 543-585.
- Angel M. V., 1983. A vertical profile of planktonic ostracods from depth of 1500-3900 m at northeast Atlantic Station. In, Maddocks R. E. (ed.). Applications of Ostracoda: Proceedings of the Eighth International Symposium on Ostracoda (July 26-29, 1982). Department of geosciences, University of Houston, Houston, 549-559.

- Angel M. V., 1984. The diel migrations and distributions within a mesopelagic community in the North East Atlantic. 3. Planktonic ostracods, a stable component in the community. *Progress of Oceanography*, 13, 319-351.
- Angel M. V. 1993. Marine planktonic ostracods. Synopsis of the British Fauna (new Series), No 48. Published for the Linnean Society of London and the Estuarine and Coastal Sciences Association by Field Studies Council. Shrewsbury. 240p.
- Angel M. V., and Fashan M. J. R., 1975. Analyses of the vertical and geographic distribution of the abundant species of planktonic ostracods in the North-East Atlantic. *Journal of the Marine Biology Association of the United Kingdom*, 55, 709-737.
- Angel M. V., Hargreaves P., Kirkpatrick P. and Domanski P., 1982. Low variability in planktonic and micronectonic populations at 1000 m depth in the vicinity of 42°N, 17°W; evidence diel migratory behavior in the majority of species. *Biological Oceanography*, 1, 287-319.
- Benassi G., Ferrari I., Rossi V., Sei S., Angel M.V., and McKenzie K. G., 1998. Distribution and taxonomy of planktonic ostracods of the Eolian Islands (Mediterranean Sea), 3 Congr. Europ. Des Ostracodologistes. Paris-Bierville 8-12 July 1996, *Bull. Centre Rech. Elf Explor. Prod.*, 3, 3-25.
- Brady G. S., 1902. On new imperfectly-known Ostracoda, chiefly from a collection in the Zoological Museum, Copenhagen. *Transaction of the Zoological Society of London*, 16, 4 (1), 179-210.
- Brady G. S., 1903. Report on dredgings and other marine research off the north-east of England in 1901. *Transaction of National Historical Society of London*, 14 (5), 87-101.
- Chavtur V. G., 1977a. [Species composition and vertical distributions of pelagic ostracods in the region of Kurile - Kamchatka Trench.]. *Trudy Instituta okeanologii AN SSSR*, 108, 136-164 [In Russian].
- Chavtur V. G., 1977b. [Fauna of pelagic ostracods of the Halocyprididae family in the northwestern part of the Pacific ocean.]. *Biologija morja*, 1, 28-38 [In Russian].
- Chavtur V. G., 1977c. Fauna of pelagic ostracods of the Halocypridinae family in the northwestern part of the Pacific ocean. *Sovetskyye Trudy Marine Biology*, 3 (3), 18-26.
- Chavtur V. G., 1978. [Species composition and feature of distribution of plankton ostracods (Ostracoda, Myodocopa) in the central part of Arctic Basin] *Zoologicheskyy zurnal*, 57 (12), 1790-1798 [In Russian].
- Chavtur V. G. 1987. Three new species of pelagic ostracods from the North Pacific. *Zoologicheskyy zhurnal*, 66 (6), 943-949 [In Russian].
- Chavtur V. G., 1991. [History - bibliographic review and analysis of study recent ostracod-myodocops of the oceans world.]. *Vladivostok Dalnevostochnoe Otdelenie Akademii Nauk SSSR*, 2 (3), 136 pp. [In Russian].
- Chavtur V. G., 1992. [Composition, structure and distribution benthic and pelagic ostracods of the superorder Myodocopa of temperate and cold waters of the Northern Hemisphere.]. *Institut Biologii morija Dalnevostochnoe Otdelenie Rossyjskoj Akademii Nauk. Vladivostok*, 155 pp. 30 figs., bibl. 96 nam. Deponent VINITI, N 3016-V92, 20.10.92. [In Russian].
- Chen Q., Yin J., and Zhang G., 1983. Studies on pelagic ostracods in the central and northern parts of the South China Sea. *Contributions on Marine Biological Research. of the South China Sea*, 3 (1), 82-132 [in Chinese].
- Chen R., Lin J., 1994a. The distribution of planktonic Ostracoda between the source of Kuroshio and west of Taiwan Strait. *Acta Oceanologica Sinica*, 13 (3), 445-452.
- Chen R., and Lin J., 1994b. Ecological study on Ostracoda in the central South China Sea. *Acta Oceanologica Sinica*, 13 (3), 413-423.
- Chen R., and Lin J., 1995. Pelagic ostracoda in China seas China. *Ocean*, 134 pp. [in Chinese].
- Cleve P. T., 1905. The plankton of the South African Seas. III. Halocyprididae, *Marinae investigations in the South Africa*. Cape Town, 4, 129-132.
- Deevey G. B., 1968. Pelagic ostracods of the Sargasso Sea off Bermuda. *Peabody Museum of Natural History Yale University. Bull.*, 26, 125 pp.
- Deevey G. B., 1970. Pelagic ostracods (Myodocopa, Halocyprididae) from the North Atlantic off Barbados. *Proceedings of the Biological Society of Washington*, 82 (62), 799-824.
- Deevey G. B., 1971. The annual cycle in quantity and composition of the zooplankton of the Sargasso Sea off Bermuda. I. The upper 500 m. *Limnology and Oceanogr*, 16 (2), 219-240.
- Deevey G. B., 1974. Pelagic ostracods collected on Hudson 70 between the equator and 55° S in the Atlantic. *Proceedings of the Biological Society Washington*, 87 (32), 351-380.

- Deevey G. B., 1978a. A taxonomic and distributional study of the planktonic ostracods collected on three cruises of the "Eltanin" in the South Pacific and the Antarctic Region of the South Pacific. Biology of the Antarctic Seas, VIII Antarctic Research, 28 (3), 43-70. (American Geophysical Union).
- Deevey G. B., 1978b. The planktonic ostracods of the Cariaco Trench and adjacent waters. Proceeding of Biological Society of Washington, 91 (1), 52-73.
- Deevey G. B., 1978c. On new and little known species of *Archiconchoecia* (Myodocopa, Halocyprididae) from the Sargasso and Caribbean Seas, with description of seven new species. Bulletin of Florida State Museum, Biological Science, 23 (2), 105-138.
- Deevey G. B., 1982. Planktonic ostracods of the north Atlantic off Barbados. Bulletin of Marine Science, 32 (2), 467-488.
- Deevey G. B., 1983. Planktonic ostracods (Myodocopa, Halocyprididae) from six Eltanin cruises in South Pacific and Antarctic waters. Journal Crustacean Biology, 3 (3), 409-412.
- Deevey G. B., and Brooks A. L., 1980. The planktonic ostracods of the Sargasso Sea off Bermuda: species composition and vertical and seasonal distribution between the surface and 2000 m. Bulletin of Florida State Museum, Biological Science, 26 (2), 37-124.
- Drapun I. E., 1981. [Pelagic ostracods from the subtropical region of the South Atlantic]. Vestnik zoologii, 3, 74-76. [In Russian].
- Drapun I. E., 1983. [Pelagic ostracods from the south-west sector of the Southatlantic subtropical circulation]. Ecologia morja, 14, 33-37. [In Russian].
- Drapun I. E., 1988. [Correlation line size with biomass of planktonic ostracods]. Ecologija morja, 28, 46-51 [In Russian].
- Ellis C. J., 1985. The effects of proximity to the continental slope sea-bed on pelagic halocyprid ostracods at 49°N-13°W. Journal of the Marine Biological Association of the United Kingdom, 65, 923-949.
- Fasham M. J. R., Angel M. V., and Roe H. S. J., 1974. An investigation of the spatial pattern of zooplankton using the Longhurst-Hardy planktonic recorder. Journal of Experimental Marine Biology and Ecology, 16: 93-112.
- Fowler G. H., 1909. Biscayan plankton collected during a cruise of H. M. S. "Research", 1900. Pt. XII. The Ostracoda. Transaction of the Linneus Society of London, Zoological Series 2, 10, 219-358.
- Granata L., and Caporiacco L., 1949. Ostracodes marins recueillis pendant les croisières du Prince Albert I-er de Monaco. Resultats Campagnes Scientifiques, Prince Albert I, 109, 1-51.
- Gamulin T., 1948. Contribution a' la connaissance du zooplancton de la zone insulaire de la *Dalmatie moyenne*. Acta Adriatica, 3 (7), 3-38. (citation on: Gooday A. J., Angel M. V., 1977).
- George J., 1979. Redescription of *Archiconchoecia striata* Muller (Ostracoda, Halocyprididae) and its distribution in the Northern Indian Ocean. Crustaceana, 36 (2), 123-128.
- George J., and Nair V. P., 1980. Planktonic ostracods of the Northern Indian Ocean. Mahasagar - Bulletin of the National Institute Oceanography, 13 (1), 29-44.
- Gonzales L.C., and Breman E., 1982. Distribution of planktonic ostracoda in the surface waters of the Bonaire Basin and adjacent seas. Crustaceana, 42 (2), 201-211.
- Gooday A. J., and Angel M. V., 1977. Distribution of planktonic Ostracoda (Halocyprididae) in the North Adriatic with the description of a new subspecies, *Conchoecia porrecta adriatica*. Crustaceana, 32 (2), 139-154.
- Grice G. D., and Hart A., 1962. The abundance seasonal occurrence and distribution of the epizooplankton between New York and Bermuda. Ecological Monographs, 32 (4), 287-309.
- Hanai T., Ikeya N., and Yajima M., 1980. Checklist Ostracoda from Southeast Asia. Bulletin of the University Museum, University of Tokyo, 17, 236 pp.
- Hillman N. S., 1969. Distribution of selected groups of marine invertebrates in waters south of 35°S latitude. Antarctic Map Folio Series, 11, Halocyprididae, 29-30.
- Hure J., 1961. Distribution saisonniere et migration journaliere verticale du zooplancton dans la region profonde de l' Adriatique. Acta Adriatica, 9 (6), 3-59.
- Juday C., 1906. Ostracoda of the San Diego Region. I. Halocyprididae. University of California Publications in Zoology, 3 (2), 13-38.
- Leveau M., 1965. Contribution a' l' etude des ostracodes et cladoceres du Golfe de Marseille. Recueil des travaux de la Station marine Endoume, Marseille, Fascicule hors serie supplement, 37 (53), 161-246.
- Leveau M., 1967. Ostracodes pelagiques du sud-ouest de l' ocean Indien (region de Tulear). Recueil des travaux de la Station marine Endoume, Marseille, Fascicule hors serie supplement, 6, 63-70.

- Leveau M., 1969. Ostracodes pelagiques recuillis lors de la 3^e campagne de l'avis "Commandant Robert Giraud". Recueil des travaux de la Station marine Endoume, Marseille, 8, 123-142.
- Martens J. M., 1979. Die pelagischen Ostracoden der Expedition Marchile I (Sudost-Pazifik), II: Systematik und Vorkommen (Crustacea: Ostracoda: Myodocopa). Mitteilungen aus dem Hamburgischen zoologischen Museum und Institut, 76, 303-366.
- Martens J. M., 1981. Die pelagischen Ostracoden der Marchile I-Expedition (Sudost-Pazifik), I: Verbreitung, zoogeographie und bedeutung als indikatoren für wasserkörper (Crust. Ostracoda: Myodocopida). Studies on Neotropical Fauna and Environment, 16, 57-97.
- McKenzie K. G., Ducasse O., Dufour E., and Peypouquet J. P., 1979. Monographie bibliographique, stratigraphique et paleocologique sur les ostracodes cenozoiques du Bessin d'Aquitaine et du Golfe de Gascogne. Bulletin de l'Institut Geology, Bassin Aquitaine Talence, 195 pp.
- Moraitou-Apostolopoulou M., 1981. Planctonic ostracods collected between 1964 and 1973 from different Greek Sea areas. Vie et Milieu, 31 (2), 171-176.
- Müller G. W., 1894. Die ostracoden des Golfes von Neapel und der angrenzenden meeresabschnitte. Fauna und flora des Golfes von Neapel. Naples. Zool. Stat. Berlin, 21, 404 pp.
- Müller G. W., 1906. Ostracoda. Wissenschaftliche Ergebnisse Deutsche Tiefsee-Expedition auf dem Dampfer "Valdivia", 8, 1-154 pp.
- Müller G. W., 1908. Die Ostracoden. Der Deutschen Sudpolar-Expedition 1901-1903. Sudpolar-Expedition 1901-1903, Zoologie, Berlin, 10 (2), 53-184.
- Müller G. W., 1912. Ostracoden. Das Tierreich, Berlin, 31, 434 pp.
- Poulsen E. M., 1969. Ostracoda -Myodocopa. Pt III A. Halocypriformes - Thaumacyprididae and Halocypridae. Dana - Reptort, 75, 100 pp.
- Poulsen E. M., 1973. Ostracoda -Myodocopa. Pt III A. Halocypriformes - Halocypridae, Conchoecinae. Dana-Reptort, 84, 224 pp.
- Puri H. S., 1963. Preliminary notes on the Ostracoda of the Gulf of Naples. Experientia, 19 (368), 1-16.
- Schweiger L., 1912. Adriatische cladoceren und plankton ostracoden. Sutzungeber.-Berlin, Akademie Wissenschaftliche, Wien, 121 (5), 239-175. (citation on: Gooday A. J., Angel M. V., 1977).
- Skogsberg T., 1931. Ostracods. In, Murray J., Hjort J. (eds.) Report of the scientific results of the "Michael Sars" North Atlantic Deep-Sea Expedition 1910, 5, 1-26.
- Tseng W. Y., 1970. Occurrence of ostracods in the neighbouring seas of Taiwan. In, Sugawara K. (ed.). The Kuroshio: Proceedings of the 2nd CSK Symposium, Tokio, 237-295.
- Vinogradov E. M. 1968. Vertical distribution of the oceanic zooplankton. Publishing House "Nauka", Moscow, 320 pp. [In Russian].
- Vucetic T., 1977. Les principales masses d'eau en Adriatique et leur influence sur les communautés pelagiques. Journées Etudes Planctonologie, Monaco, C.I.E.S.M., 105-114. (citation on: Gooday A. J., Angel M. V., 1977).