

Title	Nitric oxide-induced calcium release: activation of type 1 ryanodine receptor by endogenous nitric oxide.
Author(s)	Kakizawa, Sho; Yamazawa, Toshiko; Iino, Masamitsu
Citation	Channels (2013), 7(1): 1-5
Issue Date	2013-01-01
URL	http://hdl.handle.net/2433/179147
Right	© 2013 Landes Bioscience
Type	Journal Article
Textversion	author

Nitric oxide-induced calcium release: Activation of type 1 ryanodine receptor by endogenous nitric oxide

Sho Kakizawa^{1*}, Toshiko Yamazawa² and Masamitsu Iino³

¹Department of Biological Chemistry, Graduate School of Pharmaceutical Sciences, Kyoto University, Kyoto, Japan; ²Department of Molecular Physiology, The Jikei University School of Medicine, Tokyo, Japan;

³Department of Pharmacology, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan

*Corresponding author: Sho Kakizawa, Department of Biological Chemistry, Graduate School of Pharmaceutical Sciences, Kyoto University, Kyoto 606-8501, Japan.

Tel: +81-75-753-4552

Fax: +81-75-753-4562

E-mail: sho-kaki@pharm.kyoto-u.ac.jp

Running title: Activation of ryanodine receptor by endogenous NO

Keywords: nitric oxide, S-nitrosylation, ryanodine receptor, calcium, Purkinje cell, synaptic plasticity

Addendum to: Kakizawa S, Yamazawa T, Chen Y, Ito A, Murayama T, Oyamada H, Kurebayashi N, Sato O, Watanabe M, Mori M, Oguchi K, Sakurai T, Takeshima H, Saito N, Iino M. Nitric oxide-induced calcium release via ryanodine receptors regulates neuronal function. *EMBO J* 2012; 31: 417-428; PMID: 22036948; doi: 10.1038/emboj.2011.386

Abstract

Ryanodine receptors (RyRs), located in the sarcoplasmic/endoplasmic reticulum (SR/ER) membrane, are required for intracellular Ca^{2+} release that is involved in a wide range of cellular functions. In addition to Ca^{2+} -induced Ca^{2+} release in cardiac cells and voltage-induced Ca^{2+} release in skeletal muscle cells, we recently identified another mode of intracellular Ca^{2+} mobilization mediated by RyR, i.e., nitric oxide-induced Ca^{2+} release (NICR), in cerebellar Purkinje cells. NICR is evoked by neuronal activity, is dependent on S-nitrosylation of type 1 RyR (RyR1), and is involved in the induction of long-term potentiation (LTP) of cerebellar synapses. In this addendum, we examined whether peroxynitrite, which is produced by the reaction of nitric oxide with superoxide, may also have an effect on the Ca^{2+} release via RyR1 and the cerebellar LTP. We found that scavengers of peroxynitrite have no significant effect either on the Ca^{2+} release via RyR1 or on the cerebellar LTP. We also found that an application of a high concentration of peroxynitrite does not reproduce neuronal activity-dependent Ca^{2+} release in Purkinje cells. These results support that NICR is induced by endogenous nitric oxide produced by neuronal activity through S-nitrosylation of RyR1.

There are an increasing number of studies showing that protein S-nitrosylation, the addition of a nitric oxide (NO) group to cysteine thiols, provides important roles in a wide range of signaling pathways.^{1,2} Because the ryanodine receptor (RyR), a Ca²⁺ release channel expressed in the SR/ER, has about 100 cysteine residues per subunit and almost half of these thiols are kept in a reduced state under the resting condition,³ RyR is thought to be one of the substrate proteins for S-nitrosylation. In accordance with this notion, the open probability of RyR1 measured in lipid bilayers is increased by NO donors,⁴⁻⁷ and this increase is accompanied by an increase in S-nitrosylation of the Ca²⁺ release channel at cysteine 3635.^{4,7-9} In addition, NO-induced activation of RyR1 in lipid bilayers and in SR vesicles is influenced by pO₂ levels: RyR1 is activated by submicromolar levels of NO in the tissue pO₂ level (~10 mmHg) but not in the ambient O₂ level (~150 mmHg).⁴ In these previous studies, the redox regulations of RyRs were extensively studied using *in vitro* experimental systems, especially in lipid bilayers and SR samples isolated from skeletal muscles.^{1,10,11} Although S-nitrosylation of RyR1 is implicated in the enhancement of Ca²⁺ leak from the skeletal muscle SR in pathological conditions,^{12,13} physiological significance of S-nitrosylation-mediated modulation of RyR1 activity by endogenous NO remained elusive.

We previously found that burst stimulation (BS) of parallel fibers (PFs) generates a localized increase in NO concentration in Purkinje cell (PC) dendrites, as well as NO-dependent long-term potentiation (LTP) at the PF-PC synapse in the cerebellum.¹⁴ Because this LTP is also sensitive to the intracellular Ca²⁺ concentration, we examined a crosstalk between NO and Ca²⁺ signaling in PCs. Bath application of 1-Hydroxy-2-oxo-3-(N-methyl-3-aminopropyl)-3-methyl-1-triazene (NOC7), a NO donor, induced an increase in the intracellular Ca²⁺ concentration in PCs. The NO-induced Ca²⁺ increase was independent of Ca²⁺ influx from the extracellular medium, but was dependent on the activity of RyR1 and the sarco/endoplasmic reticulum Ca²⁺ ATPase (SERCA), suggesting that the Ca²⁺ increase is mediated by Ca²⁺ release from the intracellular store through RyR1. In addition, our study using RyR1-expressing HEK293 cells indicated that S-nitrosylation of RyR1 at cysteine 3635 is both necessary and sufficient for the NO-induced Ca²⁺ increase, which is hereafter referred to as “NO-induced Ca²⁺ release (NICR)”.¹⁵ We then showed that BS to PFs induces intracellular Ca²⁺ release in PC dendrites. When BS which induces NO-dependent-LTP was applied to PF, Ca²⁺ levels were transiently but clearly elevated in PC dendrites. This BS-induced Ca²⁺ increase was indicated to be due to Ca²⁺ release mediated by RyR1,

because the Ca^{2+} increase was abolished by the application of thapsigargin and cyclopiazonic acid, SERCA inhibitors, and dantrolene, a RyR antagonist^{16,17}. In addition, the Ca^{2+} release was inhibited by the application of *N*⁶-Nitro-L-arginine methyl ester (L-NAME), a broad inhibitor of NO synthases (NOS), and was not observed in the cerebellum of mice deficient in neuronal type NOS (nNOS).¹⁸ These results strongly suggest that BS to PFs induces NICR in the PC dendrites.

NO signaling has two types of downstream pathways: activation of soluble guanylyl cyclase (sGC) and S-nitrosylation of target proteins. Because BS-induced NICR was inhibited by intracellular application of ascorbic acid, a reducing reagent, to Purkinje cells, but not by 1H-[1,2,4]oxadiazolo[4,3-a]quinoxalin-1-one (ODQ), an inhibitor of soluble guanylyl cyclase, NICR was suggested to be dependent on S-nitrosylation of proteins including RyR1. Correspondingly, biochemical analysis demonstrated that NICR induced by the application of NOC7 is accompanied with S-nitrosylation of RyR1 in cerebellar slice preparations. Furthermore, all manipulations that we used to inhibit NICR also abolished PF-LTP. Taken together, our study indicated that NICR is a novel Ca^{2+} mobilizing mechanism in neuronal cells and is essential for the induction of PF-LTP.

However, neuronal activity in conjunction with certain forms of synaptic

plasticity may be associated with superoxide generation,¹⁹ and NO generation leads to peroxynitrite formation when superoxide is simultaneously generated.²⁰ Because peroxynitrite is known to regulate cell signaling via molecular modifications, including protein nitration,²⁰ it seems possible that NICR is affected by peroxynitrite produced from endogenous NO in the presence of superoxide. We, therefore, examined the potential role of peroxynitrite in the BS-induced Ca²⁺ release and LTP. Bath application of the peroxynitrite scavenger, uric acid,²¹ and the peroxynitrite decomposition catalyst, 5,10,15,20-Tetrakis(4-sulfonatophenyl)prophyrinato iron (III) (FeTPPS),²² had no effect on BS-induced Ca²⁺ increase (Fig. 1), although the concentration of uric acid (100 μM) and FeTPPS (10 μM) were thought to be high enough.^{23,24} We subsequently examined effects of uric acid and FeTPPS on the induction of PF-LTP. In accordance with the insensitivity of NICR to these reagents, neither uric acid nor FeTPPS impaired the induction of PF-LTP induced by BS (Fig. 2). These results do not support the involvement of peroxynitrite in NICR and PF-LTP induction, although there remains a possibility that the scavengers did not completely antagonizing the effect of peroxynitrite. We therefore examined the effect of an application of peroxynitrite on the intracellular Ca²⁺ concentration of

Purkinje cells in cerebellar slice preparations. Because our previous NO imaging experiments during BS-induced NO generation from parallel fibers indicates that the peak NO concentration in the center of response is about 5 μM ,¹⁴ we applied 10 μM peroxynitrite to the cerebellar slice preparations. However, even at this very high concentration, peroxynitrite induced only very little changes in the Ca^{2+} concentration (Fig. 1D). The slow increase in the Ca^{2+} concentration became absent in the presence of uric acid or FeTPPS. However, it was resistant to an intracellular application of 10 mM ascorbic acid, which completely abolishes BS-induced NICR.¹⁵ Therefore, the peroxynitrite-dependent slow response seems to have a mechanism different from that of NICR and might be accounted for by an irreversible effect of peroxynitrite on the endoplasmic reticulum that has been reported previously.⁵ These results suggest that the role of peroxynitrite, if any, in BS-induced Ca^{2+} release is very little.

In conclusion, in combination with the observations in our previous study,¹⁵ it is suggested that NICR is dependent on the NO-dependent S-nitrosylation of RyR1 rather than on indirect actions of peroxynitrite or cyclic GMP (Fig. 3). Biochemical analysis of RyR1 purified from the cerebellar cortex after the LTP induction may further confirm specific involvement of endogenous NO in NICR and PF-LTP.

Materials and Methods

Preparation of cerebellar slices, Ca²⁺ imaging and electrophysiological analysis were carried out essentially as described previously.^{15,25} C57BL/6 mice (postnatal days 24–32) were sacrificed by cervical dislocation under anesthesia with diethyl ether. The cerebellum was excised, and parasagittal cerebellar slices (250 μm thick) were prepared from the vermis. Whole-cell recordings were obtained from visually identified PCs at room temperature (23–25°C).

For intracellular Ca²⁺ imaging in PCs, a Ca²⁺-sensitive dye, Oregon Green 488 BAPTA-1 (100 μM), was introduced into Purkinje cells through the patch pipette. Five to nine sequential confocal images (excitation at 488 nm), obtained at 3–4 μm z-axis intervals, were acquired every 0.8 s using an upright microscope (BX51WI; Olympus) equipped with a confocal scanning unit and an argon laser (FV300, Olympus), and projected onto a plane to obtain dendrite images at 10 s intervals.

For focal stimulation of PFs, a stimulation pipette was filled with standard bathing solution and used to apply square pulses (0.1 ms in duration, 0–20 V in amplitude) to the molecular layer, specifically in the middle portion of the molecular layer approximately one-third of the

distance below the pial surface. The intensity of each stimulus was adjusted to evoke PF-EPSCs with an amplitude of 60–120 pA. The ionic current was recorded from PCs using a patch-clamp amplifier (EPC-9, HEKA, Lambrecht/Pfalz, Germany) at a holding potential of -90 or -80 mV, after compensating for liquid junction potential. The signals were filtered at 2 kHz and digitized at 20 kHz. After obtaining a stable initial recording for at least 10 min, 60 burst stimulations (1 burst stimulation: 5 pulses at 50 Hz) were repeatedly applied at 1 Hz to induce LTP. Series resistance and membrane resistance were monitored throughout the experiments, and the data were discarded when either of these resistances varied by more than 10%. The data were also discarded when the slope of PF-EPSC amplitude averaged every minute during the initial recording for 10 min was larger than 2% or when the amplitude did not become stable within 20 min after the onset of whole-cell configuration.¹⁴

References

1. Hess DT, Matsumoto A, Kim SO, Marshall HE, Stamler JS. Protein S-nitrosylation: Purview and parameters. *Nature Rev Mol Cell Biol* 2005; 6:150-66.
2. Shahani N, Sawa A. Nitric Oxide Signaling and Nitrosative Stress in Neurons: Role for S-Nitrosylation. *Antioxid Redox Signal* 2011; 14:1493-504.
3. Zalk R, Lehnart SE, Marks AR. Modulation of the ryanodine receptor and intracellular calcium. *Annu Rev Biochem* 2007; 76:367-85.
4. Eu JP, Sun JH, Xu L, Stamler JS, Meissner G. The skeletal muscle calcium release channel: Coupled O₂ sensor and NO signaling functions. *Cell* 2000; 102:499-509.
5. Stoyanovsky D, Murphy T, Anno PR, Kim YM, Salama G. Nitric oxide activates skeletal and cardiac ryanodine receptors. *Cell Calcium* 1997; 21:19-29.
6. Sun JH, Xu L, Eu JP, Stamler JS, Meissner G. Nitric oxide, NOC-12, and S-nitrosoglutathione modulate the skeletal muscle calcium release channel/ryanodine receptor by different mechanisms - An allosteric function for O₂ in S-nitrosylation of the channel. *J Biol Chem* 2003; 278:8184-9.

7. Xu L, Eu JP, Meissner G, Stamler JS. Activation of the cardiac calcium release channel (ryanodine receptor) by poly-S-nitrosylation. *Science* 1998; 279:234-7.
8. Moore CP, Zhang JZ, Hamilton SL. A role for cysteine 3635 for RYR1 in redox modulation and calmodulin binding. *J Biol Chem* 1999; 274:36831-4.
9. Sun JH, Xin CL, Eu JP, Stamler JS, Meissner G. Cysteine-3635 is responsible for skeletal muscle ryanodine receptor modulation by NO. *Proc Natl Acad Sci U S A* 2001; 98:11158-62.
10. Hidalgo C, Donoso P, Carrasco MA. The ryanodine receptors Ca²⁺ release channels: Cellular redox sensors? *IUBMB Life* 2005; 57:315-22.
11. Lanner JT, Georgiou DK, Joshi AD, Hamilton SL. Ryanodine Receptors: Structure, Expression, Molecular Details, and Function in Calcium Release. *Cold Spring Harb Perspect Biol* 2010; 2:a003996.
12. Bellinger AM, Reiken S, Carlson C, Mongillo M, Liu X, Rothman L, et al. Hypernitrosylated ryanodine receptor calcium release channels are leaky in dystrophic muscle. *Nature Med* 2009; 15:325-30.
13. Durham WJ, Aracena-Parks P, Long C, Rossi AE, Goonasekera SA, Boncompagni S, et al. RyR1 S-nitrosylation underlies environmental heat stroke and sudden death in Y522S RyR1 knockin mice. *Cell* 2008;

133:53-65.

14. Namiki S, Kakizawa S, Hirose K, Iino M. NO signalling decodes frequency of neuronal activity and generates synapse-specific plasticity in mouse cerebellum. *J Physiol* 2005; 566:849-63.
15. Kakizawa S, Yamazawa T, Chen Y, Ito A, Murayama T, Oyamada H, et al. Nitric oxide-induced calcium release via ryanodine receptors regulates neuronal function. *EMBO J* 2012; 31:417-28.
16. Kobayashi S, Yano M, Suetomi T, Ono M, Tateishi H, Mochizuki M, et al. Dantrolene, a therapeutic agent for malignant hyperthermia, markedly improves the function of failing cardiomyocytes by stabilizing interdomain interactions within the ryanodine receptor. *J Am Coll Cardiol* 2009; 53:1993-2005.
17. Zhao F, Li P, Chen SR, Louis CF, Fruen BR. Dantrolene inhibition of ryanodine receptor Ca^{2+} release channels. Molecular mechanism and isoform selectivity. *J Biol Chem* 2001; 276:13810-6.
18. Huang PL, Dawson TM, Bredt DS, Snyder SH, Fishman MC. Targeted Disruption of the Neuronal Nitric-Oxide Synthase Gene. *Cell* 1993; 75:1273-86.
19. Kishida KT, Klann E. Sources and targets of reactive oxygen species in synaptic plasticity and memory. *Antioxid Redox Signal* 2007; 9:233-44.

20. Pacher P, Beckman JS, Liaudet L. Nitric oxide and peroxynitrite in health and disease. *Physiol Rev* 2007; 87:315-424.
21. Becker BF. TOWARDS THE PHYSIOLOGICAL-FUNCTION OF URIC-ACID. *Free Radic Biol Med* 1993; 14:615-31.
22. Stern MK, Jensen MP, Kramer K. Peroxynitrite decomposition catalysts. *J Am Chem Soc* 1996; 118:8735-6.
23. Lancel S, Tissier S, Mordon S, Marechal X, Depontieu F, Scherpereel A, et al. Peroxynitrite decomposition catalysts prevent myocardial dysfunction and inflammation in endotoxemic rats. *J Am Coll Cardiol* 2004; 43:2348-58.
24. Tsukada K, Hasegawa T, Tsutsumi S, Katoh H, Kuwano H, Miyazaki T, et al. Effect of uric acid on liver injury during hemorrhagic shock. *Surgery* 2000; 127:439-46.
25. Kakizawa S, Yamasaki M, Watanabe M, Kano M. Critical period for activity-dependent synapse elimination in developing cerebellum. *J Neurosci* 2000; 20:4954-61.

Figure Legends

Figure 1. NO-induced Ca^{2+} release (NICR) is not affected by peroxynitrite.


(A-C) Ca^{2+} response, induced by a series of 60 BS at 1 Hz (BS; Gray bar) in the standard ACSF (A; data were obtained from 10 cells in 10 slices), was not inhibited by peroxynitrite scavenger, uric acid (B; 100 μM , $n = 7$ cells in 7 slices), or peroxynitrite decomposition catalyst, FeTPPS (C; 10 μM , $n = 6$ cells in 6 slices). Uric acid or FeTPPS was present in bathing solution from 30 min before the recording. (D) Ca^{2+} response to bath application of 10 μM peroxynitrite (ONOO^-) (open circle; $n = 6$ cells, 6 slices). It was abolished by 100 μM uric acid (open triangle; $n = 7$ cells, 7 slices) or 10 μM FeTPPS (closed triangle; $n = 7$ cells, 7 slices), but was insensitive to 10 mM ascorbic acid applied intracellularly through the patch pipette (closed circle; $n = 8$ cells; 8 slices). (E) Summary of data in A–D. The peak Ca^{2+} responses are shown. UA: uric acid; Fe: FeTPPS; Asc: ascorbic acid. Data are presented as Mean \pm SEM. $*p < 0.05$, ANOVA / Tukey-Kramer test.

Figure 2. Cerebellar LTP is not affected by peroxynitrite. (A-C)


Cerebellar LTP, induced by BS in the standard ACSF (A; $n = 8$ cells in 8

slices), was not inhibited by 100 μ M uric acid (B; n = 5 cells in 5 slices) or 10 μ M FeTPPS (C; n = 5 cells in 5 slices). (D) Effects of uric acid (UA) and FeTPPS on the magnitude of LTP (normalized EPSC averaged between 21-30 min after BS). Data are presented as Mean \pm SEM.


Figure 3. A schematic diagram of signaling pathways for nitric oxide-induced Ca^{2+} release (NICR) in cerebellar Purkinje cells. Nitric oxide (NO) is produced at the parallel-fiber terminal in response to neuronal activity, and diffuses into the cerebellar Purkinje cell. Then, type 1 ryanodine receptors (RyR1), located in the endoplasmic reticulum (ER) membrane, are S-nitrosylated and activated, resulting in Ca^{2+} release from the ER. Cyclic GMP and peroxynitrite, which is produced by the reaction of NO and superoxide, are unlikely to be involved in NICR. nNOS: neuronal NO synthase; $\text{IP}_3\text{R1}$: inositol 1,4,5-trisphosphate receptor; SERCA: sarco/endoplasmic reticulum Ca^{2+} ATPase.



Kakizawa *et al.* Figure 1



Kakizawa *et al.* Figure 2



Kakizawa *et al.* Figure 3