

Title	STLV-1-infected Japanese macaque as a model of HTLV-1 infection
Author(s)	Miura, Michi; Tanabe, Junko; Sugata, Kenji; Zhao, Tiejun; Ma, Guangyong; Miyazato, Paola; Yasunaga, Jun-ichiro; Matsuoka, Masao
Citation	Retrovirology (2014), 11(Suppl 1)
Issue Date	2014-01-07
URL	http://hdl.handle.net/2433/180648
Right	© 2014 Miura et al; licensee BioMed Central Ltd.
Type	Journal Article
Textversion	publisher



ORAL PRESENTATION

Open Access

STLV-1-infected Japanese macaque as a model of HTLV-1 infection

Michi Miura, Junko Tanabe, Kenji Sugata, Tiejun Zhao, Guangyong Ma, Paola Miyazato, Jun-ichiro Yasunaga, Masao Matsuoka*

From 16th International Conference on Human Retroviruses: HTLV and Related Viruses
Montreal, Canada. 26-30 June 2013

Various non-human primates are the natural hosts of simian T-cell leukemia virus type 1 (STLV-1). In the present study, we analyzed Japanese macaques naturally infected with STLV-1, and evaluated them as an animal model for HTLV-1 research. Approximately 60% of individuals in the colony are seropositive for STLV-1. Clonal proliferation of STLV-1⁺ cells was investigated by massively sequencing the provirus integration sites. We found that some clones proliferated distinctively in monkeys with higher proviral load. T lymphocytes expressing Tax in the peripheral blood were largely CD4⁺. Notably, one of the monkeys surveyed in this study developed T-cell lymphoma in the brain, indicating that STLV-1 is oncogenic in Japanese macaques. We also assessed the molecular function of STLV-1 Tax and STLV-1 bZIP factor (SBZ). STLV-1 Tax activated NFAT, AP-1, canonical Wnt and canonical NF-kappa B pathways, whereas SBZ suppressed those signaling pathways. SBZ enhanced TGF-beta signaling, but STLV-1 Tax suppressed it. These findings suggest that STLV-1 Tax and SBZ have similar functions to their counterparts of HTLV-1. In addition, we found that administration of anti-CCR4 antibody, which is currently used in Japan for the treatment of ATL patients, efficiently reduced proviral load in STLV-1-infected Japanese macaques. Our study provides the evidence that Japanese macaques naturally infected with STLV-1 correspond to HTLV-1 carriers and are a suitable animal model to investigate the pathogenesis of HTLV-1 and novel therapeutic strategies.

Published: 7 January 2014

* Correspondence: mmatsuok@virus.kyoto-u.ac.jp
Institute for Virus Research, Kyoto University, Kyoto, Japan

doi:10.1186/1742-4690-11-S1-O12

Cite this article as: Miura *et al.*: STLV-1-infected Japanese macaque as a model of HTLV-1 infection. *Retrovirology* 2014 11(Suppl 1):O12.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

