


Title	Noninvasive regional cerebral oxygen saturation for neurological prognostication of patients with out-of-hospital cardiac arrest: a prospective multicenter observational study.
Author(s)	Ito, Noritoshi; Nishiyama, Kei; Callaway, Clifton W; Orita, Tomohiko; Hayashida, Kei; Arimoto, Hideki; Abe, Mitsuru; Endo, Tomoyuki; Murai, Akira; Ishikura, Ken; Yamada, Noriaki; Mizobuchi, Masahiro; Anan, Hideki; Okuchi, Kazuo; Yasuda, Hideto; Mochizuki, Toshiaki; Tsujimura, Yuka; Nakayama, Takeo; Hatanaka, Tetsuo; Nagao, Ken
Citation	Resuscitation (2014), 85(6): 778-784
Issue Date	2014-06
URL	<a href="http://hdl.handle.net/2433/188884">http://hdl.handle.net/2433/188884</a>
Right	© 2014 The Authors. Published by Elsevier Ireland Ltd. This is an open access article under the CC BY-NC-SA license ( <a href="http://creativecommons.org/licenses/by-nc-sa/3.0/">http://creativecommons.org/licenses/by-nc-sa/3.0/</a> ).
Type	Journal Article
Textversion	publisher



## Clinical Paper

# Noninvasive regional cerebral oxygen saturation for neurological prognostication of patients with out-of-hospital cardiac arrest: A prospective multicenter observational study<sup>☆,☆☆</sup>



Noritoshi Ito <sup>a,b</sup>, Kei Nishiyama <sup>c,\*</sup>, Clifton W. Callaway <sup>d</sup>, Tomohiko Orita <sup>e</sup>, Kei Hayashida <sup>f</sup>, Hideki Arimoto <sup>g</sup>, Mitsuru Abe <sup>h</sup>, Tomoyuki Endo <sup>i</sup>, Akira Murai <sup>j</sup>, Ken Ishikura <sup>k</sup>, Noriaki Yamada <sup>l</sup>, Masahiro Mizobuchi <sup>m</sup>, Hideki Anan <sup>n</sup>, Kazuo Okuchi <sup>o</sup>, Hideto Yasuda <sup>p</sup>, Toshiaki Mochizuki <sup>q</sup>, Yuka Tsujimura <sup>r</sup>, Takeo Nakayama <sup>r</sup>, Tetsuo Hatanaka <sup>s</sup>, Ken Nagao <sup>t</sup>, for the J-POP Registry Investigators<sup>u</sup>

<sup>a</sup> Senri Cardiovascular Center, Osaka Saiseikai Senri Hospital, 1-1-6 Tsukumodai, Suita, Osaka 565-0862, Japan

<sup>b</sup> Department of Cardiovascular Medicine, Osaka University Graduate School of Medicine, 2-2 Yamadaoka, Suita, Osaka 565-0871, Japan

<sup>c</sup> Department of Primary Care and Emergency Medicine, Kyoto University Graduate School of Medicine, 54 Kawara-machi, Seigo-in, Sakyo-ku, Kyoto 606-8507, Japan

<sup>d</sup> Department of Emergency Medicine, University of Pittsburgh, Iroquois Building, Suite 400A, 3600 Forbes Avenue, Pittsburgh, PA 15261, United States

<sup>e</sup> Department of Emergency and Critical Care Medicine, Saiseikai Yokohamashi Tobu Hospital, 3-6-1 Shimosueyoshi, Tsurumi-ku, Yokohama, Kanagawa 230-0012, Japan

<sup>f</sup> Emergency and Critical Care Medicine, Keio University School of Medicine, 35 Shinanomachi Shinjuku-ku, Tokyo 160-8582, Japan

<sup>g</sup> Department of Emergency and Critical Care Medical Center, Osaka City General Hospital, 2-13-22 Miyakojima Hon-Dori, Miyakojima-ku, Osaka 534-0021, Japan

<sup>h</sup> Department of Cardiology, National Hospital Organization Kyoto Medical Center, 1-1 Fukakusumakuhatachou, Fushimi-ku, Kyoto 612-8555, Japan

<sup>i</sup> Advanced Emergency Center, Tohoku University Hospital, 1-1 Seiryo-machi, Aoba-ku, Sendai, Miyagi 980-8574, Japan

<sup>j</sup> Department of Emergency and Critical Care Medicine, Fukuoka University Hospital, 7-45-1 Nanakuma, Jyonan-ku, Fukuoka 814-0180, Japan

<sup>k</sup> Critical Care Center, Mie University Hospital, 174 Edobashi 2-chome, Tsu, Mie 514-8507, Japan

<sup>l</sup> Advanced Critical Care Center, Gifu University Hospital, 1-1 Yanagido, Gifu 501-1194, Japan

<sup>m</sup> Department of Cardiology, Kyoto Katsura Hospital, 17-Banchi, Yamada Hirao-cho, Nishikyo-ku, Kyoto 615-8256, Japan

<sup>n</sup> Division of Emergency Medicine, Fujisawa City Hospital, 2-6-1 Fujisawa, Kanagawa 251-8550, Japan

<sup>o</sup> Advanced Critical Care Center, Nara Medical University Hospital, 840 Shijocho, Kashihara, Nara 634-8522, Japan

<sup>p</sup> Emergency and Critical Care Medicine, Japanese Red Cross Musashino Hospital, 1-26-1 Sakai Minami Town, Musashino, Tokyo 180-8610, Japan

<sup>q</sup> Department of Emergency and Critical Care Medicine St Luke's International Hospital, 9-1 Akashi-cho, Chuo-ku, Tokyo 104-8560, Japan

<sup>r</sup> Department of Health Informatics, Kyoto University School of Public Health, Yoshida-Konoe-cho, Sakyo-ku, Kyoto 606-8501, Japan

<sup>s</sup> Emergency Life Saving Technique Academy, 3-8-1 Oura, Yahata-nishi-ku, Kitakyushu, Fukuoka 807-0874, Japan

<sup>t</sup> Department of Cardiology, Surugadai Nihon University Hospital, 1-8-13 Kanda-Surugadai, Chiyoda-ku, Tokyo 101-8309, Japan

## ARTICLE INFO

## Article history:

Received 5 December 2013

Received in revised form 5 February 2014

Accepted 14 February 2014

## Keywords:

Cardiac arrest

Cardiopulmonary resuscitation

Emergency department

Predictors

Regional brain oxygen saturation

## ABSTRACT

**Aim:** To investigate the association between regional brain oxygen saturation ( $rSO_2$ ) at hospital arrival and neurological outcomes at 90 days in patients with out-of-hospital cardiac arrest (OHCA).

**Methods:** The Japan-Prediction of neurological Outcomes in patients post cardiac arrest (J-POP) registry is a prospective, multicenter, cohort study to test whether  $rSO_2$  predicts neurological outcomes after OHCA. We measured  $rSO_2$  in OHCA patients immediately after hospital arrival using a near-infrared spectrometer placed on the forehead with non-blinded fashion. The primary endpoint was "neurological outcomes" at 90 days after OHCA.

**Results:** EMS providers are not permitted to terminate CPR in the field in Japan, and so most patients with OHCA who are treated by EMS personnel are transported to emergency hospitals. Among 1017 OHCA patients, 672 patients including 52 comatose patients with pulses detectable (8%) and 620 cardiac arrest patients (92%) at hospital arrival were enrolled prospectively and consecutively. Twenty-nine

<sup>☆</sup> A Spanish translated version of the summary of this article appears as Appendix in the final online version at <http://dx.doi.org/10.1016/j.resuscitation.2014.02.012>.

<sup>☆☆</sup> Previous presentation: This manuscript was presented in part at the American Heart Association Late-Breaking Resuscitation Science 2012 in Los Angeles, CA, November 4, 2012.

\* Corresponding author.

E-mail address: [keinishi@kuhp.kyoto-u.ac.jp](mailto:keinishi@kuhp.kyoto-u.ac.jp) (K. Nishiyama).

<sup>u</sup> Additional members of the J-POP Registry are listed in Appendix A.

patients with good neurological outcome had a significantly higher value of rSO<sub>2</sub> at hospital arrival than 643 patients with poor neurological outcome (mean [ $\pm$ SD] 55.6  $\pm$  20.8% vs. 19.7  $\pm$  11.0%,  $p < 0.001$ ). Receiver operating curve analysis indicated an optimal rSO<sub>2</sub> cutoff point of >42% for predicting good neurological outcome, with sensitivity 0.79 (95% confidence interval [CI], 0.60–0.92), specificity 0.95 (95% CI, 0.93–0.96), positive predictive value, 0.41 (95% CI, 0.28–0.55), negative predictive value, 0.99 (95% CI, 0.98–1.00), and area under the curve 0.90 (95% CI, 0.88–0.92).

**Conclusion:** The rSO<sub>2</sub> at hospital arrival can predict good neurological outcome at 90 days after OHCA.

© 2014 The Authors. Published by Elsevier Ireland Ltd. This is an open access article under the CC BY-NC-SA license (<http://creativecommons.org/licenses/by-nc-sa/3.0/>).

## 1. Introduction

Sudden death from cardiac arrest is a major public health problem in industrialized countries.<sup>1–4</sup> The brain is the organ most vulnerable to ischemia, and global cerebral ischemia often results in poor neurological outcome for patients with out-of-hospital cardiac arrest (OHCA).<sup>5</sup> The 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care have emphasized that cardiocerebral resuscitation (CCR) is critically important to promote the survival of patients with OHCA.<sup>1–4</sup> A real-time indicator of cerebral perfusion or function during CCR efforts would be useful for estimating the chance of favorable rehabilitation after hospital discharge.<sup>6,7</sup>

Regional cerebral oxygen saturation (rSO<sub>2</sub>) is a noninvasive indicator of cerebral perfusion. Even during cardiac arrest, rSO<sub>2</sub> is measurable because near-infrared spectroscopy (NIRS) does not require vascular pulsation.<sup>8,9</sup> Therefore, we conducted this prospective, multicenter, observational study to validate the ability of rSO<sub>2</sub> at hospital arrival to predict neurological outcomes at 90 days<sup>10</sup> after OHCA.

## 2. Methods

### 2.1. Study design

The Japan-Prediction of neurological Outcomes in patients Post-cardiac arrest (J-POP) Registry is a prospective, multicenter, cohort study. From May 15, 2011, through January 31, 2012, 13 tertiary emergency hospitals in Japan participated in this study. Of 1017 OHCA patients who were transported to the hospital, 672 were enrolled prospectively and consecutively. The inclusion criteria were unresponsive patients during and after resuscitation at hospital arrival after OHCA. The exclusion criteria were as follows: (1) trauma, (2) accidental hypothermia, (3) age under 18, (4) previous completion of the “Do Not Attempt Resuscitation” form, and (5) a Glasgow Coma Scale (GCS) score of >8 at hospital arrival. The study protocol was approved by the institutional review board or ethics committee at each participating medical institution. The requirement of written informed consent was waived according to the guidelines.<sup>11</sup>

### 2.2. Emergency medical care system in Japan

All emergency medical service (EMS) providers perform cardiopulmonary resuscitation (CPR) according to the current CPR guidelines.<sup>12,13</sup> Emergency lifesaving technicians are permitted to insert tracheal tubes and to administer intravenous epinephrine.<sup>1</sup> However, EMS providers are not permitted to terminate CPR in the field.

### 2.3. Data collection

Using the Utstein style,<sup>14</sup> data were collected prospectively. Baseline patient characteristics and in-hospital information were

collected from medical records and databases. Data management and statistical analyses were performed by an independent data coordinating center (Kyoto University Graduate School of Medicine, Kyoto, Japan).

Cardiac arrest was defined as the absence of spontaneous respiration, palpable pulse, and responsiveness to stimuli.<sup>14,15</sup> The arrest was presumed to be of cardiac origin unless it was caused by cerebrovascular disease, respiratory disease, external factors (e.g., drug overdose and asphyxia), or any other noncardiac factors. The cardiac or noncardiac origin was determined clinically by the physician in charge.

### 2.4. Procedures after hospital arrival


The CPR protocol is as described below. After arrival at the emergency department, all patients receive advanced life support (ALS) in accordance with the national guidelines for resuscitation. Patients, whose initially documented ECG rhythm was ventricular fibrillation (VF) or pulseless ventricular tachycardia (VT), receive extracorporeal CPR using extracorporeal circulatory support or cardiopulmonary bypass if sustained return of spontaneous circulation (ROSC)—restoration of palpable pulse that is sustained for at least 20 min—was not obtained by standard ALS. For patients with sustained ROSC, therapeutic hypothermia is conducted if systolic blood pressure increases above 90 mmHg and the GCS score remains  $\leq 8$ .<sup>16</sup>

### 2.5. Near-infrared spectroscopy procedures

Immediately after hospital arrival aiming within 3 min, two disposable probes of a near-infrared spectrometer (INVOS™ 5100C, Covidien, Boulder, CO, USA) were applied carefully and bilaterally onto the patient's forehead. rSO<sub>2</sub>, once stabilized in several seconds, was monitored at least for 1 min with the probes, and the lowest rSO<sub>2</sub> reading was used.<sup>8,9</sup> The cerebral oximeter emits 2 wavelengths of near-infrared rays (730 and 805 nm) into the patient's forehead, calculates spatial depth resolution by subtracting the shallow measurement from the deep, minimizes superficial signal contamination from the scalp and skull, and detects changes in oxygen saturation in the brain.<sup>8,9</sup> We could not blind investigators because the rSO<sub>2</sub> monitoring requires real-time visual confirmation during CCR efforts. Regardless of the rSO<sub>2</sub> readings, however, all patients received the best available therapy including post cardiac arrest interventions.<sup>17</sup>

### 2.6. Study endpoints

The primary endpoint was “neurological outcomes” at 90 days after OHCA according to the American Heart Association consensus statement,<sup>10</sup> which were categorized in accordance with Glasgow-Pittsburgh Cerebral Performance Categories described in the “Utstein style” guidelines<sup>14</sup>; the guidelines defined cerebral performance category (CPC) levels as follows: CPC 1 (good performance) and CPC 2 (moderate disability) as “good neurological



**Fig. 1.** Enrollment of study patients.

outcome"; and CPC 3 (severe disability), CPC 4 (vegetative state), and CPC 5 (brain death or death) as "poor neurological outcome". CPC was evaluated by more than two physicians in charge who were not aware of the rSO<sub>2</sub> readings at hospital arrival.

## 2.7. Statistical analyses

The unpaired *t*-test or the Mann–Whitney test was conducted for unpaired comparisons. The chi-square test and Fisher's exact test were made to examine differences between categorical variables. The ROC analysis was performed to evaluate the prognostic accuracy of good neurological outcome, and the optimal cutoff point was determined by estimating the Zero-One index. In patients to whom lactate and base excess (BE) were measured concurrently with rSO<sub>2</sub> at hospital arrival, we compared the AUC of rSO<sub>2</sub>, lactate, and BE, and the method of DeLong et al.<sup>18</sup> was used to for the calculation of the standard error of the AUC and the difference between two AUCs. All statistical tests were two-tailed; a *p* value less than 0.05 was considered statistically significant.

The JMP software version 10.0.0 (SAS Institute, Cary, NC, USA), the MedCalc software version 12.3.0 (MedCalc Software, Gent, Belgium), and the STATA software version 11.1 (Stata Corp., College Station, TX, USA) were used by two authors (N.I. and K.N.) to make all statistical analyses. The primary authors had full access to the data and take responsibility for its integrity. All authors have read and agreed to the manuscript as written.

## 2.8. Roles of the funding source

The funding organizations had no role in study design, data analysis, data interpretation, writing of the article, or the decision to submit it for publication. The corresponding author had full access to all the data and had final responsibility for the decision to submit for publication.

## 3. Results

### 3.1. General characteristics

During the study period, we collected data on 1017 consecutive patients with OHCA who were referred to 13 tertiary emergency hospitals. Patients were excluded because of the following causes: 111 patients, cardiac arrest caused by trauma; 19 patients, age under 18; 131 patients, the completion of the "Do Not Attempt Resuscitation Order" form; six patients, a GCS score of >8 at hospital arrival; five patients, accidental hypothermia; one patient, missing data; and 72 patients, no monitoring of rSO<sub>2</sub> due to insufficient personnel (59), shortage of disposable probes (8), or technical problems (5, i.e., no display of monitored values). Consequently, we consecutively enrolled a total of 672 unresponsive patients during and after resuscitation at hospital arrival (Fig. 1). Table 1 shows the characteristics and neurological outcomes of patients; among them, 482 (71.7%) were pronounced dead in the emergency department. Of remaining 190 patients, 152 (22.6%) died after admission to the hospital, whereas 38 (5.7%) survived at 90 days after OHCA, with CPCs 1, 2, 3, and 4 in 23 (3.4%), 6 (0.9%), 1 (0.2%), and 8 patients (1.2%), respectively.

### 3.2. Regional cerebral oxygen saturation at hospital arrival

For the enrolled 672 patients, the mean ( $\pm$ SD) rSO<sub>2</sub> at hospital arrival was  $21 \pm 14\%$ . Twenty-nine patients with good neurological outcome (CPC1, 2 at 90 days after OHCA) had a significantly higher value of rSO<sub>2</sub> at hospital arrival than 643 patients with poor neurological outcome ( $55.6 \pm 20.8\%$  vs.  $19.7 \pm 11.0\%$ ,  $p < 0.001$ ) (Fig. 2). Fifty-two comatose patients with pulses detectable at hospital arrival had a significantly higher value of rSO<sub>2</sub> at hospital arrival than 620 patients with cardiac arrest at hospital arrival ( $51.0 \pm 20.2\%$  vs.  $18.8 \pm 9.5\%$ ,  $p < 0.001$ ). One hundred eighty patients with transient or sustained ROSC after hospital arrival had a significantly higher value of rSO<sub>2</sub> at hospital arrival than 492

**Table 1**

Characteristics and neurological outcomes of 672 consecutive unresponsive patients during and after resuscitation at hospital arrival.


Characteristics	Total (n = 672)
Age—year, mean (SD)	71 (16)
Male gender—no. (%)	406 (60)
Locations of cardiac arrest—no. (%)	
Home	463 (69)
Nursing home/assisted living	61 (9)
Public building	33 (5)
Street	29 (4)
Others	86 (13)
Type of bystander-witness status—no. (%)	
No witness	361 (47)
Family members	227 (30)
EMS	48 (6)
Others	128 (17)
Bystander-initiated CPR—no. (%)	263 (39)
Types of origin—no. (%)	
Cardiac	351 (52)
Non-cardiac	321 (48)
Initially documented rhythms on the scene of cardiac arrest—no. (%)	
VF/pulseless VT	82 (12)
PEA	167 (25)
Asystole	423 (63)
Prehospital procedures—no. (%)	
Advanced airway devices	352 (52)
Intravenous epinephrine administration	126 (19)
Defibrillation	98 (15)
Emergency call to hospital arrival—min, mean (SD)	33 (12)
Rhythms at rSO <sub>2</sub> measurement—no. (%)	
VF/pulseless VT	25 (3.7)
PEA	114 (17)
Asystole	481 (71.6)
Others (pulses detectable at hospital arrival)	52 (7.7)
Procedures after hospital arrival—no. (%)	
Standard ALS	654 (97)
Extracorporeal CPR	40 (6)
Therapeutic hypothermia	81 (12)
Coronary angiography	63 (9)
Primary percutaneous coronary intervention	29 (4)
Neurological outcomes at 90 days after OHCA	
Survival (CPCs 1–4)	38 (5.6)
CPC 1, good performance	23 (3.4)
CPC 2, moderate disability	6 (0.9)
CPC 3, severe disability	1 (0.2)
CPC 4, vegetative state	8 (1.2)
Deaths (CPC 5)	634 (94.3)
Deaths pronounced in the emergency department	482 (71.7)
Deaths after hospital admission	152 (22.6)

ALS, advanced cardiac life-support; CPC, cerebral performance category; CPR, cardiopulmonary resuscitation; EMS, emergency medical service; PEA, pulseless electrical activity; rSO<sub>2</sub>, regional cerebral oxygen saturation; SD, standard deviation; VF, ventricular fibrillation; VT, ventricular tachycardia; OHCA, out-of-hospital cardiac arrest.

patients without ROSC ( $28.7 \pm 19.1\%$  vs.  $18.5 \pm 9.8\%$ ,  $p < 0.001$ ). One hundred ninety patients who survived to hospital admission had a significantly higher value of rSO<sub>2</sub> at hospital arrival than 482 patients who were pronounced dead in the emergency department ( $31.6 \pm 21.1\%$  vs.  $17.2 \pm 5.3\%$ ,  $p < 0.001$ ).

### 3.3. Relationship of regional cerebral oxygen saturation to neurological outcomes

The optimal cutoff point in the ROC analysis for predicting good neurological outcome in all 672 patients was rSO<sub>2</sub> >42%; sensitivity, 0.79 (95% CI, 0.60–0.92); specificity, 0.95 (95% CI, 0.93–0.96); positive predictive value, 0.41 (95% CI, 0.28–0.55); negative predictive value, 0.99 (95% CI, 0.98–1.00); AUC, 0.90 (95% CI, 0.88–0.92); and  $p < 0.001$  (Fig. 3A).



**Fig. 2.** Relationships between cerebral performance category and regional cerebral oxygen saturation. <sup>a</sup> Of 446 (66%) patients with an rSO<sub>2</sub> value of  $\leq 15\%$ , 441 were categorized to CPC 5. CPC, cerebral performance category; rSO<sub>2</sub>, regional cerebral oxygen saturation.

ROC analysis of subsets of patients demonstrated that the optimal cutoff point for predicting good neurological outcome was rSO<sub>2</sub> >62% in comatose patients with pulses detectable at hospital arrival ( $n = 52$ ) (AUC: 0.75 [95% CI, 0.61–0.86],  $p < 0.001$  Fig. 3B). In patients with persistent cardiac arrest at hospital arrival, on the other hand, the optimal cutoff point for predicting good neurological outcome was rSO<sub>2</sub> >21% (AUC, 0.73 [95% CI, 0.69–0.76],  $p = 0.04$ , Fig. 3C).

In a post hoc analysis, we categorized all 672 subjects into three groups based on cutoff values of rSO<sub>2</sub>  $\leq 15\%$  ("15%" reading of this device is interpreted as equivalent to no detectable cortical oxygen<sup>8,9</sup>) and rSO<sub>2</sub> of >42% (optimal cutoff in ROC analysis above): patients with an rSO<sub>2</sub>  $\leq 15\%$  ( $n = 449$ ); patients with an rSO<sub>2</sub> = 16–42% ( $n = 167$ ); and patients with an rSO<sub>2</sub> >42% ( $n = 56$ ). Good neurological outcome occurred in 3/449 (0.7%) with an rSO<sub>2</sub> value  $\leq 15\%$ ; 3/167 (1.8%) with an rSO<sub>2</sub> value of 16–42%, and 23/56 (41%) with an rSO<sub>2</sub> value >42% (Fig. 4). A significant difference ( $p < 0.001$ ) was found among categorical variables.

Among the 52 comatose patients with pulses detectable at hospital arrival, only 35 (67%) had rSO<sub>2</sub> value >42%, of whom 20 (57%) had good neurological outcome. Only 1 (6%) of 17 patients with rSO<sub>2</sub> value of  $\leq 42\%$ , had good neurological outcome (rSO<sub>2</sub> = 29%).


### 3.4. Comparisons of regional cerebral oxygen saturation, lactate, and base excess

In 633 patients (94%), lactate and BE were measured concurrently with rSO<sub>2</sub> at hospital arrival. The AUCs for rSO<sub>2</sub>, lactate, and BE to predict good neurological outcome were 0.90 (95% CI, 0.88–0.92), 0.84 (95% CI, 0.80–0.86), and 0.79 (95% CI, 0.76–0.82), respectively (Table 2). The AUC was greater with respect to the prognostication of good neurological outcome for rSO<sub>2</sub> than for BE ( $p = 0.007$ ).

## 4. Discussion

### 4.1. Regional cerebral oxygen saturation at hospital arrival and neurological outcomes at 90 days after OHCA

This study demonstrated that the rSO<sub>2</sub> at hospital arrival can predict good neurological outcome at 90 days after OHCA, with high specificity and positive predictive value (Figs. 3 and 4; Table 2). We determined that an rSO<sub>2</sub> value of >42% at hospital arrival was an optimal value for predicting good neurological outcome after OHCA. Furthermore, we note that the rSO<sub>2</sub> is a neurological



**Fig. 3.** The area under the receiver operating curve of regional cerebral oxygen saturation to predict good neurological outcome at 90 days after out-of-hospital cardiac arrest. Panel A: all unresponsive patients during and after resuscitation at hospital arrival ( $n=672$ ). Panel B: Comatose patients with pulses detectable at hospital arrival ( $n=52$ ). Panel C: Patients with cardiac arrest at hospital arrival ( $n=620$ ). CI, confidence interval; OHCA, out-of-hospital cardiac arrest; rSO<sub>2</sub>, regional cerebral oxygen saturation.

**Table 2**

Characteristics of prognostic indexes for good neurological outcome at 90 days: regional cerebral oxygen saturation, lactate, and base excess.

	Optimal cutoff	AUC (95% CI)	p-Value	Sensitivity (95% CI)	Specificity (95% CI)	PPV (95% CI)	NPV (95% CI)	p-Value (versus rSO <sub>2</sub> )
rSO <sub>2</sub>	>42%	0.90 (0.88–0.92)	<0.001	0.79 (0.60–0.92)	0.95 (0.93–0.96)	0.41 (0.28–0.55)	0.99 (0.98–1.00)	N/A
Lactate	<10.2 mmol/L	0.82 (0.79–0.85)	<0.001	0.76 (0.61–0.88)	0.76 (0.72–0.79)	0.16 (0.11–0.22)	0.98 (0.97–0.99)	0.110
Base excess	>-17.4 mmol/L	0.79 (0.76–0.82)	<0.001	0.95 (0.84–0.99)	0.55 (0.51–0.58)	0.11 (0.08–0.15)	0.99 (0.98–1.00)	0.007

AUC, area under the curve; CI, confidence interval; PPV, positive predictive value; N/A, not applicable; NPV, negative predictive value; rSO<sub>2</sub>, regional cerebral oxygen saturation.

prognostic index with advantages over lactate and base excess because of its noninvasiveness and immediate availability, as well as its specificity, positive predictive value, and AUC (Table 2). The AUC was greater for rSO<sub>2</sub> than for BE which had been shown to be associated with poor neurological outcome.<sup>19,20</sup> We could measure rSO<sub>2</sub> non-invasively immediately after hospital arrival, however it took at least 10 min to obtain the BE and lactate values at the hospital,<sup>21,22</sup> because of technical difficulties associated with blood sampling in patients with OHCA.


Even when patients had pulses detectable at hospital arrival, they had a very low chance of obtaining good neurological outcome unless achieving the rSO<sub>2</sub> >42% at hospital arrival. These data suggest that in patients without the rSO<sub>2</sub> value of >42%, regardless of the presence or absence of pulses detectable at hospital arrival,

poor cerebral perfusion may persist from the onset of cardiac arrest through to the monitoring of rSO<sub>2</sub>. Consequently, irreversible cerebral ischemia may preclude good recovery.

There is also increasing awareness that systematic post cardiac arrest intensive care can improve the likelihood of patient survival with good quality of life.<sup>2</sup> American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care emphasized that early prognostication of neurological outcome is an essential component of post-cardiac arrest care.<sup>17,23–25</sup> Our data suggest that rSO<sub>2</sub> may help predict good neurological outcome at 90 days in OHCA patients, which suggest rSO<sub>2</sub> monitoring was promising tool to select good candidates for post cardiac arrest intensive care containing therapeutic mild hypothermia, percutaneous coronary intervention, and extracorporeal cardiopulmonary resuscitation. On the other hands, in our data we found three patients with good neurological outcome at 90 days and low rSO<sub>2</sub> (<15%). So even though this large cohort showed that low rSO<sub>2</sub> (<15%) at hospital arrival indicate the high rate of poor neurological outcome at 90 days, we concluded from this study that this index could not be used alone for termination of resuscitative efforts because of the possibility of false prediction.

#### 4.2. Limitations

This study has several limitations. First, although it would be desirable to continuously monitor rSO<sub>2</sub> prior to hospital arrival, the absence of a portable NIRS device made this impossible. Nevertheless, we consider that such monitoring of rSO<sub>2</sub> with a portable device to be available in the future would allow the real-time assessment of the effects of prehospital cardiovascular care on cerebral perfusion. Second, rSO<sub>2</sub> was monitored for a limited time after hospital arrival. We recognized that time-course changes in rSO<sub>2</sub> are an important factor for the prognostication of neurological outcomes. Therefore, we are prospectively collecting the rSO<sub>2</sub> data during postresuscitation care after hospital arrival. Of special



**Fig. 4.** Three regional cerebral oxygen saturation groups of patients with good neurological outcome at 90 days after out-of-hospital cardiac arrest. The chi-square test was used to examine differences between categorical variables rSO<sub>2</sub>, regional cerebral oxygen saturation; OHCA, out-of-hospital cardiac arrest.

note was the fact that good neurological outcome was considerably more likely in patients who had rSO<sub>2</sub> >42% at hospital arrival, although the duration of monitoring at hospital arrival was short ( $\geq 1$  min). Third, rSO<sub>2</sub> measured by NIRS reflects only cerebral perfusion at the superficial layers in the limited frontal lobe areas<sup>8,9</sup> and rSO<sub>2</sub> is not a reliable marker of brain tissue oxygen partial pressure. However, rSO<sub>2</sub> measured by NIRS is closely related to oxygen saturation in the jugular bulb that represents venous oxygenation of the whole brain.<sup>26</sup> Fourth, compared with previous studies, we found lower rate of patients whose initially documented rhythms on the scene of cardiac arrest were VT/VF and very poor clinical outcomes at 90 days, especially in patients with cardiac arrest at hospital arrival, since EMS providers are not permitted to terminate CPR in the field in Japan, most patients with OHCA who are treated by EMS personnel are transported to emergency hospitals. It is conceivable that the practice of EMS system could be modified using tools such as, rSO<sub>2</sub>, in combination with existing termination of resuscitation rules.<sup>27</sup> Therefore, these results may not be generalizable to other countries with different EMS practices. Fifth, the small number of outcome events limits the ability to perform multivariable modeling. Sixth, as with any observational study, residual confounding could still account for at least part of the association. Seventh, we could not blind investigators because the rSO<sub>2</sub> monitoring requires real-time visual confirmation during CCR efforts. In pre-specified protocol all patients received the best available therapy including post cardiac arrest interventions regardless of the rSO<sub>2</sub> readings, however we could not eliminate the possibility that low rSO<sub>2</sub> might have influence the decision of stopping the resuscitation.

## 5. Conclusions

This prospective multicenter cohort study demonstrated that the rSO<sub>2</sub> at hospital arrival can predict good neurological outcome at 90 days after OHCA. Future studies should corroborate if rSO<sub>2</sub> can serve as a real-time indicator of cerebral perfusion during CCR and postresuscitation care for their improvement and for improvement of neurological outcome.

## Funding

This work was supported in part by the Grant-in-Aid for Scientific Research (B) (No. 24390400) from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) of Japan. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## Conflict of interest statement

All authors had no conflict of interest.

## Acknowledgments

We are greatly indebted to all of the J-POP Registry investigators. The authors thank Satoshi Sakima, MD, Yasuji Doi, MD, PhD, and Tetsuhisa Kitamura, MD, MSc, DrPh, for their valuable suggestions and critical review of the manuscript.

## Appendix A. Additional members of the J-POP Registry

T. Suzuki, N. Sato, T. Kimura, and K. Koike (Kyoto University Graduate School of Medicine [Kyoto, Japan]); T. Morooka, H. Rinka, and T. Ikebara (Osaka City General Hospital [Osaka, Japan]); M. Suzuki, A. Shirishita-Takeshita, and S. Hori (Keio University School of Medicine [Tokyo, Japan]); T. Unoki, S. Beppu, and I. Kaneko

(National Hospital Organization Kyoto Medical Center [Kyoto, Japan]); Y. Toyoda, and M. Kitano (Saiseikai Yokohamashi Tobu Hospital [Yokohama, Japan]); M. Machida, and H. Ishikura (Fukuoka University Hospital [Fukuoka, Japan]); T. Oomura, D. Kudo, and S. Kushimoto (Tohoku University Hospital [Sendai, Japan]); T. Watanabe, M. Fujioka and T. Seki (Nara Medical University Hospital [Kashihara, Japan]); M. Otsuka, H. Yano, K. Arakawa, M. Nitta, O. Akasaka, S. Ryu, and H. Himeno (Fujisawa City Hospital [Fujisawa, Japan]); T. Hatada and H. Imai (Mie University Hospital [Tsu, Japan]); S. Nachi, H. Ushikoshi, and S. Ogura (Gifu University Hospital [Gifu, Japan]); M. Mizobuchi, T. Kobayashi, K. Shibata, and S. Nakamura (Kyoto Katsura Hospital [Kyoto, Japan]); H. Kamura, and A. Kataoka (Japanese Red Cross Musashino Hospital [Musashino, Japan]); Y. Honma, Y. Nishi, K. Niwa, T. Watanabe, T. Inohara, T. Takabayashi, and S. Ishimatsu (St Luke's International Hospital [Tokyo, Japan]); J. Kotani and A. Hashimoto (Hyogo Medical University [Nishinomiya, Japan]); S. Marukawa (Iseikai Hospital [Osaka, Japan]); K. Shiga, S. Asai, T. Hathaway (Seirei Mikatahara General Hospital [Hamamatsu, Japan]); K. Kikuchi, M. Tokura, S. Nishino (Dokkyo Medical University Hospital [Tochigi, Japan]).

## References

- Kitamura T, Iwami T, Kawamura T, Nagao K, Tanaka H, Hiraide A. Nationwide public-access defibrillation in Japan. *N Engl J Med* 2010;362:994–1004.
- Nagao K, Hayashi N, Kanmatsuse K, et al. Cardiopulmonary cerebral resuscitation using emergency cardiopulmonary bypass, coronary reperfusion therapy and mild hypothermia in patients with cardiac arrest outside the hospital. *J Am Coll Cardiol* 2000;36:776–83.
- Nichol G, Aufderheide TP, Eigel B, et al. Regional systems of care for out-of-hospital cardiac arrest: a policy statement from the American Heart Association. *Circulation* 2010;121:709–29.
- Field JM, Hazinski MF, Sayre MR, et al. Part 1: Executive summary: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2010;122:S640–56.
- Neumar RW, Nolan JP, Adrie C, et al. Post-cardiac arrest syndrome: epidemiology, pathophysiology, treatment, and prognostication. A consensus statement from the International Liaison Committee on Resuscitation (American Heart Association, Australian and New Zealand Council on Resuscitation, European Resuscitation Council, Heart and Stroke Foundation of Canada, InterAmerican Heart Foundation, Resuscitation Council of Asia, and the Resuscitation Council of Southern Africa); the American Heart Association Emergency Cardiovascular Care Committee; the Council on Cardiovascular Surgery and Anesthesia; the Council on Cardiopulmonary, Perioperative, and Critical Care; the Council on Clinical Cardiology; and the Stroke Council. *Circulation* 2008;118:2452–83.
- Booth CM, Boone RH, Tomlinson G, Detsky AS. Is this patient dead, vegetative, or severely neurologically impaired? Assessing outcome for comatose survivors of cardiac arrest. *JAMA* 2004;291:870–9.
- Wijdicks EF, Hijdra A, Young GB, Bassetti CL, Wiebe S. Practice parameter: prediction of outcome in comatose survivors after cardiopulmonary resuscitation (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2006;67:203–10.
- Yao FS, Tseng CC, Ho CY, Levin SK, Illner P. Cerebral oxygen desaturation is associated with early postoperative neuropsychological dysfunction in patients undergoing cardiac surgery. *J Cardiothorac Vasc Anesth* 2004;18:552–8.
- Murkin JM, Adams SJ, Novick RJ, et al. Monitoring brain oxygen saturation during coronary bypass surgery: a randomized, prospective study. *Anesth Analg* 2007;104:51–8.
- Becker LB, Aufderheide TP, Geocadin RG, et al. Primary outcomes for resuscitation science studies: a consensus statement from the American Heart Association. *Circulation* 2011;124:2158–77.
- The Ministry of Health Law, Ministry of Education, Culture, Sports, Science and Technology. Ethical guidelines for epidemiologic research; 2004.
- Berg RA, Hemphill R, Abella BS, et al. Part 5: Adult basic life support: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2010;122:S685–705.
- Morrison LJ, Deakin CD, Morley PT, et al. Part 8: Advanced life support: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science With Treatment Recommendations. *Circulation* 2010;122:S345–421.
- Jacobs I, Nadkarni V, Bahr J, et al. Cardiac arrest and cardiopulmonary resuscitation outcome reports: update and simplification of the Utstein templates for resuscitation registries: a statement for healthcare professionals from a task force of the International Liaison Committee on Resuscitation (American Heart Association, European Resuscitation Council, Australian Resuscitation Council, New Zealand Resuscitation Council, Heart and Stroke Foundation of Canada, InterAmerican Heart Foundation, Resuscitation Councils of Southern Africa). *Circulation* 2004;110:3385–97.

15. Bobrow BJ, Spaite DW, Berg RA, et al. Chest compression-only CPR by lay rescuers and survival from out-of-hospital cardiac arrest. *JAMA* 2010;304:1447–54.
16. Nolan JP, Morley PT, Vanden Hoek TL, et al. Therapeutic hypothermia after cardiac arrest: an advisory statement by the advanced life support task force of the International Liaison Committee on Resuscitation. *Circulation* 2003;108:118–21.
17. Peberdy MA, Callaway CW, Neumar RW, et al. Part 9: Post-cardiac arrest care: 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation* 2010;122:S768–86.
18. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more correlated receiver operating characteristic curves: a nonparametric approach. *Biometrics* 1988;44:837–45.
19. Takasu A, Sakamoto T, Okada Y. Arterial base excess after CPR: the relationship to CPR duration and the characteristics related to outcome. *Resuscitation* 2007;73:394–9.
20. Aduen J, Bernstein WK, Khastgir T, et al. The use and clinical importance of a substrate-specific electrode for rapid determination of blood lactate concentrations. *JAMA* 1994;272:1678–85.
21. Howanitz PJ, Cembrowski GS, Steindel SJ, Long TA. Physician goals and laboratory test turnaround times. A College of American Pathologists Q-Probes study of 2763 clinicians and 722 institutions. *Arch Pathol Lab Med* 1993;117:22–8.
22. Fermann GJ, Suyama J. Point of care testing in the emergency department. *J Emerg Med* 2002;22:393–404.
23. The Hypothermia after Cardiac Arrest Study Group. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *N Engl J Med* 2002;346:549–56.
24. Bernard SA, Gray TW, Buist MD, et al. Treatment of comatose survivors of out-of-hospital cardiac arrest with induced hypothermia. *N Engl J Med* 2002;346:557–63.
25. Dumas F, White L, Stubbs BA, Cariou A, Rea TD. Long-term prognosis following resuscitation from out of hospital cardiac arrest: role of percutaneous coronary intervention and therapeutic hypothermia. *J Am Coll Cardiol* 2012;60:21–7.
26. Daubney PE, Pilkington SN, Janke E, Charlton GA, Smith DC, Webber SA. Cerebral oxygenation measured by near-infrared spectroscopy: comparison with jugular bulb oximetry. *Ann Thorac Surg* 1996;61:930–4.
27. Sasson C, Hegg AJ, Macy M, Park A, Kellermann A, McNally B. Prehospital termination of resuscitation in cases of refractory out-of-hospital cardiac arrest. *JAMA* 2008;300:1432–8.