

Title	Dose-rate effect was observed in T98G glioma cells following BNCT.
Author(s)	Kinashi, Yuko; Okumura, Kakuji; Kubota, Yoshihisa; Kitajima, Erika; Okayasu, Ryuichi; Ono, Koji; Takahashi, Sentaro
Citation	Applied radiation and isotopes : including data, instrumentation and methods for use in agriculture, industry and medicine (2014), 88: 81-85
Issue Date	2014-06
URL	http://hdl.handle.net/2433/188913
Right	© 2013 Elsevier Ltd.
Type	Journal Article
Textversion	author

1
2
3 **Dose-rate effect was observed in T98G glioma cells following BNCT**
4
5
6
7
8

9
10 Yuko KINASHI^{1)*}, Kakuji OKUMURA^{1)*}, Yoshihisa KUBOTA²⁾, Erika KITAJIMA³⁾, Ryuichi
11
12 OKAYASU²⁾, Koji ONO¹⁾ and Sentaro TAKAHASHI¹⁾
13
14
15
16
17
18

19 1) Kyoto University Research Reactor Institute,
20
21

22 2) National Institute of Radiological Sciences,
23
24

25 3) Osaka High-technology College
26
27
28
29
30

31
32 Reprint requests to Yuko Kinashi, M.D., Ph.D.
33
34

35 Research Reactor Institute, Kyoto University,
36
37

38 Kumatori-cho, Sennann-gun, Osaka 590-0494, Japan
39
40

41 Tel.:81-724-51-2437
42
43

44 Fax.: 81-724-51-2623
45
46

47 E-mail: kinashi@rri.kyoto-u.ac.jp
48
49
50

51
52 * Authors with equal contributions
53
54

55 Keywords: dose-rate effect, BNCT, DNA-DSBs, glioma
56
57
58
59
60

1
2
3 ABSTRACT
4
5

6 Background: It is generally said that low LET radiation produce high dose-rate effect, on the
7
8
9 other hand, no significant dose rate effect is observed in high LET radiation. Although high LET
10
11
12 radiations are produced in BNCT, little is known about dose-rate effect of BNCT.
13
14

15
16 Materials and Methods: T98G cells, which were tumor cells, were irradiated by neutron mixed
17
18
19 beam with BPA. As normal tissue derived cells, Chinese hamster ovary (CHO-K1) cells and
20
21
22 DNA double strand breaks (DNA-DSBs) repair deficient cells, xrs5 cells were irradiated by the
23
24
25 neutrons (not including BPA). To DNA-DSBs analysis, T98G cells were stained
26
27
28 immunochemically with 53BP1 antibody. The number of DNA-DSBs was determined by
29
30
31 counting 53BP1 foci.
32
33

34
35 Results: There was no dose-rate effect in xrs5 cells. D_0 difference between 4cGy/min and
36
37
38 20cGy/min irradiation were 0.5 and 5.9 at the neutron and gamma-ray irradiation for CHO-K1,
39
40
41 and 0.3 at the neutron for T98G cells. D_0 difference between 20cGy/min and 80cGy/min
42
43
44 irradiation for T98G cells were 1.2 and 0.6 at neutron irradiation plus BPA and gamma-ray. The
45
46
47 differences between neutron irradiations at the dose rate in T98G cells were supported by not
48
49
50 only the cell viability but also 53BP1 foci assay at 24 hours following irradiation to monitor
51
52
53 DNA-DSBs.
54
55

56
57 Conclusion: Dose-rate effect of BNCT when T98G cells include 20ppm BPA was greater than
58
59
60

1
2
3 that of gamma-ray irradiation. Moreover, Dose-rate effect of the neutron beam when CHO-K1
4
5
6 cells did not include BPA was less than that of gamma-ray irradiation These present results may
7
8
9 suggest the importance of dose-rate effect for more efficient BNCT and the side effect
10
11
12
13 reduction.

14 15 16 17 18 19 INTRODUCTION

20
21
22 Boron neutron capture therapy (BNCT) is a treatment that can selectively destroy
23
24
25 tumor tissue by $^{10}\text{B} (n,\alpha) ^7\text{Li}$ reaction without significant damage to normal tissue. In principle,
26
27
28 ^{10}B compounds are selectively delivered to tumor cells and exposed to low-energy thermal
29
30
31 neutrons. The reaction with thermal neutron produces an alpha particle and ^7Li nucleus that have
32
33
34 high linear energy transfer (LET) values and enriched biological effect. Although it depends on
35
36
37 the kinds of ^{10}B compounds, DNA damage or decrease in cell viability produced by BNCT is
38
39
40 greater than neutron alone or gamma-ray (Kinashi et al., 2011, Dagrosa et al., 2011).
41
42
43

44
45 It is generally accepted that for cell viability, low LET radiation such as X-ray and
46
47
48 gamma-ray produce high dose-rate effect, on the other hand, no significant dose rate effect is
49
50
51 observed in high LET radiation such as neutrons, alpha particle, and heavy ion (Metting et al.,
52
53
54 1985, Ainsworth et al., 1976, Heiber et al., 1987, Yang et al., 1986, Furusawa et al., 2006,
55
56
57 Fairchild et al., 1985). However, there is no data about dose-rate effect for BNCT. It is also
58
59
60

1
2
3 difficult to estimate dose-rate effect of BNCT from the past studies because the neutron beam
4
5
6 used for BNCT is the mixed beam that contains thermal, epithermal and fast neutrons, and
7
8
9 gamma-ray. A better understanding of dose-rate effect for BNCT will help to improve the
10
11
12 efficiency of this therapy as well as to reduce the side effect in normal tissue.
13
14

15
16 DNA double strand breaks (DNA-DSBs) are potentially lethal lesions and important
17
18 biological effect of ionizing radiation. DNA-DSBs can be repaired by DNA repair proteins, such
19
20
21 as gamma-H2AX, DNA-PKcs, and 53BP1. Therefore, immunochemically stained foci for the
22
23
24 proteins have been used for a sensitive and efficient marker for DNA-DSBs. Several studies
25
26
27 using this method showed that the number of foci may reflect not only the dose but also the
28
29
30 types of radiation. Okayasu et al. reported that the disappearance of phosphorylated DNA-PKcs
31
32
33 foci in 180BR cells irradiated heavy ions were slow (Okayasu et al., 2006).
34
35
36

37
38 We investigated dose-rate effect of the neutron mixed radiation beam used for BNCT
39
40
41 in Kyoto University Research Reactor (KUR) at cell survival fraction and DNA-DSBs. We
42
43
44 report here that dose-rate effect of BNCT is observed in T98G cells that are human tumor cell
45
46
47 line at above-mentioned both biological points.
48
49

50 51 52 53 MATERIALS AND METHODS

54 55 56 57 *Cell culture*

1
2
3 The Chinese hamster ovary-derived, wild-type CHO-K1 cells and the human
4
5
6 glioblastoma cell lines, T98G were purchased from Riken BRC Cell Bank. The radiosensitive
7
8
9 xrs5 mutant cells (Ku80-deficient CHO cell line) kindly supplied by Prof. P Jeggo (Jeggo et al.,
10
11
12 1983, Kemp et al., 1984). CHO-K1 and xrs5 cells were cultured in Memo medium (Invitrogen)
13
14
15 supplemented with 10% heat-inactivated FBS (Biowest). T98G cells were grown in RPMI 1640
16
17
18 (Invitrogen) supplemented 10% heat-inactivated FBS (Biowest). These cells were maintained at
19
20
21
22 37°C in a humidified atmosphere with 5.0% CO₂.

23 24 25 *¹⁰B compound and Irradiation experiments*

26
27
28 For irradiation, the cells were trypsinized, suspended in the above mentioned medium,
29
30
31 and then aliquoted to each teflon tube. A stock solution of ¹⁰B-para-boronphenylalanine (BPA)
32
33
34 was used for BNCT experiment to T98G cells. T98G cell suspensions were incubated with BPA
35
36
37 at 20ppm concentration 1 hour before neutron irradiation. The Heavy Water Facility of the
38
39
40 Kyoto University Research Reactor (KUR) was used for neutron mixed beam irradiation
41
42
43 (approximately 4cGy/min and 20cGy/min). Neutron fluencies were measured by radioactivation
44
45
46 of gold foil and gamma-ray doses by TLD. The physical dose percentages to total dose of
47
48
49 thermal (<0.5eV), epithermal (0.5eV-10keV), fast (>10keV) neutrons, and gamma-ray were
50
51
52 approximately 30%, 3%, 22% and 45% respectively, when the reactor was operated at
53
54
55 4cGy/min or 20cGy/min (Fig. 1). Gamma-ray irradiation was carried out with a ⁶⁰Co
56
57
58
59
60
61
62
63
64
65

1
2
3 gamma-ray irradiator at dose rate of about 4cGy/min, 20cGy/min, 80cGy/min and 2Gy/min.
4
5

6 *Cell survival assay*
7

8
9 Survival fraction of CHO-K1, xrs5 and T98G cells were determined by conventional
10 colony formation assay. After irradiation, the cell suspension solutions were properly diluted,
11
12 and the known number of cells were seeded in cell culture dishes and incubated for 7-9 days
13
14 (CHO-K1 and xrs5) and 2 weeks (T98G). The cells were fixed with 70% ethanol and stained
15
16 with crystal violet solution (Wako) for counting colonies.
17
18
19
20
21
22
23
24

25 *Immunofluorescent staining*
26

27
28 Irradiated T98G cells were seeded onto 22 mm × 22 mm cover slips in 6-well micro
29 plates, incubated for 30min, 1, 3, 6 and 24 hours, and washed with cold PBS (Invitrogen). The
30
31 cells were fixed with 3.6% formalin/PBS for 15 min, permeabilized with 0.5%
32
33 TritonX-100/PBS on ice for 15 min and washed thoroughly with PBS. The cover slips were
34
35 incubated for 2 hours at 37°C with primary antibody, rabbit monoclonal anti-53BP1 antibody
36
37 (Bethyl Laboratories) in TBS-DT solution (20 mM Tris-HCl, 137 mM NaCl, containing 0.1%
38
39 Tween-20, 5% skim milk). After being washed three times with cold PBS, the cells were
40
41 incubated for 1 hour at 37°C with secondary antibody, Alexa Fluor 594 goat anti-rabbit IgG
42
43 (Invitrogen) in TBS-DT solution. After washing three times with PBS, The cover slips were
44
45 mounted on slide glasses with DAPI (4',6-diamidino-2-phenylindole) in 10% glycerol/PBS to
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 counter-stain the nuclei.
4
5

6 *Image acquisition and Analysis*
7
8

9 Images were acquired with a fluorescence microscope (KEYENCE, BZ-9000). ImageJ
10 (National Institutes of Health) were used for image processing and automated foci counting.
11
12
13
14

15 Some foci images were scored by eye.
16
17
18
19
20

21
22 **RESULTS**
23
24

25 *Dose-rate effect was observed in CHO-K1 cells at cell survival assay*
26
27

28 Fig. 2 shows cell survival fraction of CHO-K1 (referred as K1 in Fig. 2) and xrs5 cells
29 irradiated with the neutron mixed beam. A scatter plot is used because the information of the
30 dose rate was not available in real time so that the total doses were not always same in the
31 mixed beam irradiation. Each point shows the survival fraction, and black and dot lines are used
32 to fit log-linear model with Microsoft Excel software. As expected, xrs5 cells were more
33 sensitive to the mixed beam than CHO-K1 cells. There was no significant difference for the cell
34 survival at xrs5 cells between 4cGy/min and 20cGy/min irradiation (Fig. 2 and Table 1). On the
35 other hand, dose-rate effect between 4cGy/min and 20cGy/min was observed in CHO-K1 cells
36 irradiated with especially gamma-ray (Fig. 2B). The D_0 differences in CHO-K1 cells were 0.5
37 and 5.9 for neutron and gamma-ray irradiation (Table 1). The surviving fraction for CHO-K1
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 cells after 20cGy/min gamma-ray irradiation was significantly lower than that after 4cGy/min
4
5
6 irradiation.
7

8
9
10 *Dose-rate effect of BNCT was greater than that of gamma-ray at T98G cell survival assay*

11
12 Fig. 3 shows cell survival fraction of T98G cells irradiated with the neutron mixed
13
14 beam alone (referred as no BPA in Fig. 3A) or the mix beam plus 20ppm BPA (referred as plus
15
16 BPA in Fig. 3B). When T98G cells did not include BPA, D_0 between 4cGy/min and 20cGy/min
17
18 irradiation did not show a big difference (D_0 difference was 0.3). 20cGy/min irradiation plus
19
20 BPA produced a decrease in cell survival fraction, compared to 4cGy/min plus BPA (Fig. 3B)
21
22 and the D_0 difference was 1.2 (Table 2). We also researched dose-rate effect of gamma-ray in
23
24 T98G cells because it was thought that gamma-ray in the neutron beam significantly contribute
25
26 to dose-rate effect. Assuming that T98G cells include 20ppm BPA, the dose-rate of 4cGy/min
27
28 and 20cGy/min neutron irradiation is converted to 20cGy/min and 80cGy/min, due to ^{10}B (n,α)
29
30 ^7Li reaction. In Fig. 4, the difference of D_0 between 20cGy/min and 80cGy/min in plus BPA
31
32 group was larger than that in gamma-ray group.
33
34
35
36
37
38
39
40
41
42
43
44
45
46

47
48 *Induction of 53BP1 foci in T98G cells*

49
50
51 Dose-rate effect was greater in T98G cells incubated with BPA (Table 2 and Fig. 4). So,
52
53 we researched DNA-DSBs, because it is said that high LET radiation induces more DNA-DSBs
54
55 than low LET radiation. The results of 53BP1 foci in T98G cells 30min after neutron irradiation
56
57
58
59
60

1
2
3 are shown in Fig. 5. Although induction of 53BP1 foci increased with dose, the number of
4
5
6 53BP1 foci following 20cGy/min irradiation seemed to be larger than 4cGy/min irradiation,
7
8
9 regardless of existence of BPA.
10

11 12 *Loss of 53BP1 foci in T98G cells*

13
14
15
16 Fig. 6 shows time dependence of 53BP1 foci after irradiation. The number of 53BP1
17
18
19 foci in not BPA including cells rapidly decrease to about zero, compared to the cells incubated
20
21
22 BPA. When T98G cells include BPA, a lot of 53BP1 foci had remained even at 24 hours after
23
24
25 neutron irradiation. Moreover, the difference between the cells plus BPA irradiated with
26
27
28 4cGy/min and 20cGy/min was significant at 24hours post irradiation.
29
30
31
32
33
34

35 DISCUSSION

36
37
38 In this report, we showed that dose-rate effect was observed at cell survival rate in
39
40
41 T98G cells following BNCT, and the effect was greater even if it was compared with
42
43
44 gamma-ray irradiation at the same dose-rate (Fig. 4). Little dose-rate effect of gamma-ray or the
45
46
47 neutron mixed beam (not irradiated with BPA) was observed in xrs5 cells. On the other hand,
48
49
50 there was the dose-rate effect in CHO-K1 cells. This means that dose-rate effect is not observed
51
52
53 in DNA damage repair deficient cells, such as xrs5 cells. DNA-DSBs were repaired, when
54
55
56 T98G cells were irradiated with the neutron beams or X-ray (Fig. 6) (Short et al., 2007).
57
58
59
60
61
62
63
64
65

1
2
3 Therefore for T98G cells, the dose-rate effect would be observed.
4
5

6 We treated CHO-K1 and xrs5 cells derived from normal tissues, and T98G cells that
7
8 are tumor cells to investigate the dose-rate effect of BNCT. In BNCT, highly ^{10}B distributed
9
10 tumor cells are exposed to thermal neutrons. The alpha particles and ^7Li nuclei produced by ^{10}B
11
12 $(n,\alpha)^7\text{Li}$ reaction have short paths ($\sim 10\mu\text{m}$), same as a cell size (Liu et al., 2009). Therefore, it is
13
14 thought that only tumor cells are affected by BNCT and normal cells are irradiated with the
15
16 neutron mixed beam. These present results may suggest that dose-rate effect of the neutron
17
18 mixed beam for BNCT would contribute to both higher cytotoxicity reaction for tumor cells
19
20 through BNCT and relative decline of biological effect to normal cells.
21
22
23
24
25
26
27
28
29
30

31 The kinetics of the number of 53BP1 foci for DNA-DSBs detection was also further
32
33 evidence to the dose-rate effect of the neutron mixed beam and BNCT. High-LET radiation
34
35 induces DNA-DSBs that are not repaired or more difficult to be repaired compared with
36
37 low-LET radiation (Okayasu et al., 2006, Hill et al., 1999, Terato et al., 2008). Although there
38
39 did not seem to be the significant difference about induction of DNA-DSBs, we showed that the
40
41 DNA-DSBs induced by the neutron mixed beam alone was fast or easy to be repaired compared
42
43 to the mixed beam plus BPA. Moreover, we thought that the significant difference between
44
45 4cGy/min and 20cGy/min about the number of 53BP1 foci 24 hours after BNCT irradiation
46
47
48
49
50
51 might reflect the difference of the survival rate produced by dose-rate effect in the case of
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3 including BPA (Fig.3B and Fig. 6). We demonstrated the relationship between DNA-DSBs and
4
5
6 the lethal potential, provided that kinetics of 53BP1 foci correlated with DNA-DSBs repair.
7
8

9
10 There is the hypothesis that residual foci are unable to pass mitosis (Marková et al.,
11
12 2007). Although this means that few foci are observed in the cell exceeding the duration of the
13
14 cell cycle after irradiation, recent studies show that the residual 53BP1/gamma-H2AX foci may
15
16 be available for biological dosimetry. Primary induced foci and residual foci may be different
17
18 types of foci (Marková et al., 2011). It is reported that the doubling time of T98G cells are 22-26
19
20 hours (Choi et al., 1997, Nakada et al., 2005). We also calculated the doubling time to be
21
22 approximately 24 hours from their growth curve analysis. So, we concluded that the difference
23
24 of 53BP1 foci between 4cGy/min and 20cGy/min at 24 hours after BNCT irradiation was the
25
26 variance of the degree of DNA-DSBs repair.
27
28
29
30
31
32
33
34
35
36
37

38 In this study, using cell survival assay, we found that greater dose-rate effect were
39
40 observed in T98G cells underwent BNCT than gamma-ray, and that the effect were less in
41
42 CHO-K1 cells irradiated with the neutron beam alone, compared with gamma-ray. Time
43
44 dependence of 53BP1 foci was not direct evidence of dose-rate effect of BNCT because the
45
46 converted absorbed dose was not same between 4cGy/min and 20c/min irradiation. However,
47
48 the difference of 53BP1 foci reflected the difference of the survival rate results. Although we
49
50
51 have thought that dose-rate effect of BNCT was not that of alpha particles or ⁷Li nuclei
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

produced by $^{10}\text{B} (n,\alpha) ^7\text{Li}$ reaction, we are interested in this result.

1
2
3 REFERENCES
4
5
6

7 Ainsworth, E. J., Jordan, D. L., Miller, M., Cooke, E. M., Hulesch, J. S., 1976. Dose rate studies
8 with fission spectrum neutrons. *Radiat. Res.* 67, 30-45.
9

10
11
12
13
14
15 Choi, B. O., Yamaki, T., Tatewaki, K., Ibayashi, Y., Hashi, K., 1997. Deletion of complex
16
17
18 gangliosides of human glioma cells during mitotic cell division. *J. Neurooncol.* 34, 211-219.
19

20
21
22
23 Dagrosa, M. A., Crivello, M., Perona, M., Thorp, S., Santa Cruz, G. A., Pozzi, E., Casal, M.,
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Thomasz, L., Cabrini, R., Kahl, S., Juvenal, G. J., Pisarev, M. A., 2011. First evaluation of the
biologic effectiveness factors of boron neutron capture therapy (BNCT) in a human colon
carcinoma cell line. *Int. J. Radiat. Oncol. Biol. Phys.* 79, 262-268.

Fairchild, R. G., Bond, V. P., 1985. Current status of ¹⁰B-neutron capture therapy: enhancement
of tumor dose via beam filtration and dose rate, and the effects of these parameters on minimum
boron content: a theoretical evaluation. *Int. J. Radiat. Oncol. Biol. Phys.* 11, 831-840.

Furusawa, Y., Hirayama, R., Matsumoto, Y., Satoh, Y., Satoh, S., Tomitani, T., Kanazawa, M.,
2006. Dose-rate Effect of Heavy Ion on Cell Killing (In Japanese). *Natl. Inst. Radiol. Sci.* 192,
79-80.

1
2
3 Hill, M. A., 1999. Radiation damage to DNA: The importance of track structure. *Radiat. Meas.*
4
5
6 31, 15-23.
7

8
9
10 Hieber, L., Ponsel, G., Roos, H., Fenn, S., Fromke, E., Kellerer, A. M., 1987. Absence of a
11
12 dose-rate effect in the transformation of C3H 10T1/2 cells by alpha-particles. *Int. J. Radiat. Biol.*
13
14
15
16
17 *Relat. Stud. Phys. Chem. Med.* 52, 859-869.
18

19
20
21 Jeggo, P. A., Kemp, L. M., 1983. X-ray-sensitive mutants of Chinese hamster ovary cell line.
22
23
24 Isolation and cross-sensitivity to other DNA-damaging agents. *Mutat. Res.* 112, 313-327.
25

26
27
28
29 Kemp, L. M., Sedgwick, S. G., Jeggo, P. A., 1984. X-ray sensitive mutants of Chinese hamster
30
31
32 ovary cells defective in double-strand break rejoining. *Mutat. Res.* 132, 189-196.
33
34

35
36
37 Kinashi, Y., Takahashi, S., Kashino, G., Okayasu, R., Masunaga, S., Suzuki, M., Ono, K., 2011.
38
39
40 DNA double-strand break induction in Ku80-deficient CHO cells following boron neutron
41
42
43 capture reaction. *Radiat. Oncol.* 6, 106.
44

45
46
47 Liu, Y., Nagata, K., Masunaga, S.-ichiro, Suzuki, M., Kashino, G., Kinashi, Y., Tanaka, H.,
48
49
50 Sakurai, Y., Maruhashi, A., Ono, K., 2009. γ -ray irradiation enhanced boron-10 compound
51
52
53 accumulation in murine tumors. *J. Radiat. Res.* 50, 553-557.
54
55

1
2
3 Metting, N. F., Braby, L. A, Roesch, W. C., Nelson, J. M., 1985. Dose-rate evidence for two
4
5
6 kinds of radiation damage in stationary-phase mammalian cells. *Radiat. Res.* 103, 204-218.
7
8

9
10
11 Marková, E., Schultz, N., Belyaev, I. Y., 2007. Kinetics and dose-response of residual
12
13
14 53BP1/gamma-H2AX foci: co-localization, relationship with DSB repair and clonogenic
15
16
17 survival. *Int. J. Radiat. Biol.* 83, 319-329.
18
19

20
21 Marková, E., Torudd, J., Belyaev, I., 2011. Long time persistence of residual 53BP1/ γ -H2AX
22
23
24 foci in human lymphocytes in relationship to apoptosis, chromatin condensation and biological
25
26
27 dosimetry. *Int. J. Radiat. Biol.* 87, 736-745.
28
29

30
31
32 Nakada, M., Niska, J. A, Tran, N. L., McDonough, W. S., Berens, M. E., 2005. EphB2/R-Ras
33
34
35 signaling regulates glioma cell adhesion, growth, and invasion. *Am. J. Pathol.* 167, 565-576.
36
37

38
39
40 Okayasu, R., Okada, M., Okabe, A., Noguchi, M., Takakura, K., Takahashi, S., 2006. Repair of
41
42
43 DNA damage induced by accelerated heavy ions in mammalian cells proficient and deficient in
44
45
46 the non-homologous end-joining pathway. *Radiat. Res.* 165, 59-67.
47
48

49
50
51 Short, S. C., Martindale, C., Bourne, S., Brand, G., Woodcock, M., Johnston, P., 2007. DNA
52
53
54 repair after irradiation in glioma cells and normal human astrocytes. *Neuro. Oncol.* 9, 404-411.
55
56

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Terato, H., Tanaka, R., Nakaarai, Y., Nohara, T., Doi, Y., Iwai, S., Hirayama, R., Furusawa, Y.,

Ide, H., 2008. Quantitative analysis of isolated and clustered DNA damage induced by gamma-rays, carbon ion beams, and iron ion beams. *J. Radiat. Res.* 49, 133-146.

Yang, T. H., Craise, L. M., Mei M. T., and Tobias, C. A., 1986. Dose protraction studies with low- and high-LET radiations on neoplastic cell transformation in vitro. *Adv. Space. Res.* 6, 137-147.

1
2
3 **FIGURE LEGENDS**
4
5

6 Fig. 1. The physical doses percentage of thermal, epithermal, fast neutrons, and gamma-ray in
7
8
9 the neutron mixed beams, when samples were irradiated at KUR operated at 4cGy/min
10
11
12 or 20cGy/min.
13
14
15

16 Fig. 2. Dose-rate effect on survival fraction for CHO-K1 and xrs5 cells irradiated with neutron
17
18
19 mixed beam (n) (A) or gamma-ray (γ) (B).
20
21

22 Fig. 3 Dose-rate effect at survival fraction for T98G cells irradiated with neutron mixed beam in
23
24
25 case of not including BPA (A) or including BPA (B). In Fig. 3B, the horizontal axis is
26
27
28 the absorbed dose converted assuming that the concentration of BPA in T98G cells is
29
30
31 20ppm.
32
33
34

35 Fig. 4 Dose-rate dependence of D_0 for T98G cells. The dose rate of plus BPA + neutron
36
37
38 irradiation group is converted assuming that the concentration of BPA in T98G cells is
39
40
41 20ppm.
42
43
44

45 Fig. 5. The number of 53BP1 foci induced in T98G cells at 30 min after the neutron mixed
46
47
48 beam irradiation in case of not including BPA (A) or including BPA (B). The data are
49
50
51 compiled from two independent experiments. The error bars indicate the standard errors
52
53
54 of the mean.
55
56
57
58
59
60
61
62
63
64
65

1
2
3 Fig. 6. Time dependent loss of 53BP1 foci in T98G cells following approximately 50min
4
5
6 neutron mixed beam irradiation at 4cGy/min and 10min at 20cGy/min. Note that the
7
8 absorbed dose converted assuming that the concentration of BPA in T98G cells is
9
10 20ppm is different between 4cGy/min and 20cGy/min irradiation. The converted dose
11
12 was approximately 10Gy in 4cGy/min neutron irradiation plus BPA, and 8Gy in
13
14 20cGy/min plus BPA. The data are compiled from two independent experiments at each
15
16 point. The error bars indicate the standard errors of the mean.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. D_0 for CHO-K1 and xrs5 cells irradiated with neutron mixed beam or gamma-ray

D_0 (Gy)	CHO-K1	xrs5
Neutron mixed beam 4cGy/min	2.2	0.40
Neutron mixed beam 20cGy/min	1.7	0.35
Gamma-ray 4cGy/min	7.4	0.53
Gamma-ray 20cGy/min	1.5	0.55

Table 2. D_0 for T98G cells irradiated with neutron mixed beam in case of including BPA or not.

D_0 (Gy)	no BPA	plus BPA
Neutron mixed beam 4cGy/min	1.8	2.7
Neutron mixed beam 20cGy/min	1.5	1.5




Fig.1





Fig.2

Figure(s)





Fig.3




Fig.4

Figure(s)





Fig.5




Fig.6