

Title	ON FUNCTIONS STARLIKE IN ONE DIRECTION (Some inequalities concerned with the geometric function theory)
Author(s)	NUNOKAWA, MAMORU; SOKOL, JANUSZ
Citation	数理解析研究所講究録 (2014), 1878: 81-84
Issue Date	2014-04
URL	http://hdl.handle.net/2433/195600
Right	
Type	Departmental Bulletin Paper
Textversion	publisher

ON FUNCTIONS STARLIKE IN ONE DIRECTION

MAMORU NUNOKAWA AND JANUSZ SOKÓŁ

ABSTRACT. In this work we investigate the notion of starlikeness in one direction.

Let \mathcal{H} denote the class of all analytic functions in the unit disc $\mathbb{D} = \{z : |z| < 1\}$ on the complex plane \mathbb{C} . Recall that a set $E \subset \mathbb{C}$ is said to be starlike with respect to a point $w_0 \in E$ if and only if the linear segment joining w_0 to every other point $w \in E$ lies entirely in E , while a set E is said to be convex if and only if it is starlike with respect to each of its points, that is if and only if the linear segment joining any two points of E lies entirely in E .

Theorem 1. *Let $p(z) = \sum_{n=0}^{\infty} c_n z^n$ be analytic in \mathbb{D} . Assume also that $p(z) \neq 0$ in \mathbb{D} . Then there exists a number $\theta_0, 0 \leq \theta_0 \leq \pi$ for which*

$$(1) \quad \arg p(e^{i(\pi+\theta_0)}) = \arg p(e^{i\theta_0}).$$

Proof. From the hypothesis, we can define the function

$$(2) \quad \varphi(\theta) = \arg p(e^{i(\pi+\theta)}) - \arg p(e^{i\theta}),$$

where $0 \leq \theta \leq \pi$. Then we have the following three cases as

$$(i) \ \varphi(0) = 0, \quad (ii) \ \varphi(0) > 0, \quad (iii) \ \varphi(0) < 0.$$

For the case (i), it proves (1) and for the case (ii), we have

$$\begin{aligned} \varphi(\pi) &= \arg p(e^{2i\pi}) - \arg p(e^{i\pi}) \\ &= -(\arg p(e^{i\pi}) - \arg p(e^{i0})) \\ &< 0. \end{aligned}$$

Applying the intermediate value theorem, there exists a number $\theta_0, 0 < \theta_0 < \pi$ for which

$$\varphi(\theta_0) = 0$$

or

$$\arg p(e^{i(\pi+\theta_0)}) = \arg p(e^{i\theta_0}).$$

For the case (iii), applying the same method as the above, we can complete the proof of (1). □

Definition 1.

Let us call the diametal line of $p(z)$ with respect to $\theta = \theta_0$ if the straight line passes through 3 points $p(e^{i(\pi+\theta_0)})$, $w = 0$ and $p(e^{i\theta_0})$.

Definition 2.

In the Definition 1, if the diametal line of $p(z)$ with respect to $\theta = \theta_0$ has only 2 common points with the image curve of $|z| = 1$ under the mapping $w = p(z)$, then we call that $p(z)$ is starlike in the direction of diametal line with respect to $\theta = \theta_0$.

These ideas of being starlike in one direction was introduced by M. S. Robertson in [1].

Lemma 2. *Let $p(z) = \sum_{n=0}^{\infty} c_n z^n$ be analytic \mathbb{D} . Assume also that $p(z) \neq 0$ in \mathbb{D} . Then there exists at least one the diametal line of $p(z)$.*

2000 *Mathematics Subject Classification.* Primary 30C45, Secondary 30C80.

Key words and phrases. analytic functions; convex functions; convex of order alpha; univalent functions; differential subordination.

Proof. From the above argument in the proof of Theorem 1 and Definitions 1 and 2, we can easily prove Lemma 2. \square

Lemma 3. (*Carathéodory's lemma*) Let $p(z) = \sum_{n=0}^{\infty} c_n z^n$ be analytic and $\Re p(z) \geq 0$ in \mathbb{D} . Then we have

$$(3) \quad \Re \{2c_0\} \geq |c_n| \quad n \in \mathbb{N} = \{1, 2, 3, \dots\}.$$

Proof. Let us put

$$(4) \quad I = \frac{1}{2\pi i} \int_{|z|=1} p(e^{i\alpha} z) (2 - z^n - 1/z^n) \frac{dz}{z}, \quad (n\alpha = -\arg c_n).$$

Then we have

$$I = 2c_0 - |c_n|$$

and

$$\begin{aligned} \Re \{I\} &= \Re \left\{ \frac{2}{\pi} \int_0^{2\pi} p(e^{i(\alpha+\theta)}) \sin^2(n\theta/2) d\theta \right\} \\ &= \Re \{2c_0 - |c_n|\} \\ &\geq 0 \end{aligned}$$

or

$$\Re \{2c_0\} \geq |c_n| \quad n \in \mathbb{N}.$$

\square

Theorem 4. Let $p(z) = 1 + \sum_{n=1}^{\infty} c_n z^n$ be analytic in \mathbb{D} . Assume also that $p(z) \neq 0$ in \mathbb{D} , $0 < c_1$ and suppose that $p(z)$ be starlike in the direction of the diametal line of $p(z)$ with respect to $\theta = 0$ and diametal line coincides real axis. Then we have

$$(5) \quad |c_n| \leq n c_1 \quad n \in \mathbb{N} = \{1, 2, 3, \dots\}.$$




Fig. 1. A picture of Theorem 4.

Proof. From the hypothesis of Theorem 4, we can have the following

$$\begin{aligned} &\Re \left\{ \frac{1}{c_1} \left((p(z) - 1) \frac{1 - z^2}{z} \right) \right\} \\ &= \Re \left\{ 1 + \frac{c_2}{c_1} z + (c_3 - c_1) \frac{z^2}{c_1} + (c_4 - c_2) \frac{z^3}{c_1} + \dots + (c_n - c_{n-2}) \frac{z^{n-1}}{c_1} + \dots \right\} \\ &= \Re \{ (p(e^{i\theta}) - 1) (-2i \sin \theta) \} \\ &\geq 0, \end{aligned}$$

where $z = e^{i\theta}$ and $0 \leq \theta \leq \pi$. Then, from Caratheodory's lemma 3, that $\Re\{2c_0\} \geq |c_n|$, we have

$$(6) \quad \left| \frac{c_2}{c_1} \right| \leq 2$$

and

$$(7) \quad \left| \frac{c_n - c_{n-2}}{c_1} \right| \leq 2, \quad n = 3, 4, 5, \dots$$

From (6) and (7) we obtain (5). □

Theorem 5. Let $p(z) = 1 + \sum_{n=1}^{\infty} c_n z^n$ be analytic in \mathbb{D} . Assume also that $p(z) \neq 0$ in \mathbb{D} , $c_1 \neq 0$ and suppose that $p(z)$ be starlike in the direction of the diametal line of $p(z)$ with respect to $\theta = \beta$. Then we have

$$(8) \quad nc_1 \geq |c_n| \quad n \in \mathbb{N} = \{1, 2, 3, \dots\}.$$




Fig. 2. $a = p(e^{i(\pi+\beta)})$, $e = p(e^{i\beta})$.

Proof. Let us put $\alpha = \arg p(e^{i\beta})$, $\gamma = \arg c_1$ and

$$\begin{aligned} I(z) &= \frac{e^{-i\gamma}}{|c_1|} \left((p(e^{i\beta}z) - 1) \frac{1-z^2}{z} \right) \\ &= \frac{c_1 e^{i(\beta-\gamma)}}{|c_1|} + \frac{c_2 e^{i(2\beta-\gamma)}}{|c_1|} z + \frac{e^{i(\beta-\gamma)}}{|c_1|} (c_3 e^{2i\beta} - c_1) z^2 \\ &\quad + \frac{e^{i(2\beta-\gamma)}}{|c_1|} (c_4 e^{2i\beta} - c_2) z^3 + \dots \\ &\quad + \frac{e^{i((n-2)\beta-\gamma)}}{|c_1|} (c_n e^{2i\beta} - c_{n-2}) z^{n-1} + \dots \end{aligned}$$

From the hypothesis of Theorem 5, we have

$$\Re\{I(z)\} \geq 0 \quad z \in \mathbb{D}.$$

Applying Caratheodory's lemma 3, we have

$$\begin{aligned} \Re \{ 2e^{i(\beta\alpha+\gamma)} \} &\geq \left| \frac{c_2}{c_1} \right|, \\ \Re \{ 2e^{i(\beta\alpha+\gamma)} \} &\geq \left| \frac{c_3 e^{2i\beta} - c_1}{c_1} \right| \\ \Re \{ 2e^{i(\beta\alpha+\gamma)} \} &\geq \left| \frac{c_4 e^{2i\beta} - c_2}{c_1} \right| \\ &\vdots \\ \Re \{ 2e^{i(\beta\alpha+\gamma)} \} &\geq \left| \frac{c_n e^{2i\beta} - c_{n-2}}{c_1} \right| \\ &\vdots \\ &\vdots \end{aligned}$$

Applying the mathematical induction, we can easily obtain Theorem 5. □

For other results on the several problems connected to the starlikeness we refer also to the recent paper [4].

REFERENCES

- [1] M. S. Robertson, Analytic functions starlike in one direction, Amer. J. Math. 58(1936) 465–472.
- [2] M. S. Robertson, On the theory of univalent functions, Ann. Math. 37(1936) 374–408.
- [3] E. Study, Konforme Abbildung Einfachzusammenhangender Bereiche, B. C. Teubner, Leipzig und Berlin 1913.
- [4] M. Nunokawa, J. Sokół, On the Order of Strongly Starlikeness of Convex Functions of Order Alpha, Mediterranean J. Math. DOI 10.1007/s00009-013-0341-6.

UNIVERSITY OF GUNMA, HOSHIKUKI-CHO 798-8, CHUOU-WARD, CHIBA, 260-0808, JAPAN
E-mail address: mamoru_nuno@doctor.nifty.jp

DEPARTMENT OF MATHEMATICS, RZESZÓW UNIVERSITY OF TECHNOLOGY, AL. POWSTAŃCÓW WARSZAWY 12,
 35-959 RZESZÓW, POLAND
E-mail address: jsokol@prz.edu.pl