

Title	Synthesis, structure and reactivity of a 1-bromoalumole.
Author(s)	Wasano, Tatsuya; Agou, Tomohiro; Sasamori, Takahiro; Tokitoh, Norihiro
Citation	Chemical communications (2014), 50(60): 8148-8150
Issue Date	2014-05-22
URL	http://hdl.handle.net/2433/198686
Right	This journal is © The Royal Society of Chemistry 2014.
Type	Journal Article
Textversion	author

COMMUNICATION

Synthesis, structure and reactivity of a 1-bromoalumole

Cite this: DOI: 10.1039/x0xx00000x

Tatsuya Wasano, Tomohiro Agou, Takahiro Sasamori and Norihiro Tokitoh*^aReceived 00th January 2012,
Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

A stable 1-bromoalumole has been synthesized by the reaction of a 1,4-dilithio-1,3-butadiene derivative with AlBr_3 . The 1-bromoalumole was found to exist as a dimeric structure in the crystalline state. Reaction of the 1-bromoalumole with Mes^*Li afforded the corresponding Mes^* -substituted alumole ($\text{Mes}^* = 2,4,6\text{-}(t\text{-Bu})_3\text{C}_6\text{H}_2$), demonstrating the potential of the 1-bromoalumole for the functionalization of alumoles.

Heteroles of group 13 elements are expected to possess unique optical and electronic properties owing to the orbital interactions between the vacant p orbital of the group 13 elements with the π^* orbitals of the butadiene moieties.¹ In particular, boroles have received considerable attention from the viewpoint of not only fundamental chemistry but also material science.² On the other hand, the heavier element analogues of boroles, such as alumoles, remain underdeveloped so far.^{3,4} Recently, we have reported the synthesis of the first stable Lewis-base-free alumole **1** by using the bulky Mes^* group ($\text{Mes}^* = 2,4,6\text{-}(t\text{-Bu})_3\text{C}_6\text{H}_2$) (Figure 1).⁵ Alumole **1** can be reduced to the corresponding dianionic species, indicating the possible use of alumole derivatives for electron accepting materials. In order to extensively elucidate the properties of alumoles and utilize them for the key components of functional materials, facile and versatile synthetic methods of alumoles bearing various functional groups should be of importance.





Figure 1. The first Lewis base free alumole **1**.


The research groups of Braunschweig and Piers reported the usability of isolable 1-haloboroles as synthetic precursors of functionalized borole derivatives.^{6,7} For instance, the nucleophilic substitution at the boron atom of 1-chloroborole **I** afforded the corresponding 1-aminoborole **II** (Scheme 1).^{6c} Such distinctive reactivities of 1-haloboroles suggest that their aluminium analogues, *i.e.*, 1-haloalumoles, would be promising modular synthetic precursors of functionalized alumole derivatives.^{8,9} In addition, the

1-haloalumoles should exhibit unique characters owing to the electron-withdrawing and mesomeric effects of the halogeno group on the aluminium atom. Here, we report the synthesis, structure, and properties of 1-bromoalumole **2**, the first example of a stable 1-haloalumole derivative.



Scheme 1. Functionalization of a borole skeleton by using 1-chloroborole **I** as a borole source.^{6c}

1-Bromoalumole **2** was obtained as a colourless solid in 64% yield by the treatment of AlBr_3 with 1,4-dilithio-1,3-butadiene derivative **3**¹⁰ in toluene at room temperature (Scheme 2). Interestingly, Robinson et al. reported that the reaction of GaCl_3 with 1,4-dilithiotetraphenylbutadiene under similar reaction conditions afforded not the corresponding 1-chlorogallole but the first “spirogallane” derivative.¹¹ Compound **2** was found to undergo facile decomposition to give 4,5-diethyl-3,5-octadiene upon exposure to air and moisture even in the solid state. Compound **2** is thermally stable under an inert atmosphere either in the solid state (decomposed at 158 °C) or in solution (toluene-*d*₈, at 100 °C for 1 d in a sealed tube).



Scheme 2. Synthesis of 1-bromoalumole **2**.

X-Ray crystallographic analysis revealed that 1-bromoalumole **2** exists as a dimer in the crystalline state. In the unit cell, two crystallographically independent dimers of **2** were found, and each dimer unit has a crystallographic C_2 axis. The structure of one of the two independent dimers is shown in Figure 2. It should be noted that the Al1-C1 bond length (2.192(3) Å) is substantially elongated compared with those of alumole **1** ($\text{Al-C}(\text{ring})$ 1.952(2)–1.955(2) Å)⁵ and the Al1-C4 bond length (1.944(3) Å). Meanwhile, the Al1-

C1* bond length (2.052(3) Å) lies between those of the Al1–C1 and Al1–C4 bonds. The geometries at the C2, C3, and C4 atoms are nearly trigonal planar, whereas the C1 atom adopts a distorted tetrahedral environment. The C1–C2 bond length (1.388(5) Å) is slightly longer than the typical value for a double bond, suggesting somewhat weakened C1–C2 bond. The Al1–Al1* distance (2.5881(18) Å) is close to those of the dimers of triorganoaluminium compounds (e.g., Al₂Me₆: Al–Al 2.606(2) Å)¹² and those of the Al–Al single bonds of dialumanes (e.g., Bbp(Br)Al–Al(Br)Bbp (Bbp = 2,6-[CH(SiMe₃)₂]₂C₆H₃): Al–Al 2.592(3) Å).¹³ On the basis of these structural features, the bonding situation in the dimer unit can be interpreted in terms of multicentre bonding between the Al1, Al1*, C1, and C1* atoms, as in the case of triorganoaluminium compounds.¹⁴ The crystal structure of the dimer of **2** was reproduced by density functional theory (DFT) calculations (see the Electronic Supplementary Information for detail).¹⁵




Figure 2. Dimeric structure of 1-bromoalumole **2** in the crystalline state (thermal ellipsoids at the 50% probability level). One of the two crystallographically independent molecules is shown here. Hydrogen atoms are omitted for clarity. Selected bond lengths [Å] and angles [°]: Molecule **A**: Al1–Al1* 2.5881(18), Al1–Br1 2.3056(10), Al1–C1 2.192(3), C1–C2 1.388(5), C2–C3 1.490(3), C3–C4 1.352(5), Al1–C4 1.944(3), Al1–C1* 2.052(3), C1–Al1–C4 89.43(14), Al1–C1–C2 92.1(2), C1–C2–C3 120.8(3), C2–C3–C4 117.2(3), C3–C4–Al1 106.3(2), Al1–C1–Al1* 75.07(11), C1–Al1–C1* 101.15(12), C1–Al1–Br1 104.75(9), C4–Al1–Br1 122.64(11); Molecule **B**: Al1–Al1* 2.5988(19), Al1–Br1 2.3096(10), Al1–C1 2.193(3), C1–C2 1.375(5), C2–C3 1.494(5), C3–C4 1.348(5), Al1–C4 1.948(3), Al1–C1* 2.052(3), C1–Al1–C4 88.97(14), Al1–C1–C2 91.8(2), C1–C2–C3 120.6(3), C2–C3–C4 116.7(3), C3–C4–Al1 106.3(2), Al1–C1–Al1* 75.48(12), C1–Al1–C1* 100.05(13), C1–Al1–Br1 106.39(10), C4–Al1–Br1 120.93(11).

1-Bromoalumole **2** in solution was characterized by using variable-temperature (VT) NMR spectroscopy. At 25 °C, the ¹H NMR spectrum of compound **2** exhibited one sharp triplet and two broad signals (Figure 3). Cooling this sample to –60 °C resolved these signals into four triplet and seven multiplet signals. Similarly, the ¹³C NMR spectrum at –60 °C showed four signals attributable to the sp² carbon atoms in the alumole ring, these chemical shifts of which were well reproduced by the GIAO calculations on the DFT-optimized structure of the 1-bromoalumole dimer. The ¹³C NMR signal corresponding to the C2 nucleus in Figure 2 appeared in a characteristic low-field region (δ = 201.6 ppm), indicating the decrease in the electron density of the C2 atom. The vinylic carbon atoms in the β-positions to the aluminium atoms in dimeric alkenylaluminium compounds are known to exhibit characteristic lower-field shifted resonances in the ¹³C NMR spectra.¹⁶ Accordingly, the dimeric form should be kept in solution at the low temperature. At 100 °C, the ¹H NMR spectrum exhibited two triplets and two quartet signals corresponding to the methyl and methylene

protons of the ethyl substituents, respectively, and only two signals were observed for the alumole ring carbon atoms in the ¹³C NMR spectrum. These VT-NMR spectroscopic data could be accounted for by the rapid site exchange of the α-carbon atoms in the dimeric structure (i.e., C1 and C4 atoms in Figure 2) and/or by the monomer-dimer equilibrium.






Figure 3. Variable-temperature (VT) ¹H NMR spectra of 1-bromoalumole **2** in toluene-*d*₈.

At room temperature, addition of excess amount of THF to the C₆D₆ solution of 1-bromoalumole **2** quantitatively afforded 1-bromoalumole–THF adduct **4** as a colourless solid (Scheme 3). Adduct **4** was characterized by the ¹H and ¹³C NMR spectroscopies.



Scheme 3. Complexation of 1-bromoalumole **2** with THF.

Finally, a nucleophilic substitution reaction on the aluminium atom of 1-bromoalumole **2** was preliminarily examined. Reaction of **2** with equimolar quantity of Mes*Li¹⁷ in C₆D₆ was completed after 48 h at 60 °C to afford the expected Mes*-substituted alumole **1** as a sole product.⁵ Alumole **1** was isolated as colourless crystals in 48% yield after recrystallization from hexane (Scheme 4). This preliminary result demonstrated that 1-bromoalumole **2** should be a promising building block for novel functionalized alumole derivatives.



Scheme 4. Reaction of 1-bromoalumole **2** with Mes*Li.

In summary, we have succeeded in the synthesis of 1-bromoalumole **2**, the first alumole derivative bearing a halogen substituent on the aluminium atom. It was revealed that 1-bromoalumole **2** exhibits its unique dimeric structure in the crystalline state. Treatment of 1-bromoalumole **2** with Mes*Li afforded the corresponding Mes*-substituted alumole **1**, demonstrating the potential of **2** as a modular-precursor of functionalized alumole derivatives.

This work was supported by JSPS KAKENHI (Nos. 22350017, 24550048, 24655028, and 24109013), by Grants for Excellent Graduate Schools, MEXT, Japan, and by the “Molecular Systems Research” project of RIKEN Advanced Science Institute. T.A. thanks to the Kyoto Technoscience Center and the Research Institute for Production Development for the financial supports. The X-ray crystallographic analysis of 1-bromoalumole **2** was performed at the BL38B1 of the SPring-8 with the approval of the Japan Synchrotron Radiation Research Institute (JASRI; proposal No. 2013B1083). The authors are grateful to Dr. K. Miura, Dr. S. Baba, and Dr. N. Mizuno (JASRI) for the X-ray crystallographic analyses at the SPring-8.

Notes and references

^a Institute for Chemical Research, Kyoto University, Gokasho, Uji, Kyoto 611-0011 (JAPAN), Fax: (+81)774-38-3209, E-mail: tokitoh@boc.kuicr.kyoto-u.ac.jp.

† Electronic Supplementary Information (ESI) available: Experimental procedures, spectroscopic and crystallographic data, and computational data. CCDC 974827. For ESI and crystallographic data in CIF or other electronic format, see DOI: 10.1039/c000000x/.

‡ Crystal data for **1**: C₂₄H₄₀Al₂Br₂, *M*_r = 542.34, colourless block, size 0.02 × 0.01 × 0.01 mm³, λ = 0.85000 Å, *T* = -170 °C, monoclinic, space group *P*2₁/*n* (#13), *a* = 17.8044(1) Å, *b* = 8.4388(1) Å, *c* = 17.8350(1) Å, β = 100.7991(3)°, *V* = 2632.21(4) Å³, *Z* = 4, μ(λ = 0.85000 Å) = 4.991 mm⁻¹, total reflections 22610, independent reflections 4671, parameters 262, *R*_{int} = 0.0412, θ_{max} = 31.00°, *R*₁ [*I* > 2σ(*I*)] = 0.0489, w*R*₂ (all data) = 0.1333, GOF 1.050.

- (a) S. Yamaguchi and A. Wakamiya, *Pure Appl. Chem.*, 2006, **78**, 1413-1424; (b) S. Yamaguchi and K. Tamao, *Chem. Lett.*, 2005, **34**, 2-7; (c) U. Salzner, J. B. Lagowski, P. G. Pickup and R. A. Poirier, *Synth. Met.*, 1998, **96**, 177-189.
- (a) H. Braunschweig, I. Krummenacher and J. Wahler, in *Adv. Organomet. Chem.*, Vol. 61, ed. A. Hill and M. J. Fink, Academic Press, 2013, pp. 1-53; (b) H. Braunschweig and T. Kupfer, *Chem. Commun.*, 2011, **47**, 10903-10914, and references therein.
- Alumole-Lewis base complexes: (a) H. Hoberg and R. Krause-Göing, *J. Organomet. Chem.*, 1977, **127**, C29-C31; (b) C. Krüger, J. C. Sekutowski, H. Hoberg and R. Krause-Göing, *J. Organomet. Chem.*, 1977, **141**, 141-148; (c) H. Hoberg, R. Krause-Göing, C. Krüger and J. C. Sekutowski, *Angew. Chem., Int. Ed. Engl.*, 1977, **16**, 183-184.
- Examples of gallole derivatives: (a) T. Matsumoto, K. Tanaka and Y. Chujo, *J. Am. Chem. Soc.*, 2013, **135**, 4211-4214; (b) A. H. Cowley, D. S. Brown, A. Decken and S. Kamepalli, *Chem. Commun.*, 1996, 2425-2426; (c) A. H. Cowley, F. P. Gabbaï and A. Decken, *Angew. Chem., Int. Ed. Engl.*, 1994, **33**, 1370-1372.
- T. Agou, T. Wasano, P. Jin, S. Nagase and N. Tokitoh, *Angew. Chem. Int. Ed.*, 2013, **52**, 10031-10034.
- (a) H. Braunschweig, C.-W. Chiu, A. Damme, K. Ferkinghoff, K. Kraft, K. Radacki and J. Wahler, *Organometallics*, 2011, **30**, 3210-3216; C. Fan, W. E. Piers and M. Parvez, *Angew. Chem. Int. Ed.*, 2009, **48**,

2955-2958; (c) H. Braunschweig and T. Kupfer, *Chem. Commun.*, 2008, 4487-4498.

- (a) H. Braunschweig, C.-W. Chiu, D. Gamon, K. Größ, C. Hörl, T. Kupfer, K. Radacki and J. Wahler, *Eur. J. Inorg. Chem.*, 2013, 1525-1530; (b) H. Braunschweig, V. Dyakonov, J. Oscar C. Jimenez-Halla, K. Kraft, I. Krummenacher, K. Radacki, A. Sperlich and J. Wahler, *Angew. Chem. Int. Ed.*, 2012, **51**, 2977-2980; (c) H. Braunschweig, C.-W. Chiu, K. Radacki and T. Kupfer, *Angew. Chem. Int. Ed.*, 2010, **49**, 2041-2044; (d) H. Braunschweig, C.-W. Chiu, K. Radacki and P. Brenner, *Chem. Commun.*, 2010, **46**, 916-918; (e) H. Braunschweig, C.-W. Chiu, J. Wahler, K. Radacki and T. Kupfer, *Chem. Eur. J.*, 2010, **16**, 12229-12233.
- 1-Halogalole-Lewis base complexes: (a) M. Saito, T. Akiba, M. Kaneko, T. Kawamura, M. Abe, M. Hada and M. Minoura, *Chem. Eur. J.*, 2013, **19**, 16946-16953; (b) P. Wei, X.-W. Li and G. H. Robinson, *Chem. Commun.*, 1999, 1287-1288; (c) J. A. C. Clyburne, R. D. Culp, S. Kamepalli, A. H. Cowley and A. Decken, *Inorg. Chem.*, 1996, **35**, 6651-6655; (d) P. J. Fagan, W. A. Nugent and J. C. Calabrese, *J. Am. Chem. Soc.*, 1994, **116**, 1880-1889.
- Proposal of 1-haloalumoles as reactive intermediates: (a) V. A. D'yakonov, R. A. Tuktarova and U. M. Dzhemilev, *Russ. Chem. Bull.*, 2011, **60**, 1633-1639; (b) Z. Xi and P. Li, *Angew. Chem. Int. Ed.*, 2000, **39**, 2950-2952.
- L. Liu, W.-X. Zhang, Q. Luo, H. Li and Z. Xi, *Organometallics*, 2010, **29**, 278-281.
- J. Su, S. D. Goodwin, X.-W. Li and G. H. Robinson, *J. Am. Chem. Soc.*, 1998, **120**, 12994-12995.
- J. C. Huffmann and W. E. Streib, *J. Chem. Soc. D, Chem. Commun.*, 1971, 911-912.
- T. Agou, K. Nagata, H. Sakai, Y. Furukawa and N. Tokitoh, *Organometallics*, 2012, **31**, 3806-3809.
- (a) S. Aldridge, A. J. Downs and D. L. Kays, in *The Group 13 Metals Aluminium, Gallium, Indium and Thallium*, ed. S. Aldridge and A. J. Downs, Wiley, 2011, pp. 148-245; (b) C. Eisenbroich, *Organometallics*, 3rd ed., Wiley-VCH, Weinheim, Germany, 2006, pp. 87-137.
- The geometry optimizations and GIAO calculations were carried out at the B3PW91/6-311G(2df) and B3PW91/6-311+G(2df) level of theory, respectively. M. J. Frisch et al., *Gaussian 09* (Rev. C.01), Gaussian, Inc., Wallingford, Connecticut, USA, 2010 (see ESI for the complete citation).
- (a) W. Uhl, E. Er, A. Hepp, J. Kösters, M. Layh, M. Rohling, A. Vinogradov, E.-U. Würthwein and N. Ghavtadze, *Eur. J. Inorg. Chem.*, 2009, 3307-3316; (b) S. S. Karpiñiec, D. S. McGuinness, M. G. Gardiner, B. F. Yates and J. Patel, *Organometallics*, 2011, **33**, 1569-1576.
- A. Wakamiya, K. Mishima, K. Ekawa and S. Yamaguchi, *Chem. Commun.*, 2008, 579-581.

Table of Contents graphic

Title: Synthesis, structure and reactivity of a 1-bromoalumole

Authors: Tatsuya Wasano, Tomohiro Agou, Takahiro Sasamori and Norihiro Tokitoh

