

Title	Fluorescent characteristics of dissolved organic matter produced by bloom-forming coastal phytoplankton
Author(s)	Fukuzaki, K.; Imai, I.; Fukushima, K.; Ishii, K.-I.; Sawayama, S.; Yoshioka, T.
Citation	Journal of Plankton Research (2014), 36(3): 685-694
Issue Date	2014-03-03
URL	http://hdl.handle.net/2433/198835
Right	This is a pre-copyedited, author-produced PDF of an article accepted for publication in "J. Plankton Res." following peer review. The version of record [Koji Fukuzaki, Ichiro Imai, Keitaro Fukushima, Ken-Ichiro Ishii, Shigeki Sawayama, and Takahito Yoshioka Fluorescent characteristics of dissolved organic matter produced by bloom-forming coastal phytoplankton. J. Plankton Res. (May/June 2014) 36(3): 685-694 first published online March 3, 2014 doi:10.1093/plankt/fbu015] is available online at: http://plankt.oxfordjournals.org/content/36/3/685 .
Type	Journal Article
Textversion	author

1
2
3
4
5
6 1 **Fluorescent characteristics of dissolved organic matter produced by**
7
8
9 2 **bloom-forming coastal phytoplankton**
10
11
12 3

14 **Koji Fukuzaki^{a*}, Ichiro Imai^b, Keitaro Fukushima^c, Ken-Ichiro Ishii^d, Shigeki**
15
16
17 **Sawayama^d and Takahito Yoshioka^{a, e**}**
18
19
20
21 6

22
23 *^aLaboratory of Forest Information, Division of Forest and Biomaterials Science,*
24
25
26 *Graduate School of Agriculture, Kyoto University, Kitashirakawa Oiwake-cho,*
27
28
29 *Sakyo-ku, Kyoto 606-8502, Japan*
30
31

32 *^bPlankton Laboratory of Marine Biology, Graduate School of Fisheries Sciences,*
33
34
35 *Hokkaido University, 3-1-1 Minato-cho, Hakodate, Hokkaido 041-8611, Japan*
36
37

38 *^cDivision of Research Project Promotion, Field Science Education and Research Center,*
39
40
41 *Kyoto University, Kitashirakawa Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan*
42
43

44 *^dLaboratory of Marine Environmental Microbiology, Division of Applied Bioscience,*
45
46
47 *Graduate School of Agriculture, Kyoto University, Kitashirakawa Oiwake-cho,*
48
49
50 *Sakyo-ku, Kyoto 606-8502, Japan*
51

52 *^eDivision of Forest Ecosystem, Field Science Education and Research Center, Kyoto*
53
54
55 *University, Kitashirakawa Oiwake-cho, Sakyo-ku, Kyoto 606-8502, Japan*
56
57

1

2 *Corresponding author. Tel.: +81-75-753-6440; Fax: +81-75-753-6443

3 E-mail: fukuzaki@kais.kyoto-u.ac.jp (K. Fukuzaki)

4 **Corresponding author. Tel.: +81-75-753-6421; Fax: +81-75-753-6443

5 E-mail: yoshioka@kais.kyoto-u.ac.jp (T. Yoshioka)

6

7 **Keywords**

8 Fluorescent dissolved organic matter, Excitation emission matrix, Incubation
9 experiment, Phytoplankton, Spectrofluorometer, Bloom

10

11 **Abstract**

12 Dynamics and sources of fluorescent dissolved organic matter (FDOM) are important
13 for understanding the biogeochemical process in aquatic ecosystems. This study aimed
14 to analyse direct production of FDOM by marine phytoplankton cultures and reveal
15 fluorescent characteristics of exuded FDOM. Axenic cultures of eight species of
16 bloom-forming marine phytoplankton, including two diatoms; a raphidophyte; two
17 dinoflagellates; a chlorophyte; a cryptophyte and a haptophyte, were incubated in an
18 artificial medium. Excitation emission matrices (EEMs) of FDOM in the culture

1 medium were spectrofluorometrically measured. FDOM production was observed in all
2 species, and fluorescent characteristics of the exudates varied considerably among
3 species. Measured EEMs had peaks at 350 nm/450 nm (excitation/emission) for the
4 diatom *Ditylum brightwellii* and 370 nm/450–470 nm for the raphidophyte *Heterosigma*
5 *akashwo* and the chlorophyte *Oltmansiellopsis viridis*, which were previously regarded
6 as the peaks of terrestrially derived humic-like substances. Direct production of FDOM
7 by marine phytoplankton should be considered in future studies of FDOM dynamics in
8 marine systems. Species-specific features of FDOM might be used for early detection of
9 noxious bloom because this method is simple, rapid, and suitable for monitoring.

10

11 **Introduction**

12

13 Marine dissolved organic matter (DOM) is one of the major factors affecting the
14 global carbon cycles and is the largest ocean reservoir of reduced carbon (Hansell et al.,
15 2009). DOM also affects various biological processes such as bacterial respiration and
16 microalgal primary production by serving as a substrate for bacterial populations
17 (Findlay et al., 2003; Findlay and Sinsabaugh, 2003), controlling the transport and
18 availability of trace elements (van den Berg et al., 1986; Shiller et al., 2006; Laglera et

1 al., 2007; Laglera and van den Berg, 2009) and absorbing photosynthetic effective
2 sunlight (Sulzberger and Durisch-Kaiser, 2009) and harmful ultraviolet (UV) light
3 (Nielsen et al., 1995; Nielsen and Ekelund, 1995). The quality and quantity of DOM are
4 thought to influence these ecological functions of DOM and should be taken into
5 account when evaluating the biogeochemical processes in aquatic ecosystems.

6 The fluorescent spectroscopic characterisation of chromophoric DOM (CDOM), which
7 is a coloured fraction of DOM, is an excellent method for evaluating the source and
8 quality of DOM (Coble, 1996, 2007; Coble et al., 1990, 1993; Stedmon et al., 2003;
9 Jaffé et al., 2008). This method compiles individual fluorescent spectra at each
10 excitation (Ex) wavelength to generate three-dimensional excitation emission matrices
11 (EEMs). EEMs of DOM can be highly variable and are controlled by different physical,
12 chemical and biological processes; therefore, they can have important ecological
13 consequences (Maie et al., 2006; Jaffé et al., 2008). In aquatic systems, fluorescent
14 CDOM (FDOM) is assumed to be derived from biological processes in the system
15 (autochthonous) as well as from the transport of terrestrial organic matter from rivers
16 and the surrounding environment (allochthonous) (Jaffé et al., 2008; Yamashita and
17 Tanoue, 2008). Autochthonous production is thought to be mainly derived from
18 bacterial metabolic by-products (Nieto-Cid et al., 2006; Shimotori et al., 2009;

1 Yamashita and Tanoue, 2008).

2 It has been well documented that phytoplankton is one of the main sources of organic
3 matter in the sea, because it releases organic compounds such as carbohydrates and
4 polysaccharides (Biddanda and Benner, 1997). Recently, the exudates from marine
5 phytoplankton have also been shown to have fluorescent properties, and they may
6 contribute as a source of marine autochthonous FDOM (Romera-Castillo et al., 2010).

7 In coastal and estuarine environments, various phytoplankton species occasionally grow
8 massively and attain high cell densities. These algal blooms are thought to have
9 substantial impacts on DOM dynamics of a region.

10 To understand direct FDOM production by marine phytoplankton, it is essential to
11 conduct experiments using axenic cultures. However, the maintenance of axenic
12 cultures is difficult, and very few studies have analysed FDOM production by axenic
13 cultures of marine phytoplankton. Indeed, direct production of FDOM has been tested
14 in only four species in the genera *Chaetoceros*, *Skeletonema*, *Prorocentrum* and
15 *Micromonas* (Romera-Castillo et al., 2010). The optical properties of DOM varied
16 considerably among these four species (Romera-Castillo et al., 2010).

17 We aimed to evaluate direct production of FDOM by eight major bloom-forming
18 coastal phytoplankton species from diverse taxonomic groups of six classes and to

1 reveal the fluorescent characteristics of the exuded FDOM.

2

3 **Methods**

4

5 *Phytoplankton cultures*

6

7 Axenic cultures of the following species were used in the incubation experiments: the
8 diatoms *Ditylum brightwellii* and *Chaetoceros curvisetus*, the raphidophyte
9 *Heterosigma akashiwo*, the dinoflagellates *Heterocapsa circularisquama* and
10 *Alexandrium catenella*, the chlorophyte *Oltmansiellopsis viridis*, the cryptophyte
11 *Rhodomonas ovalis* and the haptophyte *Pleurochrysis roscoffensis* (Table I). These are
12 commonly found bloom-forming species in the Western Pacific (Omura et al., 2012).
13 Culture of *C. curvisetus* was obtained from sea bottom sediment and made axenic (Ishii,
14 personal communication). Axenic cultures of the other seven species were made by the
15 methods described elsewhere (Imai and Yamaguchi, 1994; Nagai et al., 1998). All the
16 cultures were axenically maintained in the modified IHN medium (Imai et al., 2004).
17 The axenic conditions of each culture were confirmed by DAPI staining and
18 epifluorescence microscopy before the incubation experiments (Imai, 1987).

1

2 *Incubation experiment*

3

4 Modified IHN medium (Imai et al., 2004) was prepared using Milli-Q water and
5 dispensed into 50 mL conical glass flasks (30 mL in each) with autoclavable plastic
6 caps (Iwaki, Tokyo, Japan). The prepared culture medium was sterilised by autoclaving
7 at 121°C for 15 min. To prevent contamination with organic compounds, the glass
8 flasks were pre-combusted at 450°C for 4 h. Well-grown strains of maintenance cultures
9 were inoculated (600 µL) into each flask in a clean bench. The inoculated cultures were
10 incubated under cool-white fluorescent light at 93–145 µmol photons m⁻² s⁻¹ with
11 14:10-h light:dark cycle until they reached the stationary phase, which was 6 days for *C.*
12 *curvisetus*, 14 days for *H. akashiwo*, 27 days for *A. catenella* and 12 days for the other
13 species. Incubation temperature was set at 25°C for *H. circularisquama* and 20°C for
14 the other species, because only the maintenance culture of *H. circularisquama* indicated
15 the better growth at 25°C while the other species showed the better growth at 20°C. An
16 autoclaved culture medium without plankton inoculation was also kept in the same
17 conditions as a control. A small amount of each culture was taken to measure the *in vivo*
18 fluorescence using a fluorometer (Model 10-AU 005, Turner Designs, Sunnyvale,

1
2
3
4
5
6 1 California, USA) at the start, middle and end of the culture experiments. The cell
7
8
9 2 densities of the cultures were determined by microscopic counting at the end of the
10
11
12 3 incubation. All the culture experiments were conducted in triplicate.
13
14
15 4

17 5 *Analysis*
18
19
20 6

21
22
23 7 After reaching the stationary phase, the cultures were filtered into pre-combusted
24
25
26 8 brown glass bottles using radiation-sterilised disposable syringes (Terumo Corp., Tokyo,
27
28
29 9 Japan) and GF/F filters (Whatman, Tokyo, Japan). The glass bottles and glass filters
30
31
32 10 were pre-combusted at 450°C for 4 h. The three-dimensional EEM spectra of the culture
33
34
35 11 filtrates were measured using a spectrofluorometer (Model F-7000, Hitachi
36
37
38 12 High-Technologies, Tokyo, Japan), which was equipped with a 150 W xenon lamp. The
39
40
41 13 corrections of the spectra were performed with Rhodamine B solution, according to the
42
43
44 14 instructions in the instrument operation manual and the method described by Yoshioka
45
46
47 15 et al. (2007). The scanning ranges were 250–400 nm for Ex and 280–480 nm for
48
49
50 16 emission (Em). Fluorescence intensity (FI) was measured at 5-nm intervals for Ex and
51
52
53 17 1-nm intervals for Em, with a scanning speed of 1200 nm min⁻¹. The bandwidths were 5
54
55
56 18 nm for both Ex and Em. The EEM spectrum of the control medium was subtracted from

1
2
3
4
5
6 1 each sample EEM spectrum to obtain the net increase in FDOM as a result of the
7
8
9 2 microalgal activities. All the sample data of FI were standardised using quinine sulphate
10
11
12 3 units (QSU), where 10 QSU correspond to the FI at 350 nm/450 nm of a 10 $\mu\text{g L}^{-1}$
13
14
15 4 quinine sulphate solution in 0.1 N H_2SO_4 . Solutions of quinine sulphate (Nacalai
16
17
18 5 Tesque, Inc., Kyoto, Japan) were measured with each set of samples. The FI data of
19
20
21 6 each species were averaged for triplicates.

22
23
24 7 The bulk dissolved organic carbon (DOC) concentration of the culture filtrate was
25
26
27 8 measured using a Shimadzu TOC-V_{CSH} total organic carbon analyzer (Shimadzu, Kyoto,
28
29
30 9 Japan). The DOC content of each sample was determined using a calibration method
31
32
33 10 based on a potassium hydrogen phthalate standard for each measurement. Each sample
34
35
36 11 was injected five times, and the three values that yielded the minimum standard
37
38
39 12 deviation were used to calculate the average DOC value for a sample. The net increase
40
41
42 13 in DOC was determined by subtracting the control DOC from DOC of each sample.
43
44
45

46 *Calculation of the index*

47
48
49
50
51
52
53
54

55 17 The biomass of the cultures at the final point was estimated using previously reported
56
57
58 18 models and equations (Sun and Liu, 2003; Menden-Deuer and Lessard, 2000) because

1 we did not directly measure the cell biomass. The cell volume was calculated by
2 approximating the complex cell shapes as simple three-dimensional geometric models
3 (Sun and Liu, 2003) based on the cell sizes obtained from microscopic measurements. *D.*
4 *brightwellii* was approximated as a prism with a triangle-based girdle view shape
5 (30-H), and *C. curvisetus* was approximated as a prism with an elliptic-based girdle
6 view shape (29-H). *H. akashiwo* and *H. circularisquama* were approximated as a cone +
7 hail sphere shape (9-H) (Sun and Liu, 2003). *A. catenella* was approximated as an
8 ellipsoid (3-H). *O. viridis*, *R. ovalis* and *C. roscoffensis* were approximated as prolate
9 spheroids (2-H) (Sun and Liu, 2003). The calculated cell volumes were converted into
10 the cell biomass using the equations described by Menden-Deuer and Lessard (2000).
11 As an indicator of the percentage of extracellular carbon released during total
12 photosynthetic production, the apparent percentage of photosynthetic extracellular
13 release (APER) values (Romera-Castillo et al., 2010) were calculated using the
14 following formula:

$$\text{APER (\%)} = \text{DOC} / (\text{DOC} + \text{Biomass}) \times 100,$$

18 where DOC and Biomass represent the net increase in DOC in the culture medium (mg

1 C L⁻¹) and the phytoplankton biomass (mg C L⁻¹), respectively.

3 **Results**

5 *Growth of the cultures and cell density attained*

7 The cell densities of the stationary phase cultures varied from $11 \pm 1 \times 10^3$ (cells mL⁻¹)
8 for *D. brightwellii* to $253 \pm 45 \times 10^3$ (cells mL⁻¹) for *R. ovalis*. *H. circularisquama*
9 produced the highest biomass of 33.1 ± 7.9 mg C L⁻¹ among the eight species examined,
10 and *R. ovalis* produced the lowest biomass of 3.5 ± 0.6 mg C L⁻¹. The average growth
11 rate in the exponential growth phase, which was calculated on the basis of the *in vivo*
12 fluorescence values, varied from 0.35 ± 0.03 day⁻¹ for *H. akashiwo* to 0.97 ± 0.07 day⁻¹
13 for *C. curvisetus* (Table II). The growth rates of these species in modified IHN medium
14 were previously reported as 0.39–1.27 day⁻¹ (Naito et al. 2008), which are comparable
15 with the values obtained in the present study.

17 *DOC and APER*

1 The increased DOC concentrations and APER values are summarised in Table III. Net
2 increase in DOC concentrations ranged from $19.3 \pm 5.1 \text{ mg C L}^{-1}$ for *H. akashiwo* to
3 $49.4 \pm 7.5 \text{ mg C L}^{-1}$ for *P. roscoffensis* during incubation periods (Table III). APER
4 values were estimated to be between 46.1% for *A. catenella* and 80.2% for *P.*
5 *roscoffensis*, with rather wide variations. There were no significant differences in APER
6 values among the eight species (one-way ANOVA, $p = 0.198$).

8 *Optical properties of DOM exudates from phytoplankton*

10 Figure 1 shows the average EEMs of the net FDOM increases for each triplicate
11 culture filtrate sample. In all the cultures, EEMs had fluorescence peaks in the
12 protein-like and humic-like regions (Table IV, Fig. 1). In the protein-like region, as
13 described by Coble (1996) at Ex/Em 275 nm/340 nm (peak T) and Ex/Em 275 nm/310
14 nm (peak B), the most prominent peaks were at 280 nm/349–357 nm (Ex/Em) for the
15 cultures of *C. curvisetus*, *H. circularisquama* and *A. catenella*. In the corresponding
16 region, *H. akashiwo*, *O. viridis* and *R. ovalis* had slightly shorter Em wavelengths in the
17 range 280 nm/324–337 nm. *D. brightwellii* and *P. roscoffensis* had peaks at 255 nm/316
18 nm and 250 nm/350 nm, respectively, although these peaks were rather uniform (Fig. 1).

1 The protein-like peaks were broad toward the longer Em wavelengths. In particular, *H.*
2 *circularisquama* had a considerably broad peak toward longer Ex/Em wavelengths in
3 the region and appeared to have an overlapping peak at 290 nm/410 nm (Fig. 1), which
4 corresponded to peak M, i.e. marine humic-like substances, as defined by Coble et al.
5 (1998). Peak M was only observed with *H. circularisquama* in the present study,
6 whereas peak M was observed with all the marine phytoplankton axenic cultures
7 examined by Romera-Castillo et al. (2010). *O. viridis* has the most prominent peak
8 among the eight species in the protein-like fluorescent region, at 280 nm/337 nm, with
9 an intensity of 2.22 ± 0.36 QSU (Fig. 1, Table IV).

10 *D. brightwellii*, *H. circularisquama*, *H. akashiwo* and *O. viridis* had the maxima of
11 humic-like fluorescence at 355 nm/454 nm, 340 nm/461 nm, 375 nm/473 nm and 375
12 nm/462 nm, respectively (humic-like peak 1 in Table IV, Fig. 1). *O. viridis* had the most
13 conspicuous peak in the fluorescent area, with an FI of 0.86 ± 0.24 QSU. *C. curvisetus*,
14 *A. catenella*, *R. ovalis* and *P. roscoffensis* did not have defined peaks in this area (Fig.
15 1). Coble (1996) referred to the peaks in the region of 350 nm/420–480 nm as peak C
16 and identified them as humic-like DOM components. The peaks observed in the
17 cultures of *D. brightwellii* and *H. circularisquama* were assumed to correspond to peak
18 C. Furthermore, the peaks resembled the peak produced by a *M. pusilla* culture at 348

1 nm/434 nm and 348 nm/436 nm, as reported by Romera-Castillo et al. (2010). *H. akashiwo* and *O. viridis* had peaks at longer Em wavelengths, i.e. 370 nm/450–470 nm (Ex/Em), and these peaks were very close to the C1 peak reported to be related to terrestrial humic substances by Yamashita et al. (2008).

Humic-like peaks were also detected in all the samples at 250–255 nm/446–471 nm (humic-like peak 2), which corresponded to peak A (260 nm/458 nm), as defined by Coble (1996). The FI of humic-like peak 2 ranged from 0.54 ± 0.04 QSU for *C. curvisetus* to 1.20 ± 0.16 QSU for *O. viridis*.

DOC-specific FIs of humic like peak 1 and 2 and protein like peak considerably varied among species (Table IV). The DOC-specific FI of humic-like peak 1 was highest for *O. viridis* (Table IV). The DOC-specific FI of humic-like peak 2 ranged from 0.011 ± 0.003 QSU L mg C⁻¹ for *P. roscoffensis* to 0.046 ± 0.016 QSU L mg C⁻¹ for *H. akashiwo* (Table IV). The DOC-specific FI of protein-like peak ranged from 0.011 ± 0.002 QSU L mg C⁻¹ for *P. roscoffensis* to 0.080 ± 0.018 QSU L mg C⁻¹ for *O. viridis* (Table IV). The cell density-specific FDOM production rates were the highest for *D. brightwellii* in terms of both the protein-like and humic-like peaks (Table V). In terms of the biomass-specific FDOM production rates (data not shown), *R. ovalis* had the highest values for both fluorescent peaks.

1

2 **Discussion**

3

4 *DOC and APER*

5

6 Sharp et al. (1977) reported that photosynthetic extracellular release (PER) values
7 obtained using the ^{14}C method reached up to 70%. APER values obtained in the present
8 study were 46%–80%, which were comparable with the previously reported values. In
9 contrast, Lancelot and Billen (1985) reported that the PER value using the culture
10 method was 0%–20%. Compared with APER values of 10%–18% obtained in a
11 previous study using axenic cultures of microalgae (Romera-Castillo et al., 2010), the
12 present study obtained rather high APER values (46%–80%). In the present study, the
13 incubation period was 6–14 days, with the exception of 27 days for *A. catenella* (Table
14 II), which was longer than the incubation period (3–6 days) used previously
15 (Romera-Castillo et al., 2010). It was previously reported that PER values of the diatom
16 *Chaetoceros affinis* increased by up to 58% during the phase of decreasing
17 photosynthetic activity because of nutrient depletion (Myklestad et al., 1989). Thus,
18 APER values appear to be affected in nutrient-limited environments (Lancelot and

1 Billen, 1985). The longer incubation period probably resulted in larger release of DOC
2 from the phytoplankton cells, although we did not measure the nutrient concentration.
3 The influence of light and nutrient stress on DOC secretion should be tested in future
4 studies because phytoplankton are likely to experience light and nutrient stresses in
5 natural environments.

6 7 *Peak assignments and possible functions of the fluorescent DOM*

8
9 Significant peaks were observed at approximately 275 nm/340 nm (Ex/Em) in the
10 culture filtrates of all the species. This peak was considered to be related to protein-like
11 substances and was previously reported as peak T (Coble 1996). Romera-Castillo et al.
12 (2010) also reported a corresponding peak in cultures of *Chaetoceros*, *Skeletonema*,
13 *Prorocentrum* and *Micromonas*. Phytoplankton are known to release extracellular
14 nitrogenous compounds such as proteins, peptides and amino acids as well as
15 carbohydrates such as polysaccharides (Goldman et al., 1992; Myklestad, 1995). Some
16 of these substances with proteinaceous aromatic structures were probably detected as
17 peak T. Similar peaks were detected in EEM measurements of coastal seawaters (Para et
18 al., 2010; Yamashita et al., 2008; Maie et al., 2007). Yamashita et al. (2008) suggested

1 that a tryptophan-like fluorescent peak in a seawater sample from Mikawa Bay, Japan
2 was derived from relatively fresh long-chain peptides, which were readily degradable.
3 Maie et al. (2007) suggested that the origin of peak T in a temperate coastal seawater
4 was a mixture of proteinaceous compounds and the phenolic structures contained in
5 humic substances.

6 *H. circularisquama* had a peak at 290 nm/410 nm, which was similar to peak M
7 designated by Coble (1996). Peak M is found at 290–310 nm/370–410 nm (Ex/Em) and
8 was first reported in seawater collected during a phytoplankton bloom in the Gulf of
9 Maine (Coble, 1996) and thereafter in samples from an upwelling region in the Arabian
10 Sea (Coble et al., 1998). This peak is considered to be related to marine humic-like
11 substances (Coble, 1996). In a recent study, peak M was detected in the culture filtrates
12 of four marine algal species: *Chaetoceros* sp., *S. costatum*, *P. minimum* and *M. pusilla*
13 (Romera-Castillo et al., 2010). These findings are consistent with FDOM produced by
14 *H. circularisquama*, which showed peak M in the present study. The absence of peak M
15 in the cultures of other species examined in the present study have been possibly due to
16 differences in the metabolic processes among the species, such as different
17 photosynthetic pigments.

18 Coble (1996, 2007) suggested that terrestrial humic-like materials produce two peaks

1 at 240–260 nm/400–460 nm (Ex/Em) (peak A) and 320–360 nm/420–460 nm (Ex/Em)
2 (peak C). All the species produced peaks at 250–255 nm/ 446–471 nm (humic-like peak
3 2), which were very close to peak A. Humic-like peak 2 is also similar to Component 1
4 (<260/458 nm) in a study that used the EEM-PARAFAC method in Ise Bay, Japan
5 (Yamashita et al., 2008). In other studies, this peak was also reported to be attributable
6 to land-derived components, e.g. Q2 (Cory and Mcknight, 2005) and Component 1
7 (Stedmon and Markager, 2005).
8 *D. brightwellii* had a peak at 350 nm/450 nm (Ex/Em), which corresponded to the
9 region of peak C (Coble, 1996). *H. akashiwo* and *O. viridis* had peaks at a slightly
10 longer wavelength of 370 nm/450–470 nm (Ex/Em), and these peaks were very close to
11 the peak attributed to terrestrial humic substances (Yoshioka et al., 2007; Yamashita et
12 al., 2008, 2011). These findings suggest that FDOM produced by phytoplankton
13 occasionally have a peak in the region previously assigned to terrestrial humic
14 substances. Thus, we should be cautious when investigating the dynamics and sources
15 of DOM in coastal areas using fluorescent analysis.

16 It is not known how and why phytoplankton release humic-like substances. Bjørnsen
17 (1988) suggested that DOM exudates are caused by the passive diffusion of metabolic
18 by-products. The low-molecular-weight compounds produced by photosynthetic

1 metabolism and by-products of the decomposition of cellular polymers are assumed to
2 be released extracellularly (Myklestad, 1995). However, it is possible that
3 phytoplankton exude FDOM with ecological functions. Many species of microbial
4 prokaryotes, fungi and some phytoplankton are known to secrete organic iron ligands,
5 known as siderophores, in iron-depleted environments (Naito et al., 2001, 2004; Vraspir
6 and Butler, 2009). Most siderophores appear to have aromatic structures, although the
7 chemical structures of the siderophores secreted by eukaryotic phytoplankton are not
8 clear at present (Naito et al., 2001; Vraspir and Butler, 2009). Humic substances also
9 have aromatic structures and the capacity for metal complexation. Naito et al. (2001)
10 also suggested that *R. ovalis* secretes siderophores; this species had the highest
11 biomass-specific production of FDOM in the present study. Thus, it is possible that
12 siderophores are involved with the release of humic-like fluorescence by phytoplankton.
13 Therefore, it is necessary to investigate the biological roles of fluorescent exudates in
14 the future.

16 *Fluorescence in natural environments and bloom formation*

18 To evaluate the contributions of phytoplankton FDOM in natural environments, we

1
2
3
4
5
6 1 extrapolated the values obtained in this study to a naturally occurring *H.*
7
8
9 2 *circularisquama* bloom. The cell density of *H. circularisquama* reached >10,000 cells
10
11
12 3 mL⁻¹ during the bloom period in Japanese coastal water (Kamiyama et al., 2001). For
13
14
15 4 this bloom, the FI of humic-like peak 1 was estimated to be 1.6×10^{-2} QSU using the
16
17
18 5 data for the FI per cell density (Table V). The possible FIs of humic-like peak 2 and
19
20
21 6 protein-like peak were also estimated in the same manner. Humic-like peak 2 and
22
23
24 7 protein-like peak would have been approximately 6.60×10^{-2} QSU and 1.72×10^{-1}
25
26
27 8 QSU, respectively. These data suggest that it is possible to detect phytoplankton-derived
28
29
30 9 FDOM in the natural aquatic environment, although these are rough estimates and high
31
32
33 10 intensity FDOM in the same region derived from other sources could mask these peaks.

34
35 11 Species-specific peaks are considered to be novel indicators that could facilitate
36
37
38 12 better understanding of the contribution of microalgal activities to FDOM production.
39
40
41 13 For example, Suksomjit et al. (2009) observed significant increase of tyrosine-like,
42
43
44 14 tryptophan-like and humic-like fluorescence, which were centred at 225 nm/305 nm,
45
46
47 15 280 nm/350 nm and 230 nm/395 nm respectively, in coastal seawater during
48
49
50 16 *Heterosigma akshiwo* and *Chaetoceros* sp. bloom. While axenic cultures of *H. akshiwo*
51
52
53 17 and *C. curvisetus* were indicated to produce the corresponding peaks such as
54
55
56 18 protein-like peak and humic-like peak 2 in our study. The absence of humic-like peak 1

1 in the natural seawater was possibly due to the relatively weak FI of the peak. Moreover,
2 it was suggested that the fluorescent peaks detected in EEM of an axenic *Micromonas*
3 culture, which were centred at 275 nm/345 nm and 348 nm/436 nm, were consistent
4 with the peak detected in natural seawater during a *Micromonas* bloom
5 (Romera-Castillo et al., 2010). We observed species-specific strong peaks such as at
6 375 nm/462 nm for *O. viridis*, which are likely to be identified in natural environment.
7 It is suggested that the DOC-specific FIs are also indicators, although we should note
8 that photo-degradability of FDOM and DOC were considered to vary (Mostofa et al.,
9 2007). We should also note that heterotrophic bacteria were known to alter the FDOM
10 property. For instance, they consume peak M to produce peak C (Romera-Castillo et al.,
11 2011). The three-dimensional fluorescence method does not require any special
12 techniques during the pre-treatment procedure; therefore, it is suitable for monitoring. It
13 suggests that early detection of noxious algal blooms is possible using the EEM
14 technique. To test this possibility, we need to monitor time-series of fluorescent EEMs
15 in seawater during the course of a bloom event and compare the optical characteristics
16 of seawater samples and those of the axenic culture filtrates of the noxious
17 phytoplankton species. In the present study, we investigated the fluorescence properties
18 of DOM exuded by eight species of axenic phytoplankton using an artificial medium.

1 Our knowledge about FDOM production by pure cultures of marine phytoplankton is
2 still quite limited. For example, the study does not involve cyanobacterial species,
3 which often form nuisance bloom in coastal areas. Further studies are required to
4 understand the biogeochemical and ecological role of FDOM and its relative abundance
5 in the natural environments.

6 7 **Acknowledgements**

8
9 We are grateful to H. Shirasawa and T. Fuji at Kyoto University for their kind support
10 on data analysis and advice during the preparation of this manuscript. We also
11 acknowledge the insightful comments of three anonymous reviewers.

12 13 **Funding**

14
15 This work was conducted as part of the the research project of the connectivity of hills,
16 humans and oceans on the local society based on ecosystem services in the forested
17 watershed environment, FSERC, Kyoto University. This research was also partly
18 supported by a Sasakawa Scientific Research Grant from The Japan Science Society.

1

2 **References**

3

4 Biddanda, B. and Benner, R. (1997) Carbon, nitrogen, and carbohydrate fluxes during
5 the production of particulate and dissolved organic matter by marine
6 phytoplankton. *Limnol. Oceanogr.* **42**, 506--518.

7 Bjørnsen, P. K. (1988) Phytoplankton exudation of organic matter—why do healthy
8 cells do it. *Limnol. Oceanogr.* **33**, 151--154.

9 Coble, P. G. (1996) Characterization of marine and terrestrial DOM in seawater using
10 excitation emission matrix spectroscopy. *Mar. Chem.* **51**, 325--346.

11 Coble, P. G. (2007) Marine optical biogeochemistry: The chemistry of ocean color.
12 *Chem. Rev.* **107**, 402--418.

13 Coble, P. G., Del Castillo, C. E. and Avril, B. (1998) Distribution and optical properties
14 of CDOM in the Arabian Sea during the 1995 Southwest Monsoon. *Deep-Sea*
15 *Res. Part II-Top. Stud. Oceanogr.* **45**, 2195--2223.

16 Coble, P. G., Green, S. A., Blough, N. V. and Gagosian, R. B. (1990) Characterization
17 of dissolved organic-matter in the black-sea by fluorescence spectroscopy.
18 *Nature* **348**, 432--435.

- 1
2
3
4
5
6 1 Coble, P. G., Schultz, C. A. and Mopper, K. (1993) Fluorescence contouring analysis of
7
8
9 2 DOC intercalibration experiment samples - a comparison of techniques. *Mar.*
10
11
12 3 *Chem.* **41**, 173--178.
13
14
15 4 Cory, R. M. and Mcknight, D. M. (2005) Fluorescence spectroscopy reveals ubiquitous
16
17
18 5 presence of oxidized and reduced quinones in dissolved organic matter. *Environ.*
19
20
21 6 *Sci. Technol.* **39**, 8142--8149.
22
23
24 7 Findlay, S. E. G. and Sinsabaugh, R. L. (eds) (2003) Aquatic ecosystems: Interactivity
25
26
27 8 of dissolved organic matter. Academic Press, California, USA
28
29
30 9 Findlay, S. E. G., Sinsabaugh, R. L., Sobczak, W. V. and Hoostal, M. (2003) Metabolic
31
32
33 10 and structural response of hyporheic microbial communities to variations in
34
35
36 11 supply of dissolved organic matter. *Limnol. Oceanogr.* **48**, 1608--1617.
37
38
39 12 Goldman, J. C., Hansell, D. A. and Dennett, M. R. (1992) Chemical characterization of
40
41
42 13 3 large oceanic diatoms — potential impact on water column chemistry. *Mar.*
43
44
45 14 *Ecol.-Prog. Ser.* **88**, 257--270.
46
47
48 15 Hansell, D. A., Carlson, C. A., Repeta, D. J. and Schlitzer, R. (2009) Dissolved organic
49
50
51 16 matter in the ocean a controversy stimulates new insights. *Oceanography* **22**,
52
53
54 17 202--211.
55
56 18 Imai, I. (1987) Size distribution, number and biomass of bacteria in intertidal sediments

- 1 and seawater of Ohmi Bay, Japan. *Bull. Jpn. Soc. Microb. Ecol.* **2**, 1--11.
- 2 Imai, I. and Yamaguchi, M. (1994) A simple technique for establishing axenic cultures
3 of phytoflagellates. *Bull. Jpn. Soc. Microb. Ecol.* **9**, 15--17.
- 4 Imai, I., Hatano, M. and Naito, K. (2004) Development of a chemically defined artificial
5 medium for marine red tide-causing raphidophycean flagellates. *Plankton Biol.*
6 *Ecol.* **51**, 95--102.
- 7 Jaffé, R., McKnight, D., Maie, N., Cory, R., McDowell, W. H. and Campbell, J. L.
8 (2008) Spatial and temporal variations in DOM composition in ecosystems: The
9 importance of long-term monitoring of optical properties. *J. Geophys.*
10 *Res.-Biogeo.* **113**.
- 11 Kamiyama, T., Takayama, H., Nishii, Y. and Uchida, T. (2001) Grazing impact of the
12 field ciliate assemblage on a bloom of the toxic dinoflagellate *Heterocapsa*
13 *circularisquama*. *Plankton Biol. Ecol.* **48**, 10--18.
- 14 Laglera, L. M., Battaglia, G. and van den Berg, C. M. G. (2007) Determination of
15 humic substances in natural waters by cathodic stripping voltammetry of their
16 complexes with iron. *Analytica Chimica Acta* **599**, 58--66.
- 17 Laglera, L. M. and van den Berg, C. M. G. (2009) Evidence for geochemical control of
18 iron by humic substances in seawater. *Limnol. Oceanogr.* **54**, 610--619.

- 1
2
3
4
5
6 1 Lancelot, C. and Billen, G. (1985) Carbon- nitrogen relationships in nutrient
7
8
9 2 metabolism of coastal marine ecosystems. *Adv. Aquat. Microbiol.* **3**, 263--321.
10
11
12 3 Maie, N., Boyer, J. N., Yang, C. Y. and Jaffé, R. (2006) Spatial, geomorphological, and
13
14 4 seasonal variability of CDOM in estuaries of the Florida Coastal Everglades.
15
16
17 5 *Hydrobiologia* **569**, 135--150.
18
19
20 6 Maie, N., Scully, N. M., Pisani, O. and Jaffé, R. (2007) Composition of a protein-like
21
22 7 fluorophore of dissolved organic matter in coastal wetland and estuarine
23
24 8 ecosystems. *Water Res.* **41**, 563--570.
25
26
27 9 Menden-Deuer, S. and Lessard, E. J. (2000) Carbon to volume relationships for
28
29 10 dinoflagellates, diatoms, and other protist plankton. *Limnol. Oceanogr.* **45**,
30
31 11 569--579.
32
33
34 12 Mostofa, K.M.G., Yoshioka, T., Konohira, E. and Tanoue, E. (2007) Photodegradation
35
36 13 of fluorescent dissolved organic matter in river waters. *Geochem. J.* **41**,
37
38 14 323--331.
39
40
41 15 Mykkestad, S., Holmhansen, O., Varum, K. M. and Volcani, B. E. (1989) Rate of
42
43 16 release of extracellular amino—acids and carbohydrates from the marine diatom
44
45
46 17 *Chaetoceros affinis*. *J. Plankton Res.* **11**, 763--773.
47
48
49 18 Mykkestad, S. M. (1995) Release of extracellular products by phytoplankton with
50
51
52
53
54
55
56
57
58
59
60

- 1 special emphasis on polysaccharides. *Sci. Total Environ.* **165**, 155--164.
- 2 Nagai, S., Imai, I. and Manabe, T. (1998) A simple and quick technique for establishing
3 axenic cultures of the centric diatom *Coscinodiscus wailesii* Gran. *J. Plankton*
4 *Res.* **20**, 1417--1420.
- 5 Naito, K., Imai, I. and Nakahara, H. (2008) Complexation of iron by microbial
6 siderophores and effects of iron chelates on the growth of marine microalgae
7 causing red tides. *Phycol. Res.* **56**, 58--67.
- 8 Naito, K., Suzuki, M., Matsui, M. and Imai, I. (2004). Secretion of iron-complexing
9 ligands from *Closterium aciculare* (Charophyceae, Chlorophyta) under
10 iron-deficient conditions. *Phycologia* **43**, 632--634.
- 11 Naito, K., Suzuki, M., Mito, S., Hasegawa, H., Imai, I., Sohrin, Y. and Matsui, M.
12 (2001) The pursuit of siderophore secreted by marine phytoplankton
13 *Rhodomonas ovalis*. *Anal. Sci. S.* **17**, i817.
- 14 Nielsen, T., Björn, L. O. and Ekelund, N. G. A. (1995) Impact of natural and artificial
15 UVB radiation on motility and growth rate of marine dinoflagellates. *J.*
16 *Photochem. Photobiol. B: Biol.* **27**, 73--79.
- 17 Nielsen, T. and Ekelund, N. G. A. (1995) Influence of solar ultraviolet radiation on
18 photosynthesis and motility of marine phytoplankton. *FEMS Microbiol. Ecol.* **18**,

- 1 281--288.
- 2 Nieto-Cid, M., Alvarez-Salgado, X. A. and Perez, F. F. (2006) Microbial and
3 photochemical reactivity of fluorescent dissolved organic matter in a coastal
4 upwelling system. *Limnol. Oceanogr.* **51**, 1391--1400.
- 5 Omura, T., Iwataki, M., Borja, V. M., Takayama, H. and Fukuyo, Y. (eds) (2012)
6 Marine phytoplankton of the Western Pacific. Kouseisha Kouseikaku, Tokyo,
7 Japan
- 8 Para, J., Coble, P. G., Charriere, B., Tedetti, M., Fontana, C. and Sempere, R. (2010)
9 Fluorescence and absorption properties of chromophoric dissolved organic
10 matter (CDOM) in coastal surface waters of the northwestern Mediterranean Sea,
11 influence of the Rhone River. *Biogeosciences* **7**, 4083--4103.
- 12 Romera-Castillo, C., Sarmiento, H., Alvarez-Salgado, X. A., Gasol, J. M. and Marrase,
13 C. (2010) Production of chromophoric dissolved organic matter by marine
14 phytoplankton. *Limnol. Oceanogr.* **55**, 446--454.
- 15 Romera-Castillo, C., Sarmiento, H., Alvarez-Salgado, X. A., Gasol, J. M. and Marrase,
16 C. (2011) Net production and consumption of fluorescent colored dissolved
17 organic matter by natural bacterial assemblages growing on marine
18 phytoplankton exudates. *Appl. Environ. Microbiol.* **77**, 7490--7498.

- 1 Sharp, J. H. (1977) Excretion of organic—matter by marine—phytoplankton - healthy
2 cells do it. *Limnol. Oceanogr.* **22**, 381--399.
- 3 Shiller, A. M., Duan, S. W., van Erp, P. and Bianchi, T. S. (2006) Photo-oxidation of
4 dissolved organic matter in river water and its effect on trace element speciation.
5 *Limnol. Oceanogr.* **51**, 1716--1728.
- 6 Shimotori, K., Omori, Y. and Hama, T. (2009) Bacterial production of marine
7 humic-like fluorescent dissolved organic matter and its biogeochemical
8 importance. *Aquat. Microb. Ecol.* **58**, 55--66.
- 9 Stedmon, C. A. and Markager, S. (2005) Resolving the variability in dissolved organic
10 matter fluorescence in a temperate estuary and its catchment using PARAFAC
11 analysis. *Limnol. Oceanogr.* **50**, 686--697.
- 12 Stedmon, C. A., Markager, S. and Bro, R. (2003) Tracing dissolved organic matter in
13 aquatic environments using a new approach to fluorescence spectroscopy. *Mar.*
14 *Chem.* **82**, 239--254.
- 15 Suksomjit, M., Nagao, S., Ichimi, K., Yamada, T., Tada, K. (2009) Variation of
16 dissolved organic matter and fluorescence characteristics before, during and
17 after phytoplankton bloom. *J. Oceanogr.* **65**, 835--846.
- 18 Sulzberger, B. and Durisch-Kaiser, E. (2009) Chemical characterization of dissolved

- 1 organic matter (DOM): A prerequisite for understanding UV-induced changes of
2 DOM absorption properties and bioavailability. *Aquat. Sci.* **71**, 104--126.
- 3 Sun, J. and Liu, D. Y. (2003) Geometric models for calculating cell biovolume and
4 surface area for phytoplankton. *J. Plankton Res.* **25**, 1331--1346.
- 5 Traganza, E. D. (1969) Fluorescence excitation and emission spectra of dissolved
6 organic matter in sea water. *Bull. Mar. Sci.* **19**, 897--904.
- 7 van den Berg, C. M. G., Buckley, P. J. M., Huang, Z. Q. and Nimmo, M. (1986) An
8 electrochemical study of the speciation of copper, zinc and iron in two estuaries
9 in England. *Est. Coast. Shelf Sci.* **22**, 479--486.
- 10 Vraspir, J. M., and Butler, A. (2009) Chemistry of marine ligands and siderophores.
11 *Annu. Rev. Mar. Sci.* **1**, 43--63.
- 12 Yamashita, Y., Jaffé, R., Maie, N. and Tanoue, E. (2008) Assessing the dynamics of
13 dissolved organic matter (DOM) in coastal environments by excitation emission
14 matrix fluorescence and parallel factor analysis (EEM-PARAFAC). *Limnol.*
15 *Oceanogr.* **53**, 1900--1908.
- 16 Yamashita, Y., Panton, A., Mahaffey, C. and Jaffé, R. (2011) Assessing the spatial and
17 temporal variability of dissolved organic matter in Liverpool Bay using
18 excitation-emission matrix fluorescence and parallel factor analysis. *Ocean Dyn.*

- 1
2
3
4
5
6 1 **61**, 569--579.
7
8
9 2 Yamashita, Y. and Tanoue, E. (2008) Production of bio-refractory fluorescent dissolved
10
11
12 3 organic matter in the ocean interior. *Nat. Geosci.* **1**, 579--582.
13
14
15 4 Yamashita, Y. and Tanoue, E. (2003) Chemical characterization of protein-like
16
17
18 5 fluorophores in DOM in relation to aromatic amino acids. *Mar. Chem.* **82**,
19
20
21 6 255--271.
22
23
24 7 Yoshioka, T., Mostofa, K. M. G., Konohira, E., Tanoue, E., Hayakawa, K., Takahashi,
25
26
27 8 M., Ueda, S., Katsuyama, M., Khodzher, T., Bashenkhaeva, N., Korovyakova, I.,
28
29
30 9 Sorokovikova, L. and Gorbunova, L. (2007) Distribution and characteristics of
31
32
33 10 molecular size fractions of freshwater-dissolved organic matter in watershed
34
35
36 11 environments: its implication to degradation. *Limnology* **8**, 29--44.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6 **1 Table and Figure legends**
7
8

9 2

10
11
12 3 Table I. Coastal phytoplankton species examined.
13
14
15 4

16
17
18 5 Table II. Incubation time, growth rates (μ) during the exponential growth phase, the
19
20
21 6 final cell abundance (C) and biomass (B).
22
23
24 7

25
26 8 Table III. Net increase in the dissolved organic carbon (DOC) concentration and the
27
28
29 9 apparent percentage of net photosynthetic extracellular release (APER) for each
30
31
32 10 plankton culture. APER was calculated as follows: $\text{DOC}/(\text{DOC} + \text{Biomass}) \times$
33
34
35 11 100, where DOC and Biomass are the net increases in DOC and biomass during
36
37
38 12 the incubation period, respectively. The data shown are average values \pm
39
40
41 13 standard error (S.E.) (n = 3).
42
43
44 14

45
46 15 Table IV. Peak position, averaged fluorescence intensity (FI) standardised to quinine
47
48
49 16 sulphate unit (QSU) at each peak position and DOC-specific FI of FDOM (QSU
50
51
52 17 L mg C^{-1}) secreted by each species. Mean value \pm S.E. (n = 3).
53
54
55 18

1
2
3
4
5
6 1 Table V. DOC ($\mu\text{g C } 1000 \text{ cells}^{-1}$) and intensity of each fluorescent component (QSU
7
8
9 2 mL 1000 cells^{-1}), which were standardised on the basis of the cell density at the
10
11 3 end of the incubation, secreted by the phytoplankton cultures. Mean value \pm S.E.
12
13
14 4 ($n = 3$).
15
16

17
18 5 Fig. 1. Average excitation emission matrices of fluorescent dissolved organic matter
19
20 6 (FDOM) secreted by each plankton culture.
21
22
23
24 7
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table I. Coastal phytoplankton species examined

Bacillariophyceae

Ditylum brightwellii (T. West) Grunow, 1885

Chaetoceros curvisetus P.T. Cleve, 1889

Dinoflagellata

Heterocapsa circularisquama Horiguchi, 1995

Alexandrium catenella (Whedon & Kofoed) E. Balech, 1985

Raphidophyceae

Heterosigma akashiwo (Y. Hada) Y. Hada ex Y. Hara & M. Chihara, 1967

Chlorophyceae

Oltmannsiellopsis viridis (P.E. Hargraves & R.L. Steele) M. Chihara & I. Inouye in Chihara, Inouye & Takahata, 1986

Cryptomonadea

Rhodomonas ovalis Nygaard

Prymnesiophyceae

Pleurochrysis roscoffensis (P. Dangeard) J. Fresnel & C. Billard

Table II. Incubation time, growth rates (μ) during the exponential growth phase, the final cell abundance (C) and biomass (B)

Phytoplakton culture	Time (d)	μ (d ⁻¹)	C (cells ml ⁻¹)	B (mg C L ⁻¹)
<i>Ditylum brightwellii</i>	12	0.76 ± 0.05	11 ± 1 × 10 ³	18.8 ± 1.0
<i>Chaetoceros curvisetus</i>	6	0.97 ± 0.07	199 ± 37 × 10 ³	5.9 ± 1.1
<i>Heterocapsa circularisquama</i>	12	0.84 ± 0.01	93 ± 22 × 10 ³	33.1 ± 7.9
<i>Alexandrium catenella</i>	27	0.48 ± 0.01	22 ± 3 × 10 ³	27.6 ± 4.1
<i>Heterosigma akashiwo</i>	14	0.35 ± 0.03	121 ± 59 × 10 ³	28.7 ± 13.8
<i>Oltmannsiellopsis viridis</i>	12	0.55 ± 0.06	247 ± 95 × 10 ³	18.2 ± 7.0
<i>Rhodomonas ovalis</i>	12	0.40 ± 0.01	253 ± 45 × 10 ³	3.5 ± 0.6
<i>Pleurochrysis roscoffensis</i>	12	0.45 ± 0.03	136 ± 18 × 10 ³	11.8 ± 1.6

Table III. Net increase in the dissolved organic carbon (DOC) concentration and the apparent percentage of net photosynthetic extracellular release (APER) for each plankton culture. APER was calculated as follows: $\text{DOC}/(\text{DOC} + \text{Biomass}) \times 100$, where DOC and Biomass are the net increases in DOC and biomass during the incubation period, respectively. The data shown are average values \pm standard error (S.E.) (n = 3).

Phytoplakton culture	DOC (mgC L ⁻¹)	APER (%)
<i>Ditylum brightwellii</i>	25.1 \pm 4.9	55.9 \pm 4.5
<i>Chaetoceros curvisetus</i>	26.6 \pm 11.3	71.5 \pm 16.0
<i>Heterocapsa circularisquama</i>	38.6 \pm 14.8	51.0 \pm 14.6
<i>Alexandrium catenella</i>	24.1 \pm 5.5	46.1 \pm 9.4
<i>Heterosigma akashiwo</i>	19.3 \pm 5.1	52.4 \pm 19.6
<i>Oltmannsiellopsis viridis</i>	27.8 \pm 4.5	63.7 \pm 7.7
<i>Rhodomonas ovalis</i>	26.4 \pm 16.6	72.1 \pm 18.6
<i>Pleurochrysis roscoffensis</i>	49.9 \pm 7.5	80.2 \pm 3.7

Table IV. Peak position, averaged fluorescence intensity (FI) standardised to quinine sulphate unit (QSU) at each peak position and DOC-specific FI of FDOM (QSU L mg C⁻¹) secreted by each species. Mean value ± S.E. (n = 3).

Phytoplakton culture	Humic-like peak 1				Humic-like peak 2				Protein-like peak			
	Ex (nm)	Em (nm)	FI (QSU)	FI/DOC (QSU L mg C ⁻¹)	Ex (nm)	Em (nm)	FI (QSU)	FI/DOC (QSU L mg C ⁻¹)	Ex (nm)	Em (nm)	FI (QSU)	FI/DOC (QSU L mg C ⁻¹)
<i>Ditylum brightwellii</i>	355	454	0.26 ± 0.02	0.010 ± 0.002	250	464	0.66 ± 0.07	0.026 ± 0.006	255	316	0.58 ± 0.03	0.023 ± 0.005
<i>Chaetoceros curvisetus</i>					250	455	0.54 ± 0.04	0.020 ± 0.009	280	349	0.69 ± 0.04	0.026 ± 0.011
<i>Heterocapsa circularisquama</i>	340	461	0.15 ± 0.01	0.004 ± 0.002	255	441	0.61 ± 0.04	0.016 ± 0.006	280	357	1.60 ± 0.10	0.041 ± 0.016
<i>Alexandrium catenella</i>					250	471	0.55 ± 0.15	0.023 ± 0.008	280	352	0.75 ± 0.07	0.031 ± 0.008
<i>Heterosigma akashiwo</i>	375	473	0.31 ± 0.10	0.016 ± 0.007	250	455	0.90 ± 0.21	0.046 ± 0.016	280	335	0.44 ± 0.19	0.023 ± 0.012
<i>Oltmannsiellopsis viridis</i>	375	462	0.86 ± 0.24	0.031 ± 0.010	250	455	1.20 ± 0.16	0.043 ± 0.009	280	337	2.22 ± 0.36	0.080 ± 0.018
<i>Rhodomonas ovalis</i>					250	446	0.80 ± 0.13	0.030 ± 0.020	280	324	1.24 ± 0.61	0.047 ± 0.038
<i>Pleurochrysis roscoffensis</i>					250	455	0.56 ± 0.11	0.011 ± 0.003	250	350	0.57 ± 0.03	0.011 ± 0.002

Table V. DOC ($\mu\text{g C } 1000 \text{ cells}^{-1}$) and intensity of each fluorescent component (QSU mL 1000 cells^{-1}), which were standardised on the basis of the cell density at the end of the incubation, secreted by the phytoplankton cultures. Mean value \pm S.E. (n = 3).

Phytoplakton culture	Time (d)	DOC ($\mu\text{g C } 1000 \text{ cells}^{-1}$)	Humic-like peak 1 (QSU mL 1000 cells^{-1})	Humic-like peak 2 (QSU mL 1000 cells^{-1})	Protein-like peak (QSU mL 1000 cells^{-1})
<i>Ditylum brightwellii</i>	12	2.22 \pm 0.45	0.023 \pm 0.0024	0.058 \pm 0.007	0.051 \pm 0.004
<i>Chaetoceros curvisetus</i>	6	0.13 \pm 0.06	Not Detected	0.003 \pm 0.001	0.003 \pm 0.001
<i>Heterocapsa circularisquama</i>	12	0.42 \pm 0.19	0.0016 \pm 0.0004	0.007 \pm 0.002	0.017 \pm 0.004
<i>Alexandrium catenella</i>	27	1.09 \pm 0.29	Not Detected	0.025 \pm 0.008	0.034 \pm 0.006
<i>Heterosigma akashiwo</i>	14	0.16 \pm 0.09	0.0026 \pm 0.0015	0.007 \pm 0.004	0.004 \pm 0.002
<i>Oltmannsiellopsis viridis</i>	12	0.11 \pm 0.05	0.0035 \pm 0.0016	0.005 \pm 0.002	0.009 \pm 0.004
<i>Rhodomonas ovalis</i>	12	0.10 \pm 0.07	Not Detected	0.003 \pm 0.001	0.005 \pm 0.003
<i>Pleurochrysis roscoffensis</i>	12	0.37 \pm 0.07	Not Detected	0.004 \pm 0.001	0.004 \pm 0.001




Fig. 1. Average excitation emission matrices of fluorescent dissolved organic matter (FDOM) secreted by each plankton culture.