

Title	Functional Movement Screen for Predicting Running Injuries in 18- to 24-Year-Old Competitive Male Runners.
Author(s)	Hotta, Takayuki; Nishiguchi, Shu; Fukutani, Naoto; Tashiro, Yuto; Adachi, Daiki; Morino, Saori; Shirooka, Hidehiko; Nozaki, Yuma; Hirata, Hinako; Yamaguchi, Moe; Aoyama, Tomoki
Citation	Journal of strength and conditioning research (2015), 29(10): 2808-2815
Issue Date	2015-10
URL	<a href="http://hdl.handle.net/2433/202830">http://hdl.handle.net/2433/202830</a>
Right	© 2015 by the National Strength & Conditioning Association; This is a non-final version of an article published in final form in [Journal of Strength & Conditioning Research: October 2015, Volume 29, Issue 10, p 2808–2815, doi: 10.1519/JSC.0000000000000962]; The full-text file will be made open to the public on 1 October 2016 in accordance with publisher's 'Terms and Conditions for Self-Archiving!'; この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。 This is not the published version. Please cite only the published version.
Type	Journal Article
Textversion	author

- 1 Functional Movement Screen for Predicting Running Injuries in 18–24 Year-Old Competitive
- 2 Male Runners

**3 ABSTRACT**

4           The purpose of this study was to investigate whether the functional movement screen  
5 (FMS) could predict running injuries in competitive runners. Eighty-four competitive male  
6 runners (average age =  $20.0 \pm 1.1$  years) participated. Each subject performed the FMS,  
7 which consisted of 7 movement tests (each score range: 0–3, total score range: 0–21), during  
8 the pre-season. The incidence of running injuries (time lost due to injury  $\leq 4$  weeks) was  
9 investigated through a follow-up survey during the 6-month season. Mann–Whitney U tests  
10 were used to investigate which movement tests were significantly associated with running  
11 injuries. The receiver-operator characteristic (ROC) analysis was used to determine the  
12 cut-off. The mean FMS composite score was  $14.1 \pm 2.3$ . The ROC analysis determined the  
13 cut-off at 14/15 (sensitivity = 0.73, specificity = 0.54), suggesting that the composite score  
14 had a low predictability for running injuries. However, the total score (0–6) from the deep  
15 squat (DS) and active straight leg raise (ASLR) tests (DS & ASLR), which were significant  
16 with the U test, had relatively high predictability at the cut-off of 3/4 (sensitivity = 0.73,  
17 specificity = 0.74). Furthermore, the multivariate logistic regression analysis revealed that the  
18 DS & ASLR scores of  $\leq 3$  significantly influenced the incidence of running injuries after  
19 adjusting for subjects' characteristics (OR = 9.7, 95%CI [2.1 to 44.4]). Thus, the current  
20 study identified the DS & ASLR score as a more effective method than the composite score  
21 to screen the risk of running injuries in competitive male runners.

22 **KEY WORDS:** Distance runner, Screening, Dynamic assessment, Risk factor

## 23 INTRODUCTION

24 The functional movement screen (FMS) is a screening tool for injury risk that  
25 assesses the movement patterns of individuals, and which can evaluate mobility and stability  
26 comprehensively. The FMS consists of 7 component tests, each scored based on the  
27 movement patterns within the kinetic chain, asymmetries between the sides, and  
28 compensatory movements. The validity of the FMS as a predictor of injury risk has been  
29 confirmed in several studies (6,11,12,18). The first, by Kiesel et al. (11), examined the  
30 relationship between the FMS and serious injury in professional football players. They  
31 revealed that professional football players with an FMS score of  $\leq 14$  were at a greater risk of  
32 serious injury than those with higher scores (11). Other studies reported similar findings in  
33 other groups, such as officer candidates (12,18) and female collegiate athletes (6).

34 Recently, two studies investigated normative values of FMS scores for runners.  
35 Loudon et al (13) reported the normative value for running athletes and Agresta et al. (1)  
36 reported it for healthy runners (the mean FMS composite scores were  $13.1 \pm 1.8$  and  $15.4 \pm$   
37  $2.4$ , respectively). Additionally, Agresta et al. investigated the association between FMS  
38 scores and injury history. However, no prospective cohort studies have investigated the  
39 association between the FMS and running injuries. Running injuries are a serious problem for  
40 most runners, especially for competitive runners (9). Unfortunately, some runners are forced  
41 to retire from running due to serious running injuries. Previous studies reported some risk  
42 factors for running injuries, such as inadequate flexibility (25), muscle weakness and  
43 imbalance (17), and deficits in neuromuscular coordination (20). Cook (7,8) stated that these  
44 factors also caused poor movement patterns, which were reflected in the lower score of the  
45 FMS. Thus, runners with low FMS scores might have certain risk factors for running injury  
46 and become more prone to injury. In addition, although Parchmann and McBride (19)  
47 reported that the FMS was not significantly associated with sprinting, Chapman (5) revealed

48 that a high FMS score had a positive effect on performance in elite track and field athletes in  
49 the long view. Because athletes with a higher FMS score rarely suffered from injury, they  
50 could practice continuously and improve their performance. Therefore, we hypothesized that  
51 the FMS could predict running injuries.

52 The receiver-operator characteristic (ROC) curve is a plot of the sensitivity versus 1  
53 – specificity of a screening test; this analysis is useful in determining the cut-off where the  
54 sensitivity and specificity are maximized. In previous studies, the ROC curve was used to  
55 determine the validity of the FMS as a predictor of injury risk (3,11,18). In addition, a cut-off  
56 value allows determining more easily whether a runner has a potential injury risk simply  
57 based on the FMS scores. Therefore, the aim of the current study was to determine the cut-off  
58 value and to investigate if the FMS score during pre-season could be used to predict running  
59 injuries in young competitive runners during season.

60

## 61 **METHODS**

### 62 **Experimental Approach to the Problem**

63 This study, using a prospective cohort design, investigated whether pre-season FMS  
64 scores could predict serious running injuries during the season in 18–24-year-old competitive  
65 male runners. Figure 1 illustrates the process of this study in the form of a flow chart. The  
66 subjects performed the FMS at their college during pre-season, February 2014. To minimize  
67 the influence of fatigue on the performance, the FMS tests were conducted during the  
68 daytime on the day following a non-training day according to each team's schedule. No  
69 warm-up was included. The testing days added up to 7 days total. After the FMS test,  
70 follow-up surveys were distributed to the subjects to investigate the incidence of running  
71 injuries during the 6-month season. The follow-up surveys were conducted twice at the end  
72 of May and August 2014 to reduce a recall bias. Statistical analyses were conducted using the

73 data of the returned surveys. The ROC analysis determined the cut-off, and the logistic  
74 regression analysis determined if the FMS could be used for the prediction of running  
75 injuries.

76

## 77 **Subjects**

78 A total of 84 competitive male runners volunteered to participate in the current study  
79 (mean age =  $20.0 \pm 1.1$  years, age range = 18–24 years, height =  $171.6 \text{ cm} \pm 4.5$ , weight =  
80  $57.5 \text{ kg} \pm 4.3$ ). For inclusion, subjects had to be competitive male runners belonging to  
81 collegiate track and field teams, who were injury-free at the time of the FMS test in  
82 pre-season, whose events were middle- or long-distance, and whose running experience  
83 exceeded 1 year. The purpose and methods of this study were explained to the subjects in  
84 detail in a verbal statement, and written informed consent was obtained from the subjects.  
85 The current study did not include athletes under the age of 18 years, thus parental or guardian  
86 consent was not needed. This study was approved by the Institutional Review Board of Kyoto  
87 University (Approval No. E2023).

88

## 89 **Procedures**

90 Before the study, the physical therapists collecting data in the current study were  
91 instructed on the FMS evaluation method by an FMS specialist. The FMS scoring criteria  
92 were used as described by Cook et al. (7,8), and they discussed standardization of the testing.

93 On testing day, all subjects were questioned about their characteristics, such as age,  
94 height, weight, running experience, training sessions per week, weekly mileage, personal best  
95 time in their primary event in 2013, and injury history by questionnaire. To allow comparison  
96 between different events, performances were normalized to a percentage of collegiate  
97 Japanese record performances (as of March 31, 2013) (5). To assess injury history, we asked

98 the following question: “Have you ever suffered from musculoskeletal injury that was so  
99 severe that it required medical attention?” (6). Subsequently, all subjects were briefed on the  
100 FMS and given a demonstration of the movements. After the demonstration, all subjects  
101 performed the FMS, which consisted of 7 movement tests to comprehensively assess mobility,  
102 stability, and coordination. The 7 tests were the deep squat (DS), hurdle step (HS), in-line  
103 lunge (ILL), shoulder mobility (SM), active straight leg raise (ASLR), trunk stability push-up  
104 (TSPU), and rotary stability (RS) tests. All tests were scored using standardized scoring  
105 criteria (7,8). Each movement test was scored on a 4-point scale (0–3), and the maximal FMS  
106 score that could be achieved was 21. A score of 3 was awarded for perfect form, a score of 2  
107 was given for completing the test with compensations, a score of 1 was awarded for not  
108 completing the test accurately, and a score of 0 was given if the subjects felt any pain during  
109 the test. Each test was performed 3 times, and the highest score was used. Of the 7 tests that  
110 comprise the FMS, 5 tests (HS, ILL, SM, ASLR, and RS) were performed and scored  
111 separately for the right and left side of the body. For these bilaterally assessed tests, the lower  
112 score was used.

113         After the FMS test, follow-up surveys were distributed to all subjects through each  
114 team’s manager to investigate the incidence of running injuries during the 6-month season. If  
115 information was missing in the questionnaires, we asked the subjects to answer the omitted  
116 questions by contacting them through the team’s managers. For the current study, the  
117 definition of running injury was a musculoskeletal injury that met the following criteria: (1)  
118 the injury occurred as a result of participating in a practice or race in track and field (trauma  
119 injuries, such as sprains, were excluded), and (2) the injury was sufficiently severe to prevent  
120 participation for at least 4 weeks; this definition was based on that used in previous studies  
121 (11,18).

122

## 123 **Reliability**

124           Similar to a previous study (13), interrater reliability was assessed in a subgroup of  
125 10 subjects by 2 physical therapists. Interrater reliability was calculated for the FMS  
126 composite score using the intraclass correlation coefficient (ICC, model 2, 1). On the basis of  
127 the reliability coefficients, the standard error of measurement ( $SEM = SD \times \sqrt{1-ICC}$ ), the  
128 minimum difference to be considered real ( $MD = SEM \times 1.96 \times \sqrt{2}$ ), and the standard error  
129 of prediction ( $SEP = SD\sqrt{1-ICC^2}$ ) were calculated (24). The Bland-Altman analysis was  
130 performed to determine whether systematic error was present. The weighed kappa statistic  
131 was used to establish the interrater reliability for each movement test of the FMS.

132

## 133 **Statistical Analyses**

134           We divided the subjects into 2 groups with and without running injuries according to  
135 the follow-up survey. Comparisons between the 2 groups were made using Student's t-tests  
136 (for parametric continuous variables), Mann-Whitney U tests (for non-parametric continuous  
137 variables), or chi-squared tests (for categorical variables). The short version of the FMS was  
138 calculated from the movement tests that were significant according to the U tests. The ROC  
139 curve was calculated by pairing the FMS score with running injury to determine the cut-off  
140 on the FMS that maximized sensitivity and specificity according to previous studies  
141 (3,6,11,18). In this context, the FMS can be thought of as a screening test that determines if a  
142 runner is at risk for a running injury. An ROC curve is a plot of the sensitivity (true-positive)  
143 versus  $1 - \text{specificity}$  (false-positive) of a screening test. The area under the curve (AUC) was  
144 calculated based on the ROC curve. The optimal cut-off was determined based on the Youden  
145 index, which consists of the following formula:  $\text{Youden index} = (\text{sensitivity} + \text{specificity}) - 1$   
146 (2). Maximizing this index allows finding the optimal cut-off value. Once the cut-off value  
147 was identified, a  $2 \times 2$  contingency table was created dichotomizing those with and without



148 injury, and those above and below the cut-off on the FMS. To determine whether a runner,  
149 whose FMS score was below the cut-off, had potential injury risk during season, the  
150 multivariate logistic analysis was adjusted for each subject's characteristics including age,  
151 height, weight, running experience, training sessions per week, weekly mileage, performance  
152 level, and injury history. A value of  $p < .05$  was considered to be statistically significant for  
153 all analyses. All data were analyzed by using the Statistical Package for the Social Sciences  
154 version 20.0 (SPSS, Inc., Chicago, IL).

155

## 156 **RESULTS**

157 In pre-season, 101 runners from 7 teams participated in the FMS. Of the 84 returned  
158 the follow-up surveys (response rate was 83.2%).

159

### 160 **Reliability**

161 Interrater reliability for the FMS composite score is shown in Table 1. ICC (2, 1) was  
162 0.98 (95% confidence interval, CI [0.93, 1.00]), demonstrating excellent reliability, and the  
163 Bland-Altman analysis revealed that there was no systematic error present (both fixed bias  
164 and proportional bias). Interrater reliability (weighted kappa) for each component movement  
165 test is presented in Table 2 and shows that the majority of the FMS tests demonstrated a  
166 substantial to excellent agreement. These results were in accordance with previous studies  
167 (10,16,22) and confirmed the reliability of the procedure in the current study.

168

### 169 **FMS Score Distribution**

170 The mean FMS composite score was  $14.2 \pm 2.3$  with a range of 7–18. Of the 84  
171 subjects, 43 (51.2%) scored  $\leq 14$  on the FMS composite score, indicating that they had a high  
172 injury risk according to Kiesel et al. (11). Among all the subjects, 4 reported pain in the DS

173 and TSPU tests, 3 reported pain in the SM test, 2 reported pain in the ILL test, and 1 reported  
174 pain in the HS and RS tests, which resulted in a score of 0 for these tests.

175 The distribution of scores for each component movement test is presented in Figure  
176 2. The SM test was the movement with the highest frequency of a score of 3 (65.5%).  
177 Conversely, the RS was the movement with the highest frequency of a score of 1 (34.5%); no  
178 subject achieved a score of 3 on this test. The DS, HS, ILL, and ASLR tests had the highest  
179 frequency of a score of 2 on each test.

180

### 181 **FMS Score and Injuries**

182 Among the 84 subjects, 15 (17.9%) experienced running injuries during the season.  
183 The comparisons between groups with and without running injuries are presented in Table 3.  
184 The mean FMS composite scores were  $13.3 \pm 2.7$  and  $14.4 \pm 2.2$  for subjects with and  
185 without any injury, respectively. Although, there was a trend for the injury group to have a  
186 lower score, this difference was not significant ( $p = .07$ ). Of the 7 tests, the scores on the DS  
187 and ASLR tests were significant with the U test. Using the composite score of the 2 tests, we  
188 calculated a short version of the FMS, which was named “DS & ASLR” (score range: 0–6).  
189 Figure 3 shows the significant difference in the DS & ASLR score between the injured and  
190 non-injured groups, whose scores were  $2.9 \pm 1.0$  and  $4.1 \pm 1.1$ , respectively ( $p < .01$ ).

191 The ROC curves for the FMS composite and DS & ASLR scores are presented in  
192 Figure 4. The cut-off of the FMS composite score was determined to be 14/15, which was  
193 consistent with a previous study (11). However, the ROC curve had a relatively low AUC  
194 ( $AUC = 0.65$ ,  $p = .08$ ), and, at this point, the sensitivity was 0.73, and the specificity was 0.46.  
195 Subjects were dichotomized into groups with FMS composite scores  $\leq 14$  and  $\geq 15$ , which are  
196 presented in Table 4. Conversely, the ROC curve for the DS & ASLR score had a relatively  
197 high AUC ( $AUC = 0.79$ ,  $p = .01$ ), and it determined the cut-off to be 3/4 with a sensitivity of

198 0.73 and a specificity of 0.74 (Figure 4). Subjects were again dichotomized into groups with  
199 DS & ASLR scores  $\leq 3$  and  $\geq 4$ , which are presented in Table 5. Among the subjects with a  
200 score of  $\leq 3$ , 11 out of 29 had been injured during the season (injury rate: 37.9%), while  
201 among the subjects with a score of  $\geq 4$ , 4 out of 55 (injury rate: 7.3%) had been injured. The  
202 logistic regression analysis revealed similar results presented in Table 6. A score of  $\leq 14$  of the  
203 composite FMS did not significantly influence the incidence of running injuries (OR = 3.0,  
204 95%CI [0.8, 11.6],  $p = .10$ ). However, the same analysis revealed that a runner with a DS &  
205 ASLR score of  $\leq 3$  was significantly more likely to become injured even when adjusting for  
206 each subject's characteristics (OR = 9.7, 95%CI [2.1, 44.4],  $p < .01$ ).

207

## 208 **DISCUSSION**

209 The purpose of the current study was to investigate whether the FMS could predict  
210 running injuries in competitive male runners. The study revealed that the cut-off on the FMS  
211 was 14/15, which was in accordance with a previous study (11), but the composite score of  
212  $\leq 14$  had low predictability for running injuries. In contrast, the current study also revealed  
213 that a DS & ASLR score of  $\leq 3$  during pre-season was a more useful approach for predicting  
214 running injuries during season in 18–24 year-old competitive male runners. This is the first  
215 study to investigate the validity of the FMS as a predictor for running injuries and to establish  
216 the short version of the FMS (DS & ASLR) for screening running injuries.

217

### 218 **FMS Score Distribution**

219 The mean FMS composite score for the 18–24-year-old competitive male runners in  
220 the current study was  $14.1 \pm 2.3$ , which is similar to the results of college basketball  
221 volleyball, and soccer athletes in Warren et al.'s (23) and Chorba et al.'s (6) studies (mean

222 scores were  $14.3 \pm 2.5$  and  $14.3 \pm 1.7$ , respectively). On the other hand, Loudon et al. (13)  
223 reported a mean score for male running athletes of  $15.0 \pm 2.4$ , while Agresta et al. (1)  
224 reported a mean score for healthy male runners of  $13.1 \pm 1.7$ . Although their findings slightly  
225 differ from ours, the runners in the current study had a comparable average performance as  
226 other runners. Additionally, our scores were relatively lower than the mean composite scores  
227 for professional football players (11) and officer candidates (18) (mean scores were  $16.9 \pm$   
228  $3.0$  and  $16.6 \pm 1.7$ , respectively). These differences are expected to occur because distance  
229 running mainly requires cardiorespiratory endurance and does not involve as much stability  
230 and power as required by football players or candidate officers.

231         Considering each movement test of the FMS, Figure 2 shows that the subjects  
232 performed the best on the SM test, which required mobility of the shoulder and scapula and  
233 thoracic spine extension. Since runners need to swing their arms frequently during running,  
234 SM is needed to minimize the burden from arm swing. On the other hand, the subjects  
235 performed the worst in the RS test, which requires multi-plane trunk stability during a  
236 combined upper and lower extremity motion. This result was similar to results of previous  
237 studies (1,18,21); there were only a few subjects who scored 3 on the RS test. Thus, these  
238 findings suggest that the RS test may be one of the more difficult tests of the FMS.

239

#### 240 **FMS Score and Injuries**

241         The ROC analysis revealed that sensitivity and specificity were 0.73 and 0.74,  
242 respectively. Subsequently, the multivariate logistic regression analysis revealed that subjects  
243 with a score of  $\leq 3$  on the DS & ASLR were approximately 10 times more likely to have  
244 running injuries than those with a score  $\geq 4$  after adjusting for each subject's characteristics.  
245 The relatively strong predictability of running injuries according to the DS & ASLR score  
246 was attributed to the following reasons. First, the DS test by itself had a strong predictability

247 of injuries, which was in accordance with the result of Butler et al.'s study (3). The DS test  
248 assesses bilateral, symmetrical mobility, especially mobility of hips, ankles, and thoracic  
249 spine, and coordination in the close kinetic chain. Renström (20) reported that poor flexibility  
250 and deficit in neuromuscular coordination can cause running injuries. Additionally, excessive  
251 pronation and knee-in during testing, which was one of the causes that decreased the score on  
252 the DS test (7), was also reported to be a risk factor for injury (15). Second, the ASLR test  
253 was also found to be related to running injuries; it assesses active hamstring and  
254 gastric-soleus flexibility while maintaining a stable pelvis. This finding agreed with the study  
255 by Yagi et al. (25), who also reported that limited SLR ability increased the injury risk in high  
256 school runners. Additionally, Lysholm et al. (14) reported that flexibility of the hamstrings  
257 was a risk factor for injury. Consequently, deficits in the DS and ASLR tests are likely to  
258 induce asymmetric or compensatory movement patterns and thus result in running injuries.  
259 Thus, the FMS contains both helpful and less helpful movement tests for predicting injury  
260 risk in competitive male runners. The HS test assesses stepping ability, which requires  
261 mobility and stability of the legs as well as coordination. The ILL test requires mobility and  
262 stability in the split stance as well as coordination. Although these 2 tests seem to be relevant  
263 for running, they were not significantly associated with incidence of running injury because  
264 most subjects received a score of 2 (91.7% for HS, 86.9% for ILL). Due to their ceiling  
265 effects, these 2 tests were ineffective in screening injury risk. As a result, the FMS composite  
266 score had low predictability. For the SM, TSPU, and RS tests, there is no solid evidence that  
267 shoulder mobility and core-stability influence the incidence of running injuries.

268

### 269 **Limitation**

270 There were several limitations in the current study. The first is the definition of  
271 injury as a running injury (lost training time  $\geq 4$  weeks). Although the current study revealed

272 that the DS & ASLR could predict serious running injuries, it is unclear if it could  
273 successfully screen the risk of non-serious running injuries (lost training time <4 weeks). A  
274 second limitation was the mode of collecting injury data by a self-report questionnaire due to  
275 the absence of athletic trainers in all teams. As a result, relevant details, such as type of injury,  
276 were not collected. A third limitation was that the current study was carried out among 18–  
277 24-year-old competitive male runners in Japan. It is unclear whether the results can be  
278 extrapolated to other running populations such as female, older, or recreational runners.  
279 Therefore, further study is required to ensure the external validity of the DS & ASLR score  
280 for other runners.

281

## 282 **PRACTICAL APPLICATIONS**

283 First, the current study provided reliable normative data for FMS scores among 18–  
284 24 year-old competitive male runners. These data can be used as reference values for strength  
285 and conditioning by professional coaches when they need to assess the injury risk of similar  
286 groups using the FMS.

287 Additionally, the current study revealed that a score of  $\geq 4$  or  $\leq 3$  of the DS & ASLR  
288 was more useful for predicting running injuries than the FMS composite score in 18–24  
289 year-old competitive male runners. This finding is meaningful for the strength and  
290 conditioning professional who supports a similar group of athletes. First, injury risks can be  
291 screened easily by using the DS & ASLR as it only takes approximately 5 minutes. This is an  
292 advantage because time is often limited and rather spent on training. Second, it allows the  
293 strength and conditioning professional to prevent serious problems in younger runners that  
294 could result in retiring from running due to injuries. Timely prediction of injury risks allows  
295 initiating strategies for preventing injury. For example, performing hamstring and  
296 gastric-soleus stretches are effective in improving scores on the ASLR scores (8). As to the

297 DS test, the strength and conditioning professional or physical therapists should assess which  
298 deficit is limiting influence on this test before conducting corrective exercises. This is  
299 because the DS test is affected by many variables, such as the mobility of the hip, ankle,  
300 thoracic spine, and shoulder, the stability of the hip, and coordination (8). The current study  
301 suggests that, by improving scores on the DS & ASLR in pre-season, the incidence of  
302 running injuries in 18–24-year-old competitive male runners could be reduced.

303

### 304 **References**

- 305 1. Agresta, C, Slobodinsky, M, and Tucker, C. Functional Movement Screen™ - Normative  
306 Values in Healthy Distance Runners. *Int J Sports Med* 35: 1203–1207, 2014.
- 307 2. Akobeng, AK. Understanding diagnostic tests 3: Receiver operating characteristic curves.  
308 *Acta Paediatr* 96: 644–647, 2007.
- 309 3. Butler, RJ, Contreras, M, Burton, LC, Plisky, PJ, Goode, A, and Kiesel, K. Modifiable risk  
310 factors predict injuries in firefighters during training academies. *Work* 46: 11–17, 2013.
- 311 4. Butler, RJ, Plisky, PJ, Southers, C, Scoma, C, and Kiesel, KB. Biomechanical analysis of  
312 the different classifications of the Functional Movement Screen deep squat test. *Sports*  
313 *Biomech* 9: 270–279, 2010.
- 314 5. Chapman, RF, Laymon, AS, and Arnold, T. Functional movement scores and longitudinal  
315 performance outcomes in elite track and field athletes. *Int J Sports Physiol Perform* 9: 203–  
316 211, 2014.
- 317 6. Chorba, RS, Chorba, DJ, Bouillon, LE, Overmyer, CA, and Landis JA. Use of a functional  
318 movement screening tool to determine injury risk in female collegiate athletes. *N Am J Sports*  
319 *Phys Ther* 5: 47–54, 2010.
- 320 7. Cook, G, Burton, L, and Hoogenboom, B. Pre-participation screening: the use of  
321 fundamental movements as an assessment of function - part 1. *N Am J Sports Phys Ther* 1:

- 322 62–72, 2006.
- 323 8. Cook, G, Burton, L, and Hoogenboom, B. Pre-participation screening: the use of  
324 fundamental movements as an assessment of function - part 2. *N Am J Sports Phys Ther* 1:  
325 132–139, 2006.
- 326 9. Di Caprio, F, Buda, R, Mosca, M, Calabro, A, and Giannini, S. Foot and lower limb  
327 diseases in runners: assessment of risk factors. *J Sports Sci Med* 9: 587–596 2010.
- 328 10. Gulgin, H, and Hoogenboom, B. The functional movement screening (FMS): An  
329 inter-rater reliability study between raters of varied experience. *Int J Sports Phys Ther* 9: 14–  
330 20, 2014.
- 331 11. Kiesel, K, Plisky, PJ, and Voight, ML. Can serious injury in professional football be  
332 predicted by a preseason functional movement screen? *N Am J Sports Phys Ther* 2: 147–158,  
333 2007.
- 334 12. Lisman, P, O'Connor, FG, Deuster, PA, and Knapik, JJ. Functional movement screen and  
335 aerobic fitness predict injuries in military training. *Med Sci Sports Exerc* 45: 636–643, 2013.
- 336 13. Loudon, JK, Parkerson-Mitchell, AJ, Hildebrand, LD, and Teague, C. Functional  
337 movement screen scores in a group of running athletes. *J Strength Cond Res* 28: 909–913  
338 2014.
- 339 14. Lysholm, J, and Wiklander, J. Injuries in runners. *Am J Sports Med* 15: 168–171, 1987.
- 340 15. Messier, SP, and Pittala, KA. Etiologic factors associated with selected running injuries.  
341 *Med Sci Sports Exerc* 20: 501–505, 1988.
- 342 16. Minick, KI, Kiesel, KB, Burton, L, Taylor, A, Plisky, P, and Butler, RJ. Interrater  
343 reliability of the functional movement screen. *J Strength Cond Res* 24: 479–486, 2010.
- 344 17. Niemuth, PE, Johnson, RJ, Myers, MJ, and Thieman, TJ. Hip muscle weakness and  
345 overuse injuries in recreational runners. *Clin J Sport Med* 15: 14–21, 2005.
- 346 18. O'Connor, FG, Deuster, PA, Davis, J, Pappas, CG, Knapik, JJ. Functional movement




- 347 screening: predicting injuries in officer candidates. *Med Sci Sports Exerc* 43: 2224–2230,  
348 2011.
- 349 19. Parchmann, CJ, and McBride, JM. Relationship between functional movement screen and  
350 athletic performance. *J Strength Cond Res* 25: 3378–3384, 2011.
- 351 20. Renström, AF. Mechanism, diagnosis, and treatment of running injuries. *Instr Course Lect*  
352 42: 225–234, 1993.
- 353 21. Schneiders, AG, Davidsson, A, Horman, E, and Sullivan, SJ. Functional movement screen  
354 normative values in a young, active population. *Int J Sports Phys Ther* 6: 75–82, 2011.
- 355 22. Teyhen, DS, Shaffer, SW, Lorenson, CL, Halfpap, JP, Donofry, DF, Walker, MJ, Dugan,  
356 JL, and Childs, JD. The Functional Movement Screen: a reliability study. *J Orthop Sports*  
357 *Phys Ther* 42: 530–540, 2012.
- 358 23. Warren, M, Smith, CA, and Chimera, NJ. Association of Functional Movement Screen  
359 With Injuries in Division I Athletes. *J Sport Rehabil* Sep 8, 2014. Epub ahead of print.
- 360 24. Weir, JP. Quantifying test-retest reliability using the intraclass correlation coefficient and  
361 the SEM. *J Strength Cond Res* 19: 231–240, 2005.
- 362 25. Yagi, S, Muneta, T, and Sekiya, I. Incidence and risk factors for medial tibial stress  
363 syndrome and tibial stress fracture in high school runners. *Knee Surg Sports Traumatol*  
364 *Arthrosc* 21: 556–563, 2013.

365

## 366 **ACKNOWLEDGMENTS**


367 Special thanks to all the participants who willingly participated in this study.

368 **LEGENDS**369 **FIGURE 1.** Flow chart illustrating the process of the study.370 **FIGURE 2.** Distribution of scores for each functional movement screen (FMS) component  
371 test.372 **FIGURE 3.** Comparison of the DS & ASLR score between groups with and without injury.373 **FIGURE 4.** Receiver-operator characteristic (ROC) curves for FMS composite and DS &  
374 ASLR score.375 **TABLE 1.** Interrater reliability for the FMS composite score.376 **TABLE 2.** Interrater reliability for each FMS component test.377 **TABLE 3.** Comparison of runners with and without running injuries during the season.378 **TABLE 4.**  $2 \times 2$  contingency table: FMS composite score  $\times$  running injuries.379 **TABLE 5.**  $2 \times 2$  contingency table: DS & ASLR score  $\times$  running injuries.380 **TABLE 6.** Influence of the FMS on running injury.




1


2 **FIGURE 1.** Flow chart illustrating the process of the study.



1  
2 **FIGURE 2.** Distribution of scores for each functional movement screen (FMS) component  
3 test. DS = deep squat, HS = hurdle step, ILL = in-line lunge, SM = shoulder mobility, ASLR  
4 = active straight leg raise, TSPU = trunk stability push-up, RS = rotary stability.



1  
2 **FIGURE 3.** Comparison of DS & ASLR scores between groups with and without running  
3 injuries.



1

2 **FIGURE 4.** Receiver-operator characteristic (ROC) curves for FMS composite and DS &  
3 ASLR score.

1 **TABLE 1. Interrater reliability for the FMS composite score.**

ICC (2, 1)	95%CI	Bland-Altman plot				SE M	MD C <sub>95</sub>	SE P
		fixed bias		proportional bias				
		95%CI	presence/absence	test for no correlation	presence/absence			
0.98	0.93-1.00	-0.83-0.43	absence	r = -0.44	p = 0.90	0.31	0.87	0.44

95%CI: 95% confidence interval, SEM = standard error of measurement, MDC = minimum difference to be considered real, SEP = standard error of prediction

**TABLE 2.** Interrater reliability for each FMS component test.

Test	Kappa	Strength of agreement
Deep squat	1.000	Excellent
Hurdle step	1.000	Excellent
In-line lunge	1.000	Excellent
Shoulder mobility	1.000	Excellent
Active straight leg raise	0.831	Substantial
Trunk stability push-up	0.836	Substantial
Rotary stability	1.000	Excellent



1 **TABLE 3.** Comparison of runners with and without running injuries during the season.

Variable	Serious running injury		<i>P</i> value
	without (n = 69)	with (n = 15)	
<b>Characteristics</b>			
Age (year) <sup>†</sup>	20.1 ± 1.2	19.6 ± 0.9	0.15
Height (cm)	171.3 ± 4.3	172.7 ± 5.6	0.29
Weight (kg)	57.3 ± 4.2	58.4 ± 5.0	0.39
Running experience (year) <sup>†</sup>	6.9 ± 2.2	6.7 ± 2.4	0.64
Weekly training sessions (day/week) <sup>††</sup>	5.9 ± 0.6	5.9 ± 0.6	0.85
Weekly mileage (km/week) <sup>†</sup>	80.9 ± 53.8	98.4 ± 57.3	0.26
Performance (%)	87.6 ± 4.1	88.7 ± 3.6	0.34
Injury history, (n, %) <sup>†††</sup>	34 (49.3%)	8 (53.3%)	1.00
<b>FMS</b>			
FMS total score <sup>†</sup>	14.4 ± 2.2	13.3 ± 2.7	0.10
Deep squat <sup>††</sup>	1.8 ± 0.7	1.3 ± 0.7	0.01*
Hurdles step <sup>††</sup>	2.1 ± 0.3	2.0 ± 0.0	0.20
In-line lunge <sup>††</sup>	2.0 ± 0.4	1.9 ± 0.7	0.26
Shoulder mobility <sup>††</sup>	2.6 ± 0.8	2.5 ± 0.6	0.36
Active straight leg raise <sup>††</sup>	2.3 ± 0.6	1.6 ± 0.5	< 0.01**
Trunk stability push-up <sup>††</sup>	2.0 ± 1.0	2.5 ± 0.8	0.06
Rotary stability <sup>††</sup>	1.6 ± 0.5	1.6 ± 0.6	0.97

\**p* < .05. \*\**p* < .01

<sup>†</sup>Continuous data are expressed as the mean ± SD (tested by the student's t-tests).

<sup>††</sup>Non parametric data are expressed as the mean ± SD (tested by the Mann–Whitney U tests).

<sup>†††</sup>Categorical data are expressed as numbers (percentages) (tested by the chi-squared test).

1 **TABLE 4.**  $2 \times 2$  contingency table: FMS composite score  $\times$  running injuries.

	Running injuries	
	without	with
FMS composite score $\leq 14$	32	11
FMS composite score $\geq 15$	37	4

2

1 **TABLE 5.**  $2 \times 2$  contingency table: DS & ASLR score  $\times$  running injuries.

	Running injuries	
	without	with
DS & ASLR score $\leq 3$	18	11
DS & ASLR score $\geq 4$	51	4

2

1 **TABLE 6.** Influence of the FMS on running injury.

	univariate			multivariate*		
	OR	95%CI	<i>P</i> value	OR	95%CI	<i>P</i> value
FMS composite score $\leq 14$	3.2	0.9-11.0	0.07	3.0	0.8-11.6	0.10
DS & ASLR score $\leq 3$	7.8	2.2-27.6	<0.01**	9.7	2.1-44.4	<0.01**

\*\**p* < .01

\*Adjusted for age, height, weight, running experience, weekly training sessions, weekly mileage, performance, and injury history.