

Title	9-Oxo-10(E),12(Z),15(Z)-Octadecatrienoic Acid Activates Peroxisome Proliferator-Activated Receptor in Hepatocytes.
Author(s)	Takahashi, Haruya; Kamakari, Kosuke; Goto, Tsuyoshi; Hara, Hideyuki; Mohri, Shinsuke; Suzuki, Hideyuki; Shibata, Daisuke; Nakata, Rieko; Inoue, Hiroyasu; Takahashi, Nobuyuki; Kawada, Teruo
Citation	Lipids (2015), 50(11): 1083-1091
Issue Date	2015-11
URL	http://hdl.handle.net/2433/203050
Right	The final publication is available at Springer via http://dx.doi.org/10.1007/s11745-015-4071-3 .; The full-text file will be made open to the public on 19 September 2016 in accordance with publisher's 'Terms and Conditions for Self-Archiving'.; This is not the published version. Please cite only the published version. この論文は出版社版ではありません。引用の際には出版社版をご確認ご利用ください。
Type	Journal Article
Textversion	author

1 **9-Oxo-10(E),12(Z),15(Z)-octadecatrienoic acid activates peroxisome**
2 **proliferator-activated receptor α in hepatocytes**

3
4 Haruya Takahashi¹, Kosuke Kamakari¹, Tsuyoshi Goto^{1,2}, Hideyuki Hara¹, Shinsuke
5 Mohri¹, Hideyuki Suzuki³, Daisuke Shibata³, Rieko Nakata⁴, Hiroyasu Inoue⁴,
6 Nobuyuki Takahashi^{1,2} and Teruo Kawada^{1,2,#}

7
8 ¹*Laboratory of Molecular Function of Food, Division of Food Science and*
9 *Biotechnology, Graduate School of Agriculture, Kyoto University, Kyoto Japan*

10 ²*Research Unit for Physiological Chemistry, Kyoto University, Kyoto, Japan*

11 ³*Kazusa DNA Research Institute, Chiba, Japan*

12 ⁴*Department of Food Science and Nutrition, Nara Women's University, Nara, Japan*

13
14 #To whom correspondence should be addressed. Tel: +81-774-38-3751; Fax:
15 +81-774-38-3752; E-mail: fat@kais.kyoto-u.ac.jp

16
17 **Short title:** 9-oxo-OTA activates PPAR α in hepatocytes

18 **Abbreviations:** 9-oxo-OTA, 9-oxo-10(E),12(Z),15(Z)-octadecatrienoic acid;
19 9-oxo-ODA, 9-oxo-10,12-octadecadienoic acid; BSA, bovine serum albumin; DMEM,
20 Dulbecco's modified Eagle's medium; EtOH, ethanol; HAD, *cis*-10-heptadecenoic acid;
21 LC-MS, liquid chromatography–mass spectrometry; LNA, linoleic acid; ALA,
22 α -linolenic acid; PBS, phosphate-buffered saline; PPAR, peroxisome
23 proliferator-activated receptor; QTOFMS, quadrupole-time-of-flight MS; UPLC, ultra
24 performance LC

25

26

27 **Abstract**

28 Peroxisome proliferator-activated receptor (PPAR) α is mainly expressed in the liver
29 and plays an important role in the regulation of lipid metabolism. It has been reported
30 that PPAR α activation enhances fatty acid oxidation and reduces fat storage. Therefore,
31 PPAR α agonists are used to treat dyslipidemia. In the present study, we found that
32 9-oxo-10(*E*),12(*Z*),15(*Z*)-octadecatrienoic acid (9-oxo-OTA), which is α -linolenic acid
33 (ALA) derivative, is present in tomato (*Solanum lycopersicum*) extract. We showed that
34 9-oxo-OTA activated PPAR α and induced the mRNA expression of PPAR α target genes
35 in murine primary hepatocytes. These effects promoted fatty acid uptake and the
36 secretion of β -hydroxybutyrate, which is one of the endogenous ketone bodies. We also
37 demonstrated that these effects of 9-oxo-OTA were not observed in PPAR α -knockout
38 (KO) primary hepatocytes. To our knowledge, this is the first study to report that
39 9-oxo-OTA promotes fatty acid metabolism via PPAR α activation and discuss its
40 potential as a valuable food-derived compound for use in the management of
41 dyslipidemia.

42

43 **Key words:** PPAR α , oxylipin, hepatocytes, fatty acid metabolism, LC-MS.

44

45

46 **Introduction**

47 Dyslipidemia, which results from obesity, is a recognized risk factor for serious
48 disorders, including arteriosclerosis, hyperlipidemia, hepatic steatosis, insulin resistance,
49 and cardiovascular disease [1-4]. Therefore, avoidance of dyslipidemia is important to
50 help prevent these lifestyle diseases.

51 Peroxisome proliferator-activated receptors (PPARs) are a family of ligand-activated
52 transcription factors [5-9] comprised of three isoforms PPAR α , PPAR β/δ , and PPAR γ .
53 PPAR α is very important in the regulation of lipid metabolism [5-7] and is expressed at
54 high levels in the liver where it promotes β -oxidation, ketogenesis, and lipid transport
55 [10, 11]. PPAR α activation induces the expression of PPAR α -regulated genes (e.g.,
56 *carnitine-O-palmitoyltransferase 1* [CPT] and *acyl-CoA oxidase* [ACO]) and activates
57 β -oxidation [12-14]. These effects result in increased energy expenditure and reduced
58 fat storage [15]. It has been reported that PPAR α activation enhances fatty acid
59 metabolism and decreases the levels of circulating and cellular lipids in obese diabetic
60 patients [9, 16]. Therefore, regulation of PPAR α activity is one of the important ways to
61 manage chronic diseases related to lipid metabolism dysfunction.

62 Tomato (*Solanum lycopersicum*) is one of the most extensively consumed crops
63 worldwide. Numerous studies have shown that dietary tomato intake is associated with
64 a reduced risk of chronic diseases [17–20]. Recently, we reported that tomatoes contain
65 the oxidized linoleic acid (LNA) derivatives 9-oxo-10,12-octadecadienoic acid
66 (9-oxo-ODA) and 13-oxo-9,11-octadecadienoic acid (13-oxo-ODA), which can
67 function as PPAR α agonists [21,22]. Oxo-ODAs are oxylipins, which include fatty acid
68 hydroperoxides, hydroxyl-, oxo-, epoxy, and keto fatty acids, divinyl ethers, aldehydes,
69 and jasmonic acid [23]. Tomato is believed to contain not only oxo-ODAs but also

70 many other beneficial oxylipins that enhance lipid metabolism and suppress diseases
71 such as arteriosclerosis, hyperlipidemia, and hepatic steatosis.

72 The aim of the present study was to identify the metabolite in tomato that enhances
73 fatty acid metabolism and to determine its mechanism of action. In the present study, we
74 showed that 9-oxo-10(*E*),12(*Z*),15(*Z*)-octadecatrienoic acid (9-oxo-OTA), which is a
75 derivative of α -linolenic acid (ALA), is present in tomato fruit extract. We showed that
76 9-oxo-OTA activates PPAR α and induces the mRNA expression of PPAR α target genes,
77 which promotes fatty acid metabolism in murine hepatocytes. To the best of our
78 knowledge, this is the first study to report that 9-oxo-OTA induces fatty acid
79 metabolism through PPAR α activation, and our findings suggest that 9-oxo-OTA could
80 be a valuable food-derived compound capable of improving dyslipidemia.

81

82 **Materials and methods**

83 **Materials**

84 Authentic 9-oxo-OTA was purchased from Cayman Chemicals (Ann Arbor, MI,
85 USA). Authentic *cis*-10-heptadecenoic acid (HDA), which was used as an internal
86 standard, was purchased from Sigma (St. Louis, MO, USA). All other chemicals were
87 obtained from Sigma or Wako (Osaka, Japan). Buffers used were of high-performance
88 liquid chromatography (HPLC) or liquid chromatography-mass spectrometry (LC-MS)
89 grade. Tomatoes were obtained from a local market.

90

91 **Analysis of 9-oxo-OTA by ultra-performance liquid chromatography-quadrupole** 92 **time-of-flight mass spectrometry (UPLC-QTOF-MS)**

93 The levels of 9-oxo-OTA were assessed as previously described, using a LC-MS
94 system [24]. Briefly, each freeze-dried tomato (10 mg) was homogenized in 1 mL of
95 extraction solvent (99.5% EtOH containing 1 µg/mL HDA) with a mixer. After
96 centrifugation (15,000 rpm, 10 min, 4°C), the supernatant was collected for use as an
97 extract. The extract was filtered through a 0.2-µm-pore polyvinylidene difluoride
98 (PVDF) membrane (Whatman, Brentford, UK), and the filtrate was used for LC-MS.

99 LC-MS was performed using a Waters Acquity UPLC system (Milford, MA)
100 coupled to a Xevo QTOF-MS equipped with an electrospray source operating in
101 negative ion mode (ESI negative mode), with a lock-spray interface for accurate mass
102 measurements. Leucine enkephalin was employed as the lock-mass compound. It was
103 infused directly into the MS at a flow rate of 30 µL/min and a concentration of 200
104 µg/mL (in 50% acetonitrile, 50% water, 0.1% formic acid). The capillary, sampling cone,
105 and extraction cone voltages were set at 2700 V, 20 V, and 1 V, respectively. The source

106 and desolvation temperatures were 120°C and 450°C, respectively. The cone and
107 desolvation gas flow rates were set at 50 and 800 L/h, respectively.

108 An aliquot of the extracted sample (3 µL) was injected into an Acquity UPLC
109 BEH-C18 reversed-phase column (column size, 2.1 × 100 mm; particle size, 1.7 µm).
110 Mobile phases A (water and 0.1% formic acid) and B (acetonitrile and 0.1% formic
111 acid) were used. The column temperature was set to 40°C. The buffer gradient consisted
112 of 30% to 50% B for 0–4 min, 50% to 85% B for 4–14 min, 99% B for 14–17 min, and
113 30% B for 3 min, at a flow rate of 300 µL/min. Data were acquired with MassLynx
114 software (Waters). External mass calibration was performed following the
115 manufacturer's protocol.

116

117 **Analysis of β-hydroxybutyrate by UPLC-QTOF-MS**

118 The medium from a culture of primary hepatocytes (50 µL) was mixed with 50 µL
119 of acetonitrile. After centrifugation (15,000 rpm, 10 min, 4°C), the supernatant was
120 collected for use as an as an extract. The extract was filtered through a 0.2-µm-pore
121 polyvinylidene difluoride (PVDF) membrane (Whatman), and the filtrates were used for
122 LC-MS.

123 In the MS system, the capillary, sampling cone, and extraction cone voltages were
124 set at 2600 V, 15 V, and 3 V, respectively (ESI negative mode). The source and
125 desolvation temperatures were 120°C and 450°C, respectively. The cone and
126 desolvation gas flow rates were set at 50 and 800 L/h, respectively.

127 We used a BEH Amide column (column size, 2.1 × 100 mm; particle size, 1.7 µm).
128 The column temperature was set to 40°C. The buffer gradient consisted of 95% B for
129 0–4 min, 95% to 50% B for 4–10 min, 50% B for 10–16 min, and 95% B for 4 min, at a

130 flow rate of 300 μ L/min. Data were acquired with MassLynx software (Waters).

131

132 **Luciferase assay**

133 Luciferase assays were performed as previously described, using a GAL4/PPAR
134 chimera system [25]. We transfected p4xUASg-tk-luc (a reporter plasmid),
135 pM-hPPAR α (an expression plasmid for a chimeric protein containing the GAL4
136 DNA-binding domain and each human PPAR-ligand-binding domain), and pRL-CMV
137 (an internal control for transfection efficiency) into monkey CV1 kidney cells by using
138 Lipofectamine (Life Technologies Japan Ltd.) according to the manufacturer's protocol.
139 Luciferase activity was assayed using the dual luciferase system (Promega, Madison,
140 WI, USA) according to the manufacturer's protocol.

141

142 **Preparation of mouse primary hepatocytes**

143 All the animal experiments were approved by the Kyoto University Animal Care
144 Committee. C57BL/6J male mice (*wild type* and *PPAR α ^{-/-}*, free-feed) were anesthetized
145 with intraperitoneal administration of pentobarbital, and the liver was perfused with
146 liver perfusion medium (Life Technologies Japan Ltd.). Then, the liver was treated with
147 liver digestion medium (Life Technologies Japan Ltd.). After filtration through a
148 100- μ m nylon mesh filter, hepatocytes were isolated by repeated centrifugation at 50 \times
149 g for 3 min (3 times). The isolated hepatocytes were cultured in type-1 collagen-coated
150 12-well plates at a cell density of 2.0×10^5 cells/well. After a 5-h incubation at 37°C in
151 5% CO₂ in 20% serum DMEM (1.0g/L glucose), the hepatocytes were cultured in
152 serum-free DMEM (1.0g/L glucose) with or without 9-oxo-OTA for 5 h (mRNA assay)
153 or 8 h (LC-MS and RI assay), and then used for mRNA quantification and LC-MS

154 assays.

155

156 **Quantification of mRNA expression levels**

157 Total RNA was prepared from primary hepatocytes using Sepasol (Nacalai Tesque),
158 according to the manufacturer's protocol. Using M-MLV reverse transcriptase (Life
159 Technologies Japan Ltd.), total RNA was reverse transcribed. To determine mRNA
160 expression levels, real-time quantitative RT-PCR analysis was performed with a Light
161 Cycler System (Roche Diagnostics) using SYBR green fluorescence signals as
162 described previously [25, 26]. The oligonucleotide primer sets for mouse *36B4* (internal
163 control) and the PPAR α target genes were designed using a PCR primer selection
164 program on the website of the Virtual Genomic Center at GenBank and the sequences
165 are shown in Table 1. All mRNA expression data are presented as ratios relative to the
166 control in each experiment.

167

168 **Measurement of ^{14}C -Palmitic acid uptake and $^{14}\text{CO}_2$ production in murine** 169 **primary hepatocytes**

170 These experiments were conducted as previously described with some modifications
171 [27-29]. In the measurement of ^{14}C -palmitic acid uptake, the cells were washed with
172 PBS (500 μL /well) and then incubated in assay buffer (^{14}C -palmitic acid, 2 μCi /well;
173 fatty acid free BSA, 2.5%; and normal palmitic acid, 200 μM) for 3 min at room
174 temperature (1 mL/well). The experiment was stopped by washing the cells 3 times with
175 PBS (500 μL /well). The cells were lysed in 0.1% Triton-X solution (200 μL /well). The
176 radioactivity in the cell lysate was counted for 1 min in 2 mL of scintillation solution to
177 measure fatty acid uptake. In the measurement of $^{14}\text{CO}_2$ production, mice primary

178 hepatocytes were incubated in assay buffer (DMEM containing ^{14}C -palmitic acid, 2
179 $\mu\text{Ci}/\text{well}$; glucose, 1.0g/L; L-carnitine, 200 μM ; fatty acid free BSA, 2.5%; and normal
180 palmitic acid, 200 μM) for 8h. The medium was transferred to a 50-mL tube. An
181 uncapped-2mL sample tube containing a piece of filter paper soaked in 0.1N NaOH was
182 placed inside a 50-mL tube. After the tube was sealed, 200 μL of 70% perchloric acid
183 was added to the medium sample to release $^{14}\text{CO}_2$. The saturated filter paper containing
184 trapped $^{14}\text{CO}_2$ was assessed for radioactivity in a liquid scintillation counter (LS6500,
185 Beckman Coulter, CA, USA).

186

187 **Statistical analyses**

188 Data are presented as mean \pm standard error of the mean (SEM). Differences
189 between groups were compared with the Student's *t*-test (for two groups) and one-way
190 analysis of variance (ANOVA) followed by Tukey's test. *P* values less than 0.05 were
191 considered statistically significant.

192

193 **Results**

194 **Identification of 9-oxo-OTA in tomato fruit extract**

195 First, we explored the oxylipins, which are metabolized from ALA in plants (Fig. 1),
196 in the tomato extract using LC-MS. The results showed that the extract of tomato fruit
197 homogenate obtained under heating (37 °C, 30 min) contained 9-oxo-OTA (Fig. 2 A; Rt
198 = 7.91 min, m/z = 291.20, Fig. 2 B, C). The 9-oxo-OTA peak was detected as an
199 unfragmented deprotonation ion ($C_{18}H_{27}O_3^-$, $[M-H]^-$, Fig. 2 C) by electrospray negative
200 ionization mass spectrometry. In the tandem mass spectrometry (MS/MS) data, we
201 confirmed that the fragment ion detected in the tomato extract (m/z = 185, 125, and
202 121) matched the pure 9-oxo-OTA sample (Fig. 2D).

203 Quantitative analysis of the 9-oxo-OTA content in each part of the tomato fruit
204 homogenate under liquid nitrogen (gelatinous tissue: low limit of quantification
205 (LLOQ), sarcocarp: LLOQ, peel: LLOQ; Table 2) or heating (gelatinous tissue:
206 approximately 0.46 $\mu\text{g/g}$ of tissue weight, sarcocarp: 0.23 $\mu\text{g/g}$ of tissue weight, peel:
207 0.62 $\mu\text{g/g}$ of tissue weight; Table 2) revealed that the amount of 9-oxo-OTA in all
208 tissues was increased by heat treatment.

209

210 **9-oxo-OTA activated PPAR α and induced the mRNA expression of PPAR α target** 211 **genes**

212 9-oxo-OTA is analogous to 9-oxo-10,12-octadecadienoic acid (9-oxo-ODA, Fig. 1)
213 and 13-oxo-9,11,15-octadecatrienoic acid (13-oxo-OTA, Fig. 1). In our previous studies
214 showed that 9-oxo-ODA is involved in PPAR α activation [22] and 13-oxo-OTA
215 activates PPAR γ [24]. Therefore, we hypothesized that 9-oxo-OTA activates PPAR α
216 and/or PPAR γ . To determine the effect of 9-oxo-OTA on PPAR α and PPAR γ activity, we

217 first performed a luciferase ligand assay. In the PPAR α activation level, 9-oxo-OTA was
218 stronger than 13-oxo-OTA (Fig. 3 A). In the PPAR γ activation level, 13-oxo-OTA was
219 stronger than 9-oxo-OTA (Fig. 3 B). We also showed that 9-oxo-OTA increased
220 luciferase activity in a time dependent manner (Fig. 3 C), and this luciferase activity
221 was inhibited by PPAR α antagonist (GW6471) treatment (Fig. 3 D). PPAR α is
222 expressed in liver where it promotes β -oxidation, ketogenesis, and lipid transport [10,
223 11] and important in the regulation of lipid metabolism [5-7]. To elucidate the
224 contribution of 9-oxo-OTA to lipid metabolism in hepatocytes, murine hepatocytes were
225 cultured in medium containing 9-oxo-OTA. In murine hepatocytes, we showed that the
226 expression of PPAR α mRNA was increased by 9-oxo-OTA treatment (approximately
227 1.8-fold, Fig. 4 A). Furthermore, the mRNA expression levels of PPAR α target genes,
228 such as *Cpt1a*, *Acs*, *Hmgcs2*, and *Cd36*, were increased by 9-oxo-OTA treatment
229 (approximately 1.5-fold, 1.4-fold, 1.5-fold, and 1.4-fold, respectively, Fig. 4 B–E). We
230 also demonstrated that the 9-oxo-OTA-mediated effect on PPAR α target gene
231 expression was not observed in PPAR α -KO primary hepatocytes (Fig. 4 B–E). These
232 findings suggest that 9-oxo-OTA promotes the mRNA expression of the genes involved
233 in fatty acid oxidation via PPAR α activation.

234

235 **9-oxo-OTA promoted fatty acid oxidation in primary hepatocytes**

236 PPAR α is the major regulator of fatty acid metabolism [5-7], suggesting that
237 9-oxo-OTA induces fatty acid oxidation via PPAR α activation. We investigated whether
238 9-oxo-OTA influences not only mRNA expression levels but also lipid metabolism. The
239 results showed that ¹⁴C-palmitic acid uptake was increased by 9-oxo-OTA treatment
240 (approximately 1.4-fold, Fig. 5 A), suggesting enhanced fatty acid metabolism. We also

241 showed that $^{14}\text{CO}_2$ production was increased by 9-oxo-OTA treatment (approximately
242 1.5-fold, Fig. 5 B). β -Oxidation is one of the primary modes of fatty acid metabolism,
243 which converts fatty acids to ketone bodies. β -hydroxybutyrate, which is one of the
244 ketone bodies, is an important marker of fatty acid oxidation. We investigated whether
245 9-oxo-OTA effects the production of β -hydroxybutyrate in murine primary hepatocytes
246 and demonstrated that β -hydroxybutyrate secretion was increased by 9-oxo-OTA
247 treatment (approximately 1.5-fold, Fig. 5 C). The enhancement of fatty acid uptake and
248 secretion of $^{14}\text{CO}_2$ and β -hydroxybutyrate induced by 9-oxo-OTA were not observed in
249 PPAR α -KO primary hepatocytes (Fig. 5). These findings suggest that 9-oxo-OTA
250 promotes fatty acid metabolism in primary hepatocytes via a PPAR α -dependent
251 pathway.

252

253 **Discussion**

254 In the present study, we analyzed the metabolites in tomato extract by using LC-MS
255 and identified 9-oxo-OTA. Our previous study showed that ALA, which is a
256 9-oxo-OTA precursor, was present in tomato extracts [30]. These data suggested that
257 9-oxo-OTA can be derived from ALA via a non-enzymatic and/or enzymatic reaction. It
258 is possible that ALA is susceptible to oxidation at C9 via auto-oxidation and/or
259 lipoxygenase is involved in the production of 9-oxo-OTA in tomato fruits. Our previous
260 study also revealed that 9-oxo-ODA is localized primarily in the fruit peel [30].
261 However, here we showed that 9-oxo-OTA exists in not only peel but also in the
262 gelatinous tissue and sarcocarp. Further examinations are necessary to elucidate the
263 differences in the localization of 9-oxo-OTA and 9-oxo-ODA. We also revealed that the
264 amount of 9-oxo-OTA in tomato extract is increased by heating. This finding suggested
265 that heat processing increases the amount of 9-oxo-OTA in processed tomato foods.
266 Tomato fruit contains not only 9-oxo-OTA but also many other compounds. Therefore,
267 it is difficult that the effect of tomato on fatty acid oxidation is explained by only
268 9-oxo-OTA via PPAR α dependent pathway. To evaluate 9-oxo-OTA *in vivo*, it is
269 necessary to process selective breeding (tomato fruit having high 9-oxo-OTA content)
270 and/or to improve processing condition (e.g. tomato juice having high 9-oxo-OTA
271 content).

272 PPAR α is expressed principally in the liver, and it is a key regulator of various
273 aspects of lipid metabolism, including β -oxidation, ketogenesis, and lipid transport [10,
274 11]. PPAR α agonists are used in the treatment of dyslipidemia [9, 16]. Although we
275 previously reported that 9-oxo-ODA contributes to the activation of PPAR α [21], little
276 is known about the effect of 9-oxo-OTA on PPAR α . The structure of 9-oxo-OTA is

277 similar to that of 9-oxo-OTA. Therefore, we hypothesized that 9-oxo-OTA also has the
278 ability to activate PPAR α .

279 Here, we demonstrated for the first time that 9-oxo-OTA activates PPAR α in
280 luciferase ligand assay. There is the possibility that 9-oxo-OTA activate PPAR α both
281 directly and/or indirectly. In case of the direct action, 9-oxo-OTA acts as PPAR α ligand.
282 In case of the indirect action, the metabolites of 9-oxo-OTA act as PPAR α ligands. We
283 also showed that 9-oxo-OTA increased the mRNA expression of *Cpt1a*, *Acs*, *Hmgcs2*,
284 and *Cd36*, which are PPAR α target genes that are involved in fatty acid metabolism
285 [12-14, 31-33, 35-36]. These effects of 9-oxo-OTA on the expression of PPAR α target
286 genes were not observed in PPAR α -KO primary hepatocytes. These findings suggested
287 that 9-oxo-OTA promotes fatty acid metabolism in murine hepatocytes via a
288 PPAR α -dependent pathway. However, the expression of *Cd36* was trend toward
289 significance in PPAR α -KO primary hepatocytes treated with 9-oxo-OTA. We confirmed
290 that 9-oxo-OTA also activates PPAR γ in luciferase ligand assay. It is well known that
291 the expression of *Cd36* is induced by not only PPAR α but also PPAR γ activation
292 [34-36]. Therefore, there is the possibility that the effect of 9-oxo-OTA on PPAR γ
293 activation contributes to trend toward increment of *Cd36* expression.

294 Our present study also showed that the effect of 9-oxo-OTA on fatty acid oxidation
295 occurs through alteration of not only mRNA expression levels but also metabolism.
296 Fatty acids are taken up by hepatocytes primarily via CD36 [32]. We showed that fatty
297 acid uptake was increased by 9-oxo-OTA treatment. This effect was not observed in
298 PPAR α -KO primary hepatocytes, suggesting that 9-oxo-OTA promotes fatty acid uptake
299 via PPAR α activation. We also showed that the mRNA expression level of *Hmgcs2*
300 (3-hydroxy-3-methylglutaryl-CoA synthase 2) was increased by 9-oxo-OTA treatment.

301 *Hmgcs2* is the key enzyme involved in ketone body formation [33]. β -Hydroxybutyrate
302 is a major component of ketone bodies and a marker of fatty acid oxidation. Therefore,
303 we hypothesized that 9-oxo-OTA also promotes the production of β -hydroxybutyrate in
304 murine hepatocytes. The LC-MS analysis revealed that β -hydroxybutyrate secretion
305 from murine primary hepatocytes was increased by 9-oxo-OTA treatment. In the RI
306 experiment using ^{14}C -palmitic acid, we showed that $^{14}\text{CO}_2$ production was increased by
307 9-oxo-OTA treatment. The increment of $^{14}\text{CO}_2$ production and β -hydroxybutyrate
308 secretion suggest that 9-oxo-OTA promoted lipid metabolism in wild-type murine
309 hepatocytes. In contrast, 9-oxo-OTA treatment decreased $^{14}\text{CO}_2$ production and
310 β -hydroxybutyrate secretion in PPAR α -KO primary hepatocytes. Our present study is
311 focused on the effect of 9-oxo-OTA on PPAR α activation and its downstream targets.
312 But, there is the possibility of the effect of 9-oxo-OTA on other pathways including
313 PPAR γ activation. The previous study showed that acetoacetyl-CoA synthetase (AACS)
314 is a PPAR γ target gene [37]. AACS is a ketone body-specific ligase and might play an
315 important role in the provision of acetyl-CoA for lipogenesis [38]. 9-oxo-OTA activates
316 both PPAR α and PPAR γ . In the PPAR α KO hepatocytes treated with 9-oxo-OTA, there
317 is the possibility that fatty acid oxidation via PPAR α dependent pathway disappeared
318 and lipogenesis proceed via PPAR γ dependent pathway. In this study, we showed that
319 9-oxo-OTA promotes fatty acid metabolism via PPAR α activation and increases the
320 expression of PPAR α target genes in murine hepatocytes. Further examination is
321 necessary to determine the effect of 9-oxo-OTA on human metabolism.

322 In conclusion, 9-oxo-OTA contained in tomato fruit activates PPAR α and induces
323 the mRNA expression of PPAR α target genes. This promotes fatty acid oxidation in
324 murine hepatocytes. These findings provide the first evidence that 9-oxo-OTA induces

325 fatty acid metabolism through a PPAR α -dependent pathway and may contribute to the
326 development of improved treatments for dyslipidemia patients. Our findings suggest
327 that 9-oxo-OTA might be a valuable, food-derived compound for maintaining health.
328

329 **Acknowledgements**

330 The authors thank S. Shinoto and M. Sakai for secretarial and technical support,
331 respectively. This work was supported in part by Research and Development Projects
332 for Application in Promoting New Policies Agriculture, Forestry, and Fisheries of Japan,
333 by grants-in-aid for Scientific Research from the Ministry of Education, Culture, Sports,
334 Science, and Technology of Japan (22228001 and 24688015), and by Research
335 Fellowships of the Japan Society for the Promotion of Science for Young Scientists
336 (24521).

337

338 **References**

- 339 1. Reilly MP, Rader DJ (2003) The Metabolic Syndrome: more than the sum of its
340 parts? *Circulation* 108: 1546–1551
- 341 2. Miranda PJ, DeFronzo RA, Califf RM, Guyton JR (2005) Metabolic syndrome:
342 definition, pathophysiology, and mechanism. *Am. Heart. J.* 149: 33–45
- 343 3. Kahn BB, Flier JS (2000) Obesity and insulin resistance. *J. Clin. Invest.* 106:
344 473–481
- 345 4. Ginsberg HN (2000) Insulin resistance and cardiovascular disease. *J Clin Invest*
346 106: 453–458
- 347 5. Escher P, Wahli W (2000) Peroxisome proliferator-activated receptors: insight into
348 multiple cellular functions. *Mutat Res* 448: 121-138
- 349 6. Desvergne B, Wahli W (1999) Peroxisome proliferator-activated receptors: nuclear
350 control of metabolism. *Endocr Rev* 20: 649-688
- 351 7. Chinetti G, Fruchart JC, Staeles B (2000) Peroxisome proliferator-activated
352 receptors (PPARs): nuclear receptors at the crossroads between lipid metabolism
353 and inflammation. *Inflamm Res* 49: 497-505
- 354 8. Duval C, Chinetti G, Trottein F, Fruchart JC, Staeles B (2002) The role of PPARs in
355 atherosclerosis. *Trends Mol Med* 8: 422-430
- 356 9. Goldenberg I, Benderly M, Goldbourt U (2008) Update on the use of fibrates: focus
357 on bezafibrate. *Vasc Health Risk Manag* 4: 131-141
- 358 10. Bernal-Mizrachi C, Weng S, Feng C, Finck BN, Knutsen RH, Leone TC, Coleman T,
359 Mecham RP, Kelly DP, Semenkovich CF (2003) Dexamethasone induction of
360 hypertension and diabetes is PPAR α dependent in LDL receptor-null mice. *Nat Med*
361 9: 1069-1075

- 362 11. Reddy JK, Hashimoto T (2001) Peroxisomal β -oxidation and peroxisome
363 proliferator-activated receptor alpha: an adaptive metabolic system. *Annu Rev Nutr*
364 21: 193-230
- 365 12. Tugwood JD, Issemann I, Anderson RG, Bundell KR, McPheat WL, Green S (1992)
366 The mouse peroxisome proliferator activated receptor recognizes a response
367 element in the 5' flanking sequence of the rat acyl CoA oxidase gene. *EMBO J* 11:
368 433-439
- 369 13. Roepstorff C, Halberg N, Hillig T, Saha AK, Ruderman NB, Wojtaszewski JF,
370 Richter EA, Kiens B (2005) Malonyl-CoA and carnitine in regulation of fat
371 oxidation in human skeletal muscle during exercise. *Am J Physiol Endocrinol*
372 *Metab* 288: E133-142
- 373 14. Miller CW, Ntambi JM (1996) Peroxisome proliferators induce mouse liver
374 stearoyl-CoA desaturase 1 gene expression. *Proc Natl Acad Sci USA* 93: 9443-9448
- 375 15. Bogna GG (2014) Peroxisome proliferator-activated receptors and their ligands:
376 nutritional and clinical implications. *Nutr J* 13: 17
- 377 16. Kim S, Shin HJ, Kim SY, Kim JH, Lee YS, Kim DH, Lee MO (2004) Genistein
378 enhances expression of genes involved in fatty acid catabolism through activation
379 of PPARalpha. *Mol Cell Endocrinol* 220: 51-58
- 380 17. Pandey DK, Shekelle R, Selwyn BJ, Tangney C, Stamler J (1995) Dietary vitamin C
381 and beta-carotene and risk of death in middle-aged men. The Western Electric Study.
382 *Am J Epidemiol* 142: 1269-1278
- 383 18. Lazarus SA, Bowen K, Garg ML (2004) Tomato juice and platelet aggregation in
384 type 2 diabetes. *JAMA*. 292: 805-806
- 385 19. Giovannucci E (1999) Tomatoes, tomato-based products, lycopene, and cancer:
386 review of the epidemiologic literature. *J Natl Cancer Inst* 91: 317-331

- 387 20. Blum A, Monir M, Wirsansky I, Ben-Arzi S (2005) The beneficial effects of
388 tomatoes. *Eur J Intern Med* 16: 402–404
- 389 21. Kim YI, Hirai S, Takahashi H, Goto T, Ohyan C, Tsugane T, Konishi C, Fujii T,
390 Inai S, Iijima Y, Aoki K, Shibata D, Takahashi N, Kawada T (2011)
391 9-oxo-10(E),12(E)-Octadecadienoic acid derived from tomato is a potent PPAR α
392 agonist to decrease triglyceride accumulation in mouse primary hepatocytes. *Mol*
393 *Nutr Food Res* 55: 585-593
- 394 22. Kim YI, Hirai S, Goto T, Ohyan C, Takahashi H, Tsugane T, Konishi C, Fujii T,
395 Inai S, Iijima Y, Aoki K, Shibata D, Takahashi N, Kawada T (2012) Potent PPAR α
396 activator derived from tomato juice, 13-oxo-9,11-octadecadienoic acid, decreases
397 plasma and hepatic triglyceride in obese diabetic mice. *PLoS One* 7: e31317
- 398 23. Grechkin A (1998) Recent developments in biochemistry of the plant lipoxygenase
399 pathway. *Prog Lipid Res* 37: 317-352
- 400 24. Takahashi H, Hara H, Goto T, Kamakari K, Nomura W, Mohri S, Takahashi N,
401 Suzuki H, Shibata D, Kawada T (2015) 13-Oxo-9(Z),11(E),15(Z)-octadecatrienoic
402 Acid Activates Peroxisome Proliferator-Activated Receptor γ in Adipocytes. *Lipids*
403 50: 3-12
- 404 25. Takahashi N, Kawada T, Goto T, Yamamoto T, Taimatsu A, Matsui N, Kimura K,
405 Saito M, Hosokawa M, Miyashita K, Fushiki T (2002) Dual action of isoprenols
406 from herbal medicines on both PPAR γ and PPAR α in 3T3- L1 adipocytes and
407 HepG2 hepatocytes. *FEBS Lett* 514: 315–322
- 408 26. Goto T, Takahashi N, Kato S, Egawa K, Ebisu S, Moriyama T, Fushiki T, Kawada
409 T (2005) Phytol directly activates peroxisome proliferator-activated receptor α
410 (PPAR α) and regulates gene expression involved in lipid metabolism in
411 PPAR α -expressing HepG2 hepatocytes. *Biochem Biophys Res Commun* 337:
412 440–445

- 413 27. Takahashi N, Kang MS, Kuroyanagi K, Goto T, Hirai S, Ohyama K, Lee JY, Yu R,
414 Yano M, Sasaki T, Murakami S, Kawada T (2008) Auraptene, a citrus fruit
415 compound, regulates gene expression as a PPARalpha agonist in HepG2
416 hepatocytes. *Biofactors* 33: 25-32
- 417 28. Huynh FK, Green MF, Koves TR, Hirschey MD (2014) Measurement of fatty acid
418 oxidation rates in animal tissues and cell lines. *Methods Enzymol* 542: 391-405
- 419 29. Goto T, Lee JY., Teraminami A, Kim YI, Hirai S, Uemura T, Inoue H, Takahashi N,
420 Kawada T (2011) Activation of peroxisome proliferator-activated receptor-alpha
421 stimulates both differentiation and fatty acid oxidation in adipocytes. *J Lipid Res*
422 52: 873-884.
- 423 30. Takahashi H, Kamakari K, Suzuki H, Mohri S, Goto T, Takahashi N, Matsumura Y,
424 Shibata D, Kawada T (2014) Localization of 9- and 13-oxo-octadecadienoic acids
425 in tomato fruit. *Biosci Biotechnol Biochem* 78: 1761-1764
- 426 31. Nakamura MT, Yudell BE, Loor JJ (2014) Regulation of energy metabolism by
427 long-chain fatty acids. *Prog Lipid Res.* 53: 124-144
- 428 32. Gao M, Bu L, Ma Y, Liu D (2013) Concurrent activation of liver X receptor and
429 peroxisome proliferator-activated receptor alpha exacerbates hepatic steatosis in
430 high fat diet-induced obese mice. *PLoS One* 8: e65641
- 431 33. Lee SS, Chan WY, Lo CK, Wan DC, Tsang DS, Cheung WT (2004) Requirement of
432 PPARalpha in maintaining phospholipid and triacylglycerol homeostasis during
433 energy deprivation. *J Lipid Res* 45: 2025-2037
- 434 34. Nakamura MT, Yudell BE, Loor JJ (2014) Regulation of energy metabolism by
435 long-chain fatty acids. *Prog Lipid Res* 53: 124-144
- 436 35. Fu J, Oveisi F, Gaetani S, Lin E, Piomelli D (2005) Oleoylethanolamide, an

- 437 endogenous PPAR-alpha agonist, lowers body weight and hyperlipidemia in obese
438 rats. *Neuropharmacology* 48: 1147-1153
- 439 36. Kalinowska A, Górski J, Harasim E, Harasiuk D, Bonen A, Chabowski A (2009)
440 Differential effects of chronic, in vivo, PPAR's stimulation on the myocardial
441 subcellular redistribution of FAT/CD36 and FABPpm. *FEBS Lett* 583: 2527-2534
- 442 37. Aguiló F, Camarero N, Relat J, Marrero PF, Haro D (2010) Transcriptional
443 regulation of the human acetoacetyl-CoA synthetase gene by PPARgamma.
444 *Biochem J* 427: 255-64
- 445 38. Ito M, Fukui T, Saito T, Tomita K (1986) Acetoacetyl-CoA synthetase specific
446 activity and concentration in rat tissues. *Biochim Biophys Acta* 876: 280-287
447

448 **Figure legends**

449 **Fig. 1. Oxo-OTAs, Oxo-ODAs and their precursors (fatty acids).**

450

451 **Fig. 2. Identification of 9-oxo-10(*E*),12(*Z*),15(*Z*)-octadecatrienoic acid (9-oxo-OTA)**

452 **by LC-MS. (A)** The structure of 9-oxo-OTA. **(B)** Extracted ion chromatogram ($m/z =$

453 291.20) data and **(C)** full mass data for 9-oxo-OTA in an authentic sample and a tomato

454 extract sample. **(D)** MS/MS data for 9-oxo-OTA in an authentic sample and a tomato

455 extract sample.

456

457 **Fig. 3. 9-oxo-OTA activated PPAR α .** **(A)** Effect of 9-oxo-OTA and 13-oxo-OTA

458 (30 μ M respectively) on PPAR α activity with a luciferase reporter assay. **(B)** Effect of

459 9-oxo-OTA and 13-oxo-OTA (10 μ M respectively) on PPAR γ activity with a luciferase

460 reporter assay. **(C)** Effect of 9-oxo-OTA on PPAR α activity in a dose-dependent manner.

461 **(D)** Evaluation of PPAR α activity with a luciferase reporter assay in cells treated with

462 9-oxo-OTA (30 μ M) and/or GW6471 (PPAR α antagonist, 100 nM). Data are presented

463 as mean \pm SEM ($n = 5$). Symbol for $p < 0.05$ between the means of the 3 groups. OTA;

464 9-oxo-OTA, GW; GW6471.

465

466 **Fig. 4. 9-oxo-OTA activated the expression of PPAR α target genes in murine**

467 **hepatocytes.** Effect of 9-oxo-OTA (30 μ M) on **(A)** PPAR α expression levels in murine

468 primary hepatocytes. Effect of 9-oxo-OTA (30 μ M) on **(B)** *Cpt1a*, **(C)** *Acs*, **(D)** *Hmgcs2*,

469 and **(E)** *Cd36* expression in wild-type (WT) or PPAR $\alpha^{-/-}$ (KO) primary hepatocytes.

470 Data are presented as mean \pm SEM ($n = 5-6$). ** $p < 0.01$, *** $p < 0.001$ vs. control.

471

472 **Fig. 5. 9-oxo-OTA promoted fatty acid metabolism in murine hepatocytes.** (A) The
473 uptake of ^{14}C -palmitic acid and (B) $^{14}\text{CO}_2$ production in murine WT or KO primary
474 hepatocytes treated with 9-oxo-OTA (30 μM). (C) The secretion of β -hydroxybutyrate
475 from murine WT or KO primary hepatocytes treated with 9-oxo-OTA (30 μM). Data are
476 presented as mean \pm SEM (n = 3–5). * p < 0.05, ** p < 0.01 vs. control.








Figure 1. Takahashi H, et al.

A**B****C****D**










A**B****C****D**

Figure 3. Takahashi H, et al.

A**B****C****D****E**




Figure 5. Takahashi H, et al.

Table 1. Oligonucleotide primers used for mRNA analysis.

gene	forward primer	reverse primer
<i>Ppara</i>	TCGCGTACGGCAATGGCTTT	TCTTCATCCCCAAGCGTAGGAGG
<i>Cpt1a</i>	CTCAGTGGGAGCGACTCTCA	GGCCTCTGTGGTACACGACAA
<i>Acs</i>	ACATCCACGTGTATGAGTTCTACGC	AGTAGACGAAGTTCTCACGGTCGAT
<i>Hmgcs2</i>	AATCAGTGGAAGCAAGCTGGA	GTCCAGGGAGGCCTTCAAAA
<i>Cd36</i>	GATGTGGAACCCATAACTGGATTAC	GGTCCCAGTCTATTTAGCCACAGT
<i>36B4</i>	TCCTTCTCCAGGCTTTGGG	GACACCCTCCAGAAAGCGAG

Table 2. Quantitative analysis of 9-oxo-OTA in tomato fruit homogenate under liquid nitrogen or heating (at 37°C for 30 min; µg/g fresh weight).

tissue	liquid nitrogen	heating
gelatinous tissue	LLOQ	0.46 ± 0.10
sarcocarp tissue	LLOQ	0.23 ± 0.05
peel tissue	LLOQ	0.62 ± 0.12

LLOQ: low limit of quantification. Data are presented as mean ± SEM (n = 3).