



This is a repository copy of *The adhesion GPCR Adgrg6 (Gpr126): Insights from the zebrafish model.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/172570/>

Version: Published Version

Article:

Baxendale, S. orcid.org/0000-0002-6760-9457, Asad, A., Shahidan, N.O. et al. (2 more authors) (2021) The adhesion GPCR Adgrg6 (Gpr126): Insights from the zebrafish model. *genesis*. e23417. ISSN 1526-954X

<https://doi.org/10.1002/dvg.23417>

Reuse

This article is distributed under the terms of the Creative Commons Attribution (CC BY) licence. This licence allows you to distribute, remix, tweak, and build upon the work, even commercially, as long as you credit the authors for the original work. More information and the full terms of the licence here:
<https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

REVIEW



WILEY

The adhesion GPCR *Adgrg6* (*Gpr126*): Insights from the zebrafish model

Sarah Baxendale¹ | Anzar Asad¹ | Nahal O. Shahidan¹ |
Giselle R. Wiggin² | Tanya T. Whitfield¹

¹Department of Biomedical Science, Bateson Centre and Neuroscience Institute, University of Sheffield, Sheffield, UK

²Sosei Heptares, Steinmetz Building, Granta Park, Cambridge, UK

Correspondence

Tanya T. Whitfield, Department of Biomedical Science, University of Sheffield, Sheffield, S10 2TN, UK.

Email: t.whitfield@sheffield.ac.uk

Funding information

Biotechnology and Biological Sciences Research Council, Grant/Award Numbers: BB/S007008/1, BB/R50581X/1; Engineering and Physical Sciences Research Council, Grant/Award Number: EP/T517835/1

Summary

Adhesion GPCRs are important regulators of conserved developmental processes and represent an untapped pool of potential targets for drug discovery. The adhesion GPCR *Adgrg6* (*Gpr126*) has critical developmental roles in Schwann cell maturation and inner ear morphogenesis in the zebrafish embryo. Mutations in the human *ADGRG6* gene can result in severe deficits in peripheral myelination, and variants have been associated with many other disease conditions. Here, we review work on the zebrafish *Adgrg6* signaling pathway and its potential as a disease model. Recent advances have been made in the analysis of the structure of the *Adgrg6* receptor, demonstrating alternative structural conformations and the presence of a conserved calcium-binding site within the CUB domain of the extracellular region that is critical for receptor function. Homozygous zebrafish *adgrg6* hypomorphic mutants have been used successfully as a whole-animal screening platform, identifying candidate molecules that can influence signaling activity and rescue mutant phenotypes. These compounds offer promise for further development as small molecule modulators of *Adgrg6* pathway activity.

KEYWORDS

Adgrg6, aGPCR, drug screening, *Gpr126*, heart, inner ear, myelination, skeleton, zebrafish

1 | INTRODUCTION

Adhesion G-protein-coupled receptors (aGPCRs) are the second largest group of GPCRs, and are regulators of a wide range of developmental and physiological processes. Like other GPCRs, aGPCRs have a 7-transmembrane domain, but are specifically characterized by their very large extracellular region (ECR), which includes a GPCR autoproteolysis-inducing (GAIN) domain (see Bondarev et al., 2020; Langenhan, 2019; Morgan et al., 2019; Vizurraga, Adhikari, Yeung, Yu, & Tall, 2020 for recent reviews). *Adgrg6* (also known as *Gpr126*) is one of the better-characterized aGPCRs. Initially described in

mammals (Moriguchi et al., 2004; Stehlik, Kroismayr, Dorfleutner, Binder, & Lipp, 2004), it subsequently gained attention through the analysis of mutant phenotypes in the zebrafish (Figure 1) (Geng et al., 2013; Monk et al., 2009; Monk, Oshima, Jörs, Heller, & Talbot, 2011), originally identified through mutagenesis screens for morphology (Whitfield et al., 1996) or myelination (Pogoda et al., 2006).

Adgrg6 has a conserved role in the formation of the myelin sheath that insulates axons of the peripheral nervous system (PNS) in vertebrates (Monk et al., 2009, 2011). The PNS is myelinated by neural-crest-derived Schwann cells, and *Adgrg6* is essential for Schwann cell maturation and the initiation of myelination in both zebrafish and mammals (Monk et al., 2009, 2011). Homozygous

Sarah Baxendale and Anzar Asad contributed equally to this work.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *genesis* published by Wiley Periodicals LLC.




FIGURE 1 Otic and myelination phenotypes in zebrafish *adgrg6* mutants. (a) Phenotypically wild-type sibling (left panel, ventral view) and homozygous *adgrg6^{fr24-/-}* mutant (right panel, dorsal view), showing swollen otic vesicles (arrowhead). Anterior to the left. Note that head and eye size, pectoral fins, pigmentation, and swim bladder inflation are all normal in the mutant. Scale bar, 200 μ m. Images reproduced from Geng et al., 2013. (b) In situ hybridization to *myelin basic protein a* (*mbpa*) transcripts in the trunk (top row) and to *versican b* (*vcanb*) in the ear (bottom row). Lateral views; anterior to the left. Top row: expression of *mbpa* in Schwann cells of the posterior lateral line ganglion (pIlg, arrowhead) and posterior lateral line nerve (pIln, arrow) is reduced in the hypomorph *tb233c* allele, and lost altogether in the strong *fr24* allele. The blurred stain to the left in each image is expression in the central nervous system, which is unaffected in the mutants. Scale bar, 100 μ m. Images reproduced from Geng et al., 2013. Bottom row: strong expression of *vcanb* in the ear persists abnormally in *adgrg6* mutants. Scale bar, 50 μ m. dpf, days post fertilization. Images reproduced from Diamantopoulou et al., 2019

adgrg6^{-/-} mutant zebrafish show a reduction or loss of the Schwann cell marker gene *myelin basic protein* (*mbp*) in the periphery (Figure 1b) (Geng et al., 2013; Monk et al., 2009). In the PNS, *adgrg6* is normally expressed in Schwann cells (Monk et al., 2009): contact with the axon is thought to trigger signaling through a canonical $G\alpha_s$ cascade, leading to activation of adenylyl cyclase and the production of cAMP, in turn activating Protein Kinase A and the expression of downstream target genes, including *oct6* and *egr2* (Monk et al., 2009).

In addition to a reduction or loss of peripheral myelination, *adgrg6* mutant zebrafish also have a striking inner ear phenotype (Figure 1) (Geng et al., 2013; Monk et al., 2009; Whitfield et al., 1996). Epithelial projections in the otic vesicle, which normally initiate the formation of the semicircular canal ducts, overgrow and fail to fuse in the mutant, resulting in an easily identifiable swelling of the ear (Diamantopoulou et al., 2019; Geng et al., 2013). As in the PNS, *Adgrg6* signaling in the developing zebrafish ear appears to be triggered by cell-cell contact

and to act through cAMP, but here *adgrg6* is expressed on both sides of the fusing tissue (Geng et al., 2013). Many genes, including those coding for extracellular matrix (ECM) proteins and synthases, together with various developmental signaling pathway components, remain expressed at abnormally high levels in the *adgrg6^{-/-}* mutant ear (Geng et al., 2013). These findings suggest that *Adgrg6* signaling primarily acts (directly or indirectly) to repress gene expression in the developing ear.

The *adgrg6* gene is expressed dynamically in the developing embryo. In the zebrafish, the ear is a major site of expression, but *adgrg6* is also expressed in the neural crest and its derivatives (Schwann cells, craniofacial cartilage), the heart, mesoderm, and other tissues (Geng et al., 2013; Monk et al., 2009; reviewed in Patra, Monk, & Engel, 2014). In the 1-day-old zebrafish embryo, the expression pattern is very similar to that of the transcription factor gene *sox10*, and *adgrg6* expression in the ear and neural crest (but

not in the heart) is reduced in *sox10*^{-/-} mutants, suggesting that Sox10 may be an upstream regulator of *adgrg6* expression in these tissues (Geng et al., 2013). In mammals, *Adgrg6* is known to be expressed in multiple tissues, including in the adult (Mogha et al., 2016; Musa et al., 2019; Patra et al., 2014; Waller-Evans et al., 2010).

The zebrafish *adgrg6* mutations identified or generated to date form an allelic series (Table 1) that has helped to dissect protein function. The different variants include hypomorphic missense mutations involving single amino acid changes in the transmembrane domain, which have been exploited in chemical screening experiments (see below), and point mutations that introduce stop codons early in the coding sequence, resulting in stronger ear and myelination defects (Figure 1). However, even strong *adgrg6* mutations in zebrafish are homozygous viable, unlike in the mouse, where targeted disruption of *Adgrg6* is lethal before or soon after birth (Monk et al., 2011; Waller-Evans et al., 2010). Although this may reflect differences in *Adgrg6* function between the two species, the effects of complete deletion of the zebrafish gene have not yet been demonstrated. It is also possible that the murine gene, which is expressed in the placenta, has a role in the development of extraembryonic tissues, which may contribute to the observed lethality in mutants (Waller-Evans et al., 2010).

Given the importance of *ADGRG6* in the myelination of the PNS, and its dynamic expression during development and in adult tissues, it is not surprising that *ADGRG6* also has a role in human disease. Evidence associating *ADGRG6* variants with a wide range of human pathologies, including both rare congenital conditions and more common disorders, is accumulating from family studies, GWAS data, RNA expression profiling studies, and animal models. In this review, we highlight some of the recent advances in *Adgrg6* research, with a focus on the zebrafish model. We illustrate how the use of a suite of genetic, transgenic, imaging, and gene editing tools, together with small molecule screening approaches, make the zebrafish such a versatile model for the dissection of gene function, leading to new insights into the biology of the *Adgrg6* receptor and the disorders that can result from its loss of function.

2 | THE *ADGRG6* GENE AND PROTEIN

2.1 | *Adgrg6* gene structure and alternative splicing

Although aGPCRs are found throughout the metazoa, *Adgrg6* has no orthologue in the commonly used invertebrate model organisms *Drosophila* and *C. elegans*, the invertebrate chordates *Ciona* and amphioxus, or in jawless vertebrates such as lamprey and hagfish (Waller-Evans et al., 2010; ensembl.org GRCz11; Yates et al., 2019). The *Adgrg6* gene thus appears to be specific to jawed vertebrates, and is evolutionarily conserved between teleost fish, amphibians, reptiles, birds, and mammals (ensembl.org GRCz11). Within the aGPCR class, *Adgrg6* is most closely related to secretin-like GPCRs (reviewed in Patra et al., 2014; Scholz, Langenhan, & Schöneberg, 2019). The zebrafish *adgrg6* gene has 26 exons and spans almost 90 kb. Unlike many zebrafish genes, which have two isoforms as a result of a teleost whole-genome duplication event (Meyer & Schartl, 1999), zebrafish *adgrg6* has a 1:1 correspondence—and ~50% nucleotide identity—with the orthologous human gene *ADGRG6*. The zebrafish and human *Adgrg6* proteins also show nearly 50% identity at the amino acid level, with the highest levels of conservation in the 7-transmembrane domain and the CUB domain of the ECR (Leon et al., 2020; ensembl.org GRCz11; Yates et al., 2019).

Alternative splicing is a feature of aGPCR genes (Salzman et al., 2016). The expression of multiple isoforms, particularly of the ECR, is thought to influence mechanical interactions with binding partners, enhance receptor versatility and offer a regulatory mechanism for receptor activity (Bjarnadóttir et al., 2007). The human and zebrafish *ADGRG6* genes are known to have four main alternative splice forms (Moriguchi et al., 2004; Patra et al., 2014). These include inclusion (S1 form, also known as +ss) or exclusion (S2 form, -ss) of the short exon 6, which codes for 23 amino acids in the zebrafish *Adgrg6* ECR (Patra et al., 2014). In addition, the presence or absence of the penultimate exon results in a frameshift that alters the coding sequence for the intracellular C-terminal domain, which could potentially influence downstream signaling events. Adhesion GPCRs in general are found to have a high level of somatic mutations (Kan et al., 2010; O'Hayre et al., 2013) and the human *ADGRG6* gene in particular has numerous variants (35,447 variants; *ADGRG6*-201 transcript, Yates et al., 2019).

TABLE 1 List of key zebrafish *adgrg6* mutations

Mutation	Amino Acid	Domain	Reported phenotype	Reference
<i>stl47</i>	Δ5+3, N68K, fs*28	CUB	ear, radial sorting, PNS myelination	(Petersen et al., 2015)
<i>stl464</i>	D134A, F135A	CUB	ear & PNS myelination	(Leon et al., 2020)
<i>fr24</i>	L463*	SEA	ear & PNS myelination	(Geng et al., 2013)
<i>st49</i>	Y782*	GAIN	ear & PNS myelination	(Monk et al., 2009; Pogoda et al., 2006)
<i>vu39</i>	W804*	GPS	ear & PNS myelination	(Geng et al., 2013)
<i>stl215</i>	ΔG831, I832	<i>Stachel</i>	ear & PNS myelination	(Liebscher, Schön, et al., 2014)
<i>st63</i>	C917Y	TM2	PNS myelination	(Monk et al., 2009; Pogoda et al., 2006)
<i>tb233c</i>	I963N	TM4	ear & PNS myelination	(Geng et al., 2013; Whitfield et al., 1996)
<i>tk256a</i>	P969L	TM4	ear & PNS myelination	(Geng et al., 2013; Whitfield et al., 1996)

Amino acid sequence and numbering are based on the reference sequence NM_001163291.2, with the exception of W804, which was reported as C804 in the reference sequence (see discussion in Geng et al. 2013). N68 was reported as Q68 in Petersen et al. 2015. Italic text indicates amino acid identity or similarity with the human protein. Abbreviations: Δ, deletion; fs, frame shift; see Figure 2 for domain name abbreviations.

2.2 | Adgrg6 protein structure

Current models for the zebrafish Adgrg6 protein structure are shown in Figure 2. During biosynthesis, autoproteolytic cleavage at the GPCR Proteolysis Site (GPS) in the GAIN domain separates the protein into N- and C-terminal fragments (NTF, CTF), which can remain non-covalently associated at the membrane (Araç et al., 2012). The NTF is composed of structural domains including the Complement C1r/C1s, Uegf and Bmp1 (CUB), Pentraxin (PTX), and Hormone Receptor (HormR) domains (Araç et al., 2012; Moriguchi et al., 2004), together with a recently identified sperm protein, enterokinase and agrin (SEA) domain (Leon et al., 2020). The SEA domain contains a furin cleavage site in the human and mouse protein that is not conserved in zebrafish. Like other aGPCRs, Adgrg6 undergoes post-translational modifications including N-linked glycosylation of the ECR, which is thought to contribute to the adhesive properties of the NTF (Langenhan, Aust, & Hamann, 2013; Leon et al., 2020; Patra et al., 2014).

Determining the structure of the receptor, and in particular that of the ECR, is fundamental to understanding its mechanism of action. The crystal structure of the zebrafish Adgrg6 ECR is one of only four aGPCR protein structures determined to date (Leon et al., 2020; Ping et al., 2021; Salzman et al., 2016; Vakonakis, Langenhan, Prömel, Russ, & Campbell, 2008), and has highlighted how conformational changes of the ECR might regulate receptor activity. Crystallization of the zebrafish Adgrg6 ECR and analysis with a range of imaging techniques has revealed that the S2 form adopts a closed conformation, in which the CUB domain interacts with the HormR domain (Figure 2b) (Leon et al., 2020). This closed configuration is dependent on the presence of a proposed Ca²⁺ binding site within the CUB domain, together with a disulfide-stabilized loop between the SEA and PTX domains. The loop inserts between the CUB and HormR domains, stabilizing their interaction (Leon et al., 2020). The 23-amino-acid linker in the S1 form appears to disrupt the stability of the CUB-HormR interaction, and the S1 ECR shows a variety of conformations, indicating an ability to switch between an open and closed state (Figure 2c) (Leon et al., 2020). This mobility translates to an increase in Adgrg6 signaling in comparison with the basal activity associated with the closed isoform in an in vitro cAMP assay (Leon et al., 2020).

An intact Ca²⁺-binding site is also necessary for optimal receptor function. A conserved aspartate residue co-ordinates a calcium ion in Ca²⁺-binding CUB domains from different aGPCRs (Gregory, Thielens, Arlaud, Fontecilla-Camps, & Gaboriaud, 2003). Disruption of the site in the zebrafish Adgrg6 protein, by CRISPR/Cas9-mediated replacement of both D134 and F135 residues with alanines, induces ear and myelination defects similar to those observed in a strong loss-of-function *adgrg6* mutant (Leon et al., 2020). In the crystallized ECR, disruption of the Ca²⁺-binding site resulted in an open configuration, but despite this, the S1 Adgrg6 isoform was unable to mediate an increase in cAMP accumulation above basal levels (Leon et al., 2020). These findings propose a complex mechanism by which the ECR may regulate Adgrg6 function, with an intact Ca²⁺-binding site and the 23-amino acid linker being necessary to trigger a boost in Adgrg6 signaling.

2.3 | Adgrg6 ligands and mechanosensing

The multiple domains in the long ECRs of aGPCRs provide the opportunity to bind to many different partners. Studies in a range of model systems and contexts, including morpholino phenocopy and mutant rescue experiments in the zebrafish, indicate that the Adgrg6 NTF interacts with the ECM components Collagen IV (Paavola et al., 2014) and Laminin-211 (Petersen et al., 2015), and the flexible tail residues 23–50 of the Prion protein (PrP₂₃₋₅₀) (Küffer et al., 2016) (Figure 2d). However, it is not fully understood how Adgrg6 and these natural ligands operate at the cell membrane in vivo. Collagen IV and the Prion protein appear to mediate cAMP accumulation through binding to the Adgrg6-NTF directly (Küffer et al., 2016; Paavola et al., 2014). However, application of purified Laminin-211 to cells transfected with human ADGRG6 resulted in an unexpected decrease in cAMP accumulation, attributed to an inhibition of basal signaling activity (Petersen et al., 2015). In this experimental context, receptor activation required the application of force, by sample vibration, to mimic the in vivo environment at the cell membrane (Petersen et al., 2015). These observations suggest a variety of mechanisms through which NTF binding partners can activate Adgrg6 signaling, and a potential role for Adgrg6 in mechanosensing. Indeed, a study analyzing expression domains of a LacZ reporter for *Adgrg6* expression in the mouse has highlighted its predominant expression in mechanosensing tissues (Musa, Cazorla-Vázquez, et al., 2019).

Adapting natural ligands as agonists of the Adgrg6 signaling pathway could have exciting therapeutic potential for human diseases involving disruption of ADGRG6 function. However, recent work to develop the Prion protein (PrP) as an Adgrg6 agonist has proved challenging. The application of a stable Adgrg6-activating PrP-based molecule failed to rescue a peripheral demyelinating neuropathy in PrP knock-out mice, and transcriptomic data suggested that the rescuing agent might itself be myotoxic (Henzi et al., 2020). Further identification of NTF ligands will be critical in teasing out the underlying mechanisms leading to Adgrg6 activation. However, natural ligands may not present a viable approach for therapeutic design due to their widespread effects in vivo.

2.4 | Tethered agonist signaling: Autoactivation of the Adgrg6 receptor

Early aGPCR studies highlighted that the NTF-CTF interaction at the cell membrane provides an inhibitory control mechanism to limit receptor signaling through G proteins (Okajima, Kudo, & Yokota, 2010; Paavola, Stephenson, Ritter, Alter, & Hall, 2011; Ward et al., 2011). Utilizing cAMP assays as a measure of ADGRG6 activity, Liebscher et al. (2014) uncovered a tethered peptide agonist, terming it the “*Stachel*” (“stinger”) sequence. The peptide is located at the N-terminal end of the CTF, where it is embedded in beta sheets of the GAIN domain (Figure 2) (Araç et al., 2012; Beliu et al., 2021). Although direct binding has not yet been visualized, in vivo treatment of hypomorphic *adgrg6*^{st63} zebrafish larvae with a synthetic 16-amino-acid *Stachel*




FIGURE 2 Schematic models of zebrafish Adgrg6 protein structure and mechanism of activation. (a) Zebrafish Adgrg6 protein sequence and domain organization, drawn to scale: signal peptide (SP), Complement C1r/C1s, Uegf, BMP1 (CUB), Pentraxin (PTX), Sperm protein, Enterokinase and Agrin (SEA), hormone receptor (HormR), GPCR autoproteolysis-inducing (GAIN), *Stachel* sequence (S) and 7-pass transmembrane (7TM, magenta) domains. The sequence of Adgrg6 splice isoforms is shown below the diagram: S1, including exon 6 (+SS, pink), and the short (α) and long (β) isoforms resulting from inclusion or exclusion, respectively, of exon 25 (amino acid sequence in blue; splice site in red). The two cysteine residues (C369, C375) that form a bond in the linker between PTX and SEA domain are shown in amber. Cleavage sites are highlighted by dotted lines: GPCR autoproteolytic site (GPS, black), and position of furin cleavage site in mammalian ADGRG6 (f, green). Positions of widely studied zebrafish mutations are shown above the diagram: truncating (nonsense) mutations in red; missense mutations in black. (b) Structure of Adgrg6 S2 isoform in closed conformation (not to scale). Domains correspond to those illustrated in (b); Ca²⁺ – binding site (green) within CUB domain, cell membrane (gray). (c) Structure of Adgrg6 S1 isoform in open conformation. (d) Overview of potential Adgrg6 *Stachel* activation mechanism (not to scale). Tethered *Stachel* sequence (amber star) self-activates Adgrg6 following removal of NTF. The NTF is also thought to signal in trans (not shown). Known Adgrg6 NTF ligands (dashed shapes) include Collagen IV (pink, Col4), prion protein flexible tail (PrP^{FT}, green), and Laminin-211 (Lam211, turquoise). Signaling is transduced by various G protein α subunits activating intracellular pathways, elucidated from zebrafish Adgrg6 studies (light blue) and mouse or human ADGRG6 studies (light red). Diagrams are adapted from (Leon et al., 2020); the figure summarizes findings from this study and from (Geng et al., 2013; Küffer et al., 2016; Liebscher, Schön, et al., 2014; Lizano, Hayes, & Willard, 2021; Moriguchi et al., 2004; Paavola, Sidik, Zuchero, Eckart, & Talbot, 2014; Patra et al., 2013; Petersen et al., 2015)

peptide has been shown to mediate some restoration of Schwann cell *mbp* expression in the PNS (Liebscher, Schön, et al., 2014). A *Stachel* sequence has since been identified in multiple other aGPCRs (Demberg, Rothmund, Schöneberg, & Liebscher, 2015; Stoveken, Hajduczuk, Xu, & Tall, 2015), suggesting it may be a common mechanism of action.

Two different models have been proposed for *Stachel* activation of signaling. The first relies on the physical removal of the NTF—through ligand interactions—to expose the tethered agonist, enabling it to bind to the active site of the CTF and self-activate receptor signaling (Liebscher, Schön, et al., 2014; Petersen et al., 2015). However, a recent study using molecular dynamic simulations of aGPCRs has proposed an alternative mechanism, in which the *Stachel* sequence is transiently exposed through structural flaps within the GAIN domain, following moderate movements (Beliu et al., 2021). Although not confirmed for *Adgrg6*, this notion could explain the basal activity of *Adgrg6* that is associated with the S2 isoform (Leon et al., 2020).

Stachel-independent activation of aGPCRs has also been detected by multiple groups (Kishore, Purcell, Nassiri-Toosi, & Hall, 2016; Salzman et al., 2017). Cells expressing a mutant isoform of ADGRG1 (H381S), defective in GPS autoproteolysis, exhibit an increase in signaling following exposure to monobody ligands, a response that is unaffected by disruption of the *Stachel* sequence (Salzman et al., 2017). Furthermore, cells expressing an engineered form of ADGRB1 lacking the *Stachel* peptide exhibit differential signaling behavior, including elevated TGF α shedding, in comparison with the intact receptor (Kishore et al., 2016). Exploiting equivalent *Adgrg6* mutant isoforms may reveal further evidence of this mechanism among aGPCRs and in which tissues or developmental contexts different mechanisms apply.

2.5 | *Adgrg6* coupling to G proteins

Initial studies to identify the G protein or proteins that couple to zebrafish *Adgrg6* showed that the myelination defects in *adgrg6*^{st49} mutants could be rescued by treatment with forskolin, an adenylyl cyclase agonist that raises intracellular cAMP levels (Monk et al., 2009). The ear phenotype in *adgrg6*^{tb233c} mutants can also be ameliorated by treatment with either forskolin or 3-isobutyl-1-methylxanthine (IBMX), a phosphodiesterase inhibitor that prevents cAMP degradation (Diamantopoulou et al., 2019; Geng et al., 2013). These findings suggest that *Adgrg6* couples to a G α_s signaling pathway in both Schwann cells and in the ear (Figure 2d). Coupling to G α_s has since been confirmed by several groups by detecting changes in intracellular cAMP levels following expression of human ADGRG6 (Liebscher, Schön, et al., 2014; Paavola et al., 2014; Petersen et al., 2015). Nevertheless, this may not represent the full picture as aGPCRs are well-known to signal through multiple G-protein pathways (reviewed in Langenhan, 2019). Application of a novel enterokinase-activated tethered ligand system of ADGRG6 not only confirmed coupling to G α_s , but also demonstrated stimulation of reporters for serum response factor (SRF) and serum response element (SRE), indicating coupling to G $\alpha_{12/13}$ and G $\alpha_{q/11}$ pathways, respectively (Lizano et al., 2021). However, a study utilizing chimeric

G proteins concluded that the human receptor can couple to G α_s and G α_i , but not G α_q (Mogha et al., 2013). Given the evidence, a complex interplay of multiple signaling pathways is likely to be associated with the *Adgrg6* receptor in vivo.

2.6 | Signaling via the *Adgrg6* N-terminal fragment

The NTF produced by autoproteolytic cleavage of *Adgrg6* at the GPS, or potentially through furin cleavage of the human or mouse protein, has been shown to act in trans, independently of the CTF. Initial studies of *Adgrg6* knock-out mice uncovered a delay in radial sorting of peripheral axons by Schwann cells, suggesting a role for *Adgrg6* in this process (Monk et al., 2011). Axonal sorting deficiencies were confirmed in zebrafish by analyzing *adgrg6*^{stl47} mutants, which predict a protein with a truncated CUB domain (Petersen et al., 2015). Moreover, the *stl47* mutants were unresponsive to forskolin treatment, suggesting that axonal sorting materializes independently of the canonical G-protein-coupled signaling associated with the *Adgrg6* CTF. Laminin-211 was identified as a ligand of the *Adgrg6* NTF in axonal sorting, and multiple zebrafish *adgrg6* mutant alleles were used to characterize this interaction (Petersen et al., 2015). Given that Collagen IV can also bind the NTF (Paavola et al., 2014), and that *Col4a1* mouse mutants display peripheral radial sorting defects (Labelle-Dumais et al., 2019), it is likely that Collagen IV performs a similar function to Laminin-211 in this context in the mouse.

Further evidence for the NTF acting in a cAMP-independent manner comes from studies in mice, where conditional disruption of *Adgrg6* in chondrocyte lineages results in skeletal abnormalities that mimic human disorders, as discussed in Section 3 below. Treatment of *Adgrg6* conditional knock-out mice with the phosphodiesterase inhibitor Rolipram, which would be expected to restore cAMP signaling downstream of the receptor, was unable to rescue the skeletal phenotype, suggesting alternative signaling pathways may be required in mammalian chondrocytes (Kamer, Long, Solnica-Krezel, Monk, & Gray, 2015). In another study, over-expression of the *Adgrg6* NTF (S1 isoform) was reported to delay ossification of human mesenchymal stem cells in vitro; the authors propose that differential expression of ADGRG6 or its isoforms could contribute to the abnormal curvature of the spine in patients with Adolescent Idiopathic Scoliosis (Xu et al., 2019) (see also Section 3.4).

The *Adgrg6* NTF has also been implicated in heart development. *Adgrg6* is expressed in the mouse and zebrafish heart (Geng et al., 2013; Monk et al., 2011; Moriguchi et al., 2004; Patra et al., 2013), and homozygous mutant mice have cardiac hypotrabeulation defects (Patra et al., 2013). However, cardiovascular defects are not obvious in most of the zebrafish mutant alleles analyzed to date (Geng et al., 2013; Leon et al., 2020; Monk et al., 2009), and *adgrg6* homozygous mutant fish are adult viable. By contrast, morpholino-mediated knock-down of *adgrg6* was reported to result in a similar hypotrabeulation defect to that in the mouse mutants, which could be rescued by injection of mRNA coding for the NTF (Patra et al., 2013). The reported heart defects in zebrafish morphants could be the result of an off-target effect of the morpholino, but it is

also possible that genetic compensation—now well characterized for other zebrafish genes (El-Brolosy et al., 2019)—acts to mask any heart phenotype in the mutants. Nevertheless, the mutant alleles with no overt heart defects still predict that a truncated NTF could be produced. The most severe truncating allele, *stl47*, predicts a deletion of most of the NTF; these mutants do display heart oedema (Petersen et al., 2015), and a recent report suggests they also have hypotrabeulation (Srivastava et al., reviewed in Morgan et al., 2019). A separate report found *Adgrg6* expression downstream of Notch signaling in both trabeculation and compaction of the mouse heart, with a reduction in cardiac *adgrg6* expression in the zebrafish Notch pathway mutant *mind bomb* (*E3 ubiquitin ligase*) (D'Amato et al., 2016). In addition, injection of zebrafish embryos with a potential post-transcriptional regulator of *Adgrg6*, *miR-27b*, produced the same hypotrabeulation phenotype as in *adgrg6* morphants (Musa, Srivastava, Petzold, Cazorla-Vázquez, & Engel, 2019). Given these observations, a detailed analysis of the cardiac phenotype in the different zebrafish alleles will be important to confirm any role for *Adgrg6* in zebrafish heart trabeculation.

3 | ADGRG6 AND HUMAN DISEASE

3.1 | ADGRG6 coding sequence mutations and inherited disease

Many of the mutations in *ADGRG6* associated with human disease (Table 2) affect residues that are conserved in zebrafish. The first studies to implicate *ADGRG6* in human disease corroborated the myelination defects discovered in zebrafish and mouse mutants. Proband from three consanguineous families with Lethal Congenital Contracture Syndrome 9 (LCCS9), a rare form of Severe Arthrogryposis Multiplex Congenita (AMC), were found to harbor different homozygous recessive mutations in the *ADGRG6* coding sequence (Ravenscroft et al., 2015). AMC is known to have multiple causes that can be musculoskeletal or neuronal in origin. Contractures were observed in the limbs of mouse *Adgrg6* mutants that lack

myelination of the peripheral nerves (Monk et al., 2011). In the Ravenscroft et al. (2015) study, a lack of MBP staining was also observed in the intramuscular nerves, confirming that a myelination defect underlies the phenotype. In a separate study of an Iranian family with autosomal recessive Intellectual Disability, whole exome sequencing identified a strong candidate missense mutation in the transmembrane domain of *ADGRG6*. The two patients survived beyond puberty and displayed a number of common features with LCCS9, including contractures (Hosseini et al., 2018).

3.2 | Identification of ADGRG6 variants through GWAS and RNA sequencing

Recent Genome-Wide Association Studies (GWAS) and RNAseq studies have found many associations with variants in both coding and non-coding regions of the human *ADGRG6* locus that may provide clues to *ADGRG6* function in disease (see <https://www.ebi.ac.uk/gwas/genes/ADGRG6> for a comprehensive list). One example is a missense mutation in the *ADGRG6* transmembrane region, R1057Q, that has been linked to severe periodontitis in a cohort of Japanese patients (Kitagaki et al., 2016). In vitro studies found that the amino acid change reduced cAMP accumulation compared with wild-type *ADGRG6*; expression of downstream target genes (*Bmp2*, *ID2* and *ID4*) and cytodifferentiation were affected. This work implicates *ADGRG6* in the homeostasis of periodontal ligament tissues, making the R1057Q mutation a strong candidate as a functional cause of the disease.

The mammalian *ADGRG6* gene is widely expressed in the lung (Musa, Cazorla-Vázquez, et al., 2019) and evidence is increasing for the association of *ADGRG6* variants in both coding and intronic regions with different aspects of lung function, including chronic obstructive pulmonary disease (COPD) (Hancock et al., 2010; Jackson et al., 2016; Soler Artigas et al., 2015; Wilk et al., 2012). One of these variants, resulting in the missense mutation S123G, was also identified in Andean people living at altitude, suggesting it may be involved in the physiological adaptation to hypobaric hypoxia

TABLE 2 List of key *ADGRG6* coding variants associated with human disease conditions

Disease/condition	Amino acid	Domain	Mutation (SNP reference)	Reference
LCCS9	R7*	SP	Nonsense homozygous	(Ravenscroft et al., 2015)
LCCS9	Q716T, fs*16	GAIN	Duplication c2144dup homozygous	(Ravenscroft et al., 2015)
LCCS9	V769E	GAIN	Missense homozygous	(Ravenscroft et al., 2015)
Intellectual disability	W1088C	TM6–7 extracellular loop	Missense homozygous	(Hosseini et al., 2018)
Periodontitis	R1057Q	TM5–6 intracellular loop	Missense (rs536714306)	(Kitagaki et al., 2016)
COPD	S123G, K230Q	CUB, PTX	Missense (rs17280293)	(Terzikhan et al., 2018)
Hypobaric hypoxia adaptation	S123G	CUB	Missense (rs17280293)	(Eichstaedt et al., 2017)
Pulmonary function	K230Q	PTX	Missense (rs11155242)	(Hancock et al., 2010)

Note: Amino acid sequence and numbering are based on the reference sequence NM_198569.3. Italic text indicates amino acid identity or similarity with the zebrafish protein.

Abbreviations: COPD, chronic obstructive pulmonary disease; fs, frame shift. See Figure 2 for domain name abbreviations.

(Eichstaedt et al., 2017). A separate analysis of this variant found that expression of *ADGRG6* in human lung tissue was decreased in COPD patients and in individuals with decreased pulmonary ventilatory function, suggesting this variant may also be functional (Terzikhan et al., 2018).

ADGRG6 variants and transcriptional changes have also been associated with various human cancers. A transcriptomic analysis of 772 GPCRs in 148 acute myeloid leukemia (AML) samples, encompassing different subgroups, identified *ADGRG6* as one of 19 down-regulated GPCRs, with others including the closely related *ADGRG1* and *SMO* (Maiga et al., 2016). However, *ADGRG6* was up-regulated in specific MLL translocations, suggesting *ADGRG6* as a candidate disease-specific therapeutic target in these sub-groups (Maiga et al., 2016). The *ADGRG6* locus also contains a highly mutable palindromic motif in intron 6 that is found in 2.7% of breast cancers (Nik-Zainal et al., 2016) and in more than 45% of bladder cancers (Garinet et al., 2019). *ADGRG6* is expressed in the bladder and mutations in the intron 6 motif correlated with an increase in *ADGRG6* expression and a poor prognosis in one study (Wu et al., 2019). Increased *ADGRG6* expression also correlated with an increase in tumor angiogenesis (Wu et al., 2019), supporting previous evidence that *ADGRG6* may have a role more generally in angiogenesis via the VEGF signaling pathway (Cui et al., 2014; Stehlik et al., 2004).

3.3 | Role of *ADGRG6* in the skeleton

Noncoding variants at the human *ADGRG6* locus are associated with different aspects of musculoskeletal development, including height (Soranzo et al., 2009; J. Zhao et al., 2010) and adolescent idiopathic scoliosis (AIS) (Karner et al., 2015; Kou et al., 2013, 2018; Qin et al., 2017; J. F. Xu et al., 2015). AIS is very common, affecting up to 3% of children in the UK (Lenssinck et al., 2005). In mice, *Adgrg6* is expressed in chondrogenic lineages of the axial skeleton (Liu et al., 2019). *Adgrg6* knock-out mice are shorter than their WT littermates and have spinal abnormalities (Monk et al., 2011). Although this suggests a functional role in skeletal development, scoliosis can occur secondary to neuropathy or can be co-morbid with joint contractures, as is found in AMC (Ravenscroft et al., 2015).

To address whether skeletal defects are the underlying cause of the AIS phenotype in *Adgrg6* mutants, two studies used tissue-specific deletion of *Adgrg6* in mice (Karner et al., 2015; Sun et al., 2020). Conditional knock-out of *Adgrg6* in osteoblasts did not result in scoliosis; instead, mutant mice showed delayed ossification and reduced growth (Sun et al., 2020), supporting a role for *ADGRG6* in influencing body length. This study identified Collagen IV as the *Adgrg6* ligand and found that treatment with forskolin could partially rescue the phenotype (Sun et al., 2020). In contrast to the deletion of *Adgrg6* in osteoblasts, deletion of *Adgrg6* in chondrocyte lineages of the spine in mice resulted in abnormalities of the intervertebral disc (IVD), including mechanical stiffening and eventual disc herniation, or scoliosis, depending on the transgenic driver used (Karner et al., 2015; Liu

et al., 2019). Expression of *Stat3*, a gene known to be upregulated in disc degeneration and osteoarthritis, was increased (Liu et al., 2019). A STAT3 inhibitor used to treat osteoarthritis protected against the defect, suggesting that *ADGRG6* is a potential therapeutic target for IVD degeneration. Other studies also implicate *Adgrg6* in AIS pathology (Xu et al., 2019; Xu, Lin, et al., 2019), but analysis of the *ADGRG6* transcript expression profile in both normal and AIS patient tissue is needed to confirm these results. In the zebrafish, *adgrg6* is expressed in the developing craniofacial and axial skeleton (Geng et al., 2013; Glenn & Talbot, 2013). Delayed ossification and a slight reduction in body length has also been reported in zebrafish *adgrg6* morphants (Kou et al., 2013); however, a difference in body length has not been reported in any of the zebrafish *adgrg6* mutant alleles. Further work is needed to determine if *Adgrg6* plays a role in body length and IVD development in fish.

3.4 | Role of *Adgrg6* in remyelination and regeneration

Unlike mammals, zebrafish are able to regenerate a wide range of adult tissues following injury, including the spinal cord, fin and heart (reviewed in Cigliola, Becker, & Poss, 2020; Roehl, 2018; Sanz-Morejón & Mercader, 2020). To date, most studies of *Adgrg6* have focused on a role for the receptor in the initial development of Schwann cell myelination, rather than the maintenance or repair of myelination at adult stages (Glenn & Talbot, 2013). However, expression of *Adgrg6* is maintained and required in adult mammalian and zebrafish Schwann cells (Mogha et al., 2013; Monk et al., 2009), raising the possibility of a role for the receptor in peripheral nerve remyelination and regeneration following injury. Mogha et al. (2016) generated a Schwann-cell-specific, tamoxifen-inducible *Adgrg6* knock-out mouse model and performed a crush injury of the sciatic nerve. Twenty-one days after nerve crush, the tamoxifen-injected mice showed significantly impaired sciatic nerve remyelination and persistent myelin debris, in comparison with control-injected animals (Mogha et al., 2016). These findings suggested that *Adgrg6* is needed autonomously in mammalian Schwann cells for remyelination, but is also required non-autonomously for recruitment of macrophages to clear damaged tissue (Mogha et al., 2016). Furthermore, length measurements of the longest regenerating axons (from the crush site) revealed that axon regeneration is greatly impaired in mutant mice compared with controls, proposing another non-autonomous role for *Adgrg6* in peripheral axon regeneration following nerve injury (Mogha et al., 2016). *Adgrg6* is both expressed and required in non-myelinating terminal Schwann cells for re-innervation following nerve crush at the neuromuscular junction, where it also has a non-cell-autonomous role in the immune response after injury (Jablonka-Shariff, Lu, Campbell, Monk, & Snyder-Warwick, 2020). In the zebrafish, a study using the *stl47* allele suggests that *Adgrg6* is not required for the initial changes in Schwann cell morphology in response to peripheral motor nerve transection (Ducommun Priest, Navarro, Bremer, & Granato, 2019). However, a full analysis of nerve regeneration in zebrafish *adgrg6* mutants has not yet been performed, to our knowledge.

4 | ADGRG6 AND DRUG DISCOVERY

4.1 | GPCRs as druggable targets

Compounds targeting GPCRs constitute a major class in the global market share of therapeutic drugs, reflecting the diverse roles of these receptors in cellular physiology, accessibility at the cell surface, and the presence of binding pockets within their structure (Manglik & Kruse, 2017). Adhesion GPCRs, however, are not represented as targets for licensed drugs, but offer similar potential, given their role in human disease. There are many challenges for aGPCR drug discovery, in part due to the multimodal nature of aGPCRs, which have functions in different tissues, and the possibility of polypharmacology, due to structural similarities between aGPCRs, with any drug identified potentially acting on multiple aGPCRs. In particular, the gedunin class of partial agonists can also act on multiple receptors (Stoveken, Larsen, Smrcka, & Tall, 2018).

Knowledge of the structure of the Adgrg6 ECR (Leon et al., 2020) should facilitate drug design; however, the structure of the CTF, and in particular of the pocket where the *Stachel* peptide binds, are still uncharacterised. The Adgrg6 ECR has structural similarities with the extendable ECR of the epidermal growth factor receptor (EGFR) (Leon et al., 2020). Cetuximab, an anti-cancer monoclonal antibody drug, targets the ECR of EGFR, preventing its extension to the active isoform (Li et al., 2005). Leon et al. (2020) speculate that the dynamic Adgrg6 S1 ECR may be druggable through an equivalent approach to modulate mechanosensory and/or signal transducing functions. Although there are challenges for targeting ADGRG6, small molecule modulators have potential for therapeutic use against many ADGRG6-disease linked conditions, including cancers where ADGRG6 overexpression may contribute to the disease (see Section 3).

4.2 | Small modulators of GPCRs as biological tools

In addition to therapeutics, chemical ligands that bind directly with the Adgrg6 protein could provide valuable tools to stabilize the receptor in an active or inactive conformation, which could facilitate isolation of the receptor in its in vivo conformation, or for the manipulation of pathway activity. Agonist and antagonist compounds are widely utilized as pharmacological tools to modulate the activity of key developmental signaling pathways, including those of the Hedgehog (Hh) and Wnt proteins, which signal via Smoothed and Frizzled family GPCRs, respectively (reviewed in Agostino & Pohl, 2020). A precedent for the translation of small molecules originally identified through developmental biological research to the clinic is the Smoothed inhibitor vismodegib, now a treatment for basal cell carcinoma (Ingham, 2018).

4.3 | Zebrafish screens for small molecule modulators of Adgrg6 pathway activity

Zebrafish present an excellent whole-animal model for precision medicine and are amenable to medium- and high-throughput small

molecule screening approaches (reviewed in Baxendale, van Eeden, & Wilkinson, 2017; Lam & Peterson, 2019). The multimodal nature of aGPCRs and the potential for polypharmacology present problems for traditional in vitro drug discovery pipelines. Adhesion GPCR function is very likely to be context-dependent, due to, for example, interactions with the extracellular matrix and mechanosensing functions. In vivo screening naturally provides these physiological contexts, which are less likely to be recapitulated in an in vitro cell line-based screening platform. The use of zebrafish drug screening approaches also has the advantage of eliminating compounds with toxic or off-target effects in the primary screen.

Several features have made the zebrafish Adgrg6 pathway particularly advantageous for interrogation with small-molecule-based screens. Firstly, hypomorphic alleles—those with weak phenotypes—are amenable to modulation in two directions: rescue of the phenotype or its further exacerbation (Figure 3). Hypomorphs may also exhibit higher sensitivity to small-molecule-centered therapeutic approaches, as a weaker phenotype can sometimes be rescued more easily. Secondly, the available zebrafish *adgrg6* mutants are homozygous viable, allowing batches of 100% mutant embryos to be produced for screening assays. Thirdly, clear mRNA or transgene expression changes in the mutant have enabled the development of robust and reliable screening assays. Finally, the comparison of both hypomorphic and strong alleles in secondary screens has allowed the differentiation of different classes of hit compounds.

Two different screens have used these advantages in their approach (Bradley et al., 2019; Diamantopoulou et al., 2019). Using the expression of *versican* mRNA in the mutant ear as a phenotypic readout of Adgrg6 pathway activity, Diamantopoulou et al. (2019) identified several classes of small molecules that could rescue the otic phenotype in *adgrg6^{tb233c}* hypomorphic mutants, some of which also rescued the myelination defect (Figure 3). The hit compounds included colforsin (a forskolin derivative that raises cAMP levels), a group of dihydropyridines, and a cluster of gedunin derivatives (Diamantopoulou et al., 2019), some of which were independently identified as modulators of human ADGRG1 (Stoveken et al., 2018). Whereas colforsin was able to rescue both hypomorphic (*tb233c*) and strong (*fr24*) alleles, validating association of Adgrg6 to Gαs, other hit compounds were ineffective at rescuing the *fr24* allele (Diamantopoulou et al., 2019) (Figure 3). This latter class are of particular interest, as they may act directly at the level of the receptor. Further work will be needed to test these compounds in cell-based cAMP or Ca²⁺ immobilization assays to determine direct agonism for Adgrg6 (Liebscher, Ackley, et al., 2014; Lizano et al., 2021).

Bradley et al. (2019) monitored changes in *mbp*-driven transgene expression in Schwann cells for their compound screen in *adgrg6^{st63}* hypomorphic mutants. They identified apomorphine hydrochloride, a dopamine receptor agonist, for its ability to induce upregulation of *mbp*-driven GFP in *adgrg6* mutants and to mediate an increase in cAMP levels in Adgrg6-expressing cells (Bradley et al., 2019). Differences in the hit molecules identified between this and the Diamantopoulou et al. (2019) screen are likely to reflect the different commercially-available compound collections that were used, together with differences in the alleles used for




FIGURE 3 Design and proof-of-principle for a whole-animal compound screen to identify agonists of the zebrafish *Adgrg6* pathway. (a) Overview of screen pipeline. Batches of 100% homozygous embryos are dispensed into 96-well plates, with three embryos (biological replicates) per well. Embryos are incubated in control and test compounds from a library of choice at an optimum time for rescue of the mutant phenotype. Effects of compounds on gene expression are measured by semi-automated in situ hybridisation. (b) Example results from a two-part in situ hybridization screen using the hypomorphic (*adgrg6*^{tb233c-/-}) allele. Here, an example hit compound (nifedipine, a dihydropyridine) reduces *vcanb* expression in the ear (top panels) and restores *mbpa* expression in Schwann cells of the lateral line nerves in the vicinity of the ear. Several areas of *mbpa* expression are rescued, in particular that associated with the posterior lateral line ganglion (pll-g, arrowhead). Lateral views; anterior to the left, dorsal to the top. Abbreviations: allg, anterior lateral line ganglion; c, expression associated with the sensory cristae of the ear; DMSO, dimethyl sulfoxide; dpf, days post fertilization; io, infraorbital ramus of the anterior lateral line nerve; pll-d, pll-v, dorsal and ventral rami of the posterior lateral line nerve; pll-g, posterior lateral line ganglion. Scale bars, 50 μ m (top row); 50 μ m (bottom row). (c) Use of hypomorphic and strong alleles to differentiate hit compound classes in the *vcanb* expression assay. Effects of example compounds are shown. Colforsin, a forskolin derivative, rescues both alleles efficiently, and is likely to act downstream of the pathway to raise cAMP levels. Carapin-8(9)-ene rescues the hypomorph, but has no effect on the stronger *fr24* allele. Compounds such as this may interact directly with the *Adgrg6* receptor. Scale bar, 50 μ m. Images reproduced from Diamantopoulou et al., 2019

screening and details of the screening assay design, including compound concentration and exposure time. Clearly, each study has yielded a rich resource of material for validation and further study. A novel approach that could provide further support for the action of apomorphine hydrochloride or other candidate compounds as *Adgrg6* agonists is to determine coupling of mini-G proteins to *Adgrg6* following treatment (Carpenter & Tate, 2016; Nehmé et al., 2017). Such experiments would provide a direct measure of *Adgrg6* activity following small molecule treatments, as opposed to the detection of downstream effector molecules, such as cAMP, that can be modulated by potential compound agonism on adenylyl cyclase.

Taken together, the research findings illustrate a space for multi-faceted drug screening approaches that combine zebrafish phenotypic strategies with in vitro cell-based assays to overcome the individual limitations of each in identifying aGPCR modulators. The availability of tailored technologies such as automated imaging for the zebrafish

larva is improving the throughput of zebrafish small molecule screens (Bradley et al., 2019; Early et al., 2018). However, the lower-cost semi-quantitative in situ hybridization approach also remains effective (Diamantopoulou et al., 2019).

5 | CONCLUSIONS AND FUTURE PERSPECTIVES

Recent research on *Adgrg6* has led to a better understanding of this complex receptor, including its structure, mechanism of action, and roles in development and disease. Work in the zebrafish model has made significant contributions to this research effort. Nevertheless, many questions remain. For instance, we still need to know more about upstream regulators, downstream targets, ligand-receptor binding and the different modes of signaling mediated by the tethered agonist and the NTF. However, there are many areas for future

Adgrg6 research that play to zebrafish strengths in imaging, chemical screening, and gene editing.

One area where recent advances in technology can contribute is to understand the role that different *adgrg6* splice forms play in different developing tissues and how they lead to Adgrg6 protein diversity. At least four splice variants have been reported for the human gene, and it is likely RNAseq data will identify more. Determining when and where different isoforms are expressed, and finding any changes in expression in disease situations, is challenging. Use of single molecule fluorescent in situ hybridization (smFISH) techniques, which are being employed successfully in zebrafish (Soto et al., 2020; Stapel et al., 2016) will enable a detailed analysis of different splice forms and provide tissue-specific cellular localization of the transcripts.

Another area requiring further investigation is the potential role of Adgrg6 as a mechano-sensor, and whether mechanosensation is a key feature of Adgrg6 function in different tissues. It is interesting to speculate that Adgrg6-mediated mechanosensing might have a similar role in the outgrowth and fusion of epithelial projections in the developing ear, as in skeletal development or cardiac trabeculation. Many similar genes and ECM molecules are expressed in all three organ systems and all are subject to mechanical forces. However, both heart and cartilage formation are thought to require the Adgrg6 NTF and do not appear to signal through the canonical cAMP pathway, whereas a role for the NTF alone has not been determined in the inner ear. Zebrafish are well suited to studying the mechanical contributions to tissue morphogenesis in vivo; a recent relevant example tackles the contribution of tension heterogeneity to cardiac trabeculation (Priya et al., 2020). The development and use of tools for live imaging and the measurement of mechanical forces in the zebrafish embryo are pushing boundaries in this important area (Serwane et al., 2017; Shah et al., 2019; Tlili et al., 2019; Tsai et al., 2020).

The pharmacological tools available for aGPCRs in general and Adgrg6 in particular are currently very limited. Hit compounds identified from existing or new chemical screens could be used to target different aspects of Adgrg6 biology using zebrafish in vivo assays, and could be developed into specific small molecule modulators of the receptor. Here, existing studies have already provided promising starting points, such as the gedunin class of molecules (Diamantopoulou et al., 2019; Stoveken et al., 2018). Virtual screening and in silico approaches used in conjunction with the existing and future receptor structures will also be important in identifying novel Adgrg6-binding molecules.

Optimization of CRISPR/Cas9 technologies for zebrafish continues apace, with recent success in high-efficiency targeted integration through homology-directed repair using DNA or single-stranded oligonucleotide templates (Prykhozij et al., 2018; Wierison et al., 2020) or by direct base editing (see, for example, Rosello et al., 2021; Zhao, Shang, Ying, Cheng, & Zhou, 2020). Such approaches, as already exemplified by targeted disruption of the zebrafish Adgrg6 Ca²⁺-binding site (Leon et al., 2020), can be used to introduce specific human disease mutations into the zebrafish gene or to create a humanized Adgrg6 gene in zebrafish, which will provide a valuable resource for ADGRG6 disease modeling and drug validation. Together, the multitude of

approaches possible in the zebrafish offers great promise for furthering our understanding of the Adgrg6 signaling pathway, its roles in the developing embryo, and its significance for human disease.

CONFLICT OF INTEREST

GRW is an employee and shareholder of Sosei Heptares. The other authors declare no competing interests.

AUTHOR CONTRIBUTIONS

Sarah Baxendale, Anzar Asad, and Tanya T. Whitfield wrote the article, with contributions from Nahal O. Shahidan, and Giselle R. Wiggin. Anzar Asad, Sarah Baxendale, and Tanya T. Whitfield prepared the figures.

DATA AVAILABILITY STATEMENT

Data sharing is not applicable to this article as no new data were created or analysed in this study.

ORCID

Sarah Baxendale  <https://orcid.org/0000-0002-6760-9457>

Anzar Asad  <https://orcid.org/0000-0002-3278-9689>

Nahal O. Shahidan  <https://orcid.org/0000-0002-0253-9582>

Giselle R. Wiggin  <https://orcid.org/0000-0003-4436-9208>

Tanya T. Whitfield  <https://orcid.org/0000-0003-1575-1504>

REFERENCES

- Agostino, M., & Pohl, S. Ö.-G. (2020). The structural biology of canonical Wnt signalling. *Biochemical Society Transactions*, 48(4), 1765–1780. <https://doi.org/10.1042/BST20200243>
- Araç, D., Aust, G., Calebiro, D., Engel, F. B., Formstone, C., Goffinet, A., ... Langenhan, T. (2012). Dissecting signaling and functions of adhesion G protein-coupled receptors. *Annals of the New York Academy of Sciences*, 1276(1), 1–25. <https://doi.org/10.1111/j.1749-6632.2012.06820.x>
- Baxendale, S., van Eeden, F., & Wilkinson, R. (2017). The Power of Zebrafish in Personalised Medicine. In El-Khamisy, S. (eds), *Advances in experimental medicine and biology* (Vol. 1007, pp. 179–197). Springer, Cham. https://doi.org/10.1007/978-3-319-60733-7_10
- Beliu, G., Altrichter, S., Guixà-González, R., Hemberger, M., Brauer, I., Dahse, A.-K., ... Langenhan, T. (2021). Tethered agonist exposure in intact adhesion/class B2 GPCRs through intrinsic structural flexibility of the GAIN domain. *Molecular Cell*, 81, 905–921. <https://doi.org/10.1016/j.molcel.2020.12.042>
- Bjarnadóttir, T. K., Geirardsdóttir, K., Ingemansson, M., Mirza, M. A. I., Fredriksson, R., & Schiöth, H. B. (2007). Identification of novel splice variants of adhesion G protein-coupled receptors. *Gene*, 387(1–2), 38–48. <https://doi.org/10.1016/j.gene.2006.07.039>
- Bondarev, A. D., Attwood, M. M., Jonsson, J., Chubarev, V. N., Tarasov, V. V., & Schiöth, H. B. (2020). Opportunities and challenges for drug discovery in modulating adhesion G protein-coupled receptor (GPCR) functions. *Expert Opinion on Drug Discovery*, 15(11), 1291–1307. <https://doi.org/10.1080/17460441.2020.1791075>
- Bradley, E. C., Cunningham, R. L., Wilde, C., Morgan, R. K., Klug, E. A., Letcher, S. M., ... Petersen, S. C. (2019). In vivo identification of small molecules mediating Gpr126/Adgrg6 signaling during Schwann cell development. *Annals of the New York Academy of Sciences*, 1456(1), 44–63. <https://doi.org/10.1111/nyas.14233>
- Carpenter, B., & Tate, C. G. (2016). Engineering a minimal G protein to facilitate crystallisation of G protein-coupled receptors in their active

- conformation. *Protein Engineering, Design and Selection*, 29(12), 583–593. <https://doi.org/10.1093/protein/gzw049>
- Cigliola, V., Becker, C. J., & Poss, K. D. (2020). Building bridges, not walls: Spinal cord regeneration in zebrafish. *Disease Models & Mechanisms*, 13(5), dmm044131. <https://doi.org/10.1242/dmm.044131>
- Cui, H., Wang, Y., Huang, H., Yu, W., Bai, M., Zhang, L., ... Liu, M. (2014). GPR126 protein regulates developmental and pathological angiogenesis through modulation of VEGFR2 receptor signaling. *Journal of Biological Chemistry*, 289(50), 34871–34885. <https://doi.org/10.1074/jbc.M114.571000>
- D'Amato, G., Luxán, G., del Monte-Nieto, G., Martínez-Poveda, B., Torroja, C., Walter, W., ... de la Pompa, J. L. (2016). Sequential notch activation regulates ventricular chamber development. *Nature Cell Biology*, 18(1), 7–20. <https://doi.org/10.1038/ncb3280>
- Demberg, L. M., Rothmund, S., Schöneberg, T., & Liebscher, I. (2015). Identification of the tethered peptide agonist of the adhesion G protein-coupled receptor GPR64/ADGRG2. *Biochemical and Biophysical Research Communications*, 464(3), 743–747. <https://doi.org/10.1016/j.bbrc.2015.07.020>
- Diamantopoulou, E., Baxendale, S., de la Vega de León, A., Asad, A., Holdsworth, C. J., Abbas, L., ... Whitfield, T. T. (2019). Identification of compounds that rescue otic and myelination defects in the zebrafish *adgrg6* (*gpr126*) mutant. *eLife*, 8, 8. <https://doi.org/10.7554/eLife.44889>
- Ducommun Priest, M., Navarro, M. F., Bremer, J., & Granato, M. (2019). Dynein promotes sustained axonal growth and Schwann cell remodeling early during peripheral nerve regeneration. *PLoS Genetics*, 15(2), e1007982. <https://doi.org/10.1371/journal.pgen.1007982>
- Early, J. J., Cole, K. L. H., Williamson, J. M., Swire, M., Kamadurai, H., Muskavitch, M., & Lyons, D. A. (2018). An automated high-resolution in vivo screen in zebrafish to identify chemical regulators of myelination. *eLife*, 7, 7. <https://doi.org/10.7554/eLife.35136>
- Eichstaedt, C. A., Pagani, L., Antao, T., Inchley, C. E., Cardona, A., Mörseburg, A., ... Kivisild, T. (2017). Evidence of early-stage selection on EPAS1 and GPR126 genes in Andean high altitude populations. *Scientific Reports*, 7(1), 13042. <https://doi.org/10.1038/s41598-017-13382-4>
- El-Brolosy, M. A., Kontarakis, Z., Rossi, A., Kuenne, C., Günther, S., Fukuda, N., ... Stainier, D. Y. R. (2019). Genetic compensation triggered by mutant mRNA degradation. *Nature*, 568(7751), 193–197. <https://doi.org/10.1038/s41586-019-1064-z>
- Garinet, S., Pignot, G., Vacher, S., Le Goux, C., Schnitzler, A., Chemlali, W., ... Bieche, I. (2019). High prevalence of a hotspot of noncoding somatic mutations in intron 6 of GPR126 in bladder cancer. *Molecular Cancer Research*, 17(2), 469–475. <https://doi.org/10.1158/1541-7786.MCR-18-0363>
- Geng, F.-S., Abbas, L., Baxendale, S., Holdsworth, C. J., Swanson, A. G., Slanchev, K., ... Whitfield, T. T. (2013). Semicircular canal morphogenesis in the zebrafish inner ear requires the function of *gpr126* (*lauscher*), an adhesion class G protein-coupled receptor gene. *Development*, 140(21), 4362–4374. <https://doi.org/10.1242/dev.098061>
- Glenn, T. D., & Talbot, W. S. (2013). Analysis of *Gpr126* function defines distinct mechanisms controlling the initiation and maturation of myelin. *Development*, 140(15), 3167–3175. <https://doi.org/10.1242/dev.093401>
- Gregory, L. A., Thielens, N. M., Arlaud, G. J., Fontecilla-Camps, J. C., & Gaboriaud, C. (2003). X-ray structure of the Ca²⁺-binding interaction domain of C1s. Insights into the assembly of the C1 complex of complement. *Journal of Biological Chemistry*, 278(34), 32157–32164. <https://doi.org/10.1074/jbc.M305175200>
- Hancock, D. B., Eijgelsheim, M., Wilk, J. B., Gharib, S. A., Loehr, L. R., Marcianti, K. D., ... London, S. J. (2010). Meta-analyses of genome-wide association studies identify multiple loci associated with pulmonary function. *Nature Genetics*, 42(1), 45–52. <https://doi.org/10.1038/ng.500>
- Henzi, A., Senatore, A., Lakkaraju, A. K. K., Scheckel, C., Mühle, J., Reimann, R., ... Aguzzi, A. (2020). Soluble dimeric prion protein ligand activates *Adgrg6* receptor but does not rescue early signs of demyelination in PrP-deficient mice. *PLoS One*, 15(11), e0242137. <https://doi.org/10.1371/journal.pone.0242137>
- Hosseini, M., Fattahi, Z., Abedini, S. S., Hu, H., Ropers, H., Kalscheuer, V. M., ... Kahrizi, K. (2018). GPR126 : A novel candidate gene implicated in autosomal recessive intellectual disability. *American Journal of Medical Genetics Part A*, 179(1), ajmg.a.40531. <https://doi.org/10.1002/ajmg.a.40531>
- Ingham, P. W. (2018). From Drosophila segmentation to human cancer therapy. *Development*, 145(21), dev168898. <https://doi.org/10.1242/dev.168898>
- Jablonka-Shariff, A., Lu, C., Campbell, K., Monk, K. R., & Snyder-Warwick, A. K. (2020). *Gpr126/Adgrg6* contributes to the terminal Schwann cell response at the neuromuscular junction following peripheral nerve injury. *Glia*, 68(6), 1182–1200. <https://doi.org/10.1002/glia.23769>
- Jackson, V. E., Ntalla, I., Sayers, I., Morris, R., Whincup, P., Casas, J.-P., ... Wain, L. V. (2016). Exome-wide analysis of rare coding variation identifies novel associations with COPD and airflow limitation in MOCS3, IFIT3 and SERPINA12. *Thorax*, 71(6), 501–509. <https://doi.org/10.1136/thoraxjnl-2015-207876>
- Kan, Z., Jaiswal, B. S., Stinson, J., Janakiraman, V., Bhatt, D., Stern, H. M., ... Seshagiri, S. (2010). Diverse somatic mutation patterns and pathway alterations in human cancers. *Nature*, 466(7308), 869–873. <https://doi.org/10.1038/nature09208>
- Karner, C. M., Long, F., Solnica-Krezel, L., Monk, K. R., & Gray, R. S. (2015). *Gpr126/Adgrg6* deletion in cartilage models idiopathic scoliosis and pectus excavatum in mice. *Human Molecular Genetics*, 24(15), 4365–4373. <https://doi.org/10.1093/hmg/ddv170>
- Kishore, A., Purcell, R. H., Nassiri-Toosi, Z., & Hall, R. A. (2016). Stalk-dependent and stalk-independent signaling by the adhesion G protein-coupled receptors GPR56 (ADGRG1) and BAI1 (ADGRB1). *Journal of Biological Chemistry*, 291(7), 3385–3394. <https://doi.org/10.1074/jbc.M115.689349>
- Kitagaki, J., Miyauchi, S., Asano, Y., Imai, A., Kawai, S., Michikami, I., ... Murakami, S. (2016). A putative association of a single nucleotide polymorphism in GPR126 with aggressive periodontitis in a Japanese population. *PLoS One*, 11(8), e0160765. <https://doi.org/10.1371/journal.pone.0160765>
- Kou, I., Takahashi, Y., Johnson, T. A., Takahashi, A., Guo, L., Dai, J., ... Ikegawa, S. (2013). Genetic variants in GPR126 are associated with adolescent idiopathic scoliosis. *Nature Genetics*, 45(6), 676–679. <https://doi.org/10.1038/ng.2639>
- Kou, I., Watanabe, K., Takahashi, Y., Momozawa, Y., Khanshour, A., Grauers, A., ... Ikegawa, S. (2018). A multi-ethnic meta-analysis confirms the association of rs6570507 with adolescent idiopathic scoliosis. *Scientific Reports*, 8(1), 11575. <https://doi.org/10.1038/s41598-018-29011-7>
- Küffer, A., Lakkaraju, A. K. K., Mogha, A., Petersen, S. C., Airich, K., Doucerain, C., ... Aguzzi, A. (2016). The prion protein is an agonistic ligand of the G protein-coupled receptor *Adgrg6*. *Nature*, 536(7617), 464–468. <https://doi.org/10.1038/nature19312>
- Labelle-Dumais, C., Schuitema, V., Hayashi, G., Hoff, K., Gong, W., Dao, D. Q., ... Gould, D. B. (2019). COL4A1 mutations cause neuromuscular disease with tissue-specific mechanistic heterogeneity. *American Journal of Human Genetics*, 104(5), 847–860. <https://doi.org/10.1016/j.ajhg.2019.03.007>
- Lam, P.-Y., & Peterson, R. T. (2019). Developing zebrafish disease models for in vivo small molecule screens. *Current Opinion in Chemical Biology*, 50, 37–44. <https://doi.org/10.1016/j.cbpa.2019.02.005>
- Langenhan, T. (2019). Adhesion G protein-coupled receptors—Candidate metabotropic mechanosensors and novel drug targets. *Basic & Clinical*

- Pharmacology & Toxicology*, 126(S6), 5–16. <https://doi.org/10.1111/bcpt.13223>
- Langenhan, T., Aust, G., & Hamann, J. (2013). Sticky signaling—Adhesion class G protein-coupled receptors take the stage. *Science Signaling*, 6, re3–re3. <https://doi.org/10.1126/scisignal.2003825>
- Lenssinck, M. L. B., Frijlik, A. C., Berger, M. Y., Bierma-Zeinstra, S. M. A., Verkerk, K., & Verhagen, A. P. (2005). Effect of bracing and other conservative interventions in the treatment of idiopathic scoliosis in adolescents: A systematic review of clinical trials. *Physical Therapy*, 85, 1329–1339. <https://doi.org/10.1093/ptj/85.12.1329>
- Leon, K., Cunningham, R. L., Riback, J. A., Feldman, E., Li, J., Sosnick, T. R., ... Araç, D. (2020). Structural basis for adhesion G protein-coupled receptor Gpr126 function. *Nature Communications*, 11(1), 194. <https://doi.org/10.1038/s41467-019-14040-1>
- Li, S., Schmitz, K. R., Jeffrey, P. D., Wiltzius, J. J. W., Kussie, P., & Ferguson, K. M. (2005). Structural basis for inhibition of the epidermal growth factor receptor by cetuximab. *Cancer Cell*, 7(4), 301–311. <https://doi.org/10.1016/j.ccr.2005.03.003>
- Liebscher, I., Ackley, B., Araç, D., Ariestanti, D. M., Aust, G., Bae, B., ... Piao, X. (2014). New functions and signaling mechanisms for the class of adhesion G protein-coupled receptors. *Annals of the New York Academy of Sciences*, 1333(1), 43–64. <https://doi.org/10.1111/nyas.12580>
- Liebscher, I., Schön, J., Petersen, S. C., Fischer, L., Auerbach, N., Demberg, L. M., ... Schöneberg, T. (2014). A tethered agonist within the ectodomain activates the adhesion G protein-coupled receptors GPR126 and GPR133. *Cell Reports*, 9(6), 2018–2026. <https://doi.org/10.1016/j.celrep.2014.11.036>
- Liu, Z., Easson, G. W. D., Zhao, J., Makki, N., Ahituv, N., Hilton, M. J., ... Gray, R. S. (2019). Dysregulation of STAT3 signaling is associated with endplate-oriented herniations of the intervertebral disc in Adgrg6 mutant mice. *PLoS Genetics*, 15(10), e1008096. <https://doi.org/10.1371/journal.pgen.1008096>
- Lizano, E., Hayes, J. L., & Willard, F. S. (2021). A synthetic method to assay adhesion-family G-protein coupled receptors. Determination of the G-protein coupling profile of ADGRG6(GPR126). *Biochemical and Biophysical Research Communications*, 534, 317–322. <https://doi.org/10.1016/j.bbrc.2020.11.086>
- Maiga, A., Lemieux, S., Pabst, C., Lavallée, V.-P., Bouvier, M., Sauvageau, G., & Hébert, J. (2016). Transcriptome analysis of G protein-coupled receptors in distinct genetic subgroups of acute myeloid leukemia: Identification of potential disease-specific targets. *Blood Cancer Journal*, 6(6), e431–e431. <https://doi.org/10.1038/bcj.2016.36>
- Manglik, A., & Kruse, A. C. (2017). Structural basis for G protein-coupled receptor activation. *Biochemistry*, 56, 5628–5634. <https://doi.org/10.1021/acs.biochem.7b00747>
- Meyer, A., & Schartl, M. (1999). Gene and genome duplications in vertebrates: The one-to-four (–to-eight in fish) rule and the evolution of novel gene functions. *Current Opinion in Cell Biology*, 11(6), 699–704. [https://doi.org/10.1016/s0955-0674\(99\)00039-3](https://doi.org/10.1016/s0955-0674(99)00039-3)
- Mogha, A., Benesh, A. E., Patra, C., Engel, F. B., Schoneberg, T., Liebscher, I., & Monk, K. R. (2013). Gpr126 functions in Schwann cells to control differentiation and myelination via G-protein activation. *Journal of Neuroscience*, 33(46), 17976–17985. <https://doi.org/10.1523/JNEUROSCI.1809-13.2013>
- Mogha, A., Harty, B. L., Carlin, D., Joseph, J., Sanchez, N. E., Suter, U., ... Monk, K. R. (2016). Gpr126/Adgrg6 has Schwann cell autonomous and nonautonomous functions in peripheral nerve injury and repair. *The Journal of Neuroscience*, 36(49), 12351–12367. <https://doi.org/10.1523/JNEUROSCI.3854-15.2016>
- Monk, K. R., Naylor, S. G., Glenn, T. D., Mercurio, S., Perlin, J. R., Dominguez, C., ... Talbot, W. S. (2009). A G protein-coupled receptor is essential for Schwann cells to initiate myelination. *Science*, 325(5946), 1402–1405. <https://doi.org/10.1126/science.1173474>
- Monk, K. R., Oshima, K., Jörs, S., Heller, S., & Talbot, W. S. (2011). Gpr126 is essential for peripheral nerve development and myelination in mammals. *Development (Cambridge, England)*, 138(13), 2673–2680. <https://doi.org/10.1242/dev.062224>
- Morgan, R. K., Anderson, G. R., Araç, D., Aust, G., Balenga, N., Boucard, A., ... Mogha, A. (2019). The expanding functional roles and signaling mechanisms of adhesion G protein-coupled receptors. *Annals of the New York Academy of Sciences*, 1456(1), 5–25. <https://doi.org/10.1111/nyas.14094>
- Moriguchi, T., Haraguchi, K., Ueda, N., Okada, M., Furuya, T., & Akiyama, T. (2004). DREG, a developmentally regulated G protein-coupled receptor containing two conserved proteolytic cleavage sites. *Genes to Cells*, 9(6), 549–560. <https://doi.org/10.1111/j.1356-9597.2004.00743.x>
- Musa, G., Cazorla-Vázquez, S., Amerongen, M. J., Stemmler, M. P., Eckstein, M., Hartmann, A., ... Engel, F. B. (2019). Gpr126 (Adgrg6) is expressed in cell types known to be exposed to mechanical stimuli. *Annals of the New York Academy of Sciences*, 1456(1), 96–108. <https://doi.org/10.1111/nyas.14135>
- Musa, G., Srivastava, S., Petzold, J., Cazorla-Vázquez, S., & Engel, F. B. (2019). miR-27a/b is a posttranscriptional regulator of Gpr126 (Adgrg6). *Annals of the New York Academy of Sciences*, 1456(1), 109–121. <https://doi.org/10.1111/nyas.14245>
- Nehmé, R., Carpenter, B., Singhal, A., Strega, A., Edwards, P. C., White, C. F., ... Tate, C. G. (2017). Mini-G proteins: Novel tools for studying GPCRs in their active conformation. *PLoS One*, 12(4), e0175642. <https://doi.org/10.1371/journal.pone.0175642>
- Nik-Zainal, S., Davies, H., Staaf, J., Ramakrishna, M., Glodzik, D., Zou, X., ... Stratton, M. R. (2016). Landscape of somatic mutations in 560 breast cancer whole-genome sequences. *Nature*, 534(7605), 47–54. <https://doi.org/10.1038/nature17676>
- O'Hayre, M., Vázquez-Prado, J., Kufareva, I., Stawiski, E. W., Handel, T. M., Seshagiri, S., & Gutkind, J. S. (2013). The emerging mutational landscape of G proteins and G-protein-coupled receptors in cancer. *Nature Reviews Cancer*, 13(6), 412–424. <https://doi.org/10.1038/nrc3521>
- Okajima, D., Kudo, G., & Yokota, H. (2010). Brain-specific angiogenesis inhibitor 2 (BAI2) may be activated by proteolytic processing. *Journal of Receptors and Signal Transduction*, 30(3), 143–153. <https://doi.org/10.3109/10799891003671139>
- Paavola, K. J., Sidik, H., Zuchero, J. B., Eckart, M., & Talbot, W. S. (2014). Type IV collagen is an activating ligand for the adhesion G protein-coupled receptor GPR126. *Science Signaling*, 7(338), ra76–ra76. <https://doi.org/10.1126/scisignal.2005347>
- Paavola, K. J., Stephenson, J. R., Ritter, S. L., Alter, S. P., & Hall, R. A. (2011). The N terminus of the adhesion G protein-coupled receptor GPR56 controls receptor signaling activity. *Journal of Biological Chemistry*, 286(33), 28914–28921. <https://doi.org/10.1074/jbc.M111.247973>
- Patra, C., Monk, K. R., & Engel, F. B. (2014). The multiple signaling modalities of adhesion G protein-coupled receptor GPR126 in development. *Receptors & Clinical Investigation*, 1(3), 79. <https://doi.org/10.14800/rci.79>
- Patra, C., van Amerongen, M. J., Ghosh, S., Ricciardi, F., Sajjad, A., Novoyatleva, T., ... Engel, F. B. (2013). Organ-specific function of adhesion G protein-coupled receptor GPR126 is domain-dependent. *Proceedings of the National Academy of Sciences of the United States of America*, 110(42), 16898–16903. <https://doi.org/10.1073/pnas.1304837110>
- Petersen, S. C., Luo, R., Liebscher, I., Giera, S., Jeong, S.-J., Mogha, A., ... Monk, K. R. (2015). The adhesion GPCR GPR126 has distinct, domain-dependent functions in Schwann cell development mediated by interaction with Laminin-211. *Neuron*, 85(4), 755–769. <https://doi.org/10.1016/j.neuron.2014.12.057>
- Ping, Y.-Q., Mao, C., Xiao, P., Zhao, R.-J., Jiang, Y., Yang, Z., ... Sun, J.-P. (2021). Structures of the glucocorticoid-bound adhesion receptor GPR97-go complex. *Nature*, 589(7843), 620–626. <https://doi.org/10.1038/s41586-020-03083-w>
- Pogoda, H.-M., Sternheim, N., Lyons, D. A., Diamond, B., Hawkins, T. A., Woods, I. G., ... Talbot, W. S. (2006). A genetic screen identifies genes

- essential for development of myelinated axons in zebrafish. *Developmental Biology*, 298(1), 118–131. <https://doi.org/10.1016/j.ydbio.2006.06.021>
- Priya, R., Allanki, S., Gentile, A., Mansingh, S., Uribe, V., Maischein, H.-M., & Stainier, D. Y. R. (2020). Tension heterogeneity directs form and fate to pattern the myocardial wall. *Nature*, 588(7836), 130–134. <https://doi.org/10.1038/s41586-020-2946-9>
- Prykhodzhiy, S. V., Fuller, C., Steele, S. L., Veinotte, C. J., Razaghi, B., Robitaille, J. M., ... Berman, J. N. (2018). Optimized knock-in of point mutations in zebrafish using CRISPR/Cas9. *Nucleic Acids Research*, 46(17), e102–e102. <https://doi.org/10.1093/nar/gky512>
- Qin, X., Xu, L., Xia, C., Zhu, W., Sun, W., Liu, Z., ... Zhu, Z. (2017). Genetic variant of GPR126 gene is functionally associated with adolescent idiopathic scoliosis in Chinese population. *Spine*, 42(19), E1098–E1103. <https://doi.org/10.1097/BRS.0000000000002123>
- Ravenscroft, G., Nolent, F., Rajagopalan, S., Meireles, A. M., Paavola, K. J., Gaillard, D., ... Laing, N. G. (2015). Mutations of GPR126 are responsible for severe arthrogyposis multiplex Congenita. *The American Journal of Human Genetics*, 96(6), 955–961. <https://doi.org/10.1016/j.ajhg.2015.04.014>
- Roehl, H. H. (2018). Linking wound response and inflammation to regeneration in the zebrafish larval fin. *The International Journal of Developmental Biology*, 62(6–7–8), 473–477. <https://doi.org/10.1387/ijdb.170331hr>
- Rosello, M., Vougnay, J., Czarny, F., Mione, M. C., Concordet, J.-P., Albadri, S., & Del Bene, F. (2021). Precise base editing for the in vivo study of developmental signaling and human pathologies in zebrafish. *eLife*, 10, 65552. <https://doi.org/10.7554/eLife.65552>
- Salzman, G. S., Ackerman, S. D., Ding, C., Koide, A., Leon, K., Luo, R., ... Araç, D. (2016). Structural basis for regulation of GPR56/ADGRG1 by its alternatively spliced extracellular domains. *Neuron*, 91(6), 1292–1304. <https://doi.org/10.1016/j.neuron.2016.08.022>
- Salzman, G. S., Zhang, S., Gupta, A., Koide, A., Koide, S., & Araç, D. (2017). Stachel-independent modulation of GPR56/ADGRG1 signaling by synthetic ligands directed to its extracellular region. *Proceedings of the National Academy of Sciences*, 114(38), 10095–10100. <https://doi.org/10.1073/pnas.1708810114>
- Sanz-Morejón, A., & Mercader, N. (2020). Recent insights into zebrafish cardiac regeneration. *Current Opinion in Genetics & Development*, 64, 37–43. <https://doi.org/10.1016/j.gde.2020.05.020>
- Scholz, N., Langenhan, T., & Schöneberg, T. (2019). Revisiting the classification of adhesion GPCRs. *Annals of the New York Academy of Sciences*, 1456(1), 80–95. <https://doi.org/10.1111/nyas.14192>
- Serwane, F., Mongera, A., Rowghanian, P., Kealhofer, D. A., Lucio, A. A., Hockenbery, Z. M., & Campàs, O. (2017). In vivo quantification of spatially varying mechanical properties in developing tissues. *Nature Methods*, 14(2), 181–186. <https://doi.org/10.1038/nmeth.4101>
- Shah, G., Thierbach, K., Schmid, B., Waschke, J., Reade, A., Hlawitschka, M., ... Huisken, J. (2019). Multi-scale imaging and analysis identify pan-embryo cell dynamics of germlayer formation in zebrafish. *Nature Communications*, 10(1), 5753. <https://doi.org/10.1038/s41467-019-13625-0>
- Soler Artigas, M., Wain, L. V., Miller, S., Kheirallah, A. K., Huffman, J. E., Ntalla, I., ... Tobin, M. D. (2015). Sixteen new lung function signals identified through 1000 genomes project reference panel imputation. *Nature Communications*, 6(1), 8658. <https://doi.org/10.1038/ncomms9658>
- Soranzo, N., Rivadeneira, F., Chinappan-Horsley, U., Malkina, I., Richards, J. B., Hammond, N., ... Deloukas, P. (2009). Meta-analysis of genome-wide scans for human adult stature identifies novel loci and associations with measures of skeletal frame size. *PLoS Genetics*, 5(4), e1000445. <https://doi.org/10.1371/journal.pgen.1000445>
- Soto, X., Biga, V., Kursawe, J., Lea, R., Doostdar, P., Thomas, R., & Papalopulu, N. (2020). Dynamic properties of noise and Her6 levels are optimized by miR-9, allowing the decoding of the Her6 oscillator. *The EMBO Journal*, 39(12), e103558. <https://doi.org/10.15252/embj.2019103558>
- Stapel, L. C., Lombardot, B., Broaddus, C., Kainmueller, D., Jug, F., Myers, E. W., & Vastenhouw, N. L. (2016). Automated detection and quantification of single RNAs at cellular resolution in zebrafish embryos. *Development*, 143(3), 540–546. <https://doi.org/10.1242/dev.128918>
- Stehlik, C., Kroismayr, R., Dorfleutner, A., Binder, B. R., & Lipp, J. (2004). VIGR - a novel inducible adhesion family G-protein coupled receptor in endothelial cells. *FEBS Letters*, 569(1–3), 149–155. <https://doi.org/10.1016/j.febslet.2004.05.038>
- Stoveken, H. M., Hajduczk, A. G., Xu, L., & Tall, G. G. (2015). Adhesion G protein-coupled receptors are activated by exposure of a cryptic tethered agonist. *Proceedings of the National Academy of Sciences*, 112(19), 6194–6199. <https://doi.org/10.1073/pnas.1421785112>
- Stoveken, H. M., Larsen, S. D., Smrcka, A. V., & Tall, G. G. (2018). Gedunin and Khivorin-derivatives are small-molecule partial agonists for adhesion G protein-coupled receptors GPR56/ADGRG1 and GPR114/ADGRG5. *Molecular Pharmacology*, 93(5), 477–488. <https://doi.org/10.1124/mol.117.111476>
- Sun, P., He, L., Jia, K., Yue, Z., Li, S., Jin, Y., ... Luo, J. (2020). Regulation of body length and bone mass by Gpr126/Adgrg6. *Science Advances*, 6(12), eaaz0368. <https://doi.org/10.1126/sciadv.aaz0368>
- Terzikhan, N., Sun, F., Verhamme, F. M., Adams, H. H. H., Loth, D., Bracke, K. R., ... O'Connor, G. T. (2018). Heritability and genome-wide association study of diffusing capacity of the lung. *European Respiratory Journal*, 52(3), 1800647. <https://doi.org/10.1183/13993003.00647-2018>
- Tlili, S., Yin, J., Rupprecht, J.-F., Mendieta-Serrano, M. A., Weissbart, G., Verma, N., ... Saunders, T. E. (2019). Shaping the zebrafish myotome by intertissue friction and active stress. *Proceedings of the National Academy of Sciences*, 116(51), 25430–25439. <https://doi.org/10.1073/pnas.1900819116>
- Tsai, T. Y.-C., Sikora, M., Xia, P., Colak-Champollion, T., Knaut, H., Heisenberg, C.-P., & Megason, S. G. (2020). An adhesion code ensures robust pattern formation during tissue morphogenesis. *Science*, 370(6512), 113–116. <https://doi.org/10.1126/science.aba6637>
- Vakonakis, I., Langenhan, T., Prömel, S., Russ, A., & Campbell, I. D. (2008). Solution structure and sugar-binding mechanism of mouse Latrophilin-1 RBL: A 7TM receptor-attached Lectin-like domain. *Structure*, 16(6), 944–953. <https://doi.org/10.1016/j.str.2008.02.020>
- Vizurraga, A., Adhikari, R., Yeung, J., Yu, M., & Tall, G. G. (2020). Mechanisms of adhesion G protein-coupled receptor activation. *Journal of Biological Chemistry*, 295(41), 14065–14083. <https://doi.org/10.1074/jbc.REV120.007423>
- Waller-Evans, H., Prömel, S., Langenhan, T., Dixon, J., Zahn, D., Colledge, W. H., ... Russ, A. P. (2010). The orphan adhesion-GPCR GPR126 is required for embryonic development in the mouse. *PLoS One*, 5(11), e14047. <https://doi.org/10.1371/journal.pone.0014047>
- Ward, Y., Lake, R., Yin, J. J., Heger, C. D., Raffeld, M., Goldsmith, P. K., ... Kelly, K. (2011). LPA receptor Heterodimerizes with CD97 to amplify LPA-initiated RHO-dependent signaling and invasion in prostate Cancer cells. *Cancer Research*, 71(23), 7301–7311. <https://doi.org/10.1158/0008-5472.CAN-11-2381>
- Whitfield, T. T., Granato, M., van Eeden, F. J., Schach, U., Brand, M., Furutani-Seiki, M., ... Nüsslein-Volhard, C. (1996). Mutations affecting development of the zebrafish inner ear and lateral line. *Development (Cambridge, England)*, 123, 241–254.
- Wierson, W. A., Welker, J. M., Almeida, M. P., Mann, C. M., Webster, D. A., Torrie, M. E., ... Essner, J. (2020). Efficient targeted integration directed by short homology in zebrafish and mammalian cells. *eLife*, 9, 1–25. <https://doi.org/10.7554/eLife.53968>
- Wilk, J. B., Shrine, N. R. G., Loehr, L. R., Zhao, J. H., Manichaikul, A., Lopez, L. M., ... Stricker, B. H. (2012). Genome-wide association studies identify CHRNA5/3 and HTR4 in the development of airflow obstruction. *American Journal of Respiratory and Critical Care Medicine*, 186(7), 622–632. <https://doi.org/10.1164/rccm.201202-0366OC>

- Wu, S., Ou, T., Xing, N., Lu, J., Wan, S., Wang, C., ... Cai, Z. (2019). Whole-genome sequencing identifies ADGRG6 enhancer mutations and FRS2 duplications as angiogenesis-related drivers in bladder cancer. *Nature Communications*, 10(1), 720. <https://doi.org/10.1038/s41467-019-08576-5>
- Xu, E., Lin, T., Jiang, H., Ji, Z., Shao, W., Meng, Y., ... Zhou, X. (2019). Asymmetric expression of GPR126 in the convex/concave side of the spine is associated with spinal skeletal malformation in adolescent idiopathic (scoliosis population) asymmetric expression of GPR126 in the convex/concave side of the spine is associated wi. *European Spine Journal*, 28, 1977–1986. <https://doi.org/10.1007/s00586-019-06001-5>
- Xu, E., Shao, W., Jiang, H., Lin, T., Gao, R., & Zhou, X. (2019). A genetic variant in GPR126 causing a decreased inclusion of exon 6 is associated with cartilage development in adolescent idiopathic scoliosis population. *BioMed Research International*, 2019, 1–8. <https://doi.org/10.1155/2019/4678969>
- Xu, J. F., Yang, G. H., Pan, X. H., Zhang, S. J., Zhao, C., Qiu, B. S., ... Wang, Y. P. (2015). Association of GPR126 gene polymorphism with adolescent idiopathic scoliosis in Chinese populations. *Genomics*, 105(2), 101–107. <https://doi.org/10.1016/j.ygeno.2014.11.009>
- Yates, A. D., Achuthan, P., Akanni, W., Allen, J., Allen, J., Alvarez-Jarreta, J., ... Flicek, P. (2019). Ensembl 2020. *Nucleic Acids Research*, 48(D1), D682–D688. <https://doi.org/10.1093/nar/gkz966>
- Zhao, J., Li, M., Bradfield, J. P., Zhang, H., Mentch, F. D., Wang, K., ... Grant, S. F. A. (2010). The role of height-associated loci identified in genome wide association studies in the determination of pediatric stature. *BMC Medical Genetics*, 11(1), 96. <https://doi.org/10.1186/1471-2350-11-96>
- Zhao, Y., Shang, D., Ying, R., Cheng, H., & Zhou, R. (2020). An optimized base editor with efficient C-to-T base editing in zebrafish. *BMC Biology*, 18(1), 190. <https://doi.org/10.1186/s12915-020-00923-z>

How to cite this article: Baxendale S, Asad A, Shahidan NO, Wiggin GR, Whitfield TT. The adhesion GPCR Adgrg6 (Gpr126): Insights from the zebrafish model. *genesis*. 2021; e23417. <https://doi.org/10.1002/dvg.23417>