



This is a repository copy of *Groups of worldview transformations implied by Einstein's special principle of relativity over arbitrary ordered fields.*

White Rose Research Online URL for this paper:
<http://eprints.whiterose.ac.uk/172887/>

Version: Accepted Version

Article:

Madarász, J.X., Stannett, M. orcid.org/0000-0002-2794-8614 and Székely, G. (2021) Groups of worldview transformations implied by Einstein's special principle of relativity over arbitrary ordered fields. *The Review of Symbolic Logic*. ISSN 1755-0203

<https://doi.org/10.1017/s1755020321000149>

© 2021 Association for Symbolic Logic. This is an author produced version of a paper subsequently published in *Review of Symbolic Logic*. Article available under the terms of the CC-BY-NC-ND license (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Reuse

This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND) licence. This licence only allows you to download this work and share it with others as long as you credit the authors, but you can't change the article in any way or use it commercially. More information and the full terms of the licence here: <https://creativecommons.org/licenses/>

Takedown

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing eprints@whiterose.ac.uk including the URL of the record and the reason for the withdrawal request.



eprints@whiterose.ac.uk
<https://eprints.whiterose.ac.uk/>

Groups of Worldview Transformations Implied by Einstein's
Special Principle of Relativity over Arbitrary Ordered Fields

JUDIT X. MADARÁSZ

Alfréd Rényi Institute of Mathematics

MIKE STANNETT

Department of Computer Science, The University of Sheffield

GERGELY SZÉKELY

Alfréd Rényi Institute of Mathematics & University of Public Service

Abstract. In 1978, Yu. F. Borisov presented an axiom system using a few basic assumptions and four explicit axioms, the fourth being a formulation of the relativity principle; and he demonstrated that this axiom system had (up to choice of units) only two models: a relativistic one in which worldview transformations are Poincaré transformations and a classical one in which they are Galilean. In this paper, we reformulate Borisov's original four axioms within an intuitively simple, but strictly formal, first-order logic framework, and convert his basic background assumptions into explicit axioms. Instead of assuming that the structure of physical quantities is the field of real numbers, we assume only that they form an ordered field. This allows us to investigate how Borisov's theorem depends on the structure of quantities.

We demonstrate (as our main contribution) how to construct Euclidean, Galilean, and Poincaré models of Borisov's axiom system over every non-Archimedean field. We also demonstrate the existence of an infinite descending chain of models and transformation groups in each of these three cases, something that is not possible over Archimedean fields.

As an application, we note that there is a model of Borisov's axioms that satisfies the relativity principle, and in which the worldview transformations are Euclidean isometries. Over the field of reals it is easy to eliminate this model using natural axioms concerning time's arrow and the absence of instantaneous motion. In the case of non-Archimedean fields, however, the Euclidean isometries appear intrinsically as worldview transformations in models of Borisov's axioms and neither the assumption of time's arrow, nor the rejection of instantaneous motion, can eliminate them.

§1. Introduction

In his famous 1905 paper, Einstein (1905) based his special theory of relativity on two explicit postulates: (1) the *principle of relativity*, according to which all inertial observers (coordinate systems) are equivalent; and (2) the *light postulate*, according to which there exists a "stationary" coordinate system in which all light signals travel with the same speed. A few years later, beginning in 1910, Ignatowsky (1910a,b, 1911) attempted to simplify the theory by removing the need for the light postulate. Assuming the principle of relativity, some ideas from electrodynamics, and various hidden assumptions, he deduced that the associated (homogeneous) coordinate system transformations must be Lorentz transformations.

Frank & Rothe (1911) then considered both Einstein's and Ignatowsky's arguments in more detail, identifying four assumptions (rather than two) made by Einstein. They argued not only that the light postulate is unnecessary, but that two more of these assumptions can also be deduced by considering how transformations compose with one another and act on points and lines. Their system was less restrictive than Ignatowsky's, because transformations were shown to split into three distinct classes: Galilean transformations; Lorentz transformations; and a rather unusual class they called "Doppler" transformations (Frank & Rothe, 1911, eqn. 129).

In 1978, Yu. F. Borisov presented the axiom system with which we shall mostly be concerned in this paper. In his axiom system, Borisov presented a few basic background assumptions and four explicit axioms, the fourth being a formulation of the relativity principle. Then he demonstrated that his axiom system had (up to choice of units) only two models: a relativistic one in which worldview transformations are Poincaré transformations (the more general, inhomogeneous, counterparts of Ignatowsky's Lorentz transformations), and a classical one in which they are Galilean (Borisov, 1978) (cf. (Guts, 1982, §10, pp. 60-61)). Gyula Dávid subsequently showed (using a different framework) the existence of a model that also satisfies the relativity principle and in which worldview transformations are Euclidean isometries. He also proved (over the field of reals) a characterization theorem stating that the principle of relativity with some auxiliary assumptions implies that the worldview transformations between inertial observers are either Euclidean isometries, or else Galilean or Poincaré transformations (Dávid, 1990). However, he eliminated the models corresponding to Euclidean isometries by adding a new assumption, that motion from one spatial location to another cannot be instantaneous. Similarly, Euclidean isometries do not appear in Borisov's models because they are eliminated by his assumption (Borisov, 1978, Axiom II) that there is an arrow of time.

All of these studies implicitly assume that coordinates and other physically observable quantities can be represented as values from the real number field (\mathbb{R}), even though this assumption is not well-founded. We have no empirical reason to make this assumption, because all practical measurements yield only rational approximations – even quantum electro-dynamics (QED), widely regarded as the most precisely tested physical theory so far, only boasts accuracies to around 12 decimal digits (Odom et al., 2006). Indeed, one of our goals in writing this paper is to investigate what properties of numbers are actually needed if we want to model certain theories. For example, two of us have previously shown elsewhere that special relativity can also be modeled over the field of rational numbers (Madarász & Székely, 2013). We will demonstrate, in fact, that special relativity theories defined over *non-Archimedean* fields are fundamentally different to those defined over Archimedean fields like \mathbb{R} , because models featuring Euclidean isometries as worldview transformations cannot be eliminated, neither by introducing Borisov's arrow of time assumption, nor by allowing Dávid's ban on instantaneous motion. Moreover, the group of worldview transformations contains an infinite descending chain of proper worldview transformation subgroups – this is in sharp contrast to the situation when the field is Archimedean, where no such descending chains are possible.

TRANSFORMATION GROUPS IMPLIED BY THE RELATIVITY PRINCIPLE 3

These results argue that non-standard analysis has an important role to play in the study of physical theories, since they show that one cannot rule out – on purely empirical grounds – the possibility that infinitesimals and other non-standard values are physically relevant; they provide strong confirmatory evidence that the common preference for using real numbers to represent scalars and coordinates is one of convention rather than necessity. While a few authors have considered the use of non-standard analysis in mathematical physics and the study of stochastic processes (e.g., Albeverio et al. (1986); Nelson (1987); Albeverio (1987); Bagarello & Valenti (1988); Capinski & Cutland (1995); Perlis (2016)), most work in the field has understandably focussed on applications within mathematics itself. Our results suggest that further investigation of non-standard applications to mathematical physics is warranted, not just in terms of its usefulness as a tool for simplifying proofs, but in its own right: if non-standard scalars are taken to be physically meaningful, how does that change our understanding of the world around us?

As this brief summary illustrates, there is long history of research addressing the logical foundations of relativity theory, with each generation of researchers discovering implicit (unstated) assumptions embedded in the work of their predecessors. We believe it is important to avoid hidden assumptions wherever possible, so as to place physical theory on secure logical and strictly mathematical foundations. Our preferred approach is to express assumptions as simple, strictly formal, first-order, explicit axioms, which are then used to support formal proof-driven investigation. This provides the clarity we need for ensuring that hidden assumptions are captured explicitly (cf. (Andréka et al., 2002, §Why FOL?) and (Székely, 2009, §11)), while at the same time making it relatively simple to verify whether the axioms are strong enough to do the job we require of them (Govindarajalulu et al., 2015; Stannett & Németi, 2014).

Contributions. In this paper:

1. We reformulate Borisov’s original four axioms within an intuitively simple, but strictly formal, first-order logic framework, and convert his basic background assumptions into explicit axioms.
2. Instead of assuming that the structure of physical quantities is the field of real numbers, we assume only that they form an ordered field. This allows us to investigate how Borisov’s theorem depends on the structure of quantities.
3. We characterize the groups of worldview transformations corresponding to models of (our reformulation of) Borisov’s axiom system over any ordered field, cf. Theorems 5.10. and 5.12.. Using this characterization, we show that Borisov’s Axiom III is not needed to prove his main theorem, cf. Theorem 4.7..
4. We demonstrate (as our main contribution) how to construct Euclidean, Galilean, and Poincaré models of Borisov’s axiom system over every non-Archimedean field, cf. Theorem 4.5.. In this theorem, we also demonstrate the existence of an infinite descending chain of models and transformation groups in each of these three cases, something that is not possible over Archimedean fields like \mathbb{R} .
5. We show, in the case of non-Archimedean fields, that the Euclidean isometries appear intrinsically as worldview transformations in models of Borisov’s axioms. Neither Borisov’s assumption of time’s arrow, nor Dávid’s rejection of instantaneous motion, can eliminate them.

§2. Basic notations and axioms

2.1. The underlying logical framework Throughout this paper we consider models of the form

$$\mathfrak{M} = \langle \text{Ev}, \text{IM}, \text{IOb}, \text{Q}, +, \cdot, \leq, \text{C}, \text{P} \rangle$$

where the various components are intended to have the following interpretations:

- Ev is a nonempty set of *events*;
- IM is a nonempty set of *inertial motions*;
- IOb is a nonempty set of *inertial observers* (reference systems);
- $\Omega = (\text{Q}, +, \cdot, \leq)$ is a structure of *quantities*;
- $\text{C} \subseteq \text{IOb} \times \text{Ev} \times \text{Q}^4$ is a relation used to express *coordinatization*; and
- $\text{P} \subseteq \text{IM} \times \text{Ev}$ is a relation used to express *participation*.

In other words, we use the following many-sorted first-order logic language:

- Ev, IM, IOb, and Q are four sorts representing different kinds of basic entities (events, inertial motions, inertial observers, and quantities);
- +, ·, and ≤ are the usual operation symbols and ordering relation, defined on the sort Q;
- C is a relation of sort $\text{IOb} \times \text{Ev} \times \text{Q}^4$, where the expression $\text{C}(k, e, \vec{p})$ represents the idea that *inertial observer* $k \in \text{IOb}$ *coordinatizes* (“sees”) *event* $e \in \text{Ev}$ at *coordinate point* $\vec{p} \in \text{Q}^4$; and
- P is a relation of sort $\text{IM} \times \text{Ev}$, where the expression $\text{P}(i, e)$ represents the idea that *inertial motion* $i \in \text{IM}$ *participates in event* $e \in \text{Ev}$, or in other words worldline of inertial motion i contains event e .

We have chosen this formal language in accordance with Borisov’s choice of basic concepts, but using some notations of the Andr eka–N emeti school to make it easier to connect the results of this paper to the school’s general project of logic based axiomatic foundations of relativity theories. Axiomatic approaches to relativity theory have extensive literature, see e.g., Andr eka et al. (2006). Most of these approaches use entirely different basic concepts, and hence it is not at all straightforward to check whether these different axioms systems capture the same physical theory. By showing the connection between two radically different approaches, Andr eka & N emeti (2014) takes an important first step in bringing these sporadic axiom systems together. See Friend (2015) and Friend & Molinini (2015) for general discussions of Andr eka–N emeti school’s project and methodology from points of view of epistemological significance and role in scientific explanation.

2.2. Borisov’s basic assumptions In his paper, Borisov declares four axioms (Axiom I–Axiom IV), which are in turn based upon a number of basic assumptions, which we here refer to as BA 1–BA 4.

BA 1 $\Omega = (\text{Q}, +, \cdot, \leq)$ is an ordered field in the sense of abstract algebra.¹

¹ In (Borisov, 1978), Ω is assumed to be the field of real numbers.

BA 2 For every inertial observer $k \in \text{IOb}$, the coordinatization² of k is a bijection $C_k: \text{Ev} \rightarrow \mathbb{Q}^4$.

Given inertial observer $k \in \text{IOb}$, the binary relation $C_k \subseteq \text{Ev} \times \mathbb{Q}^4$ between events and coordinate points described in BA 2 can be defined by

$$C_k(e, \vec{p}) \stackrel{\text{def}}{\iff} C(k, e, \vec{p}).$$

By BA 2, we can introduce the *worldview transformations*, f_{kh} , between inertial observers h and k as the composition of bijections C_k and C_h^{-1} , i.e.

$$f_{kh} \stackrel{\text{def}}{=} C_k \circ C_h^{-1}: \mathbb{Q}^4 \rightarrow \mathbb{Q}^4.$$

Again by BA 2, these worldview transformations are bijections from \mathbb{Q}^4 to \mathbb{Q}^4 , for which

$$f_{mh} = f_{mk} \circ f_{kh} \quad \text{and} \quad f_{kh}^{-1} = f_{hk} \quad (1)$$

for all inertial observers $h, k, m \in \text{IOb}$.

Let us define the *worldline*³ of inertial motion $i \in \text{IM}$ as

$$\text{wl}(i) \stackrel{\text{def}}{=} \{e \in \text{Ev}: P(i, e)\},$$

and the *worldline* of inertial motion $i \in \text{IM}$ according to inertial observer $k \in \text{IOb}$ as

$$\text{wl}_k(i) \stackrel{\text{def}}{=} C_k[\text{wl}(i)].$$

By BA 2, worldview transformations map worldlines to worldlines, i.e. for all inertial observers $h, k \in \text{IOb}$ and every inertial motion $i \in \text{IM}$,

$$f_{kh}[\text{wl}_h(i)] = \text{wl}_k(i). \quad (2)$$

The *time component* and *space component* of $\vec{p} = (p_0, p_1, p_2, p_3) \in \mathbb{Q}^4$ are defined respectively as

$$p_t \stackrel{\text{def}}{=} p_0 \quad \text{and} \quad \vec{p}_s \stackrel{\text{def}}{=} (p_1, p_2, p_3).$$

A set ℓ is called a *line* if and only if there are $\vec{p}, \vec{v} \in \mathbb{Q}^4$ with $\vec{v} \neq (0, 0, 0, 0)$ such that $\ell = \{\vec{p} + \lambda \cdot \vec{v}: \lambda \in \mathbb{Q}\}$.

The line ℓ is *vertical* if and only if $\vec{p}_s = \vec{q}_s$ for every $\vec{p}, \vec{q} \in \ell$.

BA 3 For every⁴ vertical line ℓ and inertial observer $k \in \text{IOb}$, there is an inertial motion $i \in \text{IM}$ such that the worldline of i according to k is ℓ , i.e. $\text{wl}_k(i) = \ell$.

The line ℓ is of *finite slope* if and only if there are $\vec{p}, \vec{q} \in \ell$ such that $p_t \neq q_t$.

² In (Borisov, 1978), inertial observers and their coordinatizations are identified.

³ In (Borisov, 1978), inertial motions and their worldlines are identified.

⁴ In our first-order logic language, quantifying over lines can be done by quantifying over pairs of coordinate points.




Fig. 1. Illustrating the concepts of coordinatization, worldlines and worldview transformations, and the relationship between them.

BA4 For every inertial motion $i \in \text{IM}$ and every inertial observer $k \in \text{IOb}$, worldline $wl_k(i)$ is a line of finite slope.

2.3. Borisov's Axioms We say that inertial motion $i \in \text{IM}$ is **stationary** w.r.t. inertial observer k if and only if $wl_k(i)$ is a vertical line. Then inertial observer $h \in \text{IOb}$ is **at rest** w.r.t. inertial observer $k \in \text{IOb}$, if and only if, whenever inertial motion $i \in \text{IM}$ is stationary w.r.t. h , then i is also stationary w.r.t. k .

Axiom 1 There exist inertial observers $k, h \in \text{IOb}$, where h is not at rest w.r.t. k .

Given $\vec{u} = (u_0, \dots, u_{n-1}) \in \mathbb{Q}^n$ and $\vec{v} = (v_0, \dots, v_{n-1}) \in \mathbb{Q}^n$, the **scalar product** of \vec{u} and \vec{v} is defined in the usual way:

$$\vec{u} \cdot \vec{v} \stackrel{\text{def}}{=} u_0 v_0 + \dots + u_{n-1} v_{n-1}.$$

The Euclidean length of certain vectors in \mathbb{Q}^4 may not exist because \mathbb{Q} can be any ordered field, including for example the field of rational numbers in which

TRANSFORMATION GROUPS IMPLIED BY THE RELATIVITY PRINCIPLE 7

square roots are not always defined. To avoid this problem, we use instead the **squared-Euclidean length** of \vec{v} , defined as

$$|\vec{v}|^2 \stackrel{\text{def}}{=} \vec{v} \cdot \vec{v} = v_0^2 + \dots + v_{n-1}^2.$$

We call a transformation $T: \mathbb{Q}^4 \rightarrow \mathbb{Q}^4$ a **trivial transformation** if and only if (a) it takes vertical lines to vertical lines; and (b) it is the composition of a translation and a linear transformation preserving the squared-Euclidean length (or equivalently, preserving the scalar product). We say that transformation $T: \mathbb{Q}^4 \rightarrow \mathbb{Q}^4$ is **orthochronous** if and only if $T(1, 0, 0, 0)_t > T(0, 0, 0, 0)_t$. The set of orthochronous trivial transformations is denoted by Triv^\uparrow .

Axiom II

- a) Given any inertial observer $k \in \text{IOb}$ and any⁵ orthochronous trivial transformation $\varphi \in \text{Triv}^\uparrow$, there is an inertial observer $h \in \text{IOb}$ for which the worldview transformation between h and k is φ , i.e. $f_{kh} = \varphi$.
- b) For all inertial observers $k, h \in \text{IOb}$, if h is at rest w.r.t. k , then $f_{kh} \in \text{Triv}^\uparrow$.

Axiom III

Given any inertial motion $i \in \text{IM}$, any inertial observer $k \in \text{IOb}$, and any orthochronous trivial transformation $\varphi \in \text{Triv}^\uparrow$, there is an inertial motion $i' \in \text{IM}$ such that $\text{wl}_k(i') = \varphi[\text{wl}_k(i)]$.

Axiom IV

Given any inertial observers $k, k', h \in \text{IOb}$, there is an inertial observer $h' \in \text{IOb}$ such that $f_{hh'} = f_{kk'}$.

Overall, then, we capture Borisov's axiom system formally as:

$$\boxed{\text{Borisov's Axioms}} \stackrel{\text{def}}{=} \{\text{BA 1}, \dots, \text{BA 4}, \text{Axiom I}, \dots, \text{Axiom IV}\}.$$

2.4. Axiom IV as a formulation of Einstein's Special Principle of Relativity The principle of relativity can be formalized in several different ways, see e.g., Madarász et al. (2017); Gömöri (2015); Gömöri & Szabó (2015). In Borisov's axiom system, Einstein's principle of relativity is captured by Axiom IV. This is so because in terms of worldview transformations Axiom IV tells that no inertial observer is distinguished by how its worldview can be related to those of other observers. In this section, we are going to explore some equivalent formulations of this central assumption. To do so, for each inertial observer $k \in \text{IOb}$, we define the **worldview of k** to be

$$\mathbb{W}_k \stackrel{\text{def}}{=} \{f_{kh} : h \in \text{IOb}\}.$$

⁵ Trivial transformations are affine ones and hence they can be represented by a 4×4 matrix and a 4-dimensional translation vector. Therefore, in our first-order logic language, quantifying over trivial transformations can be done by quantifying over the 20 quantity parameters representing those transformations.




Fig. 2. Illustrating Axiom IV.

and the set of **worldview transformations** by

$$\mathbb{W} \stackrel{\text{def}}{=} \bigcup_{k \in \text{IOb}} \mathbb{W}_k = \{f_{kh} : k, h \in \text{IOb}\}.$$

Where we wish to emphasize the underlying model \mathfrak{M} on which these constructs are based, we will add the model name as a suffix, and write, e.g., $\mathbb{W}_{\mathfrak{M}}$.




Fig. 3. Illustrating the worldview \mathbb{W}_k of inertial observer k .

We first show (Proposition 2.1.) that Axiom IV is equivalent to the claim that all observers have the same worldview, which is in turn equivalent to saying that each worldview is a group under composition.

PROPOSITION 2.1. *Assume BA2. Then Axiom IV is equivalent to each of the following statements (and they are all equivalent to one another):*

- (i). $\mathbb{W}_k = \mathbb{W}_h$ for all $k, h \in \text{IOb}$.
- (ii). $\mathbb{W}_k = \mathbb{W}$ for all $k \in \text{IOb}$.
- (iii). $\mathbb{W}_k = \mathbb{W}$ for some $k \in \text{IOb}$.
- (iv). \mathbb{W}_k is closed under composition for all $k \in \text{IOb}$.
- (v). \mathbb{W}_k forms a group under composition for some $k \in \text{IOb}$.




Fig. 4. Diagram illustrating the steps proving Proposition 2.1.

Proof. Axiom IV says that, given any $f_{kk'} \in \mathbb{W}_k$, there is an $f_{hh'} \in \mathbb{W}_h$ satisfying $f_{kk'} = f_{hh'}$. So statement (i) is simply a reformulation of Axiom IV in terms of worldviews.

(i) \implies (ii): follows because $\mathbb{W} = \bigcup_{h \in \text{IOb}} \mathbb{W}_h$.

(ii) \implies (iii): trivial as IOb is not empty.

Let us now prove the following:

$$\mathbb{W}_k = \mathbb{W} \implies \mathbb{W}_k \text{ is a group.} \quad (3)$$

We know that $\mathbb{W}_k = \mathbb{W}$ is closed under inverses because $f_{kh}^{-1} = f_{hk}$ by (1) (which follows from BA2). It remains to show that \mathbb{W} is closed under composition. To do so, choose any $f_{dc}, f_{ba} \in \mathbb{W}$. Since $\mathbb{W}_k = \mathbb{W}$, there are observers $a', d' \in \text{IOb}$ such that $f_{ka'} = f_{ba}$ and $f_{kd'} = f_{cd}$. Now (1) yields $f_{dc} \circ f_{ba} = f_{cd}^{-1} \circ f_{ba} = f_{kd'}^{-1} \circ f_{ka'} = f_{d'k} \circ f_{ka'} = f_{d'a'} \in \mathbb{W}$, as required.

(iii) \implies (v): follows trivially from (3).

(v) \implies (i): We have $\mathbb{W}_h = f_{hk} \circ \mathbb{W}_k$ for all $k, h \in \text{IOb}$ since, by (1), $f_{hm} = f_{hk} \circ f_{km}$ for all $m \in \text{IOb}$. Therefore, since $f_{hk} = f_{kh}^{-1}$ by (1), we have $\mathbb{W}_h = f_{hk} \circ \mathbb{W}_k = f_{kh}^{-1} \circ \mathbb{W}_k = \mathbb{W}_k$ for all $h \in \text{IOb}$ if \mathbb{W}_k is a group (as $f_{kh} \in \mathbb{W}_k$).

(ii) \implies (iv): follows trivially from (3).

(iv) \implies (v): Since IOb is not empty, we only have to prove that \mathbb{W}_k is closed under composition and inverses for some $k \in \text{IOb}$. Since, by (iv), \mathbb{W}_k is closed under composition it is enough to show that it is also closed under inverses. To prove that, let $f_{km} \in \mathbb{W}_k$ be arbitrary. We need to show that $f_{mk} = f_{km}^{-1} \in \mathbb{W}_k$. Since \mathbb{W}_m is also closed under composition by (iv), we have $f_{mk} \circ f_{mk} \in \mathbb{W}_m$. Then there is $h \in \text{IOb}$ such that $f_{mk} \circ f_{mk} = f_{mh}$. Then $f_{mk} = f_{mk}^{-1} \circ f_{mk} \circ f_{mk} = f_{km} \circ f_{mk} \circ f_{mk} = f_{km} \circ f_{mh} = f_{kh} \in \mathbb{W}_k$. \square

§3. Transformations

Throughout this section, we assume BA1, i.e. that $\Omega = (\mathbb{Q}, +, \cdot, \leq)$ is an ordered field. Let $c \in \mathbb{Q}$ and suppose $c > 0$.

We define the *c-scalar product* of vectors $\vec{p}, \vec{q} \in \mathbb{Q}^4$ as:

$$\vec{p} \odot_c \vec{q} \stackrel{\text{def}}{=} cp_t \cdot cq_t + \vec{p}_s \cdot \vec{q}_s,$$

and analogously the *c-Minkowski scalar product* as:

$$\vec{p} \diamond_c \vec{q} \stackrel{\text{def}}{=} cp_t \cdot cq_t - \vec{p}_s \cdot \vec{q}_s.$$

(Note that the 1-scalar product coincides with the usual scalar product.) As usual, we write $|v| \stackrel{\text{def}}{=} \max\{-v, v\}$ for the absolute value of $v \in \mathbb{Q}$.

We say that a function $T: \mathbb{Q}^4 \rightarrow \mathbb{Q}^4$ is a *linear c-Euclidean isometry* if and only if it is a linear transformation which preserves the *c-scalar product*, i.e.

$$T\vec{p} \odot_c T\vec{q} = \vec{p} \odot_c \vec{q}$$

for every $\vec{p}, \vec{q} \in \mathbb{Q}^4$. We call T a *c-Euclidean isometry* if and only if it is a composition of a linear *c-Euclidean isometry* and a translation.

We say that T is a *linear Galilean transformation* if and only if T is a linear transformation and, for every $\vec{p}, \vec{q} \in \mathbb{Q}^4$,

$$|(T\vec{p})_t| = |p_t| \quad \text{and} \quad p_t = q_t = 0 \implies T\vec{p} \cdot T\vec{q} = \vec{p} \cdot \vec{q}.$$

T is a *Galilean transformation* if and only if it is a composition of a linear Galilean transformation and a translation.⁶

We call T a *linear c-Poincaré transformation* if and only if it is a linear transformation which preserves the *c-Minkowski scalar product*, i.e.

$$T\vec{p} \diamond_c T\vec{q} = \vec{p} \diamond_c \vec{q},$$

for every $\vec{p}, \vec{q} \in \mathbb{Q}^4$. T is a *c-Poincaré transformation* if and only if it is a composition of a linear *c-Poincaré transformation* and a translation.

In the particular case when $c = 1$, we sometimes revert to the more familiar standard terminology, viz. T is a *Euclidean isometry* if and only if it is a 1-Euclidean isometry, and a *Poincaré transformation* if and only if it is a 1-Poincaré transformation.

Finally, we note that T is a trivial transformation if and only if it is a Euclidean isometry taking vertical lines to vertical lines.

Throughout this paper we investigate the various groups corresponding to these different transformation classes. The notation we use for each of these transformation groups is shown in Table 1.

The orthochronous variants of these sets are respectively denoted by $\text{Triv}_\Omega^\uparrow$, $\text{Eucl}_\Omega^\uparrow$, $\text{Poi}_\Omega^\uparrow$, ${}_c\text{Eucl}_\Omega^\uparrow$, ${}_c\text{Poi}_\Omega^\uparrow$ and $\text{Gal}_\Omega^\uparrow$.

As one would expect, all of these sets (except ${}_c\text{Eucl}_\Omega^\uparrow$) are groups under composition. To see that ${}_c\text{Eucl}_\Omega^\uparrow$ is *not* a group, notice that while it is closed under inverses, it is not closed under composition. For example, for the case $c = 1$, let R be a rotation of 45° in the first two coordinates which leaves the other two coordinates fixed. Then $R \in {}_c\text{Eucl}_\Omega^\uparrow$, but $R \circ R \notin {}_c\text{Eucl}_\Omega^\uparrow$.

⁶ In the literature it is customary to assume that Galilean transformations preserve time orientation, but here we would like to speak also about time reversing Galilean transformations. Hence here it is more natural to introduce them this way.

Table 1. *Transformation group notations used in this paper.*

Triv_Ω	Trivial transformations over Ω
Eucl_Ω	Euclidean isometries over Ω
Poi_Ω	Poincaré transformations over Ω
${}_c\text{Eucl}_\Omega$	c -Euclidean isometries over Ω
${}_c\text{Poi}_\Omega$	c -Poincaré transformations over Ω
Gal_Ω	Galilean Transformations over Ω

Let us also note that

$$\text{Triv}_\Omega = \bigcap_{c>0} {}_c\text{Poi}_\Omega \cap \bigcap_{c>0} {}_c\text{Eucl}_\Omega \cap \text{Gal}_\Omega$$

and hence

$$\text{Triv}_\Omega^\uparrow = \bigcap_{c>0} {}_c\text{Poi}_\Omega^\uparrow \cap \bigcap_{c>0} {}_c\text{Eucl}_\Omega^\uparrow \cap \text{Gal}_\Omega^\uparrow.$$

Let us introduce the following notations for the **squared c -Euclidean length** and **squared c -Minkowski length**:

$$\|\vec{p}\|_c^2 \stackrel{\text{def}}{=} \vec{p} \odot_c \vec{p} = c^2 p_t^2 + |\vec{p}_s|^2 \quad \text{and} \quad \|\vec{p}\|_{c,\mu}^2 \stackrel{\text{def}}{=} \vec{p} \diamond_c \vec{p} = c^2 p_t^2 - |\vec{p}_s|^2$$

We note that squared 1-Euclidean length coincides with the squared-Euclidean length.

Finally, we introduce the following notations for the standard basis of \mathbb{Q}^4 :

$$\vec{e}_T \stackrel{\text{def}}{=} (1, 0, 0, 0), \quad \vec{e}_X \stackrel{\text{def}}{=} (0, 1, 0, 0), \quad \vec{e}_Y \stackrel{\text{def}}{=} (0, 0, 1, 0), \quad \vec{e}_Z \stackrel{\text{def}}{=} (0, 0, 0, 1).$$

PROPOSITION 3.2. *Let L be a linear transformation. Then the following are equivalent:*

- (i). $L \in {}_c\text{Eucl}_\Omega$.
- (ii). *Given any $i, j \in \{T, X, Y, Z\}$,*

$$L\vec{e}_i \odot_c L\vec{e}_j = \vec{e}_i \odot_c \vec{e}_j = \begin{cases} c^2 & \text{if } i = j = T \\ 1 & \text{if } i = j \neq T \\ 0 & \text{if } i \neq j. \end{cases}$$

- (iii). *Given any $\vec{p} \in \mathbb{Q}^4$, $\|L\vec{p}\|_c^2 = \|\vec{p}\|_c^2$, i.e. L preserves the squared c -Euclidean length.*

Proof. (i) \implies (ii) and (i) \implies (iii) hold by definition (this is what we mean by c -Euclidean transformations, c -scalar product, and squared c -Euclidean length).

(ii) \implies (i) follows by direct calculation in the standard basis because L is a linear transformation and $\odot_c: \mathbb{Q}^4 \times \mathbb{Q}^4 \rightarrow \mathbb{Q}$ is a bilinear function.

(iii) \implies (i) follows because \odot_c is symmetric, whence we can write

$$\vec{p} \odot_c \vec{q} = \frac{\|\vec{p} + \vec{q}\|_c^2 - \|\vec{p}\|_c^2 - \|\vec{q}\|_c^2}{2} \quad \text{for every } \vec{p}, \vec{q} \in \mathbb{Q}^4$$

and the claim follows immediately. \square

The proofs of Proposition 3.3. and 3.4. below are analogous and we leave the details to the reader.

PROPOSITION 3.3. *Let L be a linear transformation. Then the following are equivalent:*

- (i). $L \in \text{Gal}_\Omega$.
- (ii). $(L\vec{e}_T)_t = \pm 1$, $(L\vec{e}_X)_t = (L\vec{e}_Y)_t = (L\vec{e}_Z)_t = 0$; and

$$L\vec{e}_i \cdot L\vec{e}_j = \vec{e}_i \cdot \vec{e}_j = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases} \text{ for all } i, j \in \{X, Y, Z\}.$$

- (iii). *Either $(L\vec{p})_t = p_t$ for all $\vec{p} \in \mathbb{Q}^4$ or else $(L\vec{p})_t = -p_t$ for all $\vec{p} \in \mathbb{Q}^4$; and $p_t = 0 \implies |L\vec{p}|^2 = |\vec{p}|^2$. \square*

PROPOSITION 3.4. *Let L be a linear transformation. Then the following are equivalent:*

- (i). $L \in {}_c\text{Poi}_\Omega$.
- (ii). *Given any $i, j \in \{T, X, Y, Z\}$,*

$$L\vec{e}_i \diamond_c L\vec{e}_j = \vec{e}_i \diamond_c \vec{e}_j = \begin{cases} c^2 & \text{if } i = j = T \\ -1 & \text{if } i = j \neq T \\ 0 & \text{if } i \neq j. \end{cases}$$

- (iii). *Given any $\vec{p} \in \mathbb{Q}^4$, $\|L\vec{p}\|_{c,\mu}^2 = \|\vec{p}\|_{c,\mu}^2$, i.e. L preserves the squared c -Minkowskian length. \square*

§4. Theorems

In this section we outline the various theorems that we'll be proving in this paper; the proofs follow in §5. and §6.. Our main result is Theorem 4.5., showing how Borisov's theorem changes when fields other than \mathbb{R} are considered. Whenever we speak about a set G of transformations as a group, we mean the group $\langle G, \circ \rangle$. Recall that if we assume BA2 and Axiom IV, then $\mathbb{W}_k = \mathbb{W}$ for every $k \in \text{IOb}$ and \mathbb{W} is a group (by Proposition 2.1.). Recall also what Borisov's theorem states:

BORISOV'S THEOREM (Borisov, 1978)

Suppose that Borisov's Axioms hold and that Ω is the ordered field \mathbb{R} of reals. Then there are just two possibilities: either

$$\begin{aligned} \mathbb{W} &= \text{Gal}_{\mathbb{R}}^\uparrow, \text{ or else} \\ \mathbb{W} &= {}_c\text{Poi}_{\mathbb{R}}^\uparrow \text{ for some } c > 0. \end{aligned}$$

In the first-order logic framework developed in this paper, it can be shown that Borisov's Theorem remains true if we omit the assumption that Ω is the ordered field \mathbb{R} of reals, and simply assume instead that Ω is an Archimedean ordered field in which every positive number has a square root (we omit the details). It is an open question whether the statement remains true if we omit the assumption that positive numbers have square roots. Note, however, that if we drop the assumption that $\Omega = \mathbb{R}$ without adding anything new, then Borisov's Theorem is no longer valid, because \mathbb{W} can then be a *proper* subgroup of Gal^\uparrow and ${}_c\text{Poi}^\uparrow$. Indeed, \mathbb{W} can even be an orthochronous subgroup of ${}_c\text{Eucl}$.

TRANSFORMATION GROUPS IMPLIED BY THE RELATIVITY PRINCIPLE 13

As usual, we will write $H \leq G$ to mean that H is a subgroup of G , and $H < G$ to indicate that it is a proper subgroup. Analogously, we write $\mathfrak{M} < \mathfrak{M}'$ to mean that \mathfrak{M} is a proper submodel of \mathfrak{M}' .

THEOREM 4.5. (MAIN THEOREM) *Let Ω be any non-Archimedean ordered field. Then, for every $c > 0$, there are models \mathfrak{M}^E , \mathfrak{M}^G and \mathfrak{M}^P of Borisov's Axioms over Ω such that*

$$\begin{aligned} \text{Triv}_{\Omega}^{\uparrow} &< \mathbb{W}_{\mathfrak{M}^E} \subset {}_c\text{Eucl}_{\Omega}^{\uparrow}, \\ \text{Triv}_{\Omega}^{\uparrow} &< \mathbb{W}_{\mathfrak{M}^G} < \text{Gal}_{\Omega}^{\uparrow}, \\ \text{Triv}_{\Omega}^{\uparrow} &< \mathbb{W}_{\mathfrak{M}^P} < {}_c\text{Poi}_{\Omega}^{\uparrow}. \end{aligned}$$

Moreover, in each of these three cases, there is a strictly descending countably infinite chain of models

$$\begin{aligned} \dots &< \mathfrak{M}_i^E < \dots < \mathfrak{M}_1^E < \mathfrak{M}_0^E, \\ \dots &< \mathfrak{M}_i^G < \dots < \mathfrak{M}_1^G < \mathfrak{M}_0^G, \\ \dots &< \mathfrak{M}_i^P < \dots < \mathfrak{M}_1^P < \mathfrak{M}_0^P \end{aligned}$$

over Ω of Borisov's Axioms such that

$$\begin{aligned} \text{Triv}_{\Omega}^{\uparrow} &< \dots < \mathbb{W}_{\mathfrak{M}_i^E} < \dots < \mathbb{W}_{\mathfrak{M}_1^E} < \mathbb{W}_{\mathfrak{M}_0^E} \subset {}_c\text{Eucl}_{\Omega}^{\uparrow}, \\ \text{Triv}_{\Omega}^{\uparrow} &< \dots < \mathbb{W}_{\mathfrak{M}_i^G} < \dots < \mathbb{W}_{\mathfrak{M}_1^G} < \mathbb{W}_{\mathfrak{M}_0^G} < \text{Gal}_{\Omega}^{\uparrow}, \\ \text{Triv}_{\Omega}^{\uparrow} &< \dots < \mathbb{W}_{\mathfrak{M}_i^P} < \dots < \mathbb{W}_{\mathfrak{M}_1^P} < \mathbb{W}_{\mathfrak{M}_0^P} < {}_c\text{Poi}_{\Omega}^{\uparrow}. \end{aligned}$$

□

Note the use of *subset* (\subset) – as opposed to *subgroup* ($<$) – inclusion in the cases involving ${}_c\text{Eucl}_{\Omega}^{\uparrow}$ (and likewise below); this is because, as observed on page 10, ${}_c\text{Eucl}_{\Omega}^{\uparrow}$ is not a group.

THEOREM 4.6. *Let Ω be an ordered field and let G be a group for which either*

$$\begin{aligned} \text{Triv}_{\Omega}^{\uparrow} &< G \subset {}_c\text{Eucl}_{\Omega}^{\uparrow}, \text{ or} \\ \text{Triv}_{\Omega}^{\uparrow} &< G \leq \text{Gal}_{\Omega}^{\uparrow}, \text{ or} \\ \text{Triv}_{\Omega}^{\uparrow} &< G \leq {}_c\text{Poi}_{\Omega}^{\uparrow}. \end{aligned}$$

Then there is a model \mathfrak{M} of Borisov's Axioms over Ω such that $\mathbb{W}_{\mathfrak{M}} = G$. □

We also show that Borisov's Theorem remains true if we remove Axiom III from Borisov's Axioms:

THEOREM 4.7. *Assume Borisov's Axioms \setminus {Axiom III} and that Ω is the ordered field \mathbb{R} of reals. Then there are two possibilities: either*

$$\begin{aligned} \mathbb{W} &= \text{Gal}_{\mathbb{R}}^{\uparrow}, \text{ or else} \\ \mathbb{W} &= {}_c\text{Poi}_{\mathbb{R}}^{\uparrow} \text{ for some } c > 0. \end{aligned}$$

□

Remark 4.8. *If we don't require Ω to be the field \mathbb{R} of reals, then using results in Madarász et al. (2020) the following can be proven: Assume Borisov's Axioms \setminus {Axiom III} and that every positive number has a square root in Ω . Then either*

$$\begin{aligned} & \text{Triv}_\Omega^\uparrow < \mathbb{W} \subset {}_c\text{Eucl}_\Omega^\uparrow, \text{ or} \\ & \text{Triv}_\Omega^\uparrow < \mathbb{W} \leq \text{Gal}_\Omega^\uparrow, \text{ or} \\ & \text{Triv}_\Omega^\uparrow < \mathbb{W} \leq {}_c\text{Poi}_\Omega^\uparrow \text{ for some } c > 0. \end{aligned}$$

Moreover, the statement remains true even if we replace BA 4 with the more general assumption: "For every inertial motion $i \in \text{IM}$ and every inertial observer $k \in \text{IOb}$, $\text{wl}_k(i)$ is a line". It is an open question if they remain true if we omit the assumption that positive numbers have square roots.

The next result, Corollary 4.9., is a direct consequence of Borisov's Theorem and 4.6..

COROLLARY 4.9. *Let Ω be the ordered field \mathbb{R} of reals. Then there is no group G for which*

$$\begin{aligned} & \text{Triv}_\mathbb{R}^\uparrow < G \subset {}_c\text{Eucl}_\mathbb{R}^\uparrow, \text{ or} \\ & \text{Triv}_\mathbb{R}^\uparrow < G < \text{Gal}_\mathbb{R}^\uparrow, \text{ or} \\ & \text{Triv}_\mathbb{R}^\uparrow < G < {}_c\text{Poi}_\mathbb{R}^\uparrow. \end{aligned}$$

□

Corollary 4.9. fails for more general choices of Ω because (by Theorem 4.5.) for all non-Archimedean fields Ω , there are strictly descending countably infinite chains of subgroups such that

$$\begin{aligned} & \text{Triv}_\Omega^\uparrow < \dots < G_i^E < \dots < G_1^E < G_0^E \subset {}_c\text{Eucl}_\Omega^\uparrow, \\ & \text{Triv}_\Omega^\uparrow < \dots < G_i^G < \dots < G_1^G < G_0^G < \text{Gal}_\Omega^\uparrow, \text{ and} \\ & \text{Triv}_\Omega^\uparrow < \dots < G_i^P < \dots < G_1^P < G_0^P < {}_c\text{Poi}_\Omega^\uparrow. \end{aligned}$$

§5. Worldview transformation groups and model constructions

The symmetric group over \mathbb{Q}^4 is denoted by $\text{Sym}(\mathbb{Q}^4)$. The *time-axis* is defined as

$$\mathbf{t} \stackrel{\text{def}}{=} \{(t, 0, 0, 0) \in \mathbb{Q}^4 : t \in \mathbb{Q}\}.$$

THEOREM 5.10. *Assume Borisov's Axioms \setminus {Axiom III}. Then \mathbb{W} is a group satisfying:*

- (i). $\text{Triv}_\Omega^\uparrow < \mathbb{W} \leq \text{Sym}(\mathbb{Q}^4)$;
- (ii). for every $f \in \mathbb{W}$, $f[\mathbf{t}]$ is a line of finite slope;
- (iii). if $f \in \mathbb{W}$ takes vertical lines to vertical lines, then $f \in \text{Triv}_\Omega^\uparrow$.

Proof. (i) We know from Proposition 2.1. that $\mathbb{W} \leq \text{Sym}(\mathbb{Q}^4)$, and Axiom II(a) tells us that $\text{Triv}_\Omega^\uparrow \leq \mathbb{W}$. So it is enough to show that there is some $f \in \mathbb{W} \setminus \text{Triv}_\Omega^\uparrow$. By Axiom I, there are $k, h \in \text{IOb}$ such that h is not at rest w.r.t. k , i.e. there is $i \in \text{IM}$ such that $\text{wl}_h(i)$ is a vertical line but $\text{wl}_k(i)$ is not a vertical line. Since,

TRANSFORMATION GROUPS IMPLIED BY THE RELATIVITY PRINCIPLE 15

by equation (2), f_{kh} takes $\text{wl}_h(i)$ to $\text{wl}_k(i)$ and trivial transformations take vertical lines to vertical ones, we have $f_{kh} \notin \text{Triv}_\Omega$, as required.

(ii) Let $f_{kh} \in \mathbb{W}$. By BA 3, there is $i \in \text{IM}$ such that $\text{wl}_h(i) = \mathbf{t}$. By BA 4, $\text{wl}_k(i)$ is a line of finite slope. Now, by equation (2), $f_{kh}[\mathbf{t}] = f_{kh}[\text{wl}_h(i)] = \text{wl}_k(i)$, as claimed.

(iii) Finally, suppose $f_{kh} \in \mathbb{W}$ takes vertical lines to vertical lines. We will prove that h is at rest w.r.t. k . Let i be an arbitrary inertial motion which is stationary according to h , i.e. $\text{wl}_h(i)$ is a vertical line. Since f_{kh} maps $\text{wl}_h(i)$ to $\text{wl}_k(i)$, $\text{wl}_k(i)$ is also a vertical line. Thus h is at rest w.r.t. k . Hence, by Axiom II(b), $f_{kh} \in \text{Triv}_\Omega^\uparrow$ as required. \square

For every $\Omega = (\mathbb{Q}, +, \cdot, \leq)$ and $G \subseteq \text{Sym}(\mathbb{Q}^4)$, we will construct a model $\mathfrak{M}(G)$ as follows:

$$\begin{aligned} \text{Ev} &\stackrel{\text{def}}{=} \mathbb{Q}^4, & \text{IM} &\stackrel{\text{def}}{=} \{g[\mathbf{t}]: g \in G\}, & \text{IOb} &\stackrel{\text{def}}{=} G, \\ \text{C}(k, e, \vec{p}) &\stackrel{\text{def}}{\iff} k(e) = \vec{p}, & \text{P}(i, e) &\stackrel{\text{def}}{\iff} e \in i, \\ \mathfrak{M}(G) &\stackrel{\text{def}}{=} \langle \text{Ev}, \text{IM}, \text{IOb}, \mathbb{Q}, +, \cdot, \leq, \text{C}, \text{P} \rangle. \end{aligned}$$

PROPOSITION 5.11. *Suppose $G' \subset G \subseteq \text{Sym}(\mathbb{Q}^4)$. Then $\mathfrak{M}(G') < \mathfrak{M}(G)$.*

Proof. The proposition easily follows from the definitions of $\mathfrak{M}(G)$ and $\mathfrak{M}(G')$. \square

THEOREM 5.12. *Suppose Ω is an ordered field and G is a group for which:*

- (i). $\text{Triv}_\Omega^\uparrow < G \leq \text{Sym}(\mathbb{Q}^4)$;
- (ii). *For every $g \in G$, $g[\mathbf{t}]$ is a line of finite slope;*
- (iii). *If $g \in G$ takes vertical lines to vertical lines, then $g \in \text{Triv}_\Omega^\uparrow$.*

Then $\mathfrak{M}(G)$ satisfies Borisov's Axioms. Furthermore, $\mathbb{W}_{\mathfrak{M}(G)} = G$.

Remark 5.13. *Assume that Ω is an ordered field in which every positive number has a square root and that G is a group for which the conditions of Theorem 5.12. hold. Then by Theorem 5.12. and Remark 4.8., either*

$$\begin{aligned} \text{Triv}_\Omega^\uparrow < G &\subset {}_c\text{Eucl}_\Omega^\uparrow, \text{ or} \\ \text{Triv}_\Omega^\uparrow < G &\leq \text{Gal}_\Omega^\uparrow, \text{ or} \\ \text{Triv}_\Omega^\uparrow < G &\leq {}_c\text{Poi}_\Omega^\uparrow \text{ for some } c > 0. \end{aligned}$$

\square

Proof of Thm. 5.12.. It is easy to see that $C_k = k$ and $\text{wl}(i) = i$ for every $k \in \text{IOb}$ and $i \in \text{IM}$, whence $f_{kh} = k \circ h^{-1}$ and $\text{wl}_k(i) = k[i]$.

Axioms BA 1 and BA 2 hold by construction. To prove BA 3, let ℓ be a vertical line and $k \in \text{IOb} = G$. Then, there is $f \in \text{Triv}_\Omega^\uparrow$ such that $f[\mathbf{t}] = \ell$. It follows that $k^{-1} \circ f \in G$ since $\text{Triv}_\Omega^\uparrow \subseteq G$ and G is a group. Let $i = (k^{-1} \circ f)[\mathbf{t}] \in \text{IM}$. Then $\text{wl}_k(i) = k[i] = f[\mathbf{t}] = \ell$. Thus BA 3 holds.

To prove BA 4, suppose $i \in \text{IM}$ and $k \in \text{IOb} = G$. We have to prove that $\text{wl}_k(i)$ is a line of finite slope. By the definition of IM in $\mathfrak{M}(G)$, $i = g[\mathbf{t}]$ for some $g \in G$. Since G is a group, $k \circ g \in G$. But now, by assumption (ii), $\text{wl}_k(i) = k[i] = (k \circ g)[\mathbf{t}]$ is a line of finite slope. Thus BA 4 holds.

Let us now prove Axiom I. By assumption (i), we can fix some $g \in \mathbf{G} \setminus \text{Triv}_{\Omega}^{\uparrow}$. By assumption (iii), there is a vertical line ℓ such that $g[\ell]$ is not vertical. Let us fix such an ℓ . Let Id denote the identity transformation of \mathbf{Q}^4 . Then $\text{Id}, g \in \text{IOb} = \mathbf{G}$. By the already proven BA 3, there is $i \in \text{IM}$ such that $\text{wl}_{\text{Id}}(i) = \ell$. So, by equation (2), $\text{wl}_g(i) = f_{g\text{Id}}[\text{wl}_{\text{Id}}(i)] = g[\ell]$. Therefore, i is stationary w.r.t. inertial observer Id but is not stationary w.r.t. inertial observer g , which proves Axiom I.

To prove Axiom II(a), let $k \in \text{IOb}$ and $\varphi \in \text{Triv}_{\Omega}^{\uparrow}$. We have to prove that there is $h \in \text{IOb}$ such that $f_{kh} = \varphi$. Let $h = (k^{-1} \circ \varphi)^{-1}$. Then $h \in \mathbf{G} = \text{IOb}$ since \mathbf{G} is a group and $f_{kh} = k \circ h^{-1} = \varphi$, which is what we wanted to prove.

To prove Axiom II(b), let $k, h \in \text{IOb}$ be such that h is at rest w.r.t. k . We have to prove that $f_{kh} \in \text{Triv}_{\Omega}^{\uparrow}$. Let ℓ be an arbitrary vertical line. By the already proven BA 3, there is $i \in \text{IM}$ such that $\text{wl}_h(i) = \ell$. Since ℓ is vertical, i is stationary w.r.t. h . Thus i must be stationary w.r.t. k because h is at rest w.r.t. k . Hence $\text{wl}_k(i)$ is also a vertical line. By equation (2), f_{kh} takes $\text{wl}_h(i) = \ell$ to vertical line $\text{wl}_k(i)$. Since ℓ was an arbitrary vertical line, f_{kh} takes vertical lines to vertical ones. Since \mathbf{G} is a group, we have $f_{kh} = k \circ h^{-1} \in \mathbf{G}$. Hence, by assumption (iii), we have that $f_{kh} \in \text{Triv}_{\Omega}^{\uparrow}$, which is what we wanted to prove.

To prove Axiom III, let $i \in \text{IM}$, $k \in \text{IOb}$, and $\varphi \in \text{Triv}_{\Omega}^{\uparrow}$. We have to prove that there is an inertial motion $i' \in \text{IM}$ such that $\text{wl}_k(i') = \varphi[\text{wl}_k(i)]$. Let $g \in \mathbf{G} = \text{IOb}$ such that $i = g[\mathbf{t}]$. Then $k^{-1} \circ \varphi \circ k \circ g \in \mathbf{G}$ because \mathbf{G} is a group containing $\text{Triv}_{\Omega}^{\uparrow}$. Let $i' = (k^{-1} \circ \varphi \circ k \circ g)[\mathbf{t}]$. Then $i' \in \text{IM}$ by the definition of IM in $\mathfrak{M}(\mathbf{G})$ and

$$k[i'] = k[(k^{-1} \circ \varphi \circ k \circ g)[\mathbf{t}]] = (\varphi \circ k)[g[\mathbf{t}]] = \varphi[k[i]].$$

Since $\text{wl}_k(i') = k[i']$ and $\text{wl}_k(i) = k[i]$, this proves that $\text{wl}_k(i') = \varphi[\text{wl}_k(i)]$.

To prove Axiom IV, let $k, k', h \in \text{IOb}$. We have to prove that there is h' such that $f_{kk'} = f_{hh'}$, i.e. $k \circ k'^{-1} = h \circ h'^{-1}$. Such h' exists because $\text{IOb} = \mathbf{G}$ is a group: setting $h' = k' \circ k^{-1} \circ h \in \text{IOb}$ yields $f_{kk'} = f_{hh'}$.

It remains to show that $\mathbb{W}_{\mathfrak{M}(\mathbf{G})} = \mathbf{G}$. This is straightforward because, by definition and because \mathbf{G} is a group,

$$\mathbb{W}_{\mathfrak{M}(\mathbf{G})} = \{f_{kh} : k, h \in \mathbf{G}\} = \{k \circ h^{-1} : k, h \in \mathbf{G}\} = \mathbf{G}$$

as required. \square

In the proof of Theorem 4.6., we will use the following lemma.

LEMMA 5.14. *Assume BA 1. Assume $L \in {}_c\text{EucL}_{\Omega}^{\uparrow} \cup \text{Gal}_{\Omega}^{\uparrow} \cup {}_c\text{Poi}_{\Omega}^{\uparrow}$ for some $c > 0$ and that L takes vertical lines to vertical lines. Then $L \in \text{Triv}_{\Omega}^{\uparrow}$.*

Proof. Without loss of generality, we can assume that L is linear. In this case, we only have to prove that L preserves the squared-Euclidean length, i.e. that $|L\vec{p}|^2 = |\vec{p}|^2$ for every $\vec{p} \in \mathbf{Q}^4$.

Then $(L\vec{e}_{\mathbf{T}})_t > 0$ and $L\vec{e}_{\mathbf{T}} \in \mathbf{t}$ because L is an orthochronous linear transformation that takes vertical lines to vertical lines. The only point on \mathbf{t} with positive time component and squared c -Minkowskian (squared c -Euclidean) length of c^2 is $\vec{e}_{\mathbf{T}}$. Therefore, $L\vec{e}_{\mathbf{T}} = \vec{e}_{\mathbf{T}}$ by Propositions 3.2., 3.3., and 3.4..

Let $\vec{p} = (p_t, p_x, p_y, p_z) \in \mathbf{Q}^4$ be arbitrary but fixed. We have to prove that $|L\vec{p}|^2 = |\vec{p}|^2$. By the linearity of L and $L\vec{e}_{\mathbf{T}} = \vec{e}_{\mathbf{T}}$, we have

$$L(p_t, 0, 0, 0) = (p_t, 0, 0, 0). \tag{4}$$

TRANSFORMATION GROUPS IMPLIED BY THE RELATIVITY PRINCIPLE 17

We also have

$$L(0, p_x, p_y, p_z) = (0, q_x, q_y, q_z) \quad \text{and} \quad q_x^2 + q_y^2 + q_z^2 = p_x^2 + p_y^2 + p_z^2 \quad (5)$$

for some $q_x, q_y, q_z \in \mathbb{Q}$ because of the following. If $L \in {}_c\text{Poi}_\Omega^\uparrow$, then $L(0, p_x, p_y, p_z)_t = 0$ because $L(0, p_x, p_y, p_z) \diamond_c \vec{e}_T = L(0, p_x, p_y, p_z) \diamond_c L\vec{e}_T = (0, p_x, p_y, p_z) \diamond_c \vec{e}_T = 0$ as L is a linear c -Poincaré transformation. Hence $L(0, p_x, p_y, p_z) = (0, q_x, q_y, q_z)$ for some $q_x, q_y, q_z \in \mathbb{Q}$, and $q_x^2 + q_y^2 + q_z^2 = -\|L(0, p_x, p_y, p_z)\|_{c,\mu}^2 = -\|(0, p_x, p_y, p_z)\|_{c,\mu}^2 = p_x^2 + p_y^2 + p_z^2$ by Proposition 3.4.. A completely analogous proof based on Proposition 3.2. shows that (5) holds when $L \in {}_c\text{Eucl}_\Omega^\uparrow$. And if $L \in \text{Gal}_\Omega^\uparrow$, then (5) holds by Proposition 3.3..

Thus, by linearity of L and equations (4) and (5), we have

$$\begin{aligned} |L\vec{p}|^2 &= |L(p_t, 0, 0, 0) + L(0, p_x, p_y, p_z)|^2 = |(p_t, q_x, q_y, q_z)|^2 \\ &= p_t^2 + q_x^2 + q_y^2 + q_z^2 = p_t^2 + p_x^2 + p_y^2 + p_z^2 = |\vec{p}|^2 \end{aligned}$$

as claimed. \square

Proof of Theorem 4.6.. Let \mathbf{G} be a group for which $\text{Triv}_\Omega^\uparrow < \mathbf{G}$ and either $\mathbf{G} \subset {}_c\text{Eucl}_\Omega^\uparrow$, or $\mathbf{G} \leq \text{Gal}_\Omega^\uparrow$ or $\mathbf{G} \leq {}_c\text{Poi}_\Omega^\uparrow$. Then \mathbf{G} satisfies assumptions (i) and (ii) of Theorem 5.12. because elements of \mathbf{G} are orthochronous linear transformations composed with translations. \mathbf{G} also satisfies assumption (iii) of Theorem 5.12. by Lemma 5.14.. By Theorem 5.12., $\mathfrak{M}(\mathbf{G})$ is a model of Borisov's Axioms and $\mathbb{W}_{\mathfrak{M}(\mathbf{G})} = \mathbf{G}$. \square

Proof of Theorem 4.7.. Assume Borisov's Axioms $\setminus \{\text{Axiom III}\}$ and that Ω is the ordered field \mathbb{R} of reals. Then, by Theorem 5.10., \mathbb{W} is a group satisfying assumptions (i), (ii), and (iii) of Theorem 5.12.. Hence $\mathfrak{M}(\mathbb{W})$ is a model of Borisov's Axioms and $\mathbb{W}_{\mathfrak{M}(\mathbb{W})} = \mathbb{W}$. The statement now follows by Borisov's Theorem. \square

§6. Proof of the main theorem (Theorem 4.5.)

Throughout this section, we assume BA 1, i.e. that $\Omega = (\mathbb{Q}, +, \cdot, \leq)$ is an ordered field. Let us now prove our main result, Theorem 4.5.. To do so, we first introduce some additional notation. The *set of infinitesimals* is defined as follows:

$$\mathcal{E} \stackrel{\text{def}}{=} \{x \in \mathbb{Q} : |nx| < 1 \text{ for every natural number } n\},$$

where nx is an abbreviation for $\underbrace{x + \dots + x}_{n\text{-times}}$ when $x \in \mathbb{Q}$ and $n \in \mathbb{N}$.

A quantity $x \in \mathbb{Q}$ is said to be

- **infinitesimal** if and only if $x \in \mathcal{E}$;
- **unlimited** if and only if $1/x \in \mathcal{E}$; and
- **limited** if and only if it is not unlimited.

Let us note that x is limited if and only if $|x| < n$ for some natural number n .

When Ω is an Archimedean field, we have $\mathcal{E} = \{0\}$ (and hence there are no unlimited numbers), but in non-Archimedean fields there are infinitely many unlimited and infinitesimal numbers.

We call a set \mathcal{C} a **cloud** (see Figure 5 below) if and only if the following hold:

- $\{0\} \subset \mathcal{C} \subseteq \mathcal{E}$;
- if $x \in \mathcal{C}$ and $|y| \leq |x|$, then $y \in \mathcal{C}$;
- if $x \in \mathcal{C}$, then $2x \in \mathcal{C}$.

It is easy to see that (a) there is a cloud in \mathfrak{Q} if and only if (b) \mathcal{E} is a cloud if and only if (c) \mathfrak{Q} is non-Archimedean.

LEMMA 6.15. *Suppose \mathfrak{Q} is non-Archimedean. Given any non-zero $\varepsilon \in \mathcal{E}$, there is a smallest cloud \mathcal{C}_ε containing ε . It is the intersection of all clouds containing ε , and moreover*

$$\mathcal{C}_\varepsilon = \{\alpha \in \mathfrak{Q} : |\alpha| \leq |n\varepsilon| \text{ for some } n \in \mathbb{N}\}.$$

Proof. Let $\mathbb{C}_\varepsilon = \{\mathcal{C} : \mathcal{C} \text{ is a cloud containing } \varepsilon\}$ and define $\mathcal{C}_\varepsilon^* = \bigcap \mathbb{C}_\varepsilon$. Note first that \mathcal{E} is a cloud containing ε , so \mathbb{C}_ε is non-empty and $\mathcal{C}_\varepsilon^* \subseteq \mathcal{E}$.

We know that $\mathcal{C} \in \mathbb{C}_\varepsilon \implies \{0, \varepsilon\} \subseteq \mathcal{C}$, so $\{0, \varepsilon\} \subseteq \mathcal{C}_\varepsilon^*$. Thus, because $0 \neq \varepsilon$, we have $\{0\} \subset \mathcal{C}_\varepsilon^* \subseteq \mathcal{E}$. If $x \in \mathcal{C}_\varepsilon^*$ then $x \in \mathcal{C}$ for all $\mathcal{C} \in \mathbb{C}_\varepsilon$. Thus if $|y| \leq |x|$, we have that $y \in \mathcal{C}$ for all $\mathcal{C} \in \mathbb{C}_\varepsilon$ and hence $y \in \mathcal{C}_\varepsilon^*$. The proof that $2x \in \mathcal{C}_\varepsilon^*$ whenever $x \in \mathcal{C}_\varepsilon^*$ is equally straightforward. This shows that $\mathcal{C}_\varepsilon^*$ is a cloud containing ε , whence it is the smallest such cloud.

Now define $\mathcal{C}_\varepsilon = \{\alpha : |\alpha| \leq |n\varepsilon| \text{ for some } n \in \mathbb{N}\}$. We need to show that $\mathcal{C}_\varepsilon = \mathcal{C}_\varepsilon^*$. Notice first that \mathcal{C}_ε is certainly a cloud. First, it contains both 0 and ε , and all elements of \mathcal{C}_ε are infinitesimal. Second, if $x \in \mathcal{C}_\varepsilon$, there exists some n for which $|x| \leq |n\varepsilon|$, whence $|2x| \leq |(2n)\varepsilon|$ and so $2x \in \mathcal{C}_\varepsilon$. And finally, if $|y| \leq |x|$, then $|y| \leq |x| \leq |n\varepsilon|$, so $y \in \mathcal{C}_\varepsilon$. From this it follows that \mathcal{C}_ε is a cloud containing ε , and hence $\mathcal{C}_\varepsilon^* \subseteq \mathcal{C}_\varepsilon$.

It remains to show that $\mathcal{C}_\varepsilon \subseteq \mathcal{C}_\varepsilon^*$, so choose any $\alpha \in \mathcal{C}_\varepsilon$ and any $\mathcal{C} \in \mathbb{C}_\varepsilon$. By definition, there exists $n \in \mathbb{N}$ such that $|\alpha| \leq |n\varepsilon|$. Choose $m \in \mathbb{N}$ such that $n \leq 2^m$ and observe that $2^m\varepsilon \in \mathcal{C}$, because $\varepsilon \in \mathcal{C}$ and \mathcal{C} is closed under doubling. Since $|\alpha| \leq |n\varepsilon| \leq |2^m\varepsilon|$ it follows that $\alpha \in \mathcal{C}$. Thus α belongs to every cloud containing ε , and so $\alpha \in \mathcal{C}_\varepsilon^*$. It follows that $\mathcal{C}_\varepsilon \subseteq \mathcal{C}_\varepsilon^*$, as required. \square

COROLLARY 6.16. *Every non-Archimedean field contains infinitely many clouds.*

Proof. It is enough to show that for each $\varepsilon \in \mathcal{E}$, the smallest cloud containing ε^2 does not contain ε . We argue by contradiction. If $\varepsilon \in \mathcal{C}_{\varepsilon^2}$, then Lemma 6.15. tells us that there exists $n \in \mathbb{N}$ such that $|\varepsilon| \leq |n\varepsilon^2|$. It follows that $1 \leq |n\varepsilon|$, which contradicts the assumption that $\varepsilon \in \mathcal{E}$. \square

LEMMA 6.17. *Let \mathcal{C} be a cloud. Then*

- (i). $x, y \in \mathcal{C} \implies x + y \in \mathcal{C}$,
- (ii). $x \in \mathcal{C}$ and y is limited $\implies xy \in \mathcal{C}$.

Proof. To prove (i), let $x, y \in \mathcal{C}$. Without loss of generality, we can assume that $|y| \leq |x|$. Then, by the triangle inequality, $|x+y| \leq |x|+|y| \leq 2|x|$. Hence $x+y \in \mathcal{C}$.

To prove (ii), let $x \in \mathcal{C}$ and y be limited. Without loss of generality, we can assume that $x > 0$ and $y > 0$. Since y is limited there is a natural number n such that $y < n$. Since $x > 0$, we have $xy < nx$. There is a natural number k such that $n < 2^k$. Hence $0 < xy < 2^k x$. We have $2^k x \in \mathcal{C}$ since \mathcal{C} is closed under doubling. Consequently, $xy \in \mathcal{C}$. \square




Fig. 5. Illustration for clouds \mathcal{C} and \mathcal{E} , and the balls $B_{\mathcal{C}}$ and $B_{\mathcal{E}}$

Let \mathcal{C} be a cloud. Then the \mathcal{C} -ball around \vec{e}_T is defined as follows:

$$B_{\mathcal{C}} \stackrel{\text{def}}{=} \{ \vec{p} \in \mathbb{Q}^4 : |\vec{p} - \vec{e}_T|^2 \leq r^2 \text{ for some } r \in \mathcal{C} \},$$

see Figure 5 below.

By Proposition 6.18. below, the \mathcal{C} -balls and “ \mathcal{C} -boxes” around \vec{e}_T are the same sets.

PROPOSITION 6.18. *Let \mathcal{C} be a cloud. Then*

$$(1 + t, x, y, z) \in B_{\mathcal{C}} \iff t, x, y, z \in \mathcal{C}.$$

Proof. By definition, $(1 + t, x, y, z) \in B_{\mathcal{C}}$ if and only if $t^2 + x^2 + y^2 + z^2 \leq r^2$ for some $r \in \mathcal{C}$. So, if $(1 + t, x, y, z) \in B_{\mathcal{C}}$, we have $\max\{|t|, |x|, |y|, |z|\} \leq r$ for some $r \in \mathcal{C}$. Consequently, $t, x, y, z \in \mathcal{C}$. The converse follows because $t^2 + x^2 + y^2 + z^2 \leq (2 \max\{|t|, |x|, |y|, |z|\})^2$. \square

PROPOSITION 6.19. *Let $f: \mathbb{Q}^4 \rightarrow \mathbb{Q}^4$ be a linear transformation and let \mathcal{C} be a cloud. Assume that $|f(\vec{e}_x)|^2, |f(\vec{e}_y)|^2, |f(\vec{e}_z)|^2$ are limited and $f(\vec{e}_T) \in B_{\mathcal{C}}$. Then $f[B_{\mathcal{C}}] \subseteq B_{\mathcal{C}}$.*

Proof. Suppose $(1 + t, x, y, z) \in B_{\mathcal{C}}$. We need to show that $f(1 + t, x, y, z) \in B_{\mathcal{C}}$.

Note first that $t, x, y, z \in \mathcal{C}$ by Proposition 6.18.. Since f is linear,

$$f(1 + t, x, y, z) = f(\vec{e}_T) + tf(\vec{e}_T) + xf(\vec{e}_x) + yf(\vec{e}_y) + zf(\vec{e}_z).$$

For all $\vec{p}, \vec{q} \in \mathbb{Q}^4$, we have

$$|\vec{p} + \vec{q}|^2 \leq 2|\vec{p}|^2 + 2|\vec{q}|^2 \tag{6}$$

20 JUDIT X MADARÁSZ, MIKE STANNETT & GERGELY SZÉKELY

since $(a + b)^2 \leq 2a^2 + 2b^2$ for all $a, b \in \mathbb{Q}$. Consequently, we have

$$\begin{aligned}
 & |f(1 + t, x, y, z) - f(\vec{e}_T)|^2 \\
 &= |tf(\vec{e}_T) + xf(\vec{e}_x) + yf(\vec{e}_y) + zf(\vec{e}_z)|^2 \\
 &\leq 2|tf(\vec{e}_T) + xf(\vec{e}_x)|^2 + 2|yf(\vec{e}_y) + zf(\vec{e}_z)|^2 \\
 &\leq 4|tf(\vec{e}_T)|^2 + 4|xf(\vec{e}_x)|^2 + 4|yf(\vec{e}_y)|^2 + 4|zf(\vec{e}_z)|^2 \\
 &\leq 4 \max \left\{ t^2 |f(\vec{e}_T)|^2, x^2 |f(\vec{e}_x)|^2, y^2 |f(\vec{e}_y)|^2, z^2 |f(\vec{e}_z)|^2 \right\} \\
 &\leq r_0^2
 \end{aligned}$$

for some positive $r_0 \in \mathcal{C}$ by Lemma 6.17. since $f(\vec{e}_T)$ is also limited as $f(\vec{e}_T) \in B_{\mathcal{C}}$. By inequality (6),

$$|f(1 + t, x, y, z) - \vec{e}_T|^2 \leq 2|f(1 + t, x, y, z) - f(\vec{e}_T)|^2 + 2|f(\vec{e}_T) - \vec{e}_T|^2.$$

There is a positive $r_1 \in \mathcal{C}$ such that $|f(\vec{e}_T) - \vec{e}_T|^2 \leq r_1^2$ because $f(\vec{e}_T) \in B_{\mathcal{C}}$. Therefore,

$$|f(1 + t, x, y, z) - \vec{e}_T|^2 \leq 2r_0^2 + 2r_1^2 \leq (2 \max\{r_0, r_1\})^2.$$

Since $2 \max\{r_0, r_1\} \in \mathcal{C}$, we have $f(1 + t, x, y, z) \in B_{\mathcal{C}}$, which is what we wanted to prove. \square

Given any set Trf of transformations, we will write LinTrf for the set of transformations in Trf which are linear, i.e.

$$\text{LinTrf} \stackrel{\text{def}}{=} \{f \in \text{Trf} : f \text{ is linear}\}.$$

In particular, if \mathcal{C} is a cloud we will be interested in the following sets of linear transformations:

$$\begin{aligned}
 \text{LinEucl}_{\Omega}^{\uparrow}(\mathcal{C}) &\stackrel{\text{def}}{=} \{f \in \text{LinEucl}_{\Omega}^{\uparrow} : f(\vec{e}_T) \in B_{\mathcal{C}}\}, \\
 \text{LinGal}_{\Omega}^{\uparrow}(\mathcal{C}) &\stackrel{\text{def}}{=} \{f \in \text{LinGal}_{\Omega}^{\uparrow} : f(\vec{e}_T) \in B_{\mathcal{C}}\}, \\
 \text{LinPoi}_{\Omega}^{\uparrow}(\mathcal{C}) &\stackrel{\text{def}}{=} \{f \in \text{LinPoi}_{\Omega}^{\uparrow} : f(\vec{e}_T) \in B_{\mathcal{C}}\}.
 \end{aligned}$$

We will also refer below to the following sets of affine transformations:

$$\begin{aligned}
 \text{Eucl}_{\Omega}^{\uparrow}(\mathcal{C}) &\stackrel{\text{def}}{=} \text{Tran}_{\Omega} \circ \text{LinEucl}_{\Omega}^{\uparrow}(\mathcal{C}), \\
 \text{Gal}_{\Omega}^{\uparrow}(\mathcal{C}) &\stackrel{\text{def}}{=} \text{Tran}_{\Omega} \circ \text{LinGal}_{\Omega}^{\uparrow}(\mathcal{C}), \\
 \text{Poi}_{\Omega}^{\uparrow}(\mathcal{C}) &\stackrel{\text{def}}{=} \text{Tran}_{\Omega} \circ \text{LinPoi}_{\Omega}^{\uparrow}(\mathcal{C}),
 \end{aligned}$$

where Tran_{Ω} is the set of translations.

PROPOSITION 6.20. *Let \mathcal{C} be a cloud. Then*

$$\begin{aligned}
 \text{LinEucl}_{\Omega}^{\uparrow}(\mathcal{C}) &= \{f \in \text{LinEucl}_{\Omega}^{\uparrow} : f[B_{\mathcal{C}}] = B_{\mathcal{C}}\}, \\
 \text{LinGal}_{\Omega}^{\uparrow}(\mathcal{C}) &= \{f \in \text{LinGal}_{\Omega}^{\uparrow} : f[B_{\mathcal{C}}] = B_{\mathcal{C}}\}, \\
 \text{LinPoi}_{\Omega}^{\uparrow}(\mathcal{C}) &= \{f \in \text{LinPoi}_{\Omega}^{\uparrow} : f[B_{\mathcal{C}}] = B_{\mathcal{C}}\}.
 \end{aligned}$$

Proof. Let $f \in \text{LinEucl}_{\Omega}^{\uparrow} \cup \text{LinGal}_{\Omega}^{\uparrow} \cup \text{LinPoi}_{\Omega}^{\uparrow}$ and let \mathcal{C} be a cloud. It is enough to show that

$$f(\vec{e}_T) \in B_{\mathcal{C}} \iff f[B_{\mathcal{C}}] = B_{\mathcal{C}}.$$

If $f \in \text{LinPoi}_{\Omega}^{\uparrow}$, then

$$f(\vec{e}_{\tau})_t = f^{-1}(\vec{e}_{\tau})_t \quad \text{and} \quad |f(\vec{e}_{\tau})_s|^2 = |f^{-1}(\vec{e}_{\tau})_s|^2$$

because $f(\vec{e}_{\tau})_t = \vec{e}_{\tau} \diamond_1 f(\vec{e}_{\tau}) = f^{-1}(\vec{e}_{\tau}) \diamond_1 \vec{e}_{\tau} = f^{-1}(\vec{e}_{\tau})_t$, $|\vec{p}_s|^2 = p_t^2 - \|\vec{p}\|_{1,\mu}^2$ for every $\vec{p} \in \mathbb{Q}^4$ and $\|f(\vec{e}_{\tau})\|_{1,\mu}^2 = \|\vec{e}_{\tau}\|_{1,\mu}^2 = \|f^{-1}(\vec{e}_{\tau})\|_{1,\mu}^2$ by Proposition 3.4.. Thus

$$\begin{aligned} |\vec{e}_{\tau} - f(\vec{e}_{\tau})|^2 &= (1 - f(\vec{e}_{\tau})_t)^2 + |f(\vec{e}_{\tau})_s|^2 \\ &= (1 - f^{-1}(\vec{e}_{\tau})_t)^2 + |f^{-1}(\vec{e}_{\tau})_s|^2 = |\vec{e}_{\tau} - f^{-1}(\vec{e}_{\tau})|^2. \end{aligned}$$

If $f \in \text{LinEucl}_{\Omega}^{\uparrow} \cup \text{LinGal}_{\Omega}^{\uparrow}$,

$$|\vec{e}_{\tau} - f(\vec{e}_{\tau})|^2 = |f^{-1}(\vec{e}_{\tau}) - f^{-1}(f(\vec{e}_{\tau}))|^2 = |f^{-1}(\vec{e}_{\tau}) - \vec{e}_{\tau}|^2$$

because f^{-1} is linear and preserves the Euclidean distance of \vec{e}_{τ} and $f(\vec{e}_{\tau})$ by Propositions 3.2. and 3.3..

So $|\vec{e}_{\tau} - f(\vec{e}_{\tau})|^2 = |f^{-1}(\vec{e}_{\tau}) - \vec{e}_{\tau}|^2$ in all three cases. Thus $f(\vec{e}_{\tau}) \in B_C$ if and only if $f^{-1}(\vec{e}_{\tau}) \in B_C$ and hence $\text{LinEucl}_{\Omega}^{\uparrow}(C)$, $\text{LinGal}_{\Omega}^{\uparrow}(C)$ and $\text{LinPoi}_{\Omega}^{\uparrow}(C)$ are closed under taking inverse. Therefore, it is enough to show that

$$f(\vec{e}_{\tau}) \in B_C \iff f[B_C] \subseteq B_C.$$

Hence, by Proposition 6.19., it is enough to show that $f(\vec{e}_{\tau}) \in B_C$ implies that $|f(\vec{e}_x)|^2$, $|f(\vec{e}_y)|^2$, $|f(\vec{e}_z)|^2$ are limited.

If $f \in \text{LinGal}_{\Omega}^{\uparrow}$ or $f \in \text{LinEucl}_{\Omega}^{\uparrow}$, $|f(\vec{e}_x)|^2 = |f(\vec{e}_y)|^2 = |f(\vec{e}_z)|^2 = 1$. Hence they are limited.

To complete the proof in the last remaining case, let $f \in \text{LinPoi}_{\Omega}^{\uparrow}$. We will prove that for every $i \in \{x, y, z\}$, $|f(\vec{e}_i)|^2 \leq |f(\vec{e}_{\tau})|^2$. To prove this, let $i \in \{x, y, z\}$ and let

$$\vec{p} = f(\vec{e}_{\tau}) \quad \text{and} \quad \vec{q} = f(\vec{e}_i).$$

Then $p_t \cdot q_t - \vec{p}_s \cdot \vec{q}_s = f(\vec{e}_{\tau}) \diamond_1 f(\vec{e}_i) = \vec{e}_{\tau} \diamond_1 \vec{e}_i = 0$, $p_t^2 - |\vec{p}_s|^2 = \|f(\vec{e}_{\tau})\|_{1,\mu}^2 = \|\vec{e}_{\tau}\|_{1,\mu}^2 = 1$ and $q_t^2 - |\vec{q}_s|^2 = \|f(\vec{e}_i)\|_{1,\mu}^2 = \|\vec{e}_i\|_{1,\mu}^2 = -1$ by definition of Poincaré transformations and Proposition 3.4.. Consequently,

$$p_t \cdot q_t = \vec{p}_s \cdot \vec{q}_s, \quad |\vec{p}_s|^2 = p_t^2 - 1, \quad \text{and} \quad |\vec{q}_s|^2 = q_t^2 + 1.$$

Therefore, by the Cauchy-Schwarz inequality,⁷

$$(p_t \cdot q_t)^2 = (\vec{p}_s \cdot \vec{q}_s)^2 \leq |\vec{p}_s|^2 \cdot |\vec{q}_s|^2 = (p_t^2 - 1) \cdot (q_t^2 + 1) = (p_t \cdot q_t)^2 - q_t^2 + p_t^2 - 1.$$

Thus $(p_t \cdot q_t)^2 \leq (p_t \cdot q_t)^2 - q_t^2 + p_t^2 - 1$. This is equivalent to $q_t^2 \leq p_t^2 - 1$, which is equivalent to $q_t^2 + (q_t^2 + 1) \leq p_t^2 + (p_t^2 - 1)$, which is equivalent to $q_t^2 + |\vec{q}_s|^2 \leq p_t^2 + |\vec{p}_s|^2$. Thus $|\vec{q}|^2 \leq |\vec{p}|^2$, i.e. $|f(\vec{e}_i)|^2 \leq |f(\vec{e}_{\tau})|^2$. Now $|f(\vec{e}_{\tau})|^2$ is limited because $f(\vec{e}_{\tau}) \in B_C$. Therefore, $|f(\vec{e}_x)|^2$, $|f(\vec{e}_y)|^2$ and $|f(\vec{e}_z)|^2$ are limited, too. \square

⁷ The Cauchy-Schwarz inequality over \mathbb{R} has several elementary proofs, many of which remain valid over arbitrary ordered fields.

LEMMA 6.21. *Suppose $p_t, p_x, r_t, r_x \in \mathbb{Q}$ satisfy $p_t^2 + p_x^2 = r_t^2 - r_x^2 = 1$, and let q be any value in \mathbb{Q} . Then there exist $f \in \text{Eucl}_\Omega$, $g \in \text{Gal}_\Omega$ and $h \in \text{Poi}_\Omega$ such that*

$$\begin{aligned} f(\vec{e}_T) &= (p_t, p_x, 0, 0), \\ g(\vec{e}_T) &= (1, q, 0, 0), \\ h(\vec{e}_T) &= (r_t, r_x, 0, 0). \end{aligned}$$

Proof. Let f be the linear transformation that takes $\vec{e}_T, \vec{e}_X, \vec{e}_Y, \vec{e}_Z$ to $(p_t, p_x, 0, 0), (-p_x, p_t, 0, 0), \vec{e}_Y, \vec{e}_Z$, respectively. Let g be the linear transformation that takes $\vec{e}_T, \vec{e}_X, \vec{e}_Y, \vec{e}_Z$ to $(1, q, 0, 0), \vec{e}_X, \vec{e}_Y, \vec{e}_Z$, respectively. Let h be the linear transformation that takes $\vec{e}_T, \vec{e}_X, \vec{e}_Y, \vec{e}_Z$ to $(r_t, r_x, 0, 0), (r_x, r_t, 0, 0), \vec{e}_Y, \vec{e}_Z$, respectively. Then, using Propositions 3.2., 3.3., and 3.4., it is easy to check that $f \in \text{Eucl}_\Omega$, $g \in \text{Gal}_\Omega$ and $h \in \text{Poi}_\Omega$. \square

THEOREM 6.22. *Let \mathcal{C} be a cloud. Then*

$$\begin{aligned} \text{Triv}_\Omega^\uparrow < \text{Eucl}_\Omega^\uparrow(\mathcal{C}) &\subset \text{Eucl}_\Omega^\uparrow, \\ \text{Triv}_\Omega^\uparrow < \text{Gal}_\Omega^\uparrow(\mathcal{C}) &< \text{Gal}_\Omega^\uparrow, \\ \text{Triv}_\Omega^\uparrow < \text{Poi}_\Omega^\uparrow(\mathcal{C}) &< \text{Poi}_\Omega^\uparrow. \end{aligned}$$

Moreover, there is a subcloud $\mathcal{C}' \subset \mathcal{C}$ such that

$$\begin{aligned} \text{Triv}_\Omega^\uparrow < \text{Eucl}_\Omega^\uparrow(\mathcal{C}') &< \text{Eucl}_\Omega^\uparrow(\mathcal{C}) \subset \text{Eucl}_\Omega^\uparrow, \\ \text{Triv}_\Omega^\uparrow < \text{Gal}_\Omega^\uparrow(\mathcal{C}') &< \text{Gal}_\Omega^\uparrow(\mathcal{C}) < \text{Gal}_\Omega^\uparrow, \\ \text{Triv}_\Omega^\uparrow < \text{Poi}_\Omega^\uparrow(\mathcal{C}') &< \text{Poi}_\Omega^\uparrow(\mathcal{C}) < \text{Poi}_\Omega^\uparrow. \end{aligned}$$

Proof. To prove the first part of the theorem, it is enough to show that

$$\begin{aligned} \text{LinTriv}_\Omega^\uparrow < \text{LinEucl}_\Omega^\uparrow(\mathcal{C}) &\subset \text{LinEucl}_\Omega^\uparrow, \\ \text{LinTriv}_\Omega^\uparrow < \text{LinGal}_\Omega^\uparrow(\mathcal{C}) &< \text{LinGal}_\Omega^\uparrow, \\ \text{LinTriv}_\Omega^\uparrow < \text{LinPoi}_\Omega^\uparrow(\mathcal{C}) &< \text{LinPoi}_\Omega^\uparrow, \end{aligned}$$

because $\text{Trf} = \text{Tran}_\Omega \circ \text{LinTrf}$ for each of the relevant transformation groups Trf .

Since $\text{LinGal}_\Omega^\uparrow$ and $\text{LinPoi}_\Omega^\uparrow$ are groups, we have

$$\text{LinGal}_\Omega^\uparrow(\mathcal{C}) \leq \text{LinGal}_\Omega^\uparrow \quad \text{and} \quad \text{LinPoi}_\Omega^\uparrow(\mathcal{C}) \leq \text{LinPoi}_\Omega^\uparrow$$

by Proposition 6.20.. By Lemma 6.21., there are $g \in \text{LinGal}_\Omega^\uparrow$ and $h \in \text{LinPoi}_\Omega^\uparrow$ such that $g(\vec{e}_T) = (1, 1, 0, 0)$ and $h(\vec{e}_T) = (\frac{5}{3}, \frac{4}{3}, 0, 0)$. Clearly for such g and h , we have $g \notin \text{LinGal}_\Omega^\uparrow(\mathcal{C})$ and $h \notin \text{LinPoi}_\Omega^\uparrow(\mathcal{C})$ because $(1, 1, 0, 0), (\frac{5}{3}, \frac{4}{3}, 0, 0) \notin B_{\mathcal{C}}$. Therefore, $\text{LinGal}_\Omega^\uparrow(\mathcal{C}) < \text{LinGal}_\Omega^\uparrow$ and $\text{LinPoi}_\Omega^\uparrow(\mathcal{C}) < \text{LinPoi}_\Omega^\uparrow$.

Since $\text{LinEucl}_\Omega^\uparrow$ does not form a group, we need to adopt a different approach to prove that $\text{LinEucl}_\Omega^\uparrow(\mathcal{C})$ is a group. Let $f, g \in \text{LinEucl}_\Omega^\uparrow(\mathcal{C})$. Then $f \circ g \in \text{LinEucl}_\Omega$ and $f^{-1} \in \text{LinEucl}_\Omega$. By Proposition 6.20., we have $(f \circ g)[B_{\mathcal{C}}] = B_{\mathcal{C}}$ and $f^{-1}[B_{\mathcal{C}}] = B_{\mathcal{C}}$. Thus we have $(f \circ g)(\vec{e}_T)_t > 0$ and $f^{-1}(\vec{e}_T)_t > 0$ because $(f \circ g)(\vec{e}_T) \in B_{\mathcal{C}}$, $f^{-1}(\vec{e}_T) \in B_{\mathcal{C}}$ and $\mathcal{C} \subseteq \mathcal{E}$. Hence $f \circ g \in \text{LinEucl}_\Omega^\uparrow(\mathcal{C})$ and $f^{-1} \in \text{LinEucl}_\Omega^\uparrow(\mathcal{C})$. Consequently, $\text{LinEucl}_\Omega^\uparrow(\mathcal{C})$ is a group and $\text{LinEucl}_\Omega^\uparrow(\mathcal{C}) \subseteq \text{LinEucl}_\Omega^\uparrow$. Hence $\text{LinEucl}_\Omega^\uparrow(\mathcal{C}) \subset \text{LinEucl}_\Omega^\uparrow$ as $\text{LinEucl}_\Omega^\uparrow$ is not closed under composition.

TRANSFORMATION GROUPS IMPLIED BY THE RELATIVITY PRINCIPLE 23

$\text{LinTriv}_\Omega^\uparrow$ is a subgroup of $\text{LinEucl}_\Omega^\uparrow(\mathcal{C})$, $\text{LinGal}_\Omega^\uparrow(\mathcal{C})$ and $\text{LinPoi}_\Omega^\uparrow(\mathcal{C})$ because $f(\vec{e}_T) = \vec{e}_T$ for every $f \in \text{LinTriv}_\Omega^\uparrow$.

To prove that $\text{LinTriv}_\Omega^\uparrow$ is a proper subgroup of these groups, let $\varepsilon \in \mathcal{C}$ be such that $\varepsilon > 0$. Let

$$p_t = \frac{2\varepsilon+1}{2\varepsilon^2+2\varepsilon+1}, \quad p_x = \frac{2\varepsilon^2+2\varepsilon}{2\varepsilon^2+2\varepsilon+1}, \quad r_t = \frac{2\varepsilon^2+2\varepsilon+1}{2\varepsilon+1}, \quad r_x = \frac{2\varepsilon^2+2\varepsilon}{2\varepsilon+1}.$$

and notice that $p_t^2 + p_x^2 = r_t^2 - r_x^2 = 1$, because $(2\varepsilon^2+2\varepsilon+1)^2 - (2\varepsilon^2+2\varepsilon)^2 = (2\varepsilon+1)^2$.

By Lemma 6.21., there exist transformations $f \in \text{LinEucl}_\Omega^\uparrow$, $g \in \text{LinGal}_\Omega^\uparrow$ and $h \in \text{LinPoi}_\Omega^\uparrow$ such that

$$f(\vec{e}_T) = (p_t, p_x, 0, 0), \quad g(\vec{e}_T) = (1, \varepsilon, 0, 0), \quad h(\vec{e}_T) = (r_t, r_x, 0, 0).$$

It now follows, by Proposition 6.18., that $f(\vec{e}_T), g(\vec{e}_T), h(\vec{e}_T) \in B_{\mathcal{C}}$. To see this, note that $0 < \varepsilon^2 < \varepsilon < 1$, whence

- $1 - 2\varepsilon < 1 - 2\varepsilon^2 < 1 - \frac{2\varepsilon^2}{2\varepsilon^2+2\varepsilon+1} = \frac{2\varepsilon+1}{2\varepsilon^2+2\varepsilon+1} = p_t$ and $p_t < 1$;
- $0 < p_x$ and $p_x = \frac{2\varepsilon^2+2\varepsilon}{2\varepsilon^2+2\varepsilon+1} < 2\varepsilon^2 + 2\varepsilon < 4\varepsilon$,

so that $1 - 2\varepsilon < p_t < 1$ and $0 < p_x < 4\varepsilon$. Similarly,

- $1 < r_t = \frac{2\varepsilon^2+2\varepsilon+1}{2\varepsilon+1} < 2\varepsilon^2 + 2\varepsilon + 1 < 1 + 4\varepsilon$;
- $0 < r_x = \frac{2\varepsilon^2+2\varepsilon}{2\varepsilon+1} < 2\varepsilon^2 + 2\varepsilon < 4\varepsilon$

whence $1 < r_t < 1 + 4\varepsilon$ and $0 < r_x < 4\varepsilon$.

Therefore, $f \in \text{LinEucl}_\Omega^\uparrow(\mathcal{C})$, $g \in \text{LinGal}_\Omega^\uparrow(\mathcal{C})$ and $h \in \text{LinPoi}_\Omega^\uparrow(\mathcal{C})$, but $f, g, h \notin \text{LinTriv}_\Omega^\uparrow$ because $f(\vec{e}_T)_s \neq (0, 0, 0)$, $g(\vec{e}_T)_s \neq (0, 0, 0)$ and $h(\vec{e}_T)_s \neq (0, 0, 0)$. Thus $\text{LinTriv}_\Omega^\uparrow$ is a proper subgroup of groups $\text{LinEucl}_\Omega^\uparrow(\mathcal{C})$, $\text{LinGal}_\Omega^\uparrow(\mathcal{C})$ and $\text{LinPoi}_\Omega^\uparrow(\mathcal{C})$.

Finally, let \mathcal{C}' be any cloud that does not contain ε . As we saw in the proof of Corollary 6.16. we can take, e.g., $\mathcal{C}' = \mathcal{C}_{\varepsilon^2}$. To complete the present proof, it only remains to prove that $\text{Eucl}_\Omega^\uparrow(\mathcal{C}')$, $\text{Gal}_\Omega^\uparrow(\mathcal{C}')$ and $\text{Poi}_\Omega^\uparrow(\mathcal{C}')$ are proper subgroups of $\text{Eucl}_\Omega^\uparrow(\mathcal{C})$, $\text{Gal}_\Omega^\uparrow(\mathcal{C})$ and $\text{Poi}_\Omega^\uparrow(\mathcal{C})$, respectively. Moreover, to prove this, it is enough to show that $f(\vec{e}_T), g(\vec{e}_T), h(\vec{e}_T) \notin B_{\mathcal{C}'}$.

We have

$$p_x = \frac{2\varepsilon^2 + 2\varepsilon}{2\varepsilon^2 + 2\varepsilon + 1} > \frac{2\varepsilon}{2\varepsilon^2 + 2\varepsilon + 1} > \frac{2\varepsilon}{2} > \varepsilon$$

and similarly

$$r_x = \frac{2\varepsilon^2 + 2\varepsilon}{2\varepsilon + 1} > \frac{2\varepsilon}{2\varepsilon + 1} > \frac{2\varepsilon}{2} > \varepsilon.$$

Thus $p_x, r_x \notin \mathcal{C}'$ as $\varepsilon \notin \mathcal{C}'$. Hence, by Proposition 6.18., $f(\vec{e}_T), g(\vec{e}_T), h(\vec{e}_T) \notin B_{\mathcal{C}'}$. \square

Changing the units used to express spatial distance is represented by the following **space scaling bijections**. Given any $\lambda > 0$, we write $S_\lambda: \mathbb{Q}^4 \rightarrow \mathbb{Q}^4$ for the map:

$$S_\lambda: (t, x, y, z) \mapsto (t, \lambda x, \lambda y, \lambda z),$$

and note that $S_\lambda^{-1} = S_{\frac{1}{\lambda}}$.

PROPOSITION 6.23. For every $c > 0$,

$$\begin{aligned} S_c \circ \text{Eucl}_\Omega^\uparrow \circ S_c^{-1} &= {}_c\text{Eucl}_\Omega^\uparrow, & S_c \circ \text{Poi}_\Omega^\uparrow \circ S_c^{-1} &= {}_c\text{Poi}_\Omega^\uparrow, \\ S_c \circ \text{Triv}_\Omega^\uparrow \circ S_c^{-1} &= \text{Triv}_\Omega^\uparrow, & S_c \circ \text{Gal}_\Omega^\uparrow \circ S_c^{-1} &= \text{Gal}_\Omega^\uparrow. \end{aligned}$$

Proof. Since for all $T \in \text{Tran}_\Omega$ and $c > 0$, there is $T' \in \text{Tran}_\Omega$ such that $S_c \circ T = T' \circ S_c$, it is enough to check the statements for linear transformations. For all $c > 0$ and $\vec{p} \in \mathbb{Q}^4$,

$$\|S_c \vec{p}\|_c^2 = c^2 \|\vec{p}\|_1^2 \quad \text{and} \quad \|S_c \vec{p}\|_{c,\mu}^2 = c^2 \|\vec{p}\|_{1,\mu}^2.$$

Therefore, if $L \in \text{LinEucl}_\Omega^\uparrow$, then by Proposition 3.2.,

$$\left\| S_c L S_{\frac{1}{c}} \vec{p} \right\|_c^2 = c^2 \left\| L S_{\frac{1}{c}} \vec{p} \right\|_1^2 = c^2 \left\| S_{\frac{1}{c}} \vec{p} \right\|_1^2 = \left\| S_c S_{\frac{1}{c}} \vec{p} \right\|_c^2 = \|\vec{p}\|_c^2.$$

Thus, by Proposition 3.2., $S_c L S_c^{-1} \in \text{Lin}_c \text{Eucl}_\Omega$. If we compose any orthochronous transformation from left or right with S_λ , we get an orthochronous transformation. Therefore, we also have $S_c L S_c^{-1} \in \text{Lin}_c \text{Eucl}_\Omega^\uparrow$. Therefore, $S_c \circ \text{Eucl}_\Omega^\uparrow \circ S_c^{-1} \subseteq {}_c \text{Eucl}_\Omega^\uparrow$. Similarly, by Proposition 3.2., if $L \in \text{Lin}_c \text{Eucl}_\Omega^\uparrow$, then

$$c^2 \left\| S_{\frac{1}{c}} L S_c \vec{p} \right\|_1^2 = \left\| S_c S_{\frac{1}{c}} L S_c \vec{p} \right\|_c^2 = \|L S_c \vec{p}\|_c^2 = \|S_c \vec{p}\|_c^2 = c^2 \|\vec{p}\|_1^2.$$

So $S_c^{-1} L S_c \in \text{LinEucl}_\Omega^\uparrow$ and thus $S_c^{-1} \circ {}_c \text{Eucl}_\Omega^\uparrow \circ S_c \subseteq \text{Eucl}_\Omega^\uparrow$, which is equivalent to ${}_c \text{Eucl}_\Omega^\uparrow \subseteq S_c \circ \text{Eucl}_\Omega^\uparrow \circ S_c^{-1}$. Consequently,

$$S_c \circ \text{Eucl}_\Omega^\uparrow \circ S_c^{-1} = {}_c \text{Eucl}_\Omega^\uparrow. \quad (7)$$

An analogous proof based on Proposition 3.4. using $\|\dots\|_{c,\mu}^2$ in place of $\|\dots\|_c^2$ shows that

$$S_c \circ \text{Poi}_\Omega^\uparrow \circ S_c^{-1} = {}_c \text{Poi}_\Omega^\uparrow.$$

Since S_λ takes vertical lines to vertical lines for any λ , elements of $S_c \circ \text{Triv}_\Omega^\uparrow \circ S_c^{-1}$ take vertical lines to vertical ones. On the other hand, $S_c \circ \text{Triv}_\Omega^\uparrow \circ S_c^{-1} \subseteq {}_c \text{Eucl}_\Omega^\uparrow$ by $\text{Triv}_\Omega^\uparrow \subseteq \text{Eucl}_\Omega^\uparrow$ and (7). Therefore, by Lemma 5.14., $S_c \circ \text{Triv}_\Omega^\uparrow \circ S_c^{-1} \subseteq \text{Triv}_\Omega^\uparrow$. Analogously, $S_c^{-1} \circ \text{Triv}_\Omega^\uparrow \circ S_c = S_{\frac{1}{c}} \circ \text{Triv}_\Omega^\uparrow \circ S_{\frac{1}{c}}^{-1} \subseteq \text{Triv}_\Omega^\uparrow$. Consequently,

$$S_c \circ \text{Triv}_\Omega^\uparrow \circ S_c^{-1} = \text{Triv}_\Omega^\uparrow.$$

Assume $L \in \text{LinGal}_\Omega^\uparrow$. Let $\vec{p} = (p_t, p_x, p_y, p_z) \in \mathbb{Q}^4$. Then, we have that

$$(S_c L S_c^{-1} \vec{p})_t = p_t$$

since $(L\vec{q})_t = q_t$ and $(S_c \vec{q})_t = q_t$ for any $\vec{q} \in \mathbb{Q}^4$. If $p_t = 0$, then

$$\left| S_c L S_c^{-1} (0, p_x, p_y, p_z) \right|^2 = \left| S_c L \left(0, \frac{p_x}{c}, \frac{p_y}{c}, \frac{p_z}{c} \right) \right|^2 = c^2 \left| L \left(0, \frac{p_x}{c}, \frac{p_y}{c}, \frac{p_z}{c} \right) \right|^2$$

since $L \left(0, \frac{p_x}{c}, \frac{p_y}{c}, \frac{p_z}{c} \right)_t = 0$ as $L \in \text{LinGal}_\Omega$ and $|S_c \vec{q}|^2 = c^2 |\vec{q}|^2$ for every $\vec{q} \in \mathbb{Q}^4$ with $q_t = 0$. Then

$$c^2 \left| L \left(0, \frac{p_x}{c}, \frac{p_y}{c}, \frac{p_z}{c} \right) \right|^2 = c^2 \left| \left(0, \frac{p_x}{c}, \frac{p_y}{c}, \frac{p_z}{c} \right) \right|^2 = |(0, p_x, p_y, p_z)|^2$$

since $L \in \text{LinGal}_\Omega$. Consequently,

$$p_t = 0 \implies |S_c L S_c^{-1} \vec{p}|^2 = |\vec{p}|^2.$$

TRANSFORMATION GROUPS IMPLIED BY THE RELATIVITY PRINCIPLE 25

Therefore, by Proposition 3.3., $S_c L S_c^{-1} \in \text{Gal}_\Omega^\uparrow$ and analogously $S_c^{-1} L S_c = S_{\frac{1}{c}} L S_{\frac{1}{c}}^{-1} \in \text{Gal}_\Omega^\uparrow$. Therefore,

$$S_c \circ \text{Gal}_\Omega^\uparrow \circ S_c^{-1} = \text{Gal}_\Omega^\uparrow.$$

□

For every $c > 0$ let us introduce the following sets of affine transformations:

$$\begin{aligned} {}_c\text{Eucl}_\Omega^\uparrow(\mathcal{C}) &\stackrel{\text{def}}{=} S_c \circ \text{Eucl}_\Omega^\uparrow(\mathcal{C}) \circ S_c^{-1}, \\ {}_c\text{Gal}_\Omega^\uparrow(\mathcal{C}) &\stackrel{\text{def}}{=} S_c \circ \text{Gal}_\Omega^\uparrow(\mathcal{C}) \circ S_c^{-1}, \\ {}_c\text{Poi}_\Omega^\uparrow(\mathcal{C}) &\stackrel{\text{def}}{=} S_c \circ \text{Poi}_\Omega^\uparrow(\mathcal{C}) \circ S_c^{-1}. \end{aligned}$$

THEOREM 6.24. *Let \mathcal{C} be a cloud. Then there is a subcloud $\mathcal{C}' \subset \mathcal{C}$ such that*

$$\begin{aligned} \text{Triv}_\Omega^\uparrow &< {}_c\text{Eucl}_\Omega^\uparrow(\mathcal{C}') < {}_c\text{Eucl}_\Omega^\uparrow(\mathcal{C}) < {}_c\text{Eucl}_\Omega^\uparrow, \\ \text{Triv}_\Omega^\uparrow &< {}_c\text{Gal}_\Omega^\uparrow(\mathcal{C}') < {}_c\text{Gal}_\Omega^\uparrow(\mathcal{C}) < \text{Gal}_\Omega^\uparrow, \\ \text{Triv}_\Omega^\uparrow &< {}_c\text{Poi}_\Omega^\uparrow(\mathcal{C}') < {}_c\text{Poi}_\Omega^\uparrow(\mathcal{C}) < {}_c\text{Poi}_\Omega^\uparrow. \end{aligned}$$

Proof. Immediate from Theorem 6.22. and Proposition 6.23.. □

Proof of Thm.4.5.. By Theorem 6.24., there are strictly descending countably infinite chains of subgroups such that

$$\begin{aligned} \text{Triv}_\Omega^\uparrow &< \dots < G_i^E < \dots < G_1^E < G_0^E < {}_c\text{Eucl}_\Omega^\uparrow, \\ \text{Triv}_\Omega^\uparrow &< \dots < G_i^G < \dots < G_1^G < G_0^G < \text{Gal}_\Omega^\uparrow, \\ \text{Triv}_\Omega^\uparrow &< \dots < G_i^P < \dots < G_1^P < G_0^P < {}_c\text{Poi}_\Omega^\uparrow. \end{aligned}$$

Groups G_i^E 's, G_i^G 's, and G_i^P 's satisfy assumptions (i) and (ii) of Theorem 5.12., and by Lemma 5.14., they also satisfy assumption (iii). Let $\mathfrak{M}_i^E = \mathfrak{M}(G_i^E)$, $\mathfrak{M}_i^G = \mathfrak{M}(G_i^G)$ and $\mathfrak{M}_i^P = \mathfrak{M}(G_i^P)$. Then, by Theorem 5.12. and Proposition 5.11., the various \mathfrak{M}_i^E , \mathfrak{M}_i^G and \mathfrak{M}_i^P satisfy the statement of Theorem 4.5.. □

§7. Open questions

According to Theorem 4.6., whenever Ω is an ordered field and G is a group for which either

$$\begin{aligned} \text{Triv}_\Omega^\uparrow &< G \subset {}_c\text{Eucl}_\Omega^\uparrow, \text{ or} \\ \text{Triv}_\Omega^\uparrow &< G \leq \text{Gal}_\Omega^\uparrow, \text{ or} \\ \text{Triv}_\Omega^\uparrow &< G \leq {}_c\text{Poi}_\Omega^\uparrow, \end{aligned}$$

there exists a model \mathfrak{M} of Borisov's Axioms over Ω such that $\mathbb{W}_\mathfrak{M} = G$.

By Corollary 4.9., if $\Omega = \mathbb{R}$, there are no groups strictly between $\text{Triv}_\Omega^\uparrow$ and ${}_c\text{Eucl}_\Omega^\uparrow$ ($\text{Gal}_\Omega^\uparrow$, ${}_c\text{Poi}_\Omega^\uparrow$). Contrary to this, if Ω is a non-Archimedean field, then

there are groups G_1 , G_2 , and G_3 such that

$$\begin{aligned} \text{Triv}_\Omega^\uparrow &< G_1 \subset {}_c\text{Eucl}_\Omega^\uparrow, \\ \text{Triv}_\Omega^\uparrow &< G_2 \leq \text{Gal}_\Omega^\uparrow, \text{ and} \\ \text{Triv}_\Omega^\uparrow &< G_3 \leq {}_c\text{Poi}_\Omega^\uparrow. \end{aligned}$$

Moreover, there are infinitely many such groups by Theorem 6.24.. These groups were constructed from clouds in a very specific way. It is natural to ask whether all the groups strictly between $\text{Triv}_\Omega^\uparrow$ and ${}_c\text{Eucl}_\Omega^\uparrow$ ($\text{Gal}_\Omega^\uparrow$, ${}_c\text{Poi}_\Omega^\uparrow$) can be constructed from clouds the same way. More precisely, does the following hold?

Question 1 *Suppose G_1 , G_2 , and G_3 are groups satisfying*

$$\begin{aligned} \text{Triv}_\Omega^\uparrow &< G_1 \subset {}_c\text{Eucl}_\Omega^\uparrow, \\ \text{Triv}_\Omega^\uparrow &< G_2 < \text{Gal}_\Omega^\uparrow, \text{ and} \\ \text{Triv}_\Omega^\uparrow &< G_3 < {}_c\text{Poi}_\Omega^\uparrow. \end{aligned}$$

Do there exist clouds C_1 , C_2 , and C_3 such that

$$\begin{aligned} G_1 &= {}_c\text{Eucl}_\Omega^\uparrow(C_1), \\ G_2 &= {}_c\text{Gal}_\Omega^\uparrow(C_2), \\ G_3 &= {}_c\text{Poi}_\Omega^\uparrow(C_3)? \end{aligned}$$

Even if the answer to Question 1 turns out to be negative in general, it would be interesting to discover whether such clouds exist for specific classes of ordered fields, e.g., the class of real closed fields.

It is also worth noting how the consequences of Borisov’s assumptions sometimes change when we replace \mathbb{R} with other ordered fields, and sometimes remain the same.

For example, according to Borisov’s Theorem, when Ω is the field \mathbb{R} there are only two possibilities for the set of spacetime worldview transformations: either $\mathbb{W} = \text{Gal}_\mathbb{R}^\uparrow$, or $\mathbb{W} = {}_c\text{Poi}_\mathbb{R}^\uparrow$ for some $c > 0$. As explained on page 12, however, this result remains valid if we replace \mathbb{R} with any other Archimedean field, provided every positive number has a square root.

Question 2 *It remains an open question whether the existence of square roots is actually necessary in this situation.*

On the other hand, while Theorem 4.7. shows that Borisov’s Theorem remains valid if omit Axiom III, nonetheless as explained in Remark 4.8., if we simply retain the assumption of square roots without requiring Ω to be \mathbb{R} , we can show instead that either

$$\begin{aligned} \text{Triv}_\Omega^\uparrow &< \mathbb{W} \subset {}_c\text{Eucl}_\Omega^\uparrow \text{ or} \\ \text{Triv}_\Omega^\uparrow &< \mathbb{W} \leq \text{Gal}_\Omega^\uparrow \text{ or} \\ \text{Triv}_\Omega^\uparrow &< \mathbb{W} \leq {}_c\text{Poi}_\Omega^\uparrow \text{ for some } c > 0, \end{aligned}$$

and this remains true even if we replace BA4 with the more general assumption that the worldlines of inertial motions according to inertial observers are lines (but not necessarily of finite slope).

Question 3 *Once again, it remains an open question if these results remain valid if we omit the assumption that positive numbers have square roots.*

Bibliography

- Albeverio, S. (1987). An introduction to non standard analysis and applications to quantum theory. In Blaquiere, A., Diner, S., & Lochak, G., editors, *Information Complexity and Control in Quantum Physics*, Vienna, pp. 183–208. Springer Vienna.
- Albeverio, S., Fenstad, J. E., & Hoegh-Krohn, R. (1986). *Nonstandard Methods in Stochastic Analysis and Mathematical Physics*, Volume 122 of *Pure and applied mathematics*. Academic Press.
- Andréka, H., Madarász, J. X., & Németi, I. (2006). Logical axiomatizations of space-time. Samples from the literature. In Prékopa, A. & Molnár, E., editors, *Non-Euclidean Geometries: János Bolyai Memorial Volume*, pp. 155–185. Springer Verlag.
- Andréka, H., Madarász, J. X., Németi, I., with contributions from: Andai, A., Sági, G., Sain, I., & Tóke, C. (2002). *On the logical structure of relativity theories*. Research report, Alfréd Rényi Institute of Mathematics, Hungar. Acad. Sci., Budapest.
- Andréka, H., & Németi, I. (2014). Comparing theories: the dynamics of changing vocabulary. In Baltag, A. & Smets, S., editors, *Johan van Benthem on Logic and Information Dynamics*, pp. 143–172. Springer Verlag.
- Bagarello, F., & Valenti, S. (1988). Nonstandard analysis in classical physics and quantum formal scattering. *Int. J. Theor. Phys.* **27**, 557–566.
- Borisov, Y. F. (1978, Nov). Axiomatic definition of the Galilean and Lorentz groups. *Siberian Mathematical Journal* **19**(6), 870–882.
- Capinski, M., & Cutland, N. J. (1995). *Nonstandard Methods for Stochastic Fluid Mechanics*. World Scientific Publishers.
- Dávid, G. (1990). Special relativity based on group theory. Talk at Summer School on Special Relativity, Galyatető, Hungary, 1990.
- Einstein, A. (1905). Zur Elektrodynamik bewegter Körper. *Annalen der Physik* **17**, 891–921.
- Frank, P., & Rothe, H. (1911). Über die Transformation der Raumzeitkoordinaten von ruhenden auf bewegte Systeme. *Annalen der Physik* **339**(5), 825–855. English translation (transl. Morris D. Friedman Inc.): https://archive.org/details/nasa_techdoc_19880069066.
- Friend, M. (2015). On the epistemological significance of the Hungarian project. *Synthese* **192**,7, 2035–2051.
- Friend, M., & Molinini, D. (2015). Using mathematics to explain a scientific theory. *Philosophia Mathematica* **24**(2), 185–213.
- Gömöri, M. (2015). *The Principle of Relativity—An Empiricist Analysis*. Ph. D. thesis, Eötvös University, Budapest.
- Gömöri, M., & Szabó, L. E. (2015). Formal statement of the special principle of relativity. *Synthese* **192**(7), 2053–2076.
- Govindarajalulu, N. S., Bringsjord, S., & Taylor, J. (2015, Jul). Proof verification and proof discovery for relativity. *Synthese* **192**(7), 2077–2094.
- Guts, A. K. (1982). The axiomatic theory of relativity. *Russ. Math. Surv.* **37**(2), 41–89.

- Ignatowsky, W. v. (1910a). Das Relativitätsprinzip. *Archiv der Mathematik und Physik* **17**, 1–24.
- Ignatowsky, W. v. (1910b). Einige allgemeine Bemerkungen über das Relativitätsprinzip. *Physikalische Zeitschrift* **11**, 972–976. English Wikisource translation: https://en.wikisource.org/wiki/Translation:Some_General_Remarks_on_the_Relativity_Principle.
- Ignatowsky, W. v. (1911). Das Relativitätsprinzip. *Archiv der Mathematik und Physik* **18**, 17–40.
- Madarász, J. X., & Székely, G. (2013). Special relativity over the field of rational numbers. *International Journal of Theoretical Physics* **52,5**, 1706–1718.
- Madarász, J. X., Székely, G., & Stannett, M. (2017). Three different formalisations of Einstein’s relativity principle. *The Review of Symbolic Logic* **10**(3), 530–548.
- Madarász, J. X., Székely, G., & Stannett, M. (2020). Groups of worldview transformations implied by isotropy of space. arXiv:2007.14261.
- Nelson, E. (1987). *Radically Elementary Probability Theory*. Princeton University Press. Available online: <http://www.math.princeton.edu/~nelson/books/rept.pdf>.
- Odom, B., Hanneke, D., D’Urso, B., & Gabrielse, G. (2006). New measurement of the electron magnetic moment using a one-electron quantum cyclotron. *Phys. Rev. Lett.* **97**, 030801.
- Perlis, D. (2016). Taking physical infinity seriously. In Omodeo, E. G. & Policriti, A., editors, *Martin Davis on Computability, Computational Logic, and Mathematical Foundations*, pp. 243–254. Cham: Springer International Publishing.
- Stannett, M., & Németi, I. (2014, Apr). Using Isabelle/HOL to verify first-order relativity theory. *Journal of Automated Reasoning* **52**(4), 361–378.
- Székely, G. (2009). *First-order logic investigation of relativity theory with an emphasis on accelerated observers*. Ph. D. thesis, Eötvös Loránd Univ., Budapest.

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS
P.O. BOX 127
BUDAPEST 1364, HUNGARY
E-mail: madarasz.judit@renyi.hu, szekely.gergely@renyi.hu
URL: <http://www.renyi.hu/~madarasz>, <http://www.renyi.hu/~turms>

DEPARTMENT OF COMPUTER SCIENCE
THE UNIVERSITY OF SHEFFIELD
211 PORTOBELLO, SHEFFIELD S1 4DP, UK
E-mail: m.stannett@sheffield.ac.uk
URL: <https://www.sheffield.ac.uk/dcs/people/academic/mike-stannett>

UNIVERSITY OF PUBLIC SERVICE
P.O. Box 60
BUDAPEST 1441 , HUNGARY