Utah State University

DigitalCommons@USU

Fall Student Research Symposium 2020

Fall Student Research Symposium

12-10-2020

The Effects of Estrogen and Arousal on Latent Inhibition

Michael Herron Utah State University, michael.herron.mail@gmail.com

Follow this and additional works at: https://digitalcommons.usu.edu/fsrs2020



Part of the Biology Commons

Recommended Citation

Herron, Michael, "The Effects of Estrogen and Arousal on Latent Inhibition" (2020). Fall Student Research Symposium 2020. 27.

https://digitalcommons.usu.edu/fsrs2020/27

This Book is brought to you for free and open access by the Fall Student Research Symposium at DigitalCommons@USU. It has been accepted for inclusion in Fall Student Research Symposium 2020 by an authorized administrator of DigitalCommons@USU. For more information, please contact digitalcommons@usu.edu.





The Effects of Estrogen and Arousal on Latent Inhibition

Michael Herron; Megan Raddatz; Dr. Mona Buhusi



Introduction

The behaviors of humans and animals are dependent on neurotransmitters and hormones that affect attention, alertness, and associative learning. These include the hormone estrogen (which plays an interesting role in cognition attentional processes) neurotransmitter orexin (involved in wakefulness, arousal and goal-directed behaviors). Here we evaluated the roles of orexin receptors and estrogen in Latent Inhibition (LI), a measure of attention and associative learning. Latent inhibition is a behavioral phenomenon in which an organism's ability to associate new meaning to a familiar previously inconsequential (pre-exposed) stimulus is reduced when compared to associating meaning to a novel stimulus (non pre-exposed) in a classical conditioning paradigm. We hypothesize that orexin and estrogen have synergistic effects on latent inhibition (LI).

Methods

Subjects


- Female mice, aged 3-4 months, n = 33
- All mice underwent ovariectomy surgery


Experimental Design


• Alterations to the experimental subjects included condition (non-pre-exposed and pre-exposed), genotype (wild-type vs. orexin receptor knock-out), and treatment (estrogen vs. placebo).

Latent Inhibition Paradigm

- In a 5-phase experiment, about half the mice were preexposed to a stimulus and then conditioned with a foot shock, while the others did not receive pre-exposure.
- Freezing time was then manually recorded upon subsequent exposure to the stimulus as a measure of LI.







Results and Conclusion

The average time spent freezing during the NPE and PE stimuli during the test session in animals receiving placebo treatment is shown in Figure 1. A two-way ANOVA indicates a main effect of group (F(1, 17) = 4.584, p < 0.05, η p2 = 0.215)). Under placebo, both KO and WT animals in the NPE condition showed elevated freezing times as compared to PE condition. Overall, the results trend towards a display of LI in both the KO and WT animals under placebo treatment.

The average time spent freezing during the NPE and PE stimuli during the test session in KO animals receiving placebo or estrogen is shown in Figure 2. A separate two-way ANOVA did not indicate any main effect of treatment (F(1, 14) = 0.064, p > 0.05) or group (F(1, 14) = 1.822, p > 0.05). However, as seen in Figure 1, the KO under placebo that were not preexposed froze slightly more than those who were pre-exposed to the stimuli. While this is not statistically significant, the result trend toward a display of LI in KO receiving placebo but not in KO animals receiving estrogen. While the results that were obtained are of great interest, there exists a necessity to further clarify the roles that orexin and estrogen play in a latent inhibition paradigm.

Acknowledgments

This study was conducted with funding from a USU Undergraduate Research and Creative Opportunity Grant and with funding from the USU honors program. I'd also like to give a special thank you to Mona Buhusi and Megan Raddatz for their help and support in the design and execution of this project.

