

VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

Are longitudinal reallocations of time between movement behaviours associated with adiposity among elderly women? A compositional isotemporal substitution analysis

This is the Accepted version of the following publication

Pelclová, Jana, Štefelová, Nikola, Dumuid, D, Pedisic, Zeljko, Hron, K, Gába, Aleš, Olds, Timothy, Pechová, Jana, Zając-Gawlak, Izabela and Tlučáková, Lenka (2020) Are longitudinal reallocations of time between movement behaviours associated with adiposity among elderly women? A compositional isotemporal substitution analysis. *International Journal of Obesity*, 44. pp. 857-864. ISSN 0307-0565

The publisher's official version can be found at
<https://www.nature.com/articles/s41366-019-0514-x>
Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/41958/>

1 **Are longitudinal reallocations of time between movement behaviours**
2 **associated with adiposity among elderly women?**

3 **A compositional isotemporal substitution analysis**

4
5 Jana Pelclová¹, Nikola Štefelová², Dorothea Dumuid³, Željko Pedišić⁴, Karel Hron², Aleš Gába¹, Timothy
6 Olds³, Jana Pechová¹, Izabela Zajac-Gawlak⁵, Lenka Tlučáková⁶

7
8 ¹ Faculty of Physical Culture, Palacký University Olomouc, Olomouc, Czech Republic

9 ² Faculty of Science, Palacký University Olomouc, Olomouc, Czech Republic

10 ³ Alliance for Research in Exercise, Nutrition and Activity (ARENA), University of South Australia,
11 Adelaide, South Australia, Australia

12 ⁴ Institute for Health and Sport, Victoria University, Melbourne, Australia

13 ⁵ The Jerzy Kukuczka Academy of Physical Education, Katowice, Poland

14 ⁶ Faculty of Sports, University of Presov, Presov, Slovak Republic

15
16 *Running title:* Longitudinal reallocations between movement behaviours

17 *Funding:* This research was funded by a grant from the Czech Science Foundation.

18 *Correspondence address:*

19 Jana Pelclová, PhD

20 Institute of Active Lifestyle

21 Faculty of Physical Culture

22 Palacký University Olomouc

23 Tř. Míru 117, Olomouc, 771 11, Czech Republic

24 Phone: +420 585 636 469

25 Email: jana.pelclova@upol.cz

26 **ABSTRACT**

27 **Background:** This study aimed to use compositional data analysis to: 1) investigate the prospective
28 associations between changes in daily movement behaviours and adiposity among elderly women; and
29 2) to examine how the reallocation of time between movement behaviours was associated with
30 longitudinal changes in adiposity.

31 **Subjects/Methods:** This is a 7-year longitudinal study in Central European older women (n=158,
32 baseline age 63.9±4.4 years). At baseline and follow-up, light-intensity physical activity (LIPA),
33 moderate-to-vigorous physical activity (MVPA) and sedentary behaviour were measured by
34 accelerometer and body adiposity (body mass index [BMI], body fat percentage [%BF]) was assessed
35 from measured height and weight and bioelectrical impedance analyser. Compositional regression
36 with robust estimators and compositional longitudinal isotemporal substitution analysis explored if,
37 and how, changes in movement behaviours were associated with adiposity.

38 **Results:** Over 7 years, the prevalence of obesity in the sample increased by 10.1% and 14.6% according
39 to BMI and %BF, respectively, and time spent in sedentary behaviour increased by 14%, while time
40 spent in LIPA and MVPA decreased by 14% and 21%, respectively. The increase in sedentary behaviour
41 at the expense of LIPA and MVPA during the seven-year period was associated with higher BMI and
42 %BF at follow-up (both $p < 0.01$). The increase in LIPA or MVPA at the expense of sedentary behaviour
43 was associated with reduced BMI and %BF at follow-up. In our sample, the largest change in BMI (0.75
44 kg/m^2 ; 95% confidence interval [CI]: 0.37-1.13) and %BF (1.28 units; 95% CI: 0.48-2.09) was
45 associated with longitudinal reallocation of 30 min from MVPA to sedentary behaviour.

46 **Conclusion:** We found an association between longitudinal changes in daily movement behaviours and
47 adiposity among elderly women in Central Europe. Our findings support public health programs to
48 increase or maintain time spent in higher intensity physical activity among elderly women.

49

50 INTRODUCTION

51 How people spend their time in movement-related behaviours throughout the day may influence their
52 body composition.¹ It is well accepted that spending more time in physical activity is related to
53 healthier body composition (i.e., the reduction of fat mass and increase of fat-free mass),² and that
54 spending more time in sedentary behaviour is related to less healthy body composition.³ Yet few
55 studies have explored these relationships among elderly people, and even fewer have used device-
56 based, longitudinal measures of movement behaviours. As studies that apply the compositional
57 methodology specifically to older people are scarce, there is a lack of evidence to underpin obesity
58 interventions and public health policy for older people based on reallocations of time between
59 movement behaviours. More evidence is required, as robust interventions for improved time use may
60 lead to better health and alleviate future economic costs amid an ageing population.

61 Most previous studies have considered movement behaviours such as physical activity and sedentary
62 behaviour to be independent predictors of obesity.⁴ However, movement behaviours are not
63 independent of each other – they are co-dependent.⁴⁻⁶ This is because movement behaviours take
64 place in time, and available time in a day is finite. Each day, we have 24 hours on disposal. To increase
65 the time spent in one behaviour, we must take this time from one or more other behaviours within
66 that same day.⁶⁻⁸ This means it does not make sense to explore the health associations of one
67 behaviour independently of the other behaviours. Instead, movement behaviours should be
68 considered relative to each other. This holds also for any subset of daily behaviours, e.g., waking
69 behaviours.^{9,10} Although waking behaviours do not necessarily sum to the same duration for every
70 participant (as with the 24-hour day), the data are nonetheless compositional when conceptualised as
71 scale invariant, i.e., we are interested in the relative proportions (or time shares) of behaviours rather
72 than absolute amounts. Thus, when considering the impact of changing one behaviour, we
73 simultaneously consider the impact of other behaviour(s) which are changed to compensate.
74 Accordingly, there has been a recent conceptual shift in behavioural epidemiology which moves away
75 from exploring movement behaviours as independent risk factors, towards an approach which allows

76 the influence of all behaviours to be considered relative to each other, i.e., a time-use epidemiology
77 approach.¹¹ This shift has been facilitated by the development of new analytical models based on
78 compositional data analysis.^{5,8,12}

79 Studies using a compositional approach to explore the associations between movement behaviours
80 and adiposity among older adults have, to our knowledge, all been cross-sectional.^{1,3,13} They suggest
81 that older adults who spend more time in moderate-to-vigorous physical activity (MVPA) and less time
82 in sedentary behaviour have better body composition. One study estimated body mass index (BMI) to
83 decrease by 0.7 units when 15 minutes were reallocated from sedentary behaviour to MVPA.¹³
84 Unexpectedly, this study reported the same estimated improvement (0.7 units) in BMI when 15
85 minutes of light physical activity (LIPA) were reallocated to MVPA. This suggests that in relation to
86 adiposity there is no benefit of LIPA over sedentary behaviour. However, cross-sectional reallocation
87 or isothermal substitution studies do not provide evidence on how within-person changes in
88 behaviour over time are associated with health outcomes. As such, their findings should be considered
89 cautiously when planning interventions and advising policy. Studies with longitudinal exposures are
90 required to provide evidence on how changes in time use, specifically how reallocating time between
91 movement behaviours, are associated with outcomes.

92 This study aimed to use an integrated time-use approach to: 1) investigate the prospective associations
93 between changes in daily movement behaviours and adiposity among elderly women; and 2) to
94 examine how the reallocation of time between movement behaviours was associated with longitudinal
95 changes in adiposity.

96

97 **SUBJECTS AND METHODS**

98 **Design and participants**

99 This was a longitudinal study with baseline data collected during 2009–2011 in three university cities
100 in Central Europe with very similar weather, cultural and economic conditions; namely, Olomouc in
101 Czech republic, Katowice in Poland, and Prešov in Slovakia. Older women were recruited from within

102 University of Third Age programs to participate in physical activity and body composition
103 measurements. The exclusion criteria for baseline study involvement were inability to walk without
104 any prosthetic aids and being under the age 60. Follow-up data collection was conducted during 2016–
105 2018. So that data in 2009–2011, and 7 years later in 2016–2018 were collected in the same month,
106 the exact date of follow-up data collection was individually tailored. The follow-up stage implemented
107 the same assessment methods, device settings, process and measurement conditions (measurement
108 protocol) that were used at the baseline stage.

109 At baseline, valid data were available from 325 older women. After seven years, all women were
110 approached and invited to get involved in the follow-up assessment. Out of 325 baseline participants:
111 36 died before follow-up; 57 were not able to continue participating in the study due to serious illness;
112 and 65 did not agree to complete all parts of the measurements. Thus, the follow-up sample consisted
113 of 167 women. Of these, 158 women had valid baseline and follow-up data, and were included in the
114 ensuing analyses.

115 Participation in the study was voluntary and women could withdraw from the study at any time. For
116 both baseline and follow-up measurements, all participants provided their written informed consent.
117 The study was carried out in accordance with the Declaration of Helsinki and was approved by the
118 institutional scientific ethics committee.

119

120 **Measurements**

121 *Movement behaviors: physical activity and sedentary behavior*

122 Physical activity and sedentary behavior were measured at baseline and follow-up using a uniaxial
123 ActiGraph GT1M accelerometer device (Manufacturing Technology Inc., FL, USA). The research staff
124 personally checked the fastening of the device at the right hip. The participants were instructed to
125 wear the accelerometer for eight consecutive days during waking hours with exception of bathing and
126 swimming. The accelerometer sampling interval was set at 1-minute epochs. Non-wear time was
127 defined by an interval of 60 consecutive minutes of zero counts per minute (cpm), allowing for 2

128 minutes of non-zero count interruptions. This algorithm is provided in the manufacture's software
129 (ActiGraph, LLC., Pensacola, FL, USA). For the assessment of accelerometer-derived movement
130 behaviours, a 'valid day' was defined as the one in which the participant had ≥ 10 h of wear time. To be
131 included in the analyses, the participants had to have valid data for at least 4 days (3 workdays and 1
132 weekend day) in both baseline and follow-up measurements.¹⁴ Amount of time spent in sedentary
133 behavior, LIPA and MVPA was derived for each valid day. For sedentary behavior, the cut-point of 100
134 counts/min was used as the commonly used threshold for senior populations.¹⁵ LIPA and MVPA levels
135 were defined according to Freedson cut-off points.¹⁶

136

137 *Body adiposity*

138 Body height was measured barefooted using a P-375 portable anthropometer to the nearest 0.1 cm
139 (Trystom, Olomouc, Czech Republic). Body weight (to the nearest 0.1 kg) and body fat percentage
140 (%BF) were assessed using the InBody 720 multi-frequency bioelectrical impedance analyser (Biospace
141 Co., Ltd., Seoul, Korea). All the women were required fast for at least 4 h, hydrate properly for 24 h
142 preceding the measurement. BMI was calculated as $\text{weight}/\text{height}^2$ (kg/m^2) and categorised as
143 'normal' weight ($< 25 \text{ kg}/\text{m}^2$), overweight ($25\text{--}29.9 \text{ kg}/\text{m}^2$), and obesity ($\geq 30 \text{ kg}/\text{m}^2$). Body fat
144 percentage (%BF) was classified as 'normal' ($\leq 35\%$) and obesity ($> 35\%$). Regardless of body weight and
145 physical activity level, multi-frequency bioelectrical impedance analysis has been suggested as a valid
146 method for body composition assessment in older women.¹⁷

147

148 *Statistical analyses*

149 Statistical analyses were performed using SPSS 22.0 software (SPSS Inc., an IBM Company, Chicago, IL,
150 USA) and R 3.4.2 software (R Foundation for Statistical Computing, Vienna, Austria). For baseline and
151 follow-up, the daily composition consisted of three parts of waking movement behaviour (sedentary
152 behaviour, LIPA and MVPA) and was closed to 16 h (assuming 8 h of sleep a day) for the purpose of
153 isotemporal substitution modelling. We assumed 8 h of daily sleep based on previous reports of sleep

154 duration in this age group,¹⁸ and the average non-wear time observed in this sample which included
155 sleep and potentially other activities, such as bathing, and lying awake in bed (14.1 ± 1.2 h and 13.6 ± 1.2
156 h, at baseline and follow-up, respectively). The composition for isotemporal substitution modelling
157 could be also closed to the mean wake/wear time (if such data are available). The statistical analyses
158 took the relative nature of movement behaviour data into consideration. This means that not absolute
159 values of movement behaviours but rather ratios between them formed the source of relevant
160 information.¹⁰ After ensuring there were no zero values in any compositional parts, the compositions
161 were expressed as pivot coordinates,¹⁹ being a special case of isometric log ratios (ILRs). Accordingly,
162 the ILRs were constructed in a specific way so that the first pivot coordinate included all relative
163 information regarding one dominant activity (numerator), versus the geometric mean of the remaining
164 activities (denominator). This first pivot coordinate can also be expressed as the (scaled) sum of log-
165 ratios; this is reflected by the schematic notation in Table 2. Three sets of pivot coordinates were
166 constructed, with each set treating a different activity (sedentary behaviour, LIPA and MVPA) as the
167 dominant activity.

168 The prospective associations between changes in daily movement behaviours and adiposity were
169 investigated via robust compositional regression models (with the MM-estimator of regression
170 parameters)¹⁰ in order to avoid possible influence of outlying observations,¹² with the follow-up
171 adiposity parameter as the dependent variable and differences between follow-up and baseline
172 movement behaviours (in terms of pivot coordinates) as the explanatory variables. To capture the
173 differences between follow-up and baseline for the aggregated relative effect of each compositional
174 part (sedentary behaviour, LIPA and MVPA) with respect to contributions of the remaining parts, three
175 models (one for each set of differences between the respective pivot coordinates) were conducted for
176 each adiposity parameter (BMI and %BF). Age, country, respective baseline adiposity parameter and
177 pivot coordinate representations of baseline movement behaviour composition were included as
178 covariates in each model.

179 To quantify how longitudinal reallocations of time between movement behaviours were associated
180 with changes in adiposity, the above-mentioned models were used for prediction purposes.
181 Differences between pivot coordinate representations of the hypothetical follow-up and mean
182 baseline movement behaviour compositions (that were linearly adjusted to sum to 16 hours) were
183 calculated to estimate BMI and %BF changes associated with one-to-one reallocations.⁸ The estimated
184 differences for BMI and BF, respectively, were calculated for time reallocations of 5, 15 and 30 minutes
185 and 95% confidence intervals (CI) were obtained. Significance level was set a $p < 0.05$. When the 95% CI
186 did not cover zero, the change was considered as significant.

187 A comprehensive explanation of the compositional analysis is included in Additional file.

188

189 **RESULTS**

190 Baseline and follow-up characteristics of the study sample are presented in Table 1. At baseline, the
191 average age was 63.9 years, the majority of participants were non-smokers (93%) and retired (87.3%)
192 with high prevalence (53.2%) of secondary or higher education. Over 7 years, the prevalence of obesity
193 in the sample increased by 10.1% and 14.6% according to BMI and %BF, respectively. The relative
194 difference between baseline and follow-up compositional means was 1.14, 0.86, and 0.79 which means
195 that time spent in sedentary behaviour increased by 14%, while time spent in LIPA decreased by 14%
196 and time spent in MVPA decreased by 21%.

197 The results displayed in Table 2 (Model 1, Row 1, $\beta = 1.34$ for BMI and 3.15 for %BF) indicate that the
198 increase in sedentary behaviour at the expense of LIPA and MVPA during the seven-year period was
199 associated with higher BMI and %BF at follow-up. The increase in LIPA (Model 2, Row 1) or MVPA
200 (Model 3, Row 1) at the expense of the other two behaviours was associated with reduced BMI and
201 %BF at follow-up. The aggregated relative increase of LIPA was not significant for BMI ($\beta = -0.65$,
202 $p = 0.110$), however this is not surprising because the respective pivot coordinate amalgamates log-
203 ratios with contradictory associations (see Model 1, Row 2 and Model 3, Row 2).

204 Table 3 and Figures 1 show the estimated changes in BMI and BF% associated with the change in
205 movement behaviour composition, i.e., with isothermal substitutions between behaviours. By change
206 in movement behaviour composition we mean that 0 to 30 min are reallocated from one behaviour in
207 the mean baseline composition to another behaviour in the mean follow-up composition (NB the mean
208 compositions are calculated as the geometric means of the behaviours, linearly adjusted to sum to the
209 total assumed waking time of 16 h). At significance level $p < 0.05$, the estimated changes in adiposity
210 parameters were significant for all reallocation cases apart for the change in %BF for reallocation from
211 LIPA to MVPA. The largest effect was observed when the change in movement pattern was
212 characterized by replacing the time spent in MVPA by the time spent in sedentary behaviour. We can
213 expect that a 30-min exchange from MVPA to sedentary behaviour would predict on average a
214 0.75 kg/m^2 increase in BMI and a 1.28 unit increase in %BF. We can also assume that the reverse
215 exchange of time between these behaviours would result in a 0.37 kg/m^2 decrease in BMI and a
216 0.65 unit decrease in %BF.

217

218 **DISCUSSION**

219 This longitudinal study among older women revealed that reallocations of time from a higher-intensity
220 to a lower-intensity movement behaviour were associated with higher adiposity. It also seems that
221 reallocations of the same amount of time in the opposite direction (i.e. from a lower-intensity to a
222 higher-intensity movement behaviour) may be associated with smaller reductions in adiposity.

223 Our findings of decreases in physical activity and increases in sedentary behaviour over time generally
224 align with longitudinal findings previously reported among older populations.^{20,21} This may be due to
225 increasing physical impairment, co-morbidities, and changes in work, family and social commitments
226 as people age. We are unaware of any previous studies prospectively linking changes in device-
227 measured movement behaviours with adiposity in older adults; however, the findings of this study
228 partially concur with cross-sectional evidence.^{1,13} Less time spent in MVPA in favour of other
229 movement behaviours of lower intensity (LIPA and sedentary behaviour) has consistently emerged as

230 the most detrimental factor in the association with adiposity among not only older adults, but across
231 the lifespan. This is not surprising, as MVPA requires higher energy expenditure than LIPA and
232 sedentary behaviour. Thus, replacing MVPA with LIPA and sedentary behaviour may lead to an
233 imbalance between energy intake and energy expenditure, and subsequent gain of excess fat.
234 However, our study does not provide evidence on the direction of causation. Reverse causation is also
235 plausible – as adiposity increases, time spent in MVPA is replaced by behaviours requiring lower energy
236 expenditure. It is also possible that the relationship is bidirectional. A previous study suggested that
237 obesity may lead to a subsequent increase in sedentary behaviour among middle-aged and older
238 adults.²² However, this study did not conceptualise sedentary behaviour as a part of the time-use
239 composition and examine reallocations of time between different behaviours.

240 Contrary to previous cross-sectional studies,³ we found beneficial associations of adiposity status with
241 reallocations of time from sedentary behaviour to LIPA. However, these associations were weak. The
242 reallocation of 30 minutes from sedentary behaviour to LIPA was associated with -0.10 and -0.27 units
243 change in BMI and %BF, respectively. By comparison, the reallocation of 30 minutes from sedentary
244 behaviour to MVPA was associated with a much larger change in BMI (-0.37 units; -1.4%) and %BF ($-$
245 0.65 units; -1.9%). However, increasing MVPA by 30 minutes represents an increase of 75% from
246 baseline daily MVPA. This may not be an achievable intervention goal, particularly among older adults.
247 However, the reallocation of 30 minutes from sedentary behaviour to LIPA requires a comparatively
248 small behavioural change (only a 7% increase from baseline daily LIPA). To obtain the same difference
249 in BMI units (-0.37 units) estimated for reallocating 30 minutes from sedentary behaviour to MVPA,
250 104 minutes could be reallocated from sedentary behaviour to LIPA. This suggests that, in this
251 particular context, each minute of MVPA is worth around 3.5 minutes of LIPA. Similarly for %BF, the
252 reallocation of 71 minutes from sedentary behaviour to LIPA would be needed to get the same
253 estimated difference (-0.65 units). Such a reallocation strategy may be more feasible for older adults,
254 as LIPA is incidental to daily living and can be accumulated by simple modifications to daily activities,
255 such as slow walking to visit friends rather than driving.

256 Consistent with other studies using compositional data analysis,^{6,13} we found asymmetrical responses
257 in adiposity depending on whether MVPA was increased or decreased. In our sample, the average
258 benefits estimated for the reallocation of a set duration of time to MVPA were not as large as the
259 estimated worsening of the adiposity status when the same duration was reallocated away from
260 MVPA. This asymptotic dose-response relationships between PA and health outcomes are a common
261 finding in the literature.²³ For example, the relationship between physical activity dosage and all-cause
262 mortality has consistently been found to be asymptotic.²⁴ Some studies have found an asymptotic
263 relationship also between exercise dosage and weight loss.^{25,26} It should be noted, however, that most
264 of these studies did not use compositional data analysis and account for co-dependence between time-
265 use components.^{1,4} The asymmetry of estimated responses can be observed in Figures 1 and suggests
266 that the relative benefits obtained from avoiding a quantum fall in current levels of MVPA are greater
267 than the relative benefits accrued by an increase of the same quantum. This would suggest that the
268 maintenance of MVPA is an important intervention goal, particularly as people age and their MVPA
269 levels tend to decline. It should be noted, however, that the confidence intervals for absolute values
270 of the estimated changes in adiposity for reallocations to and from MVPA were overlapping, which
271 means that we cannot generalise about the asymmetry beyond our study sample.

272 Our study provided evidence to suggest that interventions enabling elderly women to shift time from
273 lower to higher intensity behaviours have the potential to decrease adiposity. Replacing sedentary
274 behaviour with MVPA appears to be the best strategy, but larger replacements of sedentary behaviour
275 with LIPA may achieve similar gains. If increasing time spent in MVPA is not feasible, our study suggests
276 that it may be worthwhile to support elderly people to maintain their current MVPA levels. Programs
277 to create safe environments and opportunities for MVPA may be warranted. A previous study
278 suggested the role of LIPA should be an important focus for future studies.²⁷ Our findings support the
279 recommendation in the context of obesity research. These findings are particularly relevant from a
280 public health perspective, because Central Europe has an aging population, consistent with most other
281 European countries, as life expectancy is increasing. However, unlike in other European regions, the

282 overall population in Central Europe is predicted to decline²⁸ due to low birth rates, a strong emigration
283 drive, and restrictive immigration policies. Evidence to inform healthier daily movement behaviours
284 among older people is, therefore, becoming increasingly important, especially among Central
285 Europeans, who are already lagging behind other countries in terms of their obesity status and overall
286 health.²⁹

287 The strengths of this study include the repeated measures of movement behaviours spanning 7 years,
288 using identical measurement procedures, and using accelerometers. Longitudinal data of older adults'
289 movement behaviours are scarce and rarely reported. However, we only had two points of data
290 measurement, meaning patterns of change may not have been detected. Adiposity indicators were
291 measured using standardised procedures and analyses were conducted using statistical models that
292 are appropriate for the relative nature of movement behaviour data. The generalizability of the study
293 is limited due to its non-probability convenience sample with very few smokers, high prevalence of
294 higher education and participation in organized PA (57 % participating one or more times/week). In
295 addition, our sample only included women, meaning results cannot be extrapolated to men without
296 caution. It is possible that our findings are confounded by unmeasured factors such as dietary changes
297 and smoking habits. Additionally, although we used the most common cut-points in accelerometry
298 data analyses,¹⁵ different cut-points can substantially impact the classification of the proportion of
299 time spent in different movement behaviours in a sample of older women. It should be considered
300 that our measurement protocol did not include examination of sleep duration, which may have
301 confounded findings as sleep is co-dependent with movement behaviours and longer sleep appears to
302 be beneficially associated with adiposity.³⁰ It is possible that the exclusion of sleep has led to
303 overestimation of the benefits of MVPA or LIPA and conservative estimates for the unfavourable
304 influence of sedentary behaviour. Additionally, for analytical purposes and interpretability of
305 estimates, we linearly adjusted the waking-day compositions to sum to 16 hours when average wear
306 times were between 14.1±1.2 and 13.6±1.2 hours. This implies that the composition of behaviours

307 during the unmeasured period of waking time is the same as during the measured period, which may
308 not necessarily be the case.

309 In conclusion, we found an association between changes in daily movement behaviours and adiposity
310 among elderly women in Central Europe. Increases in MVPA and LIPA, and decreases in sedentary
311 behaviour were beneficially associated with adiposity indicators. Our findings support public health
312 programs to increase or maintain time spent in higher intensity physical activity among elderly women.

313

314 **ACKNOWLEDGEMENT**

315 The authors would like to thank the participants for their commitment to the study.

316 The study was supported by a research grant from the Czech Science Foundation No. 18-16423S.

317

318 **CONFLICTS OF INTEREST**

319 The authors declare no conflict of interest.

320

321 **REFERENCES**

- 322 1. Grgic J, Dumuid D, Bengoechea EG, Shrestha N, Bauman A, Olds T, et al. Health outcomes
323 associated with reallocations of time between sleep, sedentary behaviour, and physical
324 activity: A systematic scoping review of isotemporal substitution studies. *Int J Behav Nutr Phys*
325 *Act.* 2018;15(1):1–68.
- 326 2. Hill J, Wyatt H. Role of physical activity in preventing and treating obesity. *J Appl Physiol.*
327 2005;99(2):765–770.
- 328 3. Pelclová J, Štefelová N, Hodonská J, Dygrýn J, Gába A, Zajac-Gawlak I. Reallocating time from
329 sedentary behavior to light and moderate-to-vigorous physical activity: What has a stronger
330 association with adiposity in older adult women? *Int J Environ Res Public Health.*
331 2018;15(7):1444.
- 332 4. Pedišić Ž. Measurement issues and poor adjustments for physical activity and sleep

- 333 undermine sedentary behaviour research - The focus should shift to the balance between
334 sleep, sedentary behaviour, standing and activity. *Kinesiology*. 2014;46(1):135–146.
- 335 5. Dumuid D, Stanford TE, Martin-Fernández J-A, Pedišić Ž, Maher CA, Lewis LK, et al.
336 Compositional data analysis for physical activity, sedentary time and sleep research. *Stat*
337 *Methods Med Res*. 2018;27(12):3726-3738.
- 338 6. Chastin SFM, Palarea-Albaladejo J, Dontje ML, Skelton DA. Combined Effects of Time Spent in
339 Physical Activity, Sedentary Behaviors and Sleep on Obesity and Cardio-Metabolic Health
340 Markers: A Novel Compositional Data Analysis Approach. *PLoS One*. 2015;10(10):e0139984.
- 341 7. Mekary RA, Willett WC, Hu FB, Ding EL. Isotemporal substitution paradigm for physical activity
342 epidemiology and weight change. *American journal of epidemiology*, 2009;170(4):519-527.
- 343 8. Dumuid D, Pedišić Ž, Stanford TE, Martín-Fernández J-A, Hron K, Maher CA, et al. The
344 compositional isotemporal substitution model: A method for estimating changes in a health
345 outcome for reallocation of time between sleep, physical activity and sedentary behaviour.
346 *Stat Methods Med Res*. 2017;28(3):846-857.
- 347 9. Buccianti A., Pawlowsky-Glahn V. New perspectives on water chemistry and compositional
348 data analysis. *Mathematical Geology*. 2005;37(7):703-727.
- 349 10. Pawlowsky-Glahn V, Egozcue JJ, Tolosana-Delgado R. (eds). *Modeling and analysis of*
350 *compositional data*. (John Wiley & Sons, Chichester, UK, 2015).
- 351 11. Pedišić Ž, Dumuid D, Olds TS. Integrating sleep, sedentary behaviour, and physical activity
352 research in the emerging field of time-use epidemiology: Definitions, concepts, statistical
353 methods, theoretical framework, and future directions. *Kinesiology*. 2017;49(2):252–269.
- 354 12. Štefelová N, Dygrýn J, Hron K, Gába A, Rubín L, Palarea-Albaladejo J. Robust Compositional
355 Analysis of Physical Activity and Sedentary Behaviour Data. *Int J Environ Res Public Health*.
356 2018;15(10):2248.
- 357 13. Dumuid D, Lewis LK, Olds TS, Maher C, Bondarenko C, Norton L. Relationships between older
358 adults' use of time and cardio-respiratory fitness, obesity and cardio-metabolic risk: A

- 359 compositional isotemporal substitution analysis. *Maturitas*. 2018;110:104–110.
- 360 14. Hart TL, Swartz AM, Strath SJ. How many days of monitoring are needed to accurately
361 estimate physical activity in older adults. *Int J Behav Nutr Phys Act*. 2011;8:62–69.
- 362 15. Gorman E, Hanson HM, Yang PH, Khan KM, Liu-Ambrose T, Ashe MC. Accelerometry analysis
363 of physical activity and sedentary behavior in older adults: a systematic review and data
364 analysis. *European Review of Aging and Physical Activity*. 2014;11(1):35.
- 365 16. Freedson PS, Melanson E, Sirard J. Calibration of the Computer Science and Applications, Inc.
366 accelerometer. *Med Sci Sports Exerc*. 1998;30(5):777–781.
- 367 17. Gába A, Kapuš O, Cuberek R, Botek M. Comparison of multi- and single-frequency bioelectrical
368 impedance analysis with dual-energy X-ray absorptiometry for assessment of body
369 composition in post-menopausal women: effects of body mass index and accelerometer-
370 determined physical activity. *J Hum Nutr Diet*. 2015;28(4):390-400.
- 371 18. da Silva AA., de Mello RGB, Schaan CW, Fuchs FD, Redline S, Fuchs SC. Sleep duration and
372 mortality in the elderly: a systematic review with meta-analysis. *BMJ open*. 2016;6(2):
373 e008119.
- 374 19. Hron K, Filzmoser P, Thompson K. Linear regression with compositional explanatory variables.
375 *J Appl Stat*. 2012;39(5):1115–1128.
- 376 20. Smith L, Gardner B, Fisher A, Hamer M. Patterns and correlates of physical activity behaviour
377 over 10 years in older adults: prospective analyses from the English Longitudinal Study of
378 Ageing. *BMJ open*. 2015;5(4):e007423.
- 379 21. Hagströmer M, Kwak L, Oja P, Sjöström M. A 6 year longitudinal study of accelerometer-
380 measured physical activity and sedentary time in Swedish adults. *Journal of science and
381 medicine in sport*. 2015;18(5): 553-557.
- 382 22. Pedišić Ž, Grunseit A, Ding, Chau JY, Banks E, Stamatakis E, et al. High sitting time or obesity:
383 Which came first? Bidirectional association in a longitudinal study of 31,787 Australian adults.
384 *Obesity*. 2014;22(10):2126–2130.

- 385 23. Woodcock J, Franco OH, Orsini N, Roberts I. Non-vigorous physical activity and all-cause
386 mortality: systematic review and meta-analysis of cohort studies. *International journal of*
387 *epidemiology*. 2011;40(1):121-138.
- 388 24. Arem H, Moore SC, Patel A, Hartge P, De Gonzalez AB, Visvanathan K, et al. Leisure time
389 physical activity and mortality: a detailed pooled analysis of the dose-response relationship.
390 *JAMA internal medicine*. 2015;175(6):959-967.
- 391 25. Ross R, Janssen IAN. Physical activity, total and regional obesity: dose-response
392 considerations. *Medicine and science in sports and exercise*. 2001;33(6; SUPP):S521-S527.
- 393 26. Khanal S, Choi L, Innes-Hughes C, Rissel C. Dose response relationship between program
394 attendance and children's outcomes in a community based weight management program for
395 children and their families. *BMC public health*. 2019;19(1):716.
- 396 27. Chastin SFM, De Craemer M, De Cocker K, Powell L, Van Cauwenberg J, Dall P, et al. How does
397 light-intensity physical activity associate with adult cardiometabolic health and mortality?
398 Systematic review with meta-analysis of experimental and observational studies. *Br J Sports*
399 *Med*. 2019;53(6):370-376.
- 400 28. United Nations, Department of Economic and Social Affairs, Population Division. *World*
401 *Population Prospects: The 2015 Revision, Key Findings and Advance Tables*. Working Paper
402 No. ESA/P/WP.241. (United Nations, New York, 2015).
- 403 29. Sowa A, Topór-Mądry R, Tobiasz-Adamczyk B, Golinowska S. Health status of older people.
404 Evidence from Europe. *Zesz Nauk Ochr Zdr*. 2015;13(4):381-396.
- 405 30. Patel SR, Hu FB. Short sleep duration and weight gain: a systematic review. *Obesity*.
406 2008;16(3): 643-653.

407

Table 1. Baseline and follow-up characteristics of the study sample

	Baseline	Follow-up
	Mean (SD)	Mean (SD)
<i>Age and nationality</i>		
Age (years)	63.9 (4.4)	
Czech, Polish, Slovak (n (%))	63 (39.9), 62 (39.2), 33 (20.9)	
<i>Anthropometrics</i>		
Body height (cm)	160.1 (6.8)	159.7 (6.9)
Body weight (kg)	68.4 (10.7)	69.6 (11.6)
Body mass index (kg/m ²)	26.5 (4.1)	27.3 (4.3)
Body fat percentage (%)	34.3 (6.9)	36.7 (6.8)
<i>Wear time and activity composition</i>		
Wear time (h)	14.1 (1.2)	13.6 (1.2)
Compositional mean of SB, LIPA, MVPA (min) ^a	505, 415, 40	573.5, 354.9, 31.6
Compositional mean of SB, LIPA, MVPA (%)	52.6, 43.2, 4.2	59.7, 37, 3.3
<i>Weight status according to BMI, n (% of n)</i>		
'Normal' weight (<25 kg/m ²)	60 (38)	75 (47.5)
Overweight (25–29.9 kg/m ²)	69 (43.7)	38 (24.0)
Obesity (≥30 kg/m ²)	29 (18.3)	45 (28.5)
<i>Obesity status according to %BF, n (% of n)</i>		
'Normal' (≤35%)	87 (55.1)	64 (40.5)
Obesity (>35%)	71 (44.9)	94 (59.5)

SD = standard deviation; SB = sedentary behaviour; LIPA = light physical activity; MVPA = moderate-to-vigorous physical activity; BMI = body mass index; %BF = body fat percentage.

^a composition closed to 16 hours.

Table 2. Pivot coordinate compositional MM-regression estimates for models with the follow-up adiposity measures as response variables.

	Body mass index (kg/m ²)		Body fat (%)	
	β_{irr} (SE)	p-value	β_{irr} (SE)	p-value
Model 1				
SB/LIPA+SB/MVPA) difference	1.34 (0.40)	< 0.001	3.15 (0.82)	<0.001
(LIPA/MVPA) difference	0.02 (0.29)	0.940	−0.50 (0.71)	0.480
Model 2				
(LIPA/SB+LIPA/MVPA) difference	−0.65 (0.41)	0.110	−2.01 (0.96)	0.040
(SB/MVPA) difference	1.17 (0.27)	<0.001	2.48 (0.50)	<0.001
Model 3				
(MVPA/SB+MVPA/LIPA) difference	−0.69 (0.19)	<0.001	−1.14 (0.41)	0.006
(SB/LIPA) difference	1.15 (0.45)	0.010	2.98 (1.00)	0.003

Note: All models were adjusted for age, country, and movement behaviour compositions at baseline. The first pivot coordinate has been expressed as the sum of individual log-ratios for ease of interpretation (a comprehensive explanation of the compositional analysis is included in Additional file).

BMI = body mass index; BF% = body fat percentage; β = unstandardised regression coefficient; SE = standard error; SB = sedentary behaviour; LIPA = light physical activity; MVPA = moderate-to-vigorous physical activity

Table 3. Estimated changes (and their 95% confidence intervals) in follow-up BMI and %BF associated with time reallocation between baseline and follow-up movement behaviour composition.

Reallocation	Body mass index (kg/m ²)			Body fat (%)		
	5 min	15 min	30 min	5 min	15 min	30 min
SB to LIPA	-0.02 (-0.03, -0.00)	-0.05 (-0.09, -0.01)	-0.10 (-0.19, -0.02)	-0.05 (-0.08, -0.01)	-0.14 (-0.23, -0.04)	-0.27 (-0.46, -0.09)
SB to MVPA	-0.07 (-0.11, -0.04)	-0.20 (-0.30, -0.10)	-0.37 (-0.54, -0.19)	-0.13 (-0.20, -0.06)	-0.36 (-0.56, -0.16)	-0.65 (-1.01, -0.30)
LIPA to SB	0.02 (0.00, 0.03)	0.05 (0.01, 0.09)	0.10 (0.02, 0.19)	0.05 (0.01, 0.08)	0.14 (0.04, 0.23)	0.27 (0.08, 0.46)
LIPA to MVPA	-0.06 (-0.09, -0.02)	-0.15 (-0.25, -0.05)	-0.26 (-0.45, -0.07)	-0.08 (-0.17, 0.00)	-0.22 (-0.46, 0.02)	-0.37 (-0.80, 0.06)
MVPA to SB	0.08 (0.04, 0.12)	0.28 (0.14, 0.41)	0.75 (0.37, 1.13)	0.14 (0.06, 0.22)	0.48 (0.19, 0.77)	1.28 (0.48, 2.09)
MVPA to LIPA	0.06 (0.02, 0.11)	0.23 (0.08, 0.37)	0.65 (0.26, 1.04)	0.10 (0.00, 0.19)	0.35 (0.02, 0.67)	1.02 (0.14, 1.89)

BMI = body mass index; %BF = body fat percentage; SB = sedentary behaviour; LIPA = light-intensity physical activity; MVPA = moderate-to-vigorous physical activity




Figure 1. Estimated changes in follow-up body mass index and body fat percentage associated with time reallocation between baseline and follow-up movement behaviour composition.

BMI = body mass index; %BF = body fat percentage; SB = sedentary behaviour; LIPA = light-intensity physical activity; MVPA = moderate-to-vigorous physical activity