



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
Main Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2020

HRS/EHRA/APHRS/LAHRS/ACC/AHA Worldwide Practice Update for Telehealth and Arrhythmia Monitoring During and After a Pandemic

Varma, Niraj ; Marrouche, Nassir F ; Aguinaga, Luis ; Albert, Christine M ; Arbelo, Elena ; Choi, Jong-II ; Chung, Mina K ; Conte, Giulio ; Dagher, Lilas ; Epstein, Laurence M ; Ghanbari, Hamid ; Han, Janet K ; Heidbuchel, Hein ; Huang, He ; Lakkireddy, Dhanunjaya R ; Ngarmukos, Tachapong ; Russo, Andrea M ; Saad, Eduardo B ; Saenz Morales, Luis C ; Sandau, Kristin E ; Sridhar, Arun Raghav M ; Stecker, Eric C ; Varosy, Paul D

DOI: <https://doi.org/10.1161/CIRCEP.120.009007>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-195538>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Varma, Niraj; Marrouche, Nassir F; Aguinaga, Luis; Albert, Christine M; Arbelo, Elena; Choi, Jong-II; Chung, Mina K; Conte, Giulio; Dagher, Lilas; Epstein, Laurence M; Ghanbari, Hamid; Han, Janet K; Heidbuchel, Hein; Huang, He; Lakkireddy, Dhanunjaya R; Ngarmukos, Tachapong; Russo, Andrea M; Saad, Eduardo B; Saenz Morales, Luis C; Sandau, Kristin E; Sridhar, Arun Raghav M; Stecker, Eric C; Varosy, Paul D (2020). HRS/EHRA/APHRS/LAHRS/ACC/AHA Worldwide Practice Update for Telehealth and Arrhythmia Monitoring During and After a Pandemic. *Circulation. Arrhythmia and Electrophysiology*, 13(7):e009007.

DOI: <https://doi.org/10.1161/CIRCEP.120.009007>



HRS/EHRA/APHRS/LAHRs/ACC/AHA Worldwide Practice Update for Telehealth and Arrhythmia Monitoring During and After a Pandemic

Niraj Varma¹, MA, MD, PhD, FACC, FRCP; Nassir F. Marrouche, MD, FHRS; Luis Aguinaga, MD, MBA, PhD, FESC, FACC*; Christine M. Albert, MD, MPH, FHRS, FACCT; Elena Arbelo², MD, MSci, PhD†; Jong-Il Choi³, MD, PhD, MHS§; Mina K. Chung⁴, MD, FHRS, FACCT||; Giulio Conte, MD, PhD‡; Lilas Dagher⁵, MD†; Laurence M. Epstein⁶, MD, FACCT; Hamid Ghanbari, MD, FACCT; Janet K. Han, MD, FHRS, FACCT; Hein Heidbuchel⁷, MD, PhD, FESC, FEHRA‡; He Huang, MD, FACC, FESC, FEHRA§; Dhanunjaya R. Lakkireddy⁸, MD, FHRS, FACCT; Tachapong Ngarmukos, MD, FAPHRs, FACCT§; Andrea M. Russo, MD, FHRS, FACCT; Eduardo B. Saad, MD, PhD, FHRS, FESC*; Luis C. Saenz Morales, MD*; Kristin E. Sandau, PhD, RN||; Arun Raghav M. Sridhar, MD, MPH, FACCT; Eric C. Stecker, MD, MPH, FACCT¶; Paul D. Varosy, MD, FHRS, FACCT¶

Key Words: COVID-19 ■ pandemic ■ QT interval ■ remote monitoring ■ telemedicine

Coronavirus disease 2019 (COVID-19), caused by the severe acute respiratory syndrome coronavirus 2, started in the city of Wuhan late in 2019. Within a few months, the disease spread toward all parts of the world and was declared a pandemic on March 11, 2020. The current healthcare dilemma worldwide is how to sustain the capacity for quality services not only for those suffering from COVID-19 but also for non-COVID-19 patients, all while protecting physicians, nurses, and other allied healthcare workers.

The pandemic poses challenges to electrophysiologists at several levels. Hospitalized patients who are COVID-19-positive may have preexisting arrhythmias, develop new arrhythmias, or be placed at increased

arrhythmic risk from therapies for COVID-19. Cardiac arrhythmia incidence in hospitalized patients has been documented in a few published studies, with reported rates of 7.9%¹ and 16.7%² in hospitals in New York City and Wuhan, respectively, and up to 44%² in patients requiring intensive care. Life-threatening arrhythmias (ventricular tachycardia/ventricular fibrillation) can occur in up to 6% of hospitalized patients with COVID-19 infection.³ There have also been several case reports of atrioventricular block in hospitalized patients, which is otherwise rarely described during viral illness.^{4,5} Although the residual left ventricular dysfunction and arrhythmic risk are currently unknown, preliminary pathophysiological,⁶ histological,⁷ and imaging⁸ data suggest that severe

Correspondence to: Heart Rhythm Society, 1325 G Street NW, Suite 400, Washington, DC 20005. Email clinicaldocs@hrsonline.org

Developed in partnership with and endorsed by the American College of Cardiology (ACC), the American Heart Association (AHA), the Asia Pacific Heart Rhythm Society (APHRS), the European Heart Rhythm Association (EHRA), the Heart Rhythm Society (HRS), and the Latin American Heart Rhythm Society (LAHRs).

This article has been copublished in *Circulation: Arrhythmia and Electrophysiology*, *EP Europace*, the *Journal of the American College of Cardiology*, the *Journal of Arrhythmia*, and *HeartRhythm*. These articles are identical except for minor stylistic and spelling differences in keeping with each journal's style.

*Representative of the Latin American Heart Rhythm Society (LAHRs).

†Representative of the Heart Rhythm Society (HRS).

‡Representative of the European Heart Rhythm Association (EHRA).

§Representative of the Asia Pacific Heart Rhythm Society (APHRS).

||Representative of the American Heart Association (AHA).

¶Representative of the American College of Cardiology (ACC).

For Disclosures, see Appendix 1.

© 2020 The Heart Rhythm Society; American Heart Association, Inc.; European Heart Rhythm Association, a registered branch of the European Society of Cardiology; American College of Cardiology Foundation; Asia Pacific Heart Rhythm Society; the Latin American Heart Rhythm Society. Published by [Elsevier Inc./Wolters Kluwer Health/Oxford University Press/Wiley]. This article is published under the [Creative Commons CC-BY](https://creativecommons.org/licenses/by/4.0/) license.

Circulation: Arrhythmia and Electrophysiology is available at www.ahajournals.org/journal/circep

Nonstandard Abbreviations and Acronyms

CIED	cardiovascular implantable electronic device
COVID-19	coronavirus disease 2019
SARS-CoV-2	severe acute respiratory syndrome coronavirus 2

acute respiratory syndrome coronavirus 2 infection holds the potential to induce durable myocardial changes predisposing to arrhythmias or heart failure.

Electrocardiographic monitoring and inpatient monitoring services may become necessary but face the potential hurdles of limited telemetry wards, contamination of equipment and infection of healthcare personnel, and shortage of personal protective equipment.^{9,10} In parallel, there is a continued responsibility to maintain care of patients who are COVID-19-negative with arrhythmias. These pressures have led to inventive utilization and adaptation of existing telemedicine technologies as alternative options.

This document discusses how digital health may facilitate electrophysiology practice for patients with arrhythmia, whether hospitalized for COVID-19 or not. The representation of authors from some of the most severely affected countries, such as China, Spain, Italy, and the United States, is a tribute from our worldwide community to those colleagues who have worked on the front lines of the pandemic.

MONITORING STRATEGIES DURING A PANDEMIC: HERE TO STAY

In light of the current pandemic, monitoring strategies should focus on selecting high-risk patients in need of close surveillance and using alternative remote recording devices to preserve personal protective equipment and protect healthcare workers from potential contagious harm.

Inpatient

For inpatient monitoring, telemetry is reasonable when there is concern for clinical deterioration (as may be indicated by acute illness, vital signs, or sinus tachycardia), or in patients with cardiovascular risk factors and/or receiving essential QT-prolonging medications. Telemetry is generally not necessary for persons under investigation without concern for arrhythmias or clinical deterioration and in those not receiving QT-prolonging drug therapy. In situations in which a hospital's existing telemetry capacity has been exceeded by patient numbers or when conventional telemetry monitoring is not feasible, such as off-site or nontraditional hospital

units, mobile devices may be used, for example, mobile cardiac outpatient telemetry as an adjunctive approach to support inpatient care.^{11–15} The majority of mobile cardiac outpatient telemetry devices can provide continuous arrhythmia monitoring using a single-lead ECG and allow for real-time and offline analysis of long-term ECG data. Telemetry can be extended using patch monitoring.^{16,17} Smartphone ECG monitors are wireless and have also been utilized during the current pandemic. Information is limited, however, on how parameters such as QTc measured on a single- (or limited number) lead ECG can reliably substitute for 12-lead ECG information.^{18,19} In one study, QT was underestimated by smartphone single-lead ECG.²⁰

Outpatient

The principles of remote patient management, crossing geographic, social, and cultural barriers, can be extended to outpatient care and are important to maintain continuity of care for non-COVID-19 patients.^{21–23} Virtual clinics move far beyond simple telephone contacts by integrating information from photos, video, mobile heart rhythm, and mobile health devices recording ECG, and remote cardiovascular implantable electronic device (CIED) interrogations.²⁴ A variety of platforms have been developed and used specifically to provide telehealth to patients via video teleconferencing^{25,26} (Table 1). Most healthcare centers have expanded use of telemedicine, with some reporting 100% transformation of in-person clinic visits to telemedicine-based visits to maintain care for non-COVID-19 patients, thus obviating their need to come to the hospital or clinic. This supplements social distancing measures and reduces the risk of transmission, especially for the older and more vulnerable populations. It also becomes a measure to control intake into emergency rooms and outpatient facilities and to permit rapid access when necessary to subspecialists.

Electrophysiology is well placed for virtual consultations. All preobtained data, including ECGs, ambulatory ECG monitoring, cardiac imaging, and coronary angiography can be adequately reviewed electronically. Digital tools such as direct-to-consumer mobile ECG (Table 2) and wireless blood pressure devices can be used to further complement the telehealth visit without in-person contact. CIED, wearable/mobile health, and clinical data can be integrated into clinician workflow.

Additional diagnostic information might be obtained without in-person contact using home enrollment of prescribed ambulatory rhythm monitors. Patch monitors can be mailed to patient homes and easily self-affixed, unlike Holter monitors with cables and electrodes requiring placement by healthcare workers. In some cases, new or follow-up telehealth visits will require an adjunctive in-person visit to perform

a 12-lead ECG, ECG stress test, echocardiogram, or other diagnostic procedures. Occasionally, conventional clinic visits are required to accurately assess the impact of comorbidities or frailty on procedural risk, or to allow comfortable discussion with multiple family members when planning procedures with high risk. Telephone-only visits (ie, without video) may allow for a broader reach because of ease and ubiquitous accessibility as a communication strategy for immediate access for urgent matters.

There are many barriers to implementation, such as inadequate reimbursement, licensing/regulatory and privacy issues, lack of infrastructure, resistance to change, lack of access/poor internet coverage, restricted financial resources, and limited technical skills (eg, in the elderly patient population). Some telehealth and remote ECG monitoring technologies may be simply unaffordable and/or unavailable, leading to different levels of uptake within communities and across the globe. All stakeholders should collaborate to address these challenges and promote the safe and effective use of digital health during the current pandemic. In recent months, regulations have been eased to permit consults with new patients, issuing prescriptions, and obtaining consents. In that sense, the COVID-19 pandemic may serve as an opportunity to evolve current technologies into indispensable tools for our future cardiological practice.

THERAPY FOR COVID-19 AND POTENTIAL ELECTRICAL EFFECTS

No specific cure exists for COVID-19.^{28–30} Potential COVID-19 therapies, especially hydroxychloroquine and azithromycin, are being investigated in ongoing trials but also have been used off label in many parts of the world. These may exert QT-prolonging effects³¹ (Table 3) and, since recent observational data have questioned their efficacy, require a careful risk-benefit adjudication.³² Combination therapy (eg, hydroxychloroquine and azithromycin) may have synergistic effects on QT prolongation.^{33,34} In a retrospective cohort study of 1438 patients with COVID-19 hospitalized in metropolitan New York (ie, a disease epicenter), cardiac arrest was more frequent in patients who received hydroxychloroquine with azithromycin compared with patients who received neither drug.³⁵ The adjusted hazard ratio for in-hospital mortality for treatment with hydroxychloroquine alone was 1.08, for azithromycin alone was 0.56, and for combined hydroxychloroquine and azithromycin was 1.35. However, none of these hazard ratios were statistically significant. The observational design of this study may limit interpretation of these findings. In the absence of clear efficacy data, treatment options should be individualized taking into account their proarrhythmic potential for torsade de pointes, which may be enhanced by

concomitant administration of other QT-prolonging drugs (eg, antiarrhythmics, psychotropics, etc).

In COVID-19 patients receiving prior antiarrhythmic therapy, there should be a thorough consideration of risk versus benefit before initiating any QT-prolonging COVID-19 therapies, especially considering their unproven value. For instance, although some recent observational studies highlight adverse effects of hydroxychloroquine in treating this infectious disease, its use is likely to persist outside of randomized trials because of its affordability and global availability compared with, for example, remdesivir.³⁵ If one of these drugs is judged to be critical, monitoring should be initiated. If life-threatening arrhythmias (ventricular tachycardia/ventricular fibrillation) occur, the benefit of antiarrhythmic drugs, notably amiodarone, outweighs the potential harm of hydroxychloroquine or other QT-prolonging drugs targeting COVID-19, and in these cases antiarrhythmic drugs should be prioritized and used as deemed necessary. Most importantly, all modifiable predisposing factors for QTc prolongation (electrolyte disturbances, drug-to-drug interaction) that may enhance arrhythmia susceptibility should be corrected, and the small subset of individuals with an underlying genetic predisposition such as congenital long QT syndrome (in whom QTc-prolonging medications are contraindicated) should be identified. Additionally, caution must be exercised in case of subclinical or manifest myocarditis that may increase the vulnerability to proarrhythmias associated with QT-prolonging drugs.

If drugs that exert a QT-prolonging effect are to be initiated in an inpatient setting, a baseline 12-lead ECG should be acquired. Following review of the QTc, patients can be stratified into low-risk group (QTc of <500 ms or <550 ms in the setting of wide baseline QRS) or high-risk group (baseline QTc of \geq 500 ms or \geq 550 ms in the setting of wide baseline QRS, or patients who are started on combination therapies), guiding selection of telemetered versus nonmonitored beds.⁵⁰ Low-risk patients treated with QT-prolonging agents may be monitored using MCT (or another available wearable) with twice-a-day transmission of QTc measurements and any urgent alerts. High-risk patients would require more continuous monitoring and follow-up QTc measurements using telemetry preferably (but if unavailable, other remote monitoring devices). A second QTc assessment via telemetry or other remote devices after 2 doses may be helpful in identifying QTc reactors—patients who have an exaggerated response to QT-prolonging agents. An increase in QTc by \geq 60 ms or to QTc \geq 500 ms on any follow-up QT assessment is considered significant and should prompt a reassessment of risks versus benefits of continuing the drug.

In the outpatient setting, a recent statement from the US Food & Drug Administration (FDA) “cautions against use of hydroxychloroquine or chloroquine for COVID-19

Table 1. Examples of Commonly Used Platforms for Telehealth²⁵⁻²⁷

	Platform	Website
HIPAA- or PHI-secure telehealth platform	accuRx*	https://www.accurx.com
	AMD Global Telemedicine†	https://www.amdtelemedicine.com/telemedicine-products/Working-Clinic-Telemedicine.html
	American Well‡	https://business.amwell.com/solution-overview
	Attend Anywhere*	https://www.attendanywhere.com
	BlueJeans for Healthcare‡	https://www.bluejeans.com/use-cases/healthcare
	Care Connect by KRY/LIVIS§	https://www.kry.care
	Caregility‡	https://caregility.com/uhe-applications
	Cisco Webex†	https://www.webex.com/webexremotehealth.html
	ClickDoc§	https://www.cgm.com/be/index.fr.jsp
	Conexa Saúde	https://www.conexasaude.com.br
	Docobo*	https://www.docobo.co.uk
	Doctena§	https://secure.doctena.com
	Doctorlink*	https://www.doctorlink.com/video-consultation
	Doximity‡	https://www.doximity.com
	Doxy.me‡	https://doxy.me
	EMIS Health Video Consult*	https://www.emishealth.com/products/video-consultation
	Exprivia§	https://www.exprivia.it
	FaceTalk§	https://facetalk.nl
	Google G Suite†	https://gsuite.google.com/industries/healthcare
	Helena§	https://qa.helena.care
	Intersysto§	http://www.intersysto.eu
	InTouch Health‡	https://intouchhealth.com/telehealth-solutions
	pMD‡	https://www.pmd.com/secure-messaging-telemedicine
	Microsoft Teams/Skype for Business E3 and E5†	https://www.microsoft.com/en-us/microsoft-365/microsoft-teams/healthcare-solutions
	Signal§¶	https://signal.org
	Silicon Practice FootFall*	https://www.siliconpractice.co.uk/video-consultations
	Spruce‡	https://www.sprucehealth.com
	TeleMedi§	https://telemedi.be
	Updox‡	https://www.updox.com
	US Department of Veterans Affairs Video Connect‡	https://www.mobile.va.gov/app/va-video-connect
VSee‡	https://vsee.com	
Zoom for Healthcare†	https://zoom.us/healthcare	
Proprietary EMR telehealth platforms		
Video-conferencing platforms (not HIPAA- or PHI-secure)	GoToMeeting	
	Skype	
	Zoom	
Nonpublic-facing mobile communication platforms (not HIPAA- or PHI-secure)	Apple FaceTime	
	Google Duo	
	Google Hangouts	
	WhatsApp	

EMR indicates electronic medical record; HIPAA, Health Insurance Portability and Accountability Act; and PHI, protected health information.

*United Kingdom.

†Global.

‡United States.

§Europe.

||South America—Brazil.

¶Encrypted, but not specifically a telehealth platform.

Table 2. Examples of Remote ECG and Heart Rate Monitoring Devices

	Device	Type	CE Mark	FDA Clearance	Additional Features/ Notes	Website
Handheld devices	AliveCor KardiaMobile	Wireless	Yes	Yes	FDA cleared for AF (1-lead) and for QTc (6 L) for COVID-19 patients on HCQ±AZM	https://www.alivecor.com/kardiamobile
	Beurer ME 90	Wireless 1-lead ECG	Yes	No		https://www.beurer.com/web/gb/products/medical/ecg-and-pulse-oximeter/mobile-ecg-device/me-90-bluetooth.php
	Cardiac Designs ECG Check	Wireless 1-lead ECG	Yes	Yes		https://www.cardiacdesigns.com
	CardioComm Solutions HeartCheck CardiBeat and ECG Pen	Wireless 1-lead ECG	Yes	Yes		https://www.theheartcheck.com
	COALA	Wireless 1-lead ECG	Yes	Yes	Remote lung auscultation	https://www.coalalife.com
	Eko DUO	Wireless 1-lead ECG	Yes	Yes	Remote cardiac auscultation/phonocardiogram	https://www.ekohealth.com
	Omron blood pressure+EKG monitor	Wireless 1-lead ECG+BP cuff	No	Yes	United States and Canada only	https://omronhealthcare.com
	EKGraph	Wireless 1-lead ECG	No	Yes	United States	https://sonohealth.org
Mobile cardiac telemetry devices	Qardio QardioCore	Chest strap 1-lead ECG	Yes	No	ECG, HR, HRV, RR, activity	https://www.getqardio.com/qardiocore-wearable-ecg-ekg-monitor-iphone
	BardyDx CAM	Patch 1-lead ECG	Yes	Yes	Under clinical investigation for QTc monitoring in COVID-19 patients	https://www.bardydx.com
	BioTel Heart	Patch 1-lead ECG	Yes—only for extended Holter	Yes	FDA cleared for QTc monitoring	https://www.myheartmonitor.com/device/mcot-patch
	BodyGuardian MINI Family/BodyGuardian MINI PLUS	Wireless, Patch: 1-lead ECG/Wired, 3-lead ECG	Yes	Yes	ECG, HR, HRV, RR	https://www.preventicesolutions.com/hcp/body-guardian-mini-family
	iRhythm Zio patch/ Zio AT	Patch 1-lead ECG	Yes	Yes		https://www.irhythmttech.com
	InfoBionic MoMe Kardia	Wired 3-lead ECG	Yes	Yes	Remote lung auscultation	https://infobionic.com
	MediBioSense MBS HealthStream, VitalPatch, MCT	Patch 1-lead ECG	Yes	Yes	Monitors up to 8 vital signs	https://www.medibiosense.com
	MEMO Patch	Patch/watch 1-lead ECG	No	No	Asia; Korea FDA approved	https://www.huinno.com
	MedilynX PocketECG	Wired 3-lead ECG	Yes	Yes	HRV	https://www.pocketecg.com
	RhythmMedix RhythmStar	Wired 3-lead ECG	No	Yes		https://www.rhythmmedix.com
	Samsung S-patch Cardio	Patch 1-lead ECG	Yes	No	Asia; Korea FDA approved	https://www.samsungsds.com/global/en/solutions/off/cardio/cardio.html
Smartwatches	Apple Watch	1-lead ECG	Yes	Yes	FDA cleared for AF notification	https://www.apple.com/watch
	Withings Move ECG	1-lead ECG	Yes	No	Requires Health Mate app for ECG analysis/ AF detection	https://www.withings.com/us/en/move-ecg

AF indicates atrial fibrillation; AZM, azithromycin; BP, blood pressure; FDA, Food & Drug Administration; HCQ, hydroxychloroquine; HR, heart rate; HRV, heart rate variability; and RR, respiratory rate.

Table 3. Effect on QTc and Proarrhythmia of Experimental Pharmacological Therapies for COVID-19³⁶

	QTc Prolongation	TdP Risk
Chloroquine ³⁷⁻⁴⁰	Moderate ↑	Low risk of TdP
Hydroxychloroquine ⁴¹	Moderate ↑	Low risk of TdP
Azithromycin ⁴²⁻⁴⁸	Moderate ↑	Very low risk of TdP
Lopinavir/ritonavir ³⁷	Moderate ↑	Low risk of TdP
Tocilizumab ⁴⁹	Mild ↑ or ↓	NR
Fingolimod	Mild ↑	NR
Remdesivir	NR	NR
Interferon alfacon-1	NR	NR
Ribavirin	NR	NR
Methylprednisolone	NR	NR

NR indicates not reported; and TdP, torsade de pointes.

outside of the hospital setting or a clinical trial due to risk of heart rhythm problems.” (This does not affect FDA-approved uses for malaria, lupus, and rheumatoid arthritis.)⁵¹ Exceptions to this practice are acknowledged to occur in some regions, as these drugs have been used outside the United States without regulatory warnings. Under these conditions, or when these drugs are maintained after hospital discharge, consumer mobile ECG devices capable of generating QTc measurements may be used. If the QTc increases significantly, physicians can consider a change or discontinuation of medication via the phone or virtual medical services.

Electrocardiographic Monitoring During Clinical Trials

Several double- and multi-arm blind randomized controlled trials are underway worldwide for COVID-19 outpatients utilizing different medications that may prolong the QT interval.⁵²⁻⁵⁶ These drugs are being tested either alone or in various combinations and are being compared with one another, with differential dosing regimens and/or placebo. These drugs are also being tested for post-exposure prophylaxis in high-risk groups.

Mobile health using smartphone-based portable ECG devices as QTc monitoring tools is an innovative and economical solution to conduct monitoring in outpatient trials. For instance, in one trial evaluating hydroxychloroquine and azithromycin (hydroxychloroquine alone and hydroxychloroquine/azithromycin combination) against a placebo, participants receive remote training to acquire a 6-lead ECG at baseline and then at specified follow-up intervals through the trial period (Figure). These ECGs are transmitted to a remote QTc monitoring site, where the QTc is assessed and monitored over the treatment period.

THE FUTURE: DIGITAL MEDICINE CATALYZED BY THE PANDEMIC

The COVID-19 public health emergency has forced changes to traditional norms of health care access and delivery across all continents.¹⁰ It has accelerated adoption of telemedicine and all aspects of digital health, regarded as a positive development. Today’s new reality will likely define medicine going forward. Many monitoring and diagnostic testing aspects of both inpatient and outpatient care will be increasingly served by digital medicine tools.

The need for contactless monitoring for inpatients triaged to intensive care unit, telemetry, or nonconventional environments, as well as for outpatients needing continued management, has triggered novel implementation of digital health monitoring tools. Some centers have created algorithms based on predictive analytics of electronic medical record data. Centralized monitoring or mobile continuous monitoring has improved patient outcomes, reduced manpower needs, and is being utilized more commonly.⁵⁷ The use of wearables such as watches, smartphones, and smart beds (with elimination of cables and skin electrodes) for in-hospital telemetry is a novel approach. This type of wireless monitoring may be continued after discharge, permitting prolonged surveillance of rhythm and other physiological parameters.¹² Bracelet technologies may transmit multiple parameters (eg, heart rate, sleep, oxygen desaturation index, blood pressure) via a smartphone link to centralized hubs. These technologies provide a solution for intensive monitoring extending beyond the hospital environment.

Outpatient management has been revolutionized since the start of the pandemic. Social distancing measures and restricted clinic access have driven the rapid adoption of telehealth mechanisms to continue management of non-COVID-19 patients. Virtual visits that have been used for decades to reach isolated communities,⁵⁸ but less commonly utilized in advanced health systems, have now become the mainstay of ambulatory care across all subspecialties. The initial experience appears to have been positive for both patient and caregiver. Heart rhythm professionals are fortunate to have a choice of wireless technologies to relay monitored information to maintain connection.¹² Wearable and smartphone-based devices allow convenient real-time monitoring for arrhythmias on a long-term basis because of the comfort associated with their small size and ease of use while reducing patient and healthcare worker exposure. Remote CIED monitoring has existed for decades.²⁴ It is strongly endorsed by professional societies, but in practice only a fraction of its diagnostic and therapeutic capabilities has been utilized—until now.⁵⁹ Since the start of the pandemic, utilization of wireless communication with CIEDs has grown exponentially, permanently altering the future of device follow-up. Patient outcomes may be improved




Figure. ECG acquisition and transmission using a smartphone-based portable ECG monitor.

with intensive device-based monitoring compared with traditional in-clinic evaluations at regular intervals.⁶⁰ Recent data indicate that in-person CIED evaluation can be extended safely to at least biennially when daily digital connectivity is maintained.⁶¹ Remote monitoring has the potential advantage of detecting and alerting caregivers (and in the future—patients directly) about important parameter changes, enabling earlier patient hospitalization, even during a presymptomatic phase.⁶²

Connectivity permits longitudinal follow-up, with advantages ranging from individual disease management to assessment of penetration of recommended therapies into communities.^{60,63} The ability for CIED remote monitoring data to be streamed to or accessed by multiple providers can facilitate communication and cooperative treatment and should be encouraged. This will require approval by patients, regulators, and manufacturers. Lessons learned from implantable devices can be applied widely in telemedicine. Regulatory bodies have been responsive, for example, approving smartphone-based QT interval measurement and telehealth services across state lines in the United States. The pandemic experience should serve as an impetus to expedite the resolution of persistent challenges, such as validation of digital technologies, infrastructure for data management (and mechanism for relay to patient and caregiver), interoperability with electronic medical record, application of predictive analytics, cybersecurity (and with it the capability for limited forms of remote CIED programming), and reimbursement.^{64–66}

In summary, the crisis precipitated by the pandemic has catalyzed the adoption of remote patient management across many specialties and for heart rhythm professionals, in particular. This practice is here to stay—it will persist even if other less arrhythmogenic treatment

strategies evolve for COVID-19 and after the pandemic has passed. This is an opportunity to embed and grow remote services in everyday medical practice worldwide.

ARTICLE INFORMATION

Affiliations

Cleveland Clinic, OH (N.V., M.K.C.). Tulane University School of Medicine, New Orleans, LA (N.F.M., L.D.). Centro Privado de Cardiología, Tucuman, Argentina (L.A.). Cedars-Sinai Medical Center, Los Angeles, CA (C.M.A.). Arrhythmia Section, Cardiology Department, Hospital Clínic, Universitat de Barcelona, Spain. Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain (E.A.). Centro de Investigación Biomédica en Red de Enfermedades Cardiovasculares (CIBERCV), Madrid, Spain (E.A.). Korea University Medical Center, Seoul, Republic of Korea (J.-I.C.). Cardiocentro, Lugano, Switzerland (G.C.). Northwell Health, North Shore University Hospital, Manhasset, New York (L.M.E.). University of Michigan, Ann Arbor (H.G.). VA Greater Los Angeles Healthcare System and David Geffen School of Medicine at the University of California (J.K.H.). Antwerp University and University Hospital, Belgium (H. Heidebuchel). Renmin Hospital of Wuhan University, China (H. Huang). Kansas City Heart Rhythm Institute and Research Foundation, Overland Park (D.R.K.). Faculty of Medicine Ramathibodi Hospital, Mahidol University, Bangkok, Thailand (T.N.). Cooper Medical School of Rowan University, Camden, NJ (A.M.R.). Hospital Pró-Cardíaco, Rio de Janeiro, Brazil (E.B.S.). CardiolInfantil Foundation, Cardiac Institute, Bogota, Colombia (L.C.S.M.). Bethel University, St. Paul, MN (K.E.S.). University of Washington, Seattle (A.R.M.S.). Oregon Health & Science University, Portland, OR (E.C.S.). VA Eastern Colorado Health Care System and University of Colorado, Aurora (P.D.V.).

Disclosures

None.

REFERENCES

- Goyal P, Choi JJ, Pinheiro LC, Schenck EJ, Chen R, Jabri A, Sattlin MJ, Campion TR Jr, Nahid M, Ringel JB, et al. Clinical characteristics of Covid-19 in New York City. *N Engl J Med*. 2020;382(24):2372–2374. doi: 10.1056/NEJMc2010419
- Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, Wang B, Xiang H, Cheng Z, Xiong Y, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA*. 2020;323:1061–1069. doi: 10.1001/jama.2020.1585
- Guo T, Fan Y, Chen M, Wu X, Zhang L, He T, Wang H, Wan J, Wang X, Lu Z. Cardiovascular implications of fatal outcomes of patients with

- coronavirus disease 2019 (COVID-19). *JAMA Cardiol.* 2020;5:811–818. doi: 10.1001/jamacardio.2020.1017
4. Azarkish M, Laleh Far V, Eslami M, Mollazadeh R. Transient complete heart block in a patient with critical COVID-19. *Eur Heart J.* 2020;41:2131. doi: 10.1093/eurheartj/ehaa307
 5. Noll A, William A, Varma N. A young woman presenting with a viral prodrome, palpitations, dizziness, and heart block. *JAMA Cardiol.* 2020 [In press].
 6. Kochi AN, Tagliari AP, Forleo GB, Fassini GM, Tondo C. Cardiac and arrhythmic complications in patients with COVID-19. *J Cardiovasc Electrophysiol.* 2020;31:1003–1008. doi: 10.1111/jce.14479
 7. Yao XH, Li TY, He ZC, Ping YF, Liu HW, Yu SC, Mou HM, Wang LH, Zhang HR, Fu WJ, et al. [A pathological report of three COVID-19 cases by minimal invasive autopsies]. *Zhonghua Bing Li Xue Za Zhi.* 2020;49:411–417. doi: 10.3760/cma.j.cn112151-20200312-00193
 8. Inciardi RM, Lupi L, Zacccone G, Italia L, Raffo M, Tomasoni D, Cani DS, Cerini M, Farina D, Gavazzi E, et al. Cardiac involvement in a patient with coronavirus disease 2019 (COVID-19). *JAMA Cardiol.* 2020;5:819–824. doi: 10.1001/jamacardio.2020.1096
 9. Sapp JL, Alqarawi W, MacIntyre CJ, Tadros R, Steinberg C, Roberts JD, Laksman Z, Healey JS, Krahn AD. Guidance on minimizing risk of drug-induced ventricular arrhythmia during treatment of COVID-19: a statement from the Canadian Heart Rhythm Society. *Can J Cardiol.* 2020;36(6):948–951. doi: 10.1016/j.cjca.2020.04.003
 10. Udawadia ZF, Raju RS. How to protect the protectors: 10 lessons to learn for doctors fighting the COVID-19 coronavirus. *Med J Armed Forces India.* 2020;76:128–131. doi: 10.1016/j.mjafi.2020.03.009
 11. Gabriels J, Saleh M, Chang D, Epstein LM. Inpatient use of mobile continuous telemetry for COVID-19 patients treated with hydroxychloroquine and azithromycin. *Heart Rhythm Case Rep.* 2020;6:241–243. doi: 10.1016/j.hrcr.2020.03.017
 12. Steinberg JS, Varma N, Cygankiewicz I, Aziz P, Balsam P, Baranchuk A, Cantillon DJ, Dilaveris P, Dubner SJ, El-Sherif N, et al. 2017 ISHNE-HRS expert consensus statement on ambulatory ECG and external cardiac monitoring/telemetry. *Heart Rhythm.* 2017;14:e55–e96. doi: 10.1016/j.hrthm.2017.03.038
 13. Garabelli P, Stavrakis S, Albert M, Koomson E, Parwani P, Chohan J, Smith L, Albert D, Xie R, Xie Q, et al. Comparison of QT Interval Readings in Normal Sinus Rhythm Between a Smartphone Heart Monitor and a 12-Lead ECG for Healthy Volunteers and Inpatients Receiving Sotalol or Dofetilide. *J Cardiovasc Electrophysiol.* 2016;27:827–832. doi: 10.1111/jce.12976
 14. Castelletti S, Dagradi F, Goulene K, Danza AI, Baldi E, Stramba-Badiale M, Schwartz PJ. A wearable remote monitoring system for the identification of subjects with a prolonged QT interval or at risk for drug-induced long QT syndrome. *Int J Cardiol.* 2018;266:89–94. doi: 10.1016/j.ijcard.2018.03.097
 15. Gropler MRF, Dalal AS, Van Hare GF, Silva JNA. Can smartphone wireless ECGs be used to accurately assess ECG intervals in pediatrics? A comparison of mobile health monitoring to standard 12-lead ECG. *PLoS One.* 2018;13:e0204403. doi: 10.1371/journal.pone.0204403
 16. US Food & Drug Administration. VitalConnect Inc. VitalPatch: Letter of Authorization 2020. <https://www.fda.gov/media/137397/download>. Accessed May 15, 2020.
 17. VitalConnect. COVID-19. <https://vitalconnect.com/covid-19-remote-patient-monitoring>. Accessed May 15, 2020.
 18. Rimmer LK, Rimmer JD. Comparison of 2 methods of measuring the QT interval. *Am J Crit Care.* 1998;7:346–354.
 19. Rimmer LK. Bedside monitoring of the QT interval. *Am J Crit Care.* 1998;7:183–9; quiz 190.
 20. Koltowski L, Balsam P, Gilowczynska R, Rokicki JK, Peller M, Maksym J, Blicharz L, Maciejewski K, Niedziela M, Opolski G, et al. Kardia Mobile applicability in clinical practice: a comparison of Kardia Mobile and standard 12-lead electrocardiogram records in 100 consecutive patients of a tertiary cardiovascular care center [published online January 15, 2019]. *Cardiol J.* doi: 10.5603/CJ.a2019.0001. https://journals.viamedica.pl/cardiology_journal/article/view/58839.
 21. Ohannessian R. Telemedicine: potential applications in epidemic situations. *European Research in Telemedicine.* 2015;4:95–98.
 22. Hollander JE, Carr BG. Virtually Perfect? Telemedicine for Covid-19. *N Engl J Med.* 2020;382:1679–1681. doi: 10.1056/NEJMp2003539
 23. Liu S, Yang L, Zhang C, Xiang YT, Liu Z, Hu S, Zhang B. Online mental health services in China during the COVID-19 outbreak. *Lancet Psychiatry.* 2020;7:e17–e18. doi: 10.1016/S2215-0366(20)30077-8
 24. Varma N, Ricci RP. Telemedicine and cardiac implants: what is the benefit? *Eur Heart J.* 2013;34:1885–1895. doi: 10.1093/eurheartj/ehs388
 25. NHSX. Information governance. <https://www.nhs.uk/covid-19-response/data-and-information-governance/information-governance>. Accessed May 15, 2020.
 26. NHS Digital. Approved video consultation systems. <https://digital.nhs.uk/services/future-gp-it-systems-and-services/approved-econsultation-systems>. Accessed May 15, 2020.
 27. National Health Service. Procurement of pre-approved suppliers of online and video consultation systems for GP practices to support COVID-19. <https://www.england.nhs.uk/coronavirus/wp-content/uploads/sites/52/2020/03/C0079-Suppliers-of-online-and-video-consultations.pdf>. Accessed April 29, 2020.
 28. Grein J, Ohmagari N, Shin D, Diaz G, Asperges E, Castagna A, Feldt T, Green G, Green ML, Lescure F-X, et al. Compassionate use of remdesivir for patients with severe Covid-19. *N Engl J Med.* 2020;382(24):2327–2336. doi: 10.1056/NEJMoa2007016
 29. European Society of Cardiology. ESC guidance for the diagnosis and management of CV disease during the COVID-19 pandemic. <https://www.escardio.org/Education/COVID-19-and-Cardiology/ESC-COVID-19-Guidance>. Accessed April 29, 2020.
 30. Chen Z, Hu J, Zhang Z, Jiang S, Han S, Yan D, Zhuang R, Hu B, Zhang Z. Efficacy of hydroxychloroquine in patients with COVID-19: results of a randomized clinical trial. *medRxiv.* 2020.03.22.20040758. doi: 10.1101/2020.03.22.20040758
 31. Haeusler IL, Chan XHS, Guérin RJ, White NJ. The arrhythmogenic cardiotoxicity of the quinoline and structurally related antimalarial drugs: a systematic review. *BMC Med.* 2018;16:200. doi: 10.1186/s12916-018-1188-2
 32. Geleris J, Sun Y, Platt J, Zucker J, Baldwin M, Hripcsak G, Labelle A, Manson D, Kuber C, Graham Barr R, et al. Observational study of hydroxychloroquine in hospitalized patients with Covid-19. *N Engl J Med.* 2020;382:2411–2418. doi: 10.1056/NEJMoa2012410
 33. Chorin E, Wadhvani L, Magnani S, Dai M, Shulman E, Nadeau-Routhier C, Knotts R, Bar-Cohen R, Kogan E, Barbhuiya C, et al. QT interval prolongation and torsade de pointes in patients with COVID-19 treated with hydroxychloroquine/azithromycin. *Heart Rhythm.* 2020;17:1425–1433. doi: 10.1016/j.hrthm.2020.05.014
 34. Saleh M, Gabriels J, Chang D, Kim BS, Mansoor A, Mahmood E, Makker P, Ismail H, Galdner B, Willner J, et al. The effect of chloroquine, hydroxychloroquine and azithromycin on the corrected QT interval in patients with SARS-CoV-2 infection. *Circ Arrhythm Electrophysiol.* 2020;13:e008662. doi: 10.1161/CIRCEP.120.008662
 35. Rosenberg ES, Dufort EM, Udo T, Wilberschied LA, Kumar J, Tesoriero J, Weinberg P, Kirkwood J, Muse A, DeHovitz J, et al. Association of treatment with hydroxychloroquine or azithromycin with in-hospital mortality in patients with COVID-19 in New York state. *JAMA.* 2020;323:2493–2502. doi: 10.1001/jama.2020.8630
 36. CredibleMeds. COVID-19 experimental therapies and TdP risk. <https://www.crediblemeds.org/blog/covid-19-experimental-therapies-and-tdp-risk>. Accessed April 20, 2020.
 37. Vicente J, Zusterzeel R, Johannesen L, Ochoa-Jimenez R, Mason JW, Sanabria C, Kemp S, Sager PT, Patel V, Matta MK, et al. Assessment of Multi-Ion Channel Block in a Phase I Randomized Study Design: Results of the CiPA Phase I ECG Biomarker Validation Study. *Clin Pharmacol Ther.* 2019;105:943–953. doi: 10.1002/cpt.1303
 38. Mzayek F, Deng H, Mather FJ, Wasilevich EC, Liu H, Hadi CM, Chansolme DH, Murphy HA, Melek BH, Tenaglia AN, et al. Randomized dose-ranging controlled trial of AQ-13, a candidate antimalarial, and chloroquine in healthy volunteers. *PLoS Clin Trials.* 2007;2:e6. doi: 10.1371/journal.pctr.0020006
 39. Wozniacka A, Cygankiewicz I, Chudzik M, Sysa-Jedrzejowska A, Wrancik JK. The cardiac safety of chloroquine phosphate treatment in patients with systemic lupus erythematosus: the influence on arrhythmia, heart rate variability and repolarization parameters. *Lupus.* 2006;15:521–525. doi: 10.1191/0961203306lu2345oa
 40. Teixeira RA, Martinelli Filho M, Benvenuti LA, Costa R, Pedrosa AA, Nishioka SA. Cardiac damage from chronic use of chloroquine: a case report and review of the literature. *Arq Bras Cardiol.* 2002;79:85–88. doi: 10.1590/s0066-782x2002001000009
 41. McGhie TK, Harvey P, Su J, Anderson N, Tomlinson G, Touma Z. Electrocardiogram abnormalities related to anti-malarials in systemic lupus erythematosus. *Clin Exp Rheumatol.* 2018;36:545–551.
 42. Zhang M, Xie M, Li S, Gao Y, Xue S, Huang H, Chen K, Liu F, Chen L. Electrophysiologic Studies on the Risks and Potential Mechanism Underlying the Proarrhythmic Nature of Azithromycin. *Cardiovasc Toxicol.* 2017;17:434–440. doi: 10.1007/s12012-017-9401-7

43. Choi Y, Lim HS, Chung D, Choi JG, Yoon D. Risk Evaluation of Azithromycin-Induced QT Prolongation in Real-World Practice. *Biomed Res Int*. 2018;2018:1574806. doi: 10.1155/2018/1574806
44. US Food and Drug Administration. ZITHROMAX (azithromycin) for IV infusion only. Highlights of prescribing information. Reference ID: 4051690. https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/050733s043lbl.pdf. Accessed May 20, 2020.
45. Ray WA, Murray KT, Hall K, Arbogast PG, Stein CM. Azithromycin and the risk of cardiovascular death. *N Engl J Med*. 2012;366:1881–1890. doi: 10.1056/NEJMoa1003833
46. Poluzzi E, Raschi E, Motola D, Moretti U, De Ponti F. Antimicrobials and the risk of torsades de pointes: the contribution from data mining of the US FDA Adverse Event Reporting System. *Drug Saf*. 2010;33:303–314. doi: 10.2165/11531850-000000000-00000
47. Cheng YJ, Nie XY, Chen XM, Lin XX, Tang K, Zeng WT, Mei WY, Liu LJ, Long M, Yao FJ, et al. The Role of Macrolide Antibiotics in Increasing Cardiovascular Risk. *J Am Coll Cardiol*. 2015;66:2173–2184. doi: 10.1016/j.jacc.2015.09.029
48. Maisch NM, Kochupurackal JG, Sin J. Azithromycin and the risk of cardiovascular complications. *J Pharm Pract*. 2014;27:496–500. doi: 10.1177/0897190013516503
49. Grange S, Schmitt C, Banken L, Kuhn B, Zhang X. Thorough QTc study of tocilizumab after single-dose administration at therapeutic and supratherapeutic doses in healthy subjects. *Int J Clin Pharmacol Ther*. 2011;49:648–655. doi: 10.5414/cp201549
50. Giudicessi JR, Noseworthy PA, Friedman PA, Ackerman MJ. Urgent Guidance for Navigating and Circumventing the QTc-Prolonging and Torsadogenic Potential of Possible Pharmacotherapies for Coronavirus Disease 19 (COVID-19). *Mayo Clin Proc*. 2020;95:1213–1221. doi: 10.1016/j.mayocp.2020.03.024
51. US Food & Drug Administration. FDA cautions against use of hydroxychloroquine or chloroquine for COVID-19 outside of the hospital setting or a clinical trial due to risk of heart rhythm problems. <https://www.fda.gov/drugs/drug-safety-and-availability/fda-cautions-against-use-hydroxychloroquine-or-chloroquine-covid-19-outside-hospital-setting-or>. Accessed April 29, 2020.
52. ClinicalTrials.gov. COVID-19 clinical trials. <https://clinicaltrials.gov/ct2/results?cond=COVID+19&term=&cntry=&state=&city=&dist=>. Accessed April 18, 2020.
53. ClinicalTrials.gov. Efficacy of novel agents for treatment of SARS-CoV-2 infection among high-risk outpatient adults: an adaptive randomized platform trial. https://clinicaltrials.gov/ct2/show/NCT04354428?type=Intr&cond=COVID+19&map_cntry=US&map_state=US%3AWA&draw=2. Accessed April 26, 2020.
54. ClinicalTrials.gov. WU 352: open-label, randomized controlled trial of hydroxychloroquine alone or hydroxychloroquine plus azithromycin or chloroquine alone or chloroquine plus azithromycin in the treatment of SARS CoV-2 infection. https://clinicaltrials.gov/ct2/show/NCT04341727?type=Intr&cond=COVID+19&map_cntry=US&map_state=US%3AMO&draw=2. Accessed April 26, 2020.
55. ClinicalTrials.gov. Efficacy of hydroxychloroquine for post-exposure prophylaxis (PEP) to prevent severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection among adults exposed to coronavirus disease (COVID-19): a blinded, randomized study. https://clinicaltrials.gov/ct2/show/NCT04328961?type=Intr&cond=COVID+19&map_cntry=US&map_state=US%3AWA&draw=2. Accessed April 26, 2020.
56. ClinicalTrials.gov. ChemoPROPhyLaxIs For covid-19 Infectious Disease (the PROLIFIC Trial). <https://clinicaltrials.gov/ct2/show/NCT04352933?type=Intr&cond=COVID+19&draw=2>. Accessed April 26, 2020.
57. Cantillon DJ, Loy M, Burkle A, Pengel S, Brosovich D, Hamilton A, Khot UN, Lindsay BD. Association Between Off-site Central Monitoring Using Standardized Cardiac Telemetry and Clinical Outcomes Among Non-Critically Ill Patients. *JAMA*. 2016;316:519–524. doi: 10.1001/jama.2016.10258
58. Bagchi S. Telemedicine in rural India. *PLoS Med*. 2006;3:e82. doi: 10.1371/journal.pmed.0030082
59. Slotwiner D, Varma N, Akar JG, Annas G, Beardsall M, Fogel RI, Galizio NO, Glotzer TV, Leahy RA, Love CJ, et al. HRS Expert Consensus Statement on remote interrogation and monitoring for cardiovascular implantable electronic devices. *Heart Rhythm*. 2015;12:e69–100. doi: 10.1016/j.hrthm.2015.05.008
60. Hindricks G, Varma N, Kacet S, Lewalter T, Søgaard P, Guédon-Moreau L, Proff J, Gerds TA, Anker SD, Torp-Pedersen C. Daily remote monitoring of implantable cardioverter-defibrillators: insights from the pooled patient-level data from three randomized controlled trials (IN-TIME, ECOST, TRUST). *Eur Heart J*. 2017;38:1749–1755. doi: 10.1093/eurheartj/ehx015
61. Watanabe E, Yamazaki F, Goto T, Asai T, Yamamoto T, Hirooka K, Sato T, Kasai A, Ueda M, Yamakawa T, et al. Remote Management of Pacemaker Patients With Biennial In-Clinic Evaluation: Continuous Home Monitoring in the Japanese At-Home Study: A Randomized Clinical Trial. *Circ Arrhythm Electrophysiol*. 2020;13:e007734. doi: 10.1161/CIRCEP.119.007734
62. Varma N, Epstein AE, Irimpen A, Schweikert R, Love C, TRUST Investigators. Efficacy and safety of automatic remote monitoring for implantable cardioverter-defibrillator follow-up: The Lumos-T Safely Reduces Routine Office Device Follow-up (TRUST) trial. *Circulation* 2010;122:325–332.
63. Varma N, Jones P, Wold N, Stein K. How well do results from large randomized clinical trials diffuse into clinical practice? Impact of MADIT-RIT in a large cohort of implantable cardioverter defibrillator patients (ALTIUDE). *Eur Heart J*. 2014;35(Suppl 1) [Abstract].
64. Saxon LA, Varma N, Epstein LM, Ganz LI, Epstein AE. Factors Influencing the Decision to Proceed to Firmware Upgrades to Implanted Pacemakers for Cybersecurity Risk Mitigation. *Circulation*. 2018;138:1274–1276. doi: 10.1161/CIRCULATIONAHA.118.034781
65. Slotwiner DJ, Abraham RL, Al-Khatib SM, Anderson HV, Bunch TJ, Ferrara MG, Lippman N, Serwer GA, Steiner PR, Tchong JE, et al. HRS White Paper on interoperability of data from cardiac implantable electronic devices (CIEDs). *Heart Rhythm*. 2019;16:e107–e127. doi: 10.1016/j.hrthm.2019.05.002
66. Heidebuchel H, Hindricks G, Broadhurst P, Van Erven L, Fernandez-Lozano I, Rivero-Ayerza M, Malinowski K, Marek A, Romero Garrido RF, Löscher S, et al. EuroEco (European Health Economic Trial on Home Monitoring in ICD Patients): a provider perspective in five European countries on costs and net financial impact of follow-up with or without remote monitoring. *Eur Heart J*. 2015;36:158–169. doi: 10.1093/eurheartj/ehu339

Appendix 1. Author Disclosure Table

Writing Group Member	Employment	Honoraria/ Speaking/ Consulting	Speakers' Bureau	Research*	Fellowship Support*	Ownership/ Partnership/ Principal/ Majority Stockholder	Stock or Stock Options	Intellectual Property/ Royalties	Other
Niraj Varma, MA, MD, PhD, FACC, FRCP (Chair)	Cleveland Clinic, Cleveland, OH	1: BIOTRONIK; 1: EP Solutions; 1: Medtronic; 1: MicroPort	None	1: Boston Scientific; 2: Abbott	None	None	None	None	None
Nassir F. Marrouche, MD, FHRS (Vice-Chair)	Tulane University School of Medicine, New Orleans, LA	0: Biosense Webster; 0: BIOTRONIK; 0: Cardiac Design; 0: Medtronic; 1: Preventice	None	0: Abbott; 0: Boston Scientific; 0: GE Healthcare; 5: Biosense Webster	None	None	None	None	None
Luis Aguinaga, MD, MBA, PhD, FESC, FACC	Centro Privado de Cardiologia, Tucuman, Argentina	None	None	None	None	None	None	None	None
Christine M. Albert, MD, MPH, FHRS, FACC	Cedars-Sinai Medical Center, Los Angeles, CA	1: Roche Diagnostics	None	5: Abbott; 5: NIH; 5: Roche Diagnostics	None	None	None	None	None
Elena Arbelo, MD, MSci, PhD	Arrhythmia Section, Cardiology Department, Hospital Clinic, Universitat de Barcelona, Barcelona, Spain; Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain; Centro de Investigación Biomédica en Red de Enfermedades Cardiovasculares (CIBERCV), Madrid, Spain	1: Biosense Webster	None	None	None	None	None	None	None
Jong-Il Choi, MD, PhD, MHS	Korea University Medical Center, Seoul, Republic of Korea	None	None	None	None	None	None	None	None
Mina K. Chung, MD, FHRS, FACC	Cleveland Clinic, Cleveland, OH	2: ABIM	None	5: AHA; 5: NIH	None	None	None	1: Elsevier; 1: UpToDate	0: ACC (EP Section Leadership Council member); 0: AHA (Chair, ECG & Arrhythmias Committee; Member, Clinical Cardiology Leadership Committee; Member, Committee on Scientific Sessions Programming); 0: Amarin (Data monitoring committee member); 0: BIOTRONIK (EPIC Alliance steering committee member); 2: AHA (Associate Editor, <i>Circulation: Arrhythmia and Electrophysiology</i>)

(Continued)

Appendix 1. Author Disclosure Table Continued

Writing Group Member	Employment	Honoraria/ Speaking/ Consulting	Speakers' Bureau	Research*	Fellowship Support*	Ownership/ Partnership/ Principal/ Majority Stockholder	Stock or Stock Options	Intellectual Property/ Royalties	Other
Giulio Conte, MD, PhD	Cardiocentro, Lugano, Switzerland	None	None	5: Boston Scientific; 5: Swiss National Science Foundation	None	None	None	None	None
Lilas Dagher, MD	Tulane University School of Medicine, New Orleans, LA	None	None	None	None	None	None	None	None
Laurence M. Epstein, MD, FACC	Northwell Health, North Shore University Hospital, Manhasset, NY	1: Abbott; 2: Medtronic; 2: Spectranetics Corporation	None	None	None	None	None	None	2: Boston Scientific (Clinical Events Committee)
Hamid Ghanbari, MD, FACC	University of Michigan, Ann Arbor, MI	2: Preventice	None	1: BIOTRONIK; 1: Boston Scientific; 1: Medtronic; 1: Toyota	None	None	None	None	1: Preventice (Travel/ Entertainment)
Janet K. Han, MD, FHRS, FACC	VA Greater Los Angeles Healthcare System and David Geffen School of Medicine at the University of California, Los Angeles, Los Angeles, CA	1: Abbott; 1: Medtronic	None	None	None	None	None	None	None
Hein Heidbuchel, MD, PhD, FESC, FEHRA	Antwerp University and University Hospital, Antwerp, Belgium	None	None	None	None	None	None	None	None
He Huang, MD, FACC, FESC, FEHRA	Renmin Hospital of Wuhan University, Wuhan, China	None	None	None	None	None	None	None	None
Dhanunjaya R. Lakkireddy, MD, FHRS, FACC	Kansas City Heart Rhythm Institute and Research Foundation, Overland Park, KA	1: BIOTRONIK; 2: Abbott	1: Abiomed; 1: Biosense Webster; 1: Boston Scientific; 2: Janssen	None	None	None	None	None	None
Tachapong Ngarmukos, MD, FAPHRs, FACC	Faculty of Medicine Ramathibodi Hospital, Mahidol University, Bangkok, Thailand	1: Abbott; 1: Bayer; 1: Biosense Webster; 1: BIOTRONIK; 1: Boehringer Ingelheim; 1: Boston Scientific; 1: Daiichi Sankyo; 1: Medtronic; 1: Pfizer	None	None	None	None	None	None	None
Andrea M. Russo, MD, FHRS, FACC	Cooper Medical School of Rowan University, Camden, NJ	None	None	1: MediLynx; 2: Boehringer Ingelheim; 2: Boston Scientific	None	None	None	1: UpToDate	0: ABIM (Member, ABIM Cardiovascular Board); 0: Apple, Inc (Steering Committee Apple Heart Study); 0: Boston Scientific (Steering Committee, Research)

(Continued)

Appendix1. Author Disclosure Table Continued

Writing Group Member	Employment	Honoraria/ Speaking/ Consulting	Speakers' Bureau	Research*	Fellowship Support*	Ownership/ Partnership/ Principal/ Majority Stockholder	Stock or Stock Options	Intellectual Property/ Royalties	Other
Eduardo B. Saad, MD, PhD, FHRS, FESC	Hospital Pró-Cardiaco, Rio de Janeiro, Brazil	1: Biosense Webster	None	None	None	None	None	None	1: Abbott (Travel/ Entertainment)
Luis C. Saenz Morales, MD	Cardiolfantil Foundation, Cardiac Institute, Bogota, Colombia	None	None	None	None	None	None	None	None
Kristin E. Sandau, PhD, RN	Bethel University, St. Paul, MN	1: Japanese Association of Cardiovascular Nursing	None	None	None	None	None	None	1: Patient Safety Authority
Arun Raghav M. Sridhar, MD, MPH, FACC	University of Washington, Seattle, WA	None	None	None	None	None	None	None	None
Eric C. Stecker, MD, MPH, FACC	Oregon Health & Science University, Portland, OR	None	None	4: AHA	None	None	None	None	None
Paul D. Varosy, MD, FHRS, FACC	VA Eastern Colorado Health Care System and University of Colorado, Aurora, CO	None	None	4: NIH	None	None	None	None	0: ACC (Committee Chair); 0: AHA (Committee Member); 0: NCDR (Committee Chair/Member)

Number value: 0=\$0; 1=≤\$10 000; 2=≤\$10 000 to ≤\$25 000; 3=≤\$25 000 to ≤\$50 000; 4=≤\$50 000 to ≤\$100 000; 5=≤\$100 000. ABIM indicates American Board of Internal Medicine; ACC, American College of Cardiology; AHA, American Heart Association; NCDR, National Cardiovascular Data Registry; and NIH, National Institutes of Health.

*Research and fellowship support are classed as programmatic support. Sources of programmatic support are disclosed but are not regarded as a relevant relationship with industry for writing group members or reviewers.