

Contents lists available at [ScienceDirect](https://www.sciencedirect.com)

Genomics

journal homepage: www.elsevier.com/locate/ygeno

Original Article

Analysis of organellar genomes in brown algae reveals an independent introduction of similar foreign sequences into the mitochondrial genome

Shivani Rana^a, Klaus Valentin^{b,*}, Jana Riehl^a, Aurélie Blanfuné^c, Lauric Reynes^c,
Thierry Thibaut^c, Inka Bartsch^b, Ludwig Eichinger^{a,*}, Gernot Glöckner^{a,1}

^a Institute of Biochemistry I, Faculty of Medicine, University of Cologne, Cologne, Germany

^b Alfred-Wegener-Institute, Helmholtz-Center for Marine and Polar Research, Bremerhaven, Germany

^c Aix-Marseille University and University of Toulon, CNRS, IRD, Mediterranean Institute of Oceanography (MIO), Marseille, France



ARTICLE INFO

Keywords:

Brown algae
Organellar genomes
Phylogeny
Mitochondrial lateral gene transfer

ABSTRACT

Kelp species (Laminariales, Phaeophyceae) are globally widespread along temperate to Polar rocky coastal lines. Here we analyse the mitochondrial and chloroplast genomes of *Laminaria rodriguezii*, in comparison to the organellar genomes of other kelp species. We also provide the complete mitochondrial genome sequence of another endemic kelp species from a Polar habitat, the Arctic *Laminaria solidungula*. We compare phylogenetic trees derived from twenty complete mitochondrial and seven complete chloroplast kelp genomes. Interestingly, we found a stretch of more than 700 bp in the mitochondrial genome of *L. rodriguezii*, which is not present in any other yet sequenced member of the Phaeophyceae. This stretch matches a protein coding region in the mitochondrial genome from *Desmarestia viridis*, another brown seaweed. Their high similarity suggests that these sequences originated through independent introduction into the two species. Their origin could have been by infection by yet unknown similar mitoviruses, currently only known from fungi and plants.

1. Introduction

Seaweeds are multicellular photoautotrophic macroalgal species. All brown algae (Phaeophyceae) are part of this non-taxonomic classification, since so far no unicellular members of this group were found [1]. Thus, multicellularity is one of the founding factors of brown algal evolution and factors enabling this were recruited early in the evolution of the branch. Similar functions are often established by using the same toolbox [2]. Therefore, we might learn from the analysis of one group of organisms, how and why other groups evolved. Intriguingly, similar factors seem to have been used to establish multicellularity in green plants and brown algae [3]. Thus, by comparative analyses of genomes from distantly related clades we might learn about common mechanisms of differentiation, habitat occupation, or pathogen susceptibility.

Kelps are a subgroup of brown algal seaweeds, which constitute the order Laminariales in the Phaeophyceae. They form large marine forests which provide habitat and food to other species [4], analogue to land-based tree forests. They thus have a high impact on local environments and need protection as a hot spot of biodiversity. Kelps are also of

high economic value as providers of alginates and as a food source [5]. Despite their economical and ecological importance only the nuclear genomes of *Saccharina japonica* (Areschoug) C.E.Lane, C.Mayes, Druehl & G.W.Saunders [6], including different cultivars [7], and recently of *Undaria pinnatifida* (Harvey) Suringar [8] were published so far. This fact prompted the Phaeoexplorer project (<https://www.france-genomique.org/projet/phaeoexplorer/?lang=en>) aiming at whole breadth genomics of brown algae. However, several organellar genomes of brown algae have been published. This is partly due to their over-representation regarding their copy numbers in eukaryote cells compared to the nuclear genome. Thus, reconstruction of organellar genomes needs far less sequencing reads than constructing a draft nuclear genome. Moreover, the small size of organellar genomes enables their completion without the need of long read sequences from nanopore [9] or PacBio [10] sequencing. Thus, several genomes from kelp chloroplasts and mitochondria are available: 42 kelp mitochondrial genomes are stored in public databases, kelp chloroplast genomes are rarer with only yet 9 available at the NCBI nucleotide database. A number of chloroplast genomes were studied and published [11–14] yet only one

* Corresponding authors.

E-mail addresses: Klaus.Valentin@awi.de (K. Valentin), ludwig.eichinger@uni-koeln.de (L. Eichinger).

¹ Sadly Gernot Glöckner passed away a few weeks after submission of this manuscript.

<https://doi.org/10.1016/j.ygeno.2021.01.003>

Received 24 August 2020; Received in revised form 19 December 2020; Accepted 18 January 2021

Available online 22 January 2021

0888-7543/© 2021 The Author(s).

Published by Elsevier Inc.

This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

complete mitochondrial genome was so far analysed in more detail [15]. Other mitochondrial genomes were made publicly available but seemingly used for barcoding purposes only, e.g. [16,17]. In a previous study we analysed variation in kelp chloroplast genomes [11]. Only minor rearrangements took place during kelp chloroplast genome evolution and therefore available kelp chloroplast genomes appear collinear. This resembles the situation in land plants, which also mostly possess collinear chloroplast genomes [18,19]. However, kelp species are much younger than land plants with major speciation events occurring only 35 million years ago [20]. As collinearity describes the maintenance of genes in corresponding orders over time we could analyse the abundance of single nucleotide polymorphisms over the complete length of available chloroplast genomes [11]. This analysis showed that likely incomplete lineage sorting contributed to chloroplast evolution. The phylogenetic relationships of genera within the kelp lineage were resolved recently using partial organellar and nuclear genome data [20]. Additional species can easily be fitted into this phylogenetic tree as they become available. Yet it is unclear whether these data are sufficient to finally get a robust tree topology.

There are some kelp species, which are especially rare or growing in remote areas and thus are not easily collectable and readily available for analysis. Especially the endemic deep-water kelp species growing below the thermocline in deep warm-temperate to subtropical waters, such as *Laminaria rodriguezii* Bornet (Mediterranean Sea [21,22]), *Laminaria abyssalis* A.B.Joly & E.C. Oliveira (S-Atlantic) [23] or *Laminaria philippinensis* J.E.Petrov & M.V. Suchovejeva (Pacific Ocean), [24] have seldom been collected. Another habitat difficult to assess is the Arctic sublittoral where *Laminaria solidungula* (J. Agardh) thrives in shaded habitats [22]. Despite their differences in latitudinal distribution and temperature environment, these species all are subjected to especially low light conditions (e.g. [25,26]) and their environment seems to be rather constant compared to species with a wide horizontal or vertical distribution gradient (e.g. [27]).

Laminaria rodriguezii is restricted to deep water habitats stretching between 70 and 120 m and rarely up to 50 m or down to 260 m [22,27,28]. Temperatures there seem to be quite constant year round at 13–14 °C [27]. *L. solidungula* thrives in Arctic habitats where temperatures typically vary between –1.5 and 5 °C and populations at their southern distribution boundary in south east Canada subduct to zones with year round cold temperatures ([29] and references therein). Both species endure low light conditions over most of the year. Besides its endemism *L. rodriguezii* also is distinctive from all other kelp species, except *Laminaria sinclairii* (Harvey ex J.D.Hooker & Harvey) Farlow, Anderson & Eaton [30], in its ability to form stolons and produce clonal sporophytes by vegetative propagation [31] thereby being able to bypass the obligate sexual life history of Laminariales [32].

To be able to compare available chloroplast and mitochondrial genomes with those of endangered *L.rodriguezii* and *L. solidungula*, we sampled sporophytes of *L. rodriguezii* from its Mediterranean habitat via the GOMBESSA expedition on the 7-9th of July in 2019 at Banc de Magaud, a rocky ledge between 68 and 80 m depth off Le Levant Island (Provence, France) [33]. In addition, we cultivated Arctic *L. solidungula* from stock cultures to comparatively study their organellar genomes. We were interested to see whether these two species from extreme habitats and despite their phylogenetic difference [27], might share common characters or show signs of special adaptation in their organellar genomes that deviate from each other or from kelp species with broad distribution ranges.

2. Material and methods

2.1. Sampling

Cultivated sporophytes propagated from clonal gametophyte samples of *L. solidungula*, originally sampled at Spitsbergen were used for DNA extraction and all cultivation details are given in [11]. Several

samples of *L. rodriguezii* were taken from the Mediterranean Sea on the 7-9th of July 2019 at Banc de Magaud, a rocky ledge between 68 and 80 m depth off Le Levant Island (Provence, France; 6°0.67375 E - 43°0.05624 N) [33]. The *L. rodriguezii* samples were from in situ sporophytes, silica dried and then used for DNA extraction.

2.2. DNA extraction and PCR analysis

Freeze dried juvenile sporophytes of *L. solidungula* and silica dried sporophytic meristem discs of *L. rodriguezii* were grinded under liquid nitrogen to yield a fine powder, where all cell walls were destroyed. The DNA then was extracted according to Doyle and Doyle modified cetyl trimethyl ammonium bromide method (CTAB) [34]. PCR analysis with total DNA from *L. rodriguezii*, *Laminaria digitata* (Hudson) J.V.Lamouroux, and *L. solidungula* was essentially performed as described [35]. The external primers were specific for the rpl31 and 16S rRNA genes, respectively, which are located directly up- and downstream of the *L. rodriguezii* ORF. The internal primers were specific for the 5' and 3' region of the *L. rodriguezii* open reading frame (ORF), respectively. Primers are listed in Table S1.

2.3. Sequencing and assembly

Total DNA (5 µg) was converted to an Illumina sequencing library and analysed on an Illumina HiSeq machine. Read sizes in the paired end library were 150 bases with a mean distance of reads of 235 bases. Trimming and further processing were done with the Illumina software suit. De novo assembly was performed with abyss-pe [36] using kmers 40, 45, and 55. The resulting contigs were converted to a BLAST Table database and queried with the *L. digitata* chloroplast and mitochondrial genome sequences. Matching contigs were used to reconstruct the complete chloroplast and mitochondrial genomes by closing gaps with Gapfiller [37].

2.4. Annotation, alignment and phylogenetic analyses

The coding sequences (CDS) of the recently published *L. digitata* mitochondrial and plastid genomes [11] were used to find the corresponding CDS on the completed organellar genomes. Furthermore, with the algorithm implemented in tRNAscan [38] we detected the tRNA genes. Detection of the rRNA genes was done by BLASTing the respective *L. digitata* nucleotide sequences against the whole organellar genomes. We then scanned the sequence portions larger than 100 bases without any annotation for presence of additional open reading frames (ORFs). The completed annotated organellar genome annotation was converted to a GenBank file and visualized with OGDRAW [39].

Sequences of other mitochondrial and chloroplast genomes from kelp species used for phylogenetic analysis were retrieved from the NCBI database (Table 1). Collinearity of the assembled kelp chloroplast and mitochondrial genomes was tested with the nucmer tool of mummer [40]. All sequences were edited to start at the same position and then complete genomes were aligned with Clustal Omega [41,42]. Further manual inspection of the sequences was done to ensure proper alignment for phylogenetic analyses.

For phylogenetic analyses we used MEGAX [43] and MrBayes [44]. The evolutionary history was inferred by using the Maximum Likelihood method and General Time Reversible model [45]. Initial trees for the heuristic search were obtained automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise distances estimated using the Maximum Composite Likelihood (MCL) approach, and then selecting the topology with superior log likelihood value. We performed 500 bootstrap repetitions on the data sets. In case of the MrBayes analysis we set the evolutionary model to GTR with gamma distributed rate variation and a proportion of invariable sites. We used 80,000 generations to get the standard deviation of split frequencies below 0.01 and then used a burn-in of 1000.

Table 1

The organellar genomes used in this study. The IDs are the NCBI accession numbers.

	Species name	ID	Sequence length [bp]	Reference	
Mitochondrial genomes	<i>Desmarestia viridis</i>	AY500367.1	39,049	[47]	
	<i>Lessonia spicata</i>	NC_044181.1	37,097	[66]	
	<i>Nereocystis luetkeana</i>	NC_042395.1	37,399	[67]	
	<i>Macrocystis integrifolia</i>	NC_042669.1	37,366	[68]	
	<i>Laminaria solidungula</i>	MT732098	37,862	This study	
	<i>Laminaria rodriguezii</i>	MT732097	38,047	This study	
	<i>Laminaria hyperborea</i>	JN099683.1	37,976	[69]	
	<i>Laminaria digitata</i>	AJ344328.1	38,007	[15]	
	<i>Saccharina latissima</i>	KM675818.1	37,659	[70]	
	<i>Saccharina longissima</i>	JN099684.1	37,628	[69]	
	<i>Saccharina japonica</i>	MG712776.1	37,657	[71]	
	Dongfang No.6				
	<i>Saccharina japonica</i> SJAPO	AP011493	37,657	[72]	
	<i>Saccharina japonica</i> var. <i>religiosa</i>	AP011494.1	37,657	[72]	
	<i>Saccharina longipedalis</i>	AP011497.1	37,657	[72]	
	<i>Saccharina japonica</i> var. <i>diabolica</i>	AP011496.1	37,657	[72]	
	<i>Saccharina japonica</i> var. <i>ochotensis</i>	AP011495.1	37,656	[72]	
	<i>Saccharina cichorioides</i> f. <i>coriacea</i>	AP011499.1	37,500	[72]	
	<i>Saccharina angustata</i>	AP011498.1	37,605	[72]	
	<i>Saccharina sculpera</i>	KR350664.1	37,627	[73]	
	<i>Costaria costata</i>	KF384641.1	37,461	[74]	
	<i>Undaria pinnatifida</i>	KF319031.1	37,402	[75]	
	Chloroplast genomes	<i>Lessonia spicata</i>	NC_044182.1	130,305	[66]
		<i>Laminaria solidungula</i>	MH784528.1	130,784	[11]
		<i>Costaria costata</i>	NC_028502.1	129,947	[14]
		<i>Undaria pinnatifida</i>	NC_028503.1	130,383	[13]
		<i>Laminaria digitata</i>	MH784527	130,376	[11]
		<i>Saccharina japonica</i>	JQ405663.1	130,584	[12]
		<i>Laminaria rodriguezii</i>	MT732096	131,092	This study

3. Results

3.1. New organellar genome sequences

We completely reconstructed the mitochondrial genomes of *L. rodriguezii* and *L. solidungula* and the chloroplast genome of *L. rodriguezii* from short read sequencing of extracted total DNA from the two species (Fig. 1). The organellar contigs were retrieved using related sequences. We closed the remaining gaps (one gap in the *L. solidungula* mitochondrial genome; three gaps in the *L. rodriguezii* chloroplast genome) from the assembly procedure using available raw read information (see methods). All these genomes were of similar length as their

related counterparts from other kelp (Table 1).

3.2. Phylogeny of kelp species using complete mitochondrial and chloroplast genomes

After annotation of the newly assembled genomes we checked all brown algal organellar genomes from the NCBI database for collinearity and retrieved 18 of the 42 collinear kelp mitochondrial genomes and 6 of the 9 collinear kelp plastid genomes (Table 1; Tables S2 and S3). According to this analysis all Laminariales (kelp) organellar genomes are collinear. Additionally, the mitochondrial genome of the brown alga *Desmarestia viridis* (O.F.Müller) J.V. Lamouroux [46,47], which is not a kelp species, turned out to be collinear to kelp mitochondrial genomes, providing a suitable outgroup for phylogenetic analyses. Other known brown algal mitochondrial and chloroplast genomes exhibit structural recombination and are therefore not compatible with the analyses performed here. In a previous study [11], we only used the coding sequences (CDS) for phylogenetic analyses of brown algal chloroplast genomes. As we here focus on the brown algal species with collinear organellar genomes, we can make use of this collinearity and employ complete organellar genome alignments directly for a phylogenetic analysis irrespective of their coding potential, thus also including tRNAs, rRNAs, and intergenic regions. Here, the alignment of intergenic regions is crucial to the outcome of the analysis. We therefore manually checked the alignments for incorrectly aligned portions and used a refined alignment for further analyses. The alignment of the mitochondrial genomes can be found in the supplement (Supplemental alignment 1). The chloroplast phylogenetic reconstruction recapitulated the previously published topology of kelp species obtained with concatenated CDS [11], albeit without the possibility to root them with other brown algae (Fig. S1). We used 18 complete mitochondrial genomes of kelps for the calculation of the mitochondrial tree together with the newly sequenced mitochondrial genomes from *L. solidungula* and *L. rodriguezii* and *D. viridis* as an outgroup. We found no differences in species placements between the two organellar trees (Fig. 2 and Fig. S1) in cases where both trees contained the species. Thus, we could place the newly added species in the trees with high confidence. As the mitochondrial maximum likelihood tree was not well resolved in the *Saccharina* species complex due to scarce phylogenetic signal, we additionally used MrBayes (see Methods) to compare its outcome with the maximum likelihood tree. This approach confirmed the tree topology of the maximum likelihood method and delivered even higher support at the deeper branches with cumulative probability of 100% at all nodes (Fig. S2). MrBayes also yielded two alternative topologies with lower cumulative probability affecting the placement of *S. japonica* var. *diabolica* (Miyabe) and *S. japonica* var. *ochotensis* (Miyabe) [48] (not shown).

3.3. Special features in organelle genomes

Overall, all kelp species organellar genomes contain the same gene sets, i.e. no gene was missing or additionally present in any species with two exceptions: i) In a previous sequencing effort we found, that *L. solidungula*, in comparison to other Laminariales, lacks an ORF (YCF37) in its plastid genome [11]. YCF37 is also present in cyanobacteria and it was reported that a deletion mutant shows enhanced susceptibility to high light [49]. *L. solidungula* is a low light adapted kelp and one of the reasons to sequence low light adapted *L. rodriguezii* and high light tolerant *L. digitata* was to investigate whether *L. rodriguezii* also lacks the ORF. This is, however, not the case and we also found no other common markers in the organellar genomes of the two low light species, *L. solidungula* and *L. rodriguezii*, which distinguishes them from the high light tolerant ones. ii) In this study we found in *L. rodriguezii* an additional ORF in the mitochondrial genome. This ORF is embedded in an approximately 700 bp stretch of DNA, is 456 bases or 152 amino acids long and, according to IPRscan [50], contains no recognizable domain.




Fig. 1. Newly sequenced organellar genomes. A: *Laminaria solidungula* mitochondrial genome. B: *Laminaria rodriguezii* mitochondrial genome. C: *L. rodriguezii* chloroplast genome. The arrow in B indicates the position of the ORF discussed in the text. The figures were made with OGdraw [39] and the legends are depicted in D.

To exclude contamination or an assembly error we confirmed the presence of the ORF in *L. rodriguezii* and its absence in *L. solidungula* and *L. digitata* by PCR (Fig. S3). BLAST searches with the ORF against all published 90 complete mitochondrial genomes of Phaeophyceae (Table S4) did not result in any match. However, a search against all brown algae revealed significant similarity to a mitochondrial ORF in *D. viridis* [47], where the similarity covers the 3' end of the *L. rodriguezii* ORF and at the 5' end of the *D. viridis* ORF over a length of 234 bases or 78 amino acids (Fig. 3B). Also, a short stretch of 45 bases upstream of orf211 in *D. viridis* is similar to ORF 2 in *L. rodriguezii* (Fig. 3A, C). Despite overall collinearity of the two mitochondrial genomes the two ORFs do not share the same relative position within the mitochondrial genome. While the *D. viridis* ORF is located between tRNA-K and tRNA-V adjacent to the 23S rRNA gene replacing tRNA-A, the *L. rodriguezii* ORF is placed between the 16S rRNA gene and rpl31 (Fig. 1B). The foreign

sequences in both, *L. rodriguezii* and *D. viridis*, show no detectable similarity to any known sequences including brown algal or kelp nuclear genomes, bacteria associated with brown algae, or viruses. We also found no evidence for transcriptional activity of the foreign sequences as there are no similar sequences in any currently available brown algal transcriptome dataset [51–54].

4. Discussion

4.1. Phylogenetic analysis

Since currently no complete nuclear data are available for most kelp species our phylogenetic analysis was restricted to that of the organellar genomes. Only 9 kelp chloroplast genomes are yet sequenced (Table S3), but these are collinear with only minor rearrangements at the inverted




Fig. 2. Phylogenetic analysis of kelp species with complete mitochondrial genomes. Maximum likelihood tree of aligned mitochondrial genomes. The evolutionary history was inferred by using the Maximum Likelihood method and General Time Reversible model [45].


repeat (IR) regions. Similarly, all 44, including the two new ones from *L. rodriguezii* and *L. solidungula*, available mitochondrial genomes from kelp species are collinear (Table S2) and we have used 18 for the phylogeny (Table 1). Even the mitochondrial genome from *D. viridis*, a more distantly related brown alga from the sister order Desmarestiales, which was used as outgroup in the phylogenetic analysis, has the same gene order as the kelp species. Thus, we conclude that rearrangements of mitochondrial genomes in kelps are rare if not completely absent. The chloroplast and mitochondrial genome sequences can be aligned completely irrespective of coding or intergenic region. Thus, for phylogenetic analyses we have slightly more data per species at our disposal than previous studies [20,27], but limited to organellar genomes only. With this data set it was impossible to resolve the *Saccharina* species complex with the maximum likelihood method indicating weak phylogenetic signal in mitochondrial genomes within this group. The MrBayes analysis however resolved the topology with high confidence values (99% each).

We compared our mitochondrial tree with the previously published comprehensive overview on kelp radiation [20] and found no difference. Our tree includes other kelp species than Starko *et al.* [20], but the species used in both studies are placed at alike positions in the tree. We therefore are confident that the placement of *L. solidungula* and *L. rodriguezii* within our tree is also correct. Interestingly the close relationship between *L. digitata* and *L. rodriguezii* is also reflected in their temperature tolerance profiles which are more similar to each other than to *L. ochroleuca* Bachelot Pylaie, a southern European kelp species [55]. The tree topology within the Laminariaceae conforms to that, which was previously published using roughly 2 kb of data from internal transcribed spacer (ITS; nuclear), Rubisco spacer (chloroplast) and mitochondrial cytochrome *c* oxidase sequences [27]. The number of informative sites present in organellar genomes thus seems to be sufficient to resolve the kelp phylogenetic tree with high confidence. When further kelp organellar genomes are being produced they can be readily added to the alignment. The phylogenetic position of *L. rodriguezii* presented here builds the framework, on which our further interpretation of results relied.

4.2. Deviations from collinearity in kelp organellar genomes

In light of the overall collinearity of kelp mitochondrial genomes it is surprising that we observed a long stretch of DNA in the *L. rodriguezii* mitochondrial genome adjacent to the 16S rRNA gene without counterparts in 43 other kelp species and 48 other Phaeophyceae mitochondrial genomes (Table S4). We can exclude a technical (sequencing or assembly) problem here since neither the cultivating nor the sequencing laboratory ever handled *D. viridis* and the sequences in the two species differ enough to exclude an accidental co-cultivation with assembly errors afterwards. Nevertheless, we confirmed the presence of the ORF in the mitochondrial DNA of *L. rodriguezii* and its absence in *L. digitata*, and *L. solidungula* by PCR (Fig. S3). Part of the additional DNA in *L. rodriguezii* can be translated into an ORF and yields a potential protein of 152 amino acids. The encoded protein of this ORF has an overlap with high identity of 72% and a similarity of more than 88% to part of a protein encoded by an ORF in the *D. viridis* mitochondrial genome (Fig. 3B). The nucleotide identity is even higher than the amino acid sequence identity indicating mutational changes affecting all codon positions. Possible reasons for this could be fast evolution (positive selection) or degradation. The nucleotide sequence similarity of the region in which ORF211 from *D. viridis* resides, extends 5' to its start codon (Fig. 3A, C), indicating that the annotated start codon of this ORF is not the original one. Furthermore, a stretch of 45 bases in the 5' region of ORF211 has also high similarity of 80% to ORF2 of *L. rodriguezii*. These similarities, extending from the originally defined ORF, indicate that the ancient coding gene in *D. viridis* was longer than the extant one. Similarities on nucleotide level are very unlikely to have occurred by chance and should have originated from independent lateral introduction of the same gene from an unknown source. A vertical transfer between the two species is unlikely since they are separated by several other brown algae in the tree (Fig. 2). In case of vertical inheritance some traces of this transfer should have remained in at least some of the other mitochondrial genomes of kelps. Moreover, the differing locations of the two ORFs argues against a vertical transfer in the light of the otherwise strictly collinear genomes. The different location together with the only

A



B

```

L. rodriguezii  MGYTHPQGCITSSGLSSPGCIVGITNVLNIHPSGMLSKEFLFIQILEQLTE
D. viridis      -----
L. rodriguezii  SILGIIHKYETTFPSSIFKNKPSDDPILISLLEQIVQSQEDQRHFVREMS
D. viridis      -----MEQVVQSQEDQRREVKELIS
                    :*:*****:*:*:
L. rodriguezii  NRQNI LLES LNDVVMRQDKISLDLQKMFYKDNNSWYYSHLSTTLQYLQII
D. viridis      NRQNI LLES LNDVVIRQDKMSLDLQKHLSSGKSHSWHY-LSTTLQYLQVI
                    *****:*****:  .:*: * *****:
L. rodriguezii  NV-----
D. viridis      SIFTPYIAKILPTTLIENIPLVNVVFFSPGQPEVVGPMRNMETNINTL
                    .:
L. rodriguezii  -----
D. viridis      AAALNQTNQGLAGVHTQVTALNETARGLVESQTNIQRIHENGLAEQLARD
L. rodriguezii  -----
D. viridis      LNTLATQTPYTPPVPTAEVTSAVDNSTPMDRPRLQNTAHLFRRT
    
```

C

```

L. rodriguezii  ATATAATAATGTATATATAATGGGGACATATCATCTCAAGGATGTATAACGTCTTCAGG
D. viridis      ---C...C.AT..G.G..TG.AA.A...C..T.G.T.-.CATCAA..AT.TTAG..ATAT
consensus      .....*:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*
L. rodriguezii  CCTCTCTTCCTCCGGGGTGTATAGTAGGAATCACTAACGTACTTAAATATACACCCCTAGCGG
D. viridis      ..C..T.C.....C.....T.....G..A...GC...
consensus      ..*:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*
L. rodriguezii  TATGCTTAGCAAGTTTTTATTCATACAAATCCTCGAACAACTCACGGAATCTATACTAGG
D. viridis      -----T...T..T..GG.T...TGA.TTGAG.T..TC...AGA..AA
consensus      .....*:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*
L. rodriguezii  CATAATTCACAAATATGAAACTACATTC---CTTCTAGTATTTCAAGAACAAGCCTTC
D. viridis      ACCC.AC.GTCTCA.A.GCCA.TTTAAG.CCCT.A.AGTG.CAAAA.CAT.ATGAA..CA
consensus      .....*:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*
L. rodriguezii  TTECGATCCCATTCCTTATTTCCCTTTTAGAGCAAATCGTGCAGTCCCAAGAAGATCAACG
D. viridis      AP.G.....CC...G.....C.G.....G.....C.....G...
consensus      ..*:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*
L. rodriguezii  TCACCTTTGTGAGAGAAATGTCAAATAGACAAAACATTCPTTTAGAAAGCTTAAATGATGT
D. viridis      S.GG.....A.....C.....C.....T.....C.....
consensus      .:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*
L. rodriguezii  CGTAATCGTCAAGATAAGATTTCATTGGACCTTAAACAAATGTTTTATAAGGATAATAA
D. viridis      S.....C.....G..C.....CC...C..C.GGTA.G.G.C.
consensus      .....*:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*
L. rodriguezii  CTCTGGTATTACTACACCTATCTACTACCTACAATATTTACAAATAATTAATGTGTA
D. viridis      ...G..C.....T.G.....G.C...G.A.A.T
consensus      ***:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*.**:*
    
```

Fig. 3. Alignment of the gene product of ORF2 of *Laminaria rodriguezii* with its partial counterpart from *Desmarestia viridis* (orf211 in NC_007684). A: A sketch of the two ORFs. ORFs are drawn as open rectangles, the unique regions of both species are represented by red lines. The two overlaps are depicted as grey boxes. B: The alignment of the translated protein sequences of the two ORFs. Asterisks denote conserved residues, colons conserved substitutions, and dots semi conserved substitutions. C: The section comprising the two overlaps aligned on nucleotide level. The overlapping parts from A are framed with red lines, the start codon of orf211 is highlighted by a green open box. Asterisks denote conserved residues in the consensus. Identical bases in *D. viridis* are represented by dots. The alignments were done with clustalw [65]. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

partial overlap suggests independent introductions of the two ORFs into the genomes and thus the overlapping conserved region.

In our opinion, the conservation of nucleotide sequences could have been achieved by two reasons: Functional conservation, i.e. purifying selection, or recent introductions from a similar source into the two species, so that only a few mutations could accumulate. Yet, in the overlapping stretch the nucleotide conservation is higher than the amino acid conservation, which points to loss of functional constraint and rapid degradation. We thus prefer the idea, that relatively recent independent events introduced these sequences into the mitochondrial genomes of *L. rodriguezii* and *D. viridis*, respectively. The origin of such sequences could lay in the nuclear genome, yet we could not find such a conserved gene in the published genome of a *Saccharina* species [7] and in our draft assembly of *L. digitata*. Thus, a horizontal gene transfer (HTG) of the genes from external sources is currently the best explanation for the presence of these sequences. Since we found no similarities to other sequences the source of this HTG remains unknown. We can, however, speculate, where this sequence came from. A HTG is often achieved via a vector, be it plasmid or virus. We therefore searched for potential candidates of the observed HTG, and found that in certain evolutionary branches so-called mitoviruses exist, which are exclusively found in mitochondria. Such mitoviruses replicate in mitochondria only and encode a single protein on their RNA(+) strand, an RNA-dependent RNA polymerase. They are so far known from fungi [56] and plants [57] only. In fungi they are transmitted via spores, mating or cytoplasmic mixing, but can also be transmitted between distantly related species by unknown means [56]. The viral RNA polymerases are generally not well conserved [58,59] and therefore it is not surprising that we found no match of the brown algal ORF to publicly available mitovirus sequences. Even the complete mitovirus sequences from plants and fungi cannot be aligned together and branch specific alignments (fungi or plants) yield only a few conserved residues [57]. Moreover, the apparent degradation of the ORFs likely affected the identifiability of the scarce conserved amino acids in RNA-dependent RNA polymerases. Nevertheless, mitoviruses are reasonable candidates for the independent infestation of the mitochondria of the two brown algae. Consequently, we also searched brown algal genome and transcriptome data for similar sequences but without positive match. One explanation for this situation could be that mitoviruses normally do not integrate into DNA and thus are mainly present in transcript data [57]. However, such data are currently scarce for brown algae and unfortunately focussed on polyA+ mRNAs only, which renders the probability to detect RNA viruses marginal.

From gene and virus integration assays it is known that such sequences preferentially integrate into transcriptionally highly active portions of a genome [60,61]. From the analysis of mammalian and plant mitochondrial genomes it is also known that the rRNA genes are normally expressed at a much higher level than the rest of the mitochondrial genome, which seems to be transcribed as one large RNA [62,63], however, sometimes post transcriptional regulation may

account for different levels of steady state levels of RNA [64]. Taken together it is intriguing to speculate that the observed locations of the two similar sequences in the vicinity of rRNA genes are due to the higher propensity of DNA pieces to integrate at highly expressed sequences.

5. Conclusion

Here we present three newly sequenced organellar genomes from kelp species together with a thorough analysis of their specific features and phylogenetic information content in comparison to other kelp organellar genomes. Collinearity of the organellar genomes within the kelp species together with their considerable length and thus informative positions make them suitable as very robust phylogenetic markers for the entire order. Our finding of an independent introduction of homologous nucleotide sequences into two brown algal mitochondrial genomes, potentially of mitovirus origin, is unprecedented and should be the beginning of endeavours to search for similar sequences in other brown algae, e.g. via the PHAEOEXPLORER project, and ultimately target complete mitovirus genomes from kelp.

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ygeno.2021.01.003>.

Author contributions

GG conceived the study and wrote the manuscript. SR and JR carried out the experiments and analysed the data. LE contributed to the experiments and to the final version of the manuscript. IB, AB, LR, and TT contributed material and to the final version of the manuscript. KV contributed to the collection of *L. rodriguezii* and discussions of the results. All authors provided critical feedback.

Data accessibility

The newly described organellar genome sequences have been deposited in the NCBI database under the Accession numbers MT732096, MT732097, and MT732098.

Author statement

We declare, that we have no competing financial interests.

Acknowledgement

We hereby confirm that the sampling of *Laminaria rodriguezii* was in compliance with the french ABS legislation. We thank Andreas Wagner for his efforts to participate in the collection of *L. rodriguezii* and the preparation of samples. We are also indebted to the Cologne Center for Genomics (CCG) for conducting the sequencing of our DNA preparations. The study was conducted in the framework of the European

Biodiversa call MARFOR (<http://www.biodiversa.org/1019>) with the DFG project number GL235/21-1.

We thank Andromede Oceanologie in the Frame of the Operation Gombessa for collecting materials and technical support.

References

- [1] T. Silberfeld, J.W. Leigh, H. Verbruggen, C. Cruaud, B. de Reviers, F. Rousseau, A multi-locus time-calibrated phylogeny of the brown algae (Heterokonta, Ochrophyta, Phaeophyceae): investigating the evolutionary nature of the “brown algal crown radiation”, *Mol. Phylogenet. Evol.* 56 (2010) 659–674.
- [2] F. Hillmann, G. Forbes, S. Novohradská, I. Ferling, K. Riege, M. Groth, M. Westermann, M. Marz, T. Spaller, T. Winckler, P. Schaap, G. Glockner, Multiple roots of fruiting body formation in Amoebozoa, *Genome Biol. Evol.* 10 (2018) 591–606.
- [3] A. Arun, S.M. Coelho, A.F. Peters, S. Bourdareaux, L. Peres, D. Scornet, M. Strittmatter, A.P. Lipinska, H. Yao, O. Godfroy, G.J. Montecinos, K. Avia, N. Macaisne, C. Troadec, A. Bendahmane, J.M. Cock, Convergent recruitment of TALE homeodomain life cycle regulators to direct sporophyte development in land plants and brown algae, *Elife* 8 (2019).
- [4] R.S. Steneck, M.H. Graham, B.J. Bourque, D. Corbett, J.M. Erlandson, J.A. Estes, M. J. Tegner, Kelp forest ecosystems: biodiversity, stability, resilience and future, *Environ. Conserv.* (2002) 436–459.
- [5] S. Bennett, T. Wernberg, S.D. Connell, A.J. Hobday, C.R. Johnson, E. S. Poloczanska, The ‘great southern reef’: social, ecological and economic value of Australia’s neglected kelp forests, *Mar. Freshw. Res.* 67 (2016) 47–56.
- [6] C.E. Lane, C. Mayes, L.D. Druel, G.W. Saunders, A multi-gene molecular investigation of the kelp (Laminariales, Phaeophyceae) supports substantial taxonomic re-organization 1, *J. Phycol.* 42 (2006) 493–512.
- [7] N. Ye, X. Zhang, M. Miao, X. Fan, Y. Zheng, D. Xu, J. Wang, L. Zhou, D. Wang, Y. Gao, Saccharina genomes provide novel insight into kelp biology, *Nat. Commun.* 6 (2015) 1–11.
- [8] T. Shan, J. Yuan, L. Su, J. Li, X. Leng, Y. Zhang, H. Gao, S. Pang, First genome of the Brown alga *Undaria pinnatifida*: chromosome-level assembly using PacBio and Hi-C technologies, *Front. Genet.* 11 (2020) 140.
- [9] H. Lu, F. Giordano, Z. Ning, Oxford Nanopore MinION sequencing and genome assembly, *Genom. Proteomics Bioinformatics* 14 (2016) 265–279.
- [10] A. Rhoads, K.F. Au, PacBio sequencing and its applications, *Genom. Proteomics Bioinformatics* 13 (2015) 278–289.
- [11] S. Rana, K. Valentin, I. Bartsch, G. Glöckner, Loss of a chloroplast encoded function could influence species range in kelp, *Ecol. Evol.* 9 (2019) 8759–8770.
- [12] X. Wang, Z. Shao, W. Fu, J. Yao, Q. Hu, D. Duan, Chloroplast genome of one brown seaweed, *Saccharina japonica* (Laminariales, Phaeophyta): its structural features and phylogenetic analyses with other photosynthetic plastids, *Mar. Genomics* 10 (2013) 1–9.
- [13] L. Zhang, X. Wang, T. Liu, G. Wang, S. Chi, C. Liu, H. Wang, Complete plastid genome sequence of the brown alga *Undaria pinnatifida*, *PLoS One* 10 (2015), e0139366.
- [14] L. Zhang, X. Wang, T. Liu, H. Wang, G. Wang, S. Chi, C. Liu, Complete plastid genome of the brown alga *Costaria costata* (Laminariales, Phaeophyceae), *PLoS One* 10 (2015), e0140144.
- [15] M.-P. Oudot-Le Secq, B. Kloareg, S. Loiseaux-De Goër, The mitochondrial genome of the brown alga *Laminaria digitata*: a comparative analysis, *Eur. J. Phycol.* 37 (2002) 163–172.
- [16] F.C. Küpper, A.F. Peters, D.M. Shewring, M.D. Sayer, A. Mystikou, H. Brown, E. Azzopardi, O. Dargent, M. Strittmatter, D. Brennan, Arctic marine phytobenthos of northern Baffin Island, *J. Phycol.* 52 (2016) 532–549.
- [17] M.D. Rothman, L. Mattio, R.J. Anderson, J.J. Bolton, A phylogeographic investigation of the kelp genus *Laminaria* (Laminariales, Phaeophyceae), with emphasis on the South Atlantic Ocean, *J. Phycol.* 53 (2017) 778–789.
- [18] Y. Huo, L. Gao, B. Liu, Y. Yang, S. Kong, Y. Sun, Y. Yang, X. Wu, Complete chloroplast genome sequences of four *Allium* species: comparative and phylogenetic analyses, *Sci. Rep.* 9 (2019) 1–14.
- [19] V.A. Thode, L.G. Lohmann, Comparative chloroplast genomics at low taxonomic levels: a case study using *Amphilophium* (Bignoniaceae, Bignoniaceae), *Front. Plant Sci.* 10 (2019) 796.
- [20] S. Starko, M.S. Gomez, H. Darby, K.W. Demes, H. Kawai, N. Yotsukura, S. C. Lindstrom, P.J. Keeling, S.W. Graham, P.T. Martone, A comprehensive kelp phylogeny sheds light on the evolution of an ecosystem, *Mol. Phylogenet. Evol.* 136 (2019) 138–150.
- [21] M.E. Bornet, Note sur une nouvelle espèce de Laminaria (*Laminaria rodriguezii*) de la Méditerranée, *Bulletin de la société botanique de France* 35 (1888) 361–366.
- [22] K. Lüning, *Seaweeds: Their Environment, Biogeography, and Ecophysiology*, John Wiley & Sons, 1990.
- [23] A.B. Joly, Two Brazilian laminarians, *Inst. Pesq. Mar.* 4 (1967) 1–13.
- [24] J. Petrov, M. Suchovejeva, G. Avdejev, Species generis *Laminaria* Lam. e mari philippinensi nova, *Novosti Sistematiki Nizshikh Rastenii* 10 (1973) 59–61.
- [25] K. Dunton, Growth and production in *Laminaria solidungula*: relation to continuous underwater light levels in the Alaskan high Arctic, *Mar. Biol.* 106 (1990) 297–304.
- [26] S. Joher, E. Ballesteros, E. Cebrian, N. Sánchez, C. Rodríguez-Prieto, Deep-water macroalgal-dominated coastal detritic assemblages on the continental shelf off Mallorca and Menorca (Balearic Islands, Western Mediterranean), *Bot. Mar.* 55 (2012) 485–497.
- [27] A. Žuljević, A.F. Peters, V. Nikolić, B. Antolić, M. Despalatović, I. Cvitković, I. Isajlović, H. Mihanović, S. Matijević, D.M. Shewring, The Mediterranean deep-water kelp *Laminaria rodriguezii* is an endangered species in the Adriatic Sea, *Mar. Biol.* 163 (2016) 69.
- [28] A. Ercegović, La végétation des algues sur les fonds pécheux de l’Adriatique, Institut za Oceanografiju i Ribarstvo, 1960.
- [29] R. Müller, T. Laepple, I. Bartsch, C. Wiencke, Impact of oceanic warming on the distribution of seaweeds in polar and cold-temperate waters, *Bot. Mar.* 52 (2009) 617–638.
- [30] W. Farlow, C. Anderson, D. Eaton, *Algae exsiccatae Americae borealis*, Boston, Massachusetts 2, 1878, pp. 51–100.
- [31] H. Huvé, Présence de *Laminaria rodriguezii* Bornet sur les côtes françaises de la Méditerranée, *REC. Stat. Mar. Endoume*, Fr 15 (1955) 73–91.
- [32] M.J. Wynne, H. Bold, *Introduction to the Algae: Structure and Reproduction*, Prentice-Hall, Incorporated, 1985.
- [33] J. Deter, J.-B. Juhel, E. Boulanger, N. Guellati, S. Maun, F. Holon, D. Mouillot, R. Hocdé, **Gombessa 5 cruise: CTD profiles in western Mediterranean, July 2019.**, SEANO, 2020, doi.org/10.17882/71814.
- [34] J.J. Doyle, J.L. Doyle, Isolation of plant DNA from fresh tissue, *Focus* 12 (1990) 39–40.
- [35] P. Farbrother, C. Wagner, J. Na, B. Tunggal, T. Morio, H. Urushihara, Y. Tanaka, M. Schleicher, M. Steinert, L. Eichinger, Dictyostelium transcriptional host cell response upon infection with *Legionella*, *Cell. Microbiol.* 8 (2006) 438–456.
- [36] J.T. Simpson, K. Wong, S.D. Jackman, J.E. Schein, S.J. Jones, I. Birol, ABYSS: a parallel assembler for short read sequence data, *Genome Res.* 19 (2009) 1117–1123.
- [37] M. Boetzer, W. Pirovano, Toward almost closed genomes with GapFiller, *Genome Biol.* 13 (2012) R56.
- [38] T.M. Lowe, S.R. Eddy, tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence, *Nucleic Acids Res.* 25 (1997) 955–964.
- [39] S. Greiner, P. Lehwork, R. Bock, OrganellarGenomeDRAW (OGDRAW) version 1.3.1: expanded toolkit for the graphical visualization of organellar genomes, *Nucleic Acids Res.* 47 (2019) W59–W64.
- [40] S. Kurtz, A. Phillippy, A.L. Delcher, M. Smoot, M. Shumway, C. Antonescu, S. L. Salzberg, Versatile and open software for comparing large genomes, *Genome Biol.* 5 (2004) R12.
- [41] H. McWilliam, W. Li, M. Uludag, S. Squizzato, Y.M. Park, N. Buso, A.P. Cowley, R. Lopez, Analysis tool web services from the EMBL-EBI, *Nucleic Acids Res.* 41 (2013) W597–W600.
- [42] F. Sievers, A. Wilm, D. Dineen, T.J. Gibson, K. Karplus, W. Li, R. Lopez, H. McWilliam, M. Remmert, J. Söding, Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal omega, *Mol. Syst. Biol.* 7 (2011) 539.
- [43] S. Kumar, G. Stecher, M. Li, C. Knyaz, K. Tamura, MEGA X: molecular evolutionary genetics analysis across computing platforms, *Mol. Biol. Evol.* 35 (2018) 1547–1549.
- [44] F. Ronquist, M. Teslenko, P. Van Der Mark, D.L. Ayres, A. Darling, S. Höhna, B. Larget, L. Liu, M.A. Suchard, J.P. Huelsenbeck, MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space, *Syst. Biol.* 61 (2012) 539–542.
- [45] M. Nei, S. Kumar, *Molecular Evolution and Phylogenetics*, Oxford university press, 2000.
- [46] J.V. Lamouroux, Essai sur les genres de la famille des thalassiphytes non articulées, De l’imprimerie de A. Belin, 1813.
- [47] M.-P. Oudot-Le Secq, S. Loiseaux-de Goer, W.T. Stam, J.L. Olsen, Complete mitochondrial genomes of the three brown algae (Heterokonta: Phaeophyceae) *Dictyota dichotoma*, *Fucus vesiculosus* and *Desmarestia viridis*, *Curr. Genet.* 49 (2006) 47–58.
- [48] N. Yotsukura, A systematic re-examination of four *Laminaria* species: *L. japonica*, *L. religiosa*, *L. ochotensis* and *L. diabolica*, *J. Jpn. Bot.* 83 (2008) 165–176.
- [49] U. Dühring, K.D. Irrgang, K. Lunser, J. Kehr, A. Wilde, Analysis of photosynthetic complexes from a cyanobacterial ycf37 mutant, *Biochim. Biophys. Acta* 1757 (2006) 3–11.
- [50] P. Jones, D. Binns, H.-Y. Chang, M. Fraser, W. Li, C. McAnulla, H. McWilliam, J. Maslen, A. Mitchell, G. Nuka, InterProScan 5: genome-scale protein function classification, *Bioinformatics* 30 (2014) 1236–1240.
- [51] H. Li, C. Monteiro, S. Heinrich, I. Bartsch, K. Valentin, L. Harms, G. Glöckner, E. Corre, K. Bischof, Responses of the kelp *Saccharina latissima* (Phaeophyceae) to the warming Arctic: from physiology to transcriptomics, *Physiol. Plant.* 168 (2020) 5–26.
- [52] C. Monteiro, S. Heinrich, I. Bartsch, K. Valentin, E. Corre, J. Collén, L. Harms, G. Glöckner, K. Bischof, Temperature modulates sex-biased gene expression in the gametophytes of the kelp *Saccharina latissima*, *Front. Mar. Sci.* 6 (2019) 769.
- [53] C. Monteiro, S. Heinrich, I. Bartsch, K. Valentin, E. Corre, J. Collén, L. Harms, G. Glöckner, K. Bischof, Temperature dependent sex-biased gene expression in the gametophytes of the kelp *Saccharina latissima*, *bioRxiv* (2019) 750455.
- [54] C.M.M. Monteiro, H. Li, K. Bischof, I. Bartsch, K.U. Valentin, E. Corre, J. Collén, L. Harms, G. Glöckner, S. Heinrich, Is geographical variation driving the transcriptomic responses to multiple stressors in the kelp *Saccharina latissima*? *BMC Plant Biol.* 19 (2019) 1–15.
- [55] I. Tom Dieck, North Pacific and North Atlantic digitate *Laminaria* species (Phaeophyta): hybridization experiments and temperature responses, *Phycologia* 31 (1992) 147–163.
- [56] B.I. Hillman, G. Cai, The family *Narnaviridae*: simplest of RNA viruses, in: *Advances in Virus Research*, Elsevier, 2013, pp. 149–176.

- [57] M.L. Nibert, M. Vong, K.K. Fugate, H.J. Debat, Evidence for contemporary plant mitoviruses, *Virology* 518 (2018) 14–24.
- [58] A. Bartholomäus, D. Wibberg, A. Winkler, A. Pühler, A. Schlüter, M. Varrelmann, Deep sequencing analysis reveals the mycoviral diversity of the virome of an avirulent isolate of *Rhizoctonia solani* AG-2-2 IV, *PLoS One* 11 (2016), e0165965.
- [59] E.V. Koonin, The phylogeny of RNA-dependent RNA polymerases of positive-strand RNA viruses, *J. Gen. Virol.* 72 (1991) 2197–2206.
- [60] I.K. Christiansen, G.K. Sandve, M. Schmitz, M. Dürst, E. Hovig, Transcriptionally active regions are the preferred targets for chromosomal HPV integration in cervical carcinogenesis, *PLoS One* 10 (2015), e0119566.
- [61] S.A. O'Brien, K. Lee, H.Y. Fu, Z. Lee, T.S. Le, C.S. Stach, M.G. McCann, A.Q. Zhang, M.J. Smsanski, N.V. Somia, Single copy transgene integration in a transcriptionally active site for recombinant protein synthesis, *Biotechnol. J.* 13 (2018) 1800226.
- [62] P.M. Finnegan, G.G. Brown, Transcriptional and post-transcriptional regulation of RNA levels in maize mitochondria, *Plant Cell* 2 (1990) 71–83.
- [63] C.M. Gustafsson, M. Falkenberg, N.-G. Larsson, Maintenance and expression of mammalian mitochondrial DNA, *Annu. Rev. Biochem.* 85 (2016) 133–160.
- [64] P. Giegé, M. Hoffmann, S. Binder, A. Brennicke, RNA degradation buffers asymmetries of transcription in *Arabidopsis* mitochondria, *EMBO Rep.* 1 (2000) 164–170.
- [65] M.A. Larkin, G. Blackshields, N.P. Brown, R. Chenna, P.A. McGettigan, H. McWilliam, F. Valentin, I.M. Wallace, A. Wilm, R. Lopez, Clustal W and Clustal X version 2.0, *Bioinformatics* 23 (2007) 2947–2948.
- [66] D. Tineo, K.B. Rubio, J.B. Melendez, J.E. Mendoza, J.O. Silva, J. Perez, E. Esquerre, M. Perez-Alania, S.L. Fernandez, S.E. Aguilar, Analysis of the complete organellar genomes of the economically valuable kelp *Lessonia spicata* (Lessoniaceae, Phaeophyceae) from Chile, *Mitochondrial DNA Part B* 4 (2019) 2581–2582.
- [67] Z. Zheng, H. Chen, H. Wang, W. Jiang, Q. Cao, N. Du, Characterization of the complete mitochondrial genome of bull kelp, *Nereocystis luetkeana*, *Mitochondrial DNA Part B* 4 (2019) 630–631.
- [68] J. Chen, Y. Zang, S. Shang, X. Tang, The complete mitochondrial genome of the brown alga *Macrocystis integrifolia* (Laminariales, Phaeophyceae), *Mitochondrial DNA Part B* 4 (2019) 635–636.
- [69] J. Zhang, X. Wang, C. Liu, Y. Jin, T. Liu, The complete mitochondrial genomes of two brown algae (Laminariales, Phaeophyceae) and phylogenetic analysis within *Laminaria*, *J. Appl. Phycol.* 25 (2013) 1247–1253.
- [70] S. Wang, X. Fan, Z. Guan, D. Xu, X. Zhang, D. Wang, Y. Miao, N. Ye, Sequencing of complete mitochondrial genome of *Saccharina latissima* ye-C14, *Mitochondrial DNA Part A* 27 (2016) 4037–4038.
- [71] J. Zhang, N. Liu, T. Liu, The complete mitogenome of *Saccharina* cultivar 'Dongfang no. 6' and its phylogenetic analysis, *Mitochondrial DNA Part B* 4 (2019) 1161–1162.
- [72] N. Yotsukura, T. Shimizu, T. Katayama, L.D. Druehl, Mitochondrial DNA sequence variation of four *Saccharina* species (Laminariales, Phaeophyceae) growing in Japan, *J. Appl. Phycol.* 22 (2010) 243–251.
- [73] L. Zhang, X. Wang, T. Liu, G. Wang, S. Chi, C. Liu, H. Wang, Complete mitochondrial genome of *Kjellmaniella crassifolia* (Laminariaceae, Phaeophyceae): *Laminaria* and *Saccharina* are distinct genus, *Mitochondrial DNA Part A* 27 (2016) 4592–4594.
- [74] J.-Q. Qu, C. Liu, X.-M. Wang, Z.-B. Zhang, S. Chi, T. Liu, Complete mitochondrial genome of *Costaria costata* shows conservative evolution in Laminariales, *Mitochondrial DNA* 26 (2015) 919–920.
- [75] T.-Y. Li, J.-Q. Qu, Y.-J. Feng, C. Liu, S. Chi, T. Liu, Complete mitochondrial genome of *Undaria pinnatifida* (Alariaceae, Laminariales, Phaeophyceae), *Mitochondrial DNA* 26 (2015) 953–954.