

Blunted leptin sensitivity during hedonic overeating can be reinstated by activating galanin 2 receptors (Gal2R) in the lateral hypothalamus

Este Leidmaa^{1,2,3}  | Mary Gazea¹  | Alexandre V. Patchev¹ | Anna Pissioti¹ | Nils Christian Gassen¹ | Mayumi Kimura¹ | Zsolt Liposits⁴ | Imre Kallo⁴  | Osborne F. X. Almeida¹ 

¹Max Planck Institute of Psychiatry, Munich, Germany

²Graduate School of Systems Neuroscience, Munich University, Planegg-Martinsried, Germany

³Institute of Molecular Psychiatry, Bonn, Germany

⁴Institute of Experimental Medicine, Hungarian Academy of Sciences, Budapest, Hungary

Correspondence

Osborne F. X. Almeida, Max Planck Institute of Psychiatry – NeuroAdaptations, Munich, Germany
Email: osa@psych.mpg.de

Funding information

European Union (FP7 Health), Grant/Award Number: 259772 to OFXA; National Science Foundation of Hungary (Országos Tudományos Kutatási Alapprogramok), Grant/Award Number: K101326 to IK

Abstract

Aim: Since foods with high hedonic value are often consumed in excess of energetic needs, this study was designed to identify the mechanisms that may counter anorexigenic signalling in the presence of hedonic foods in lean animals.

Methods: Mice, in different states of satiety (fed/fasted, or fed/fasted and treated with ghrelin or leptin, respectively), were allowed to choose between high-fat/high-sucrose and standard foods. Intake of each food type and the activity of hypothalamic neuropeptidergic neurons that regulate appetite were monitored. In some cases, food choice was monitored in leptin-injected fasted mice that received microinjections of galanin receptor agonists into the lateral hypothalamus.

Results: Appetite-stimulating orexin neurons in the lateral hypothalamus are rapidly activated when lean, satiated mice consume a highly palatable food (PF); such activation (upregulated c-Fos expression) occurred even after administration of the anorexigenic hormone leptin and despite intact leptin signalling in the hypothalamus. The ability of leptin to restrain PF eating is restored when a galanin receptor 2 (Gal2R) agonist is injected into the lateral hypothalamus.

Conclusion: Hedonically-loaded foods interrupt the inhibitory actions of leptin on orexin neurons and interfere with the homeostatic control of feeding. Overeating of palatable foods can be curtailed in lean animals by activating Gal2R in the lateral hypothalamus.

KEYWORDS

galanin, ghrelin, lateral hypothalamus, leptin, orexin, overeating

1 | INTRODUCTION

The abundance and convenience of energy-rich foods with high hedonic value poses a homeostatic challenge for modern

humans and predisposes them to obesity and associated cardiometabolic and psychiatric diseases.¹ While the maladaptive peripheral and central responses to chronic overeating have been extensively explored,^{2–4} present knowledge of the

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Acta Physiologica* published by John Wiley & Sons Ltd on behalf of Scandinavian Physiological Society

proximal neurobiological mechanisms that initiate overeating is poor. In light of the “hedonia hypothesis of obesity” that posits that foods imbued with pleasurable signals override (and/or reset) homeostatic set-points,⁵ we here aimed at identifying mechanisms that may counter anorexigenic signalling when lean mice encounter hedonic foods.

The orexigenic peptide ghrelin stimulates appetite; it is the only “hunger hormone” known to date.^{6,7} Ghrelin levels are downregulated in the obese state, making its direct involvement in the pathophysiology of obesity unlikely.^{8,9} However, the early role of ghrelin in triggering obesity should not be underestimated since the start and termination of the feeding cycle depends on a fine balance between the orexigenic effects of ghrelin and those of anorexigenic hormones such as leptin. Leptin, released into the blood as energy reserves are replenished, reduces appetite and stimulates energy expenditure¹⁰; thus, leptin serves as a major satiety signal. While the ability of leptin to act as a physiological brake on eating is compromised in the obese state because of the development of leptin resistance,^{11,12} it seems plausible that transient leptin resistance may explain why metabolically healthy individuals indulge in foods of high hedonic value. The hedonic properties of palatable foods (PF) can be mainly ascribed to their high content of energy-dense fats and sugars that stimulate the senses of taste, smell and texture.

Ghrelin and leptin receptors are prominently expressed in hypothalamic nuclei (arcuate nucleus [Arc] and lateral hypothalamic area [LHA]) that are critical for the regulation of energy balance^{13–15} where these orexigenic (ghrelin) and anorexigenic (leptin) hormones exert opposing effects. Ghrelin stimulates orexigenic neuropeptide Y/agouti-related peptide (NPY) neurons in the Arc^{16,17} and orexin (OX/hypocretin) neurons that are expressed exclusively in the LHA.^{7,18–20} Leptin activates anorexigenic pro-opiomelanocortin (POMC) neurons and inhibits orexigenic NPY neurons^{3,21} which ultimately innervate OX neurons in the LHA. Besides increasing appetite, OX is also implicated in reward-related behaviour and as a driver of hedonic feeding.^{22,23} It is important to note that OX neurons that are subject to inhibition by leptin^{24,25} do not express LepRb^{14,26} but in addition to their POMC inputs, receive dense inputs from neighbouring LepRb-positive neurons. The latter co-express galanin (Gal), neurotensin (Nts) and gamma-aminobutyric acid (GABA).^{14,24,26–28} There is evidence that Gal mediates the inhibitory actions of leptin on OX neurons from studies using acute hypothalamic slices and knockout mouse models.^{24,25}

In this study, using lean, adult male mice, we tested the hypothesis that overeating of PF results from disruption of the mechanisms through which leptin inhibits OX neurons. To begin dissecting how different elements of this complex neurocircuitry contribute to hedonic overeating, we first tested whether acute PF affects the efficacy of leptin or ghrelin on food intake. First, we compared food consumption

in mice that received peripheral injections of either saline, ghrelin or leptin before being given access to standard chow (SC, here considered to be hedonically neutral) for 1 h vs saline-, ghrelin- or leptin-injected mice exposed to a so-called “choice paradigm” in which mice could choose between SC and PF (hedonically loaded) or SC over 1 h. Brain areas possibly contributing to hedonic eating were mapped using immediate early gene c-Fos, and c-Fos was monitored in first- and second-order neurons (POMC, OX); in addition, we examined the efficacy of leptin signalling (induction of pSTAT3)^{11,12} when leptin-treated mice were confronted with hedonically-loaded PF. Lastly, we examined whether intra-LHA injections of Gal can restore the ability of leptin to reduce hedonic eating.

Briefly, the present experiments demonstrated that exposure of mice to PF triggers hyperphagia, a phenomenon that can be attributed to an escape of OX neurons from the inhibitory actions of leptin. Furthermore, we show that the anorexigenic effects of leptin can be pharmacologically reinstated by stimulation of the Gal receptor 2 (Gal2R) in the LHA. These observations suggest that descending hedonic information impinges on Gal neurons which, in turn, gate the access of peripheral signals of satiety to OX neurons.

2 | RESULTS

2.1 | Palatable food challenges homeostatic control mechanisms in LHA

In order to study the efficacy of homeostatic signals during initial acute exposure to hedonic foods, non-obese mice were injected with either ghrelin to induce hunger or leptin to mimic satiety, before testing their intake of standard chow (SC) in a no-choice paradigm (SC/SC) vs palatable food (PF) in a choice paradigm (SC/PF), as shown schematically in Figure S1A,B. These tests were conducted at ZT3 in the light-phase (resting period) to increase the level of conflict between the homeostatic state of the animal and the hedonic motivation to eat.

In lean, fasted mice, independent experiments using within-subject (Figure 1A) and between-subject (Figure 1B) designs showed that systemic leptin injections are more effective at suppressing SC ingestion than PF within the first hour of food presentation (for normalized data, see Figure S1C). Pre-treatment with leptin before exposure to SC/SC suppressed eating, as expected. In contrast, leptin-pretreated mice that were subsequently given access to SC/PF choice escaped from the anorexigenic effects of the peptide (Figure 1B). As shown in Figure S1D, the amount of SC ingested in the SC/PF choice paradigm was negligible in both saline- and leptin-treated mice. The same experiment was repeated in the dark phase (ZT16) when similar results were obtained (data not shown).




FIGURE 1 Overeating of highly palatable food (PF) is enhanced by ghrelin but is not suppressed by leptin. A, Leptin (3 mg/kg ip, after 24 h fast) suppresses consumption of SC (within-subject design; $n = 18$); $F_{1,34} = 50.2$, $P < .0001$; $time \times drug$ interaction: $F_{3,34} = 33.1$, $P < .0001$, *Sidak's post hoc $P < .05$) but only partially reduces PF intake ($F_{1,34} = 7.1$, $P < .05$; $time \times drug$ interaction: $F_{3,34} = 24.9$, $P < .0001$, § $P < .05$). B, Leptin initially (during 1 h) suppresses SC but not PF intake (between-subject design; $n = 20$ -22/group; $leptin$ effect: $F_{1,81} = 4.564$, $P = .036$; $food$ type $\times leptin$ interaction: $F_{1,81} = 2.00$, $P = .16$, *Tukey's post hoc $P < .05$). C, In an independent set of satiated mice exposed to the SC/PF choice paradigm, ghrelin (Ghr) injections (0.3 mg/kg) stimulated the cumulative intake of palatable food (PF) over 24 h ($F_{1,34} = 12.6$, *Sidak's post-hoc $P < .01$); Ghr did not increase SC consumption ($F_{1,34} = 6.6$, $P < .05$, §post-hoc $P < .05$), within-subject design ($n = 18$). (D) Ghr stimulates PF ingestion and not that of SC (between-subject design, $n = 10$ /group; measurements 1 h-post treatment; $drug$ effect: $F_{1,35} = 10.2$, $P = .003$, *Tukey's post-hoc $P < .05$)

A converse set of experiments showed that ghrelin (Ghr) administration to mice that had been previously satiated on SC stimulated only PF and not SC consumption (compare Control SC/SC vs Choice SC/PF) (Figure 1C within-subject and Figure 1D between-subject experiment, normalized data in Figure S1F).

We next sought to identify the initial neural correlates of hedonic eating (ZT16, 4 h into the dark phase) when mice would have satiated their homeostatic energy needs. In this experiment, SC-satiated mice that were allowed to choose between a palatable milk solution and SC, preferentially consumed milk (0.52 ± 0.1 kcal/g BW) as compared to SC (0.02 ± 0.02 kcal/g BW) during the first hour. Neuronal activity in brain areas likely involved in the physiological and behavioural regulation of hedonic eating was monitored by measuring c-Fos (mRNA and protein) expression.^{29,30} After 1 h of exposure to milk, increased activity was observed in brain regions implicated in the processing of reward/hedonic information (Figure 2A), eg piriform cortex (olfaction), insular cortex (taste), ventral pallidum (hedonic, reward)

orbitofrontal cortex, lateral septum (association) and amygdala (emotional learning and reward valuation). Hedonic eating was also accompanied by strong activation of two satiety centres, the ventromedial and dorsomedial hypothalamic nuclei, as well of the lateral hypothalamic area (LHA). Orexin (OX) neurons located in the LHA^{19,31} are normally activated during hunger to stimulate food intake. In the present study, evaluation of pre-pro-OX mRNA transcript levels in the LHA of SC-satiated mice revealed that OX mRNA levels are up-regulated within 1 h of consuming milk and return to baseline levels after 24 h. Interestingly, OX mRNA was again up-regulated when mice, previously satiated on milk for 24 h, were offered a different hedonic stimulus (solid PF comprised of high-fat/high-sucrose), as shown in Figure 2B. These data imply that, in addition to hunger per se, hedonic foods increase the activity of OX neurons in the LHA. Under both conditions (presentation of milk for 1 h and presentation of solid PF for 1 h to mice previously satiated with milk), the increases in OX mRNA levels were accompanied by elevations in serum corticosterone (CORT) levels (Figure 2C).

(A)

Function	Region of interest	c-Fos mRNA expression
Sensory	Agranular insular cortex, <i>AID, AIV</i>	++
	Piriform cortex, <i>Pir</i>	+++
Motor	Dorsal striatum, <i>CPu</i>	no change
	Motor cortex, <i>M1, M2</i>	+
	Substantia Nigra, <i>SNR, SNC</i>	no change
Association	Orbitofrontal cortex, <i>VO, LO, MO</i>	+
	Prefrontal cortex, <i>IL, PL</i>	+
	Anterior cingulate cortex, <i>Cg1, Cg2</i>	no change
	Habenula, <i>LHbM, MHb</i>	no change
Emotion and memory (limbic areas)	Bed nucleus of stria terminalis, <i>BNST</i>	++
	Amygdala, <i>BLA, BMA, LA, CeA</i>	++
	Lateral septum, <i>LSI</i>	+++
Hedonic, reward	Nucleus accumbens, <i>NacSh, NacC</i>	no change
	Ventral pallidum, <i>VP</i>	++
	Ventral tegmental area, <i>VTA, PBP</i>	no change
Endocrine	Hypothalamus, <i>ARC, VMH, DMH</i>	++
	Lateral hypothalamus, <i>LH, PeF</i>	++
Brainstem	Locus Coeruleus, <i>LC</i>	no change
	Substantia Nigra, <i>SN</i>	no change
	Parabrachial nucleus, <i>PBN</i>	no change
	Nucleus of Solitary tract <i>NTS</i>	no change
	Pontine nuclei, <i>Pn</i>	++







FIGURE 2 Hedonic eating upregulates c-Fos and orexin expression in the LHA. A, Qualitative relative scores of c-Fos mRNA expression in various relevant brain areas of mice after they had consumed either a hedonic solution (milk) or water for 1 h; the mice were already satiated on standard chow (SC) (liquids co-presented with SC). Mice consumed significantly greater amounts of milk vs water (0.52 ± 0.1 and 0.02 ± 0.02 kcal/g BW respectively). B, Acute access (1 h) to hedonic foods (either milk or a solid high-fat-containing food [PF]) during a state of satiety (ZT 16) also upregulates pre-pro-OX mRNA expression in the LHA ($n = 2-3$ /group). C, Serum corticosterone (CORT) levels were increased after PF presentation for 1 h. Serum was collected at ZT 16 from control (SC-exposed) mice ($n = 10$), mice that had ingested milk for 1 h ($n = 10$) or 24 h ($n = 6$), and mice that were satiated (24 h) on milk before being allowed to eat a novel palatable food (PF) for 1 h ($n = 4$). D, Ingestion of PF (high-fat/high sucrose food) by fasted mice increases mRNA transcript levels of orexin (OX) ($F_{1,20} = 16.92$, $P = .0005$), but not pro-MCH. Importantly, exogenous leptin (3 mg/kg ip) does not influence OX expression in either SC/SC- or SC/PF-exposed animals. Numerical data are mean \pm SEM ($n = 6-7$ /group). *indicates $P < .05$

2.2 | Orexin neurons in LHA escape inhibition by leptin during eating


Since PF intake is not decreased by leptin (Figure 1A,B) and upregulates OX expression (Figure 2B), we next measured the effects of leptin on orexigenic peptide expression in the LHA. As observed in SC-satiated mice (Figure 2B), PF upregulated OX expression (Figure 2D). Unexpectedly,^{4,32} administration of exogenous leptin to previously fasted mice did not influence OX mRNA expression in the LHA (Figure 2D). Notably, neither PF nor leptin altered mRNA expression of another LHA orexigenic peptide, melanin-concentrating hormone (MCH) (Figure 2D). Nevertheless, leptin reduced the number of c-FOS/OX co-expressing LHA neurons in mice offered SC/SC. Consistent with the results of previous

electrophysiological studies,¹⁹ our observations indicate that leptin suppresses OX-ergic activity when mice are exposed to control food (SC). Interestingly, however, leptin did not affect OX neuron activity in mice given the SC/PF choice (Figure 3A,B); this result matches the observation that leptin does not curtail the eating of PF (Figure 1A,B).

2.3 | Initial contact with PF does not interrupt leptin signalling in LHA

Effective dosage of leptin was confirmed by assaying serum levels of leptin (Figure 4A) and of insulin and Ghr (Figure S2). Moreover, measurements of pSTAT3 (Tyr705), which mediates the actions of leptin,^{2,12} revealed that acute PF ingestion does not disrupt leptin signalling in the brain. Briefly,

(A) Anti-c-FOS/OREXIN



(B)




FIGURE 3 Leptin fails to suppress OX neuron activity in the LHA during hedonic eating. A, Double immune-labelling of c-FOS (black) and orexin (brown) neurons in the LHA. The high-power inset shows neurons that express either c-FOS (arrowheads) or, orexin (open arrow) and neurons that display colocalized c-FOS and orexin (arrows) immunoreactivity. B, Leptin treatment significantly decreased the number c-FOS/OX co-expressing neurons in the LHA of mice on the SC/SC diet. In contrast, the number of c-FOS/OX double-labelled neurons was not reduced when leptin was administered to mice given the SC/PF choice (drug effect: $F_{1,21} = 27.4$, $P < .0001$; food type effect: $F_{1,21} = 13.9$, $P = .0012$; drug \times food type interaction: $F_{1,21} = 12.2$, $P = .02$). Scale bar represents 50 μ m. Semi-quantitative analysis; mean \pm SEM; $n = 6-7$ /group; *Tukey's post-hoc $P < .05$





FIGURE 4 Hedonic PF eating does not interrupt proximal leptin signalling in hypothalamus. Leptin was injected (ip 3 mg/kg at ZT 3) 30 min before access to SC/SC and PF/SC choice PF over 1 h. A, Serum levels of leptin 1.5 h post-injection ($n = 8-10/\text{group}$). B, Representative results for total and pSTAT3 in hypothalamic lysates, as measured by capillary electrophoresis. C, Semi-quantitative analysis, based on capillary electrophoretic detection, of phosphorylated STAT3 (pSTAT3, Tyr705) in the hypothalamus after leptin injection. Leptin induced pSTAT3 expression to a similar extent in SC/SC- and SC/PF-exposed groups (*main leptin effect*: $F_{1,20} = 12.38$, $P = .0022$, *food type effect*: $F_{1,20} = 0.51$, $P = .86$). D, Number of pSTAT3-positive nuclei in the LHA (*main leptin and food type effects*: $F_{1,21} = 25.3$, $P < .0001$ and $F_{1,21} = 0.0149$, $P = .90$ respectively). E and F, Immunostaining of pSTAT3 in the Arc (Bregma -1.46) and LHA (Bregma -1.82) respectively. Scale bar: 50 μ m. * $P < .05$

exogenous leptin induced pSTAT3 levels in the hypothalamus (Figure 4B,C), including the Arc (Figure 4E) and LHA (Figure 4D,F).

Leptin is known to stimulate anorexigenic POMC-expressing neurons in the Arc which, in turn, inhibit orexiogenic NPY neurons in that nucleus.^{3,33,34} We here found that

FIGURE 5 Neurons in the hypothalamic arcuate nucleus (Arc) remain responsive to leptin during hedonic eating. A, Immunofluorescent images of the Arc of leptin-treated mice that could choose between either SC/SC or SC/PF show co-expression of POMC (red) and c-FOS (green); the upregulation of c-FOS expression in a subset of POMC-positive neurons is shown at high power in the *inset*. Scale bar: 20 μ m. B, Semi-quantitative analysis of c-FOS and POMC double-immunostained Arc neurons. Data (mean \pm SEM; n = 6-7/group; *Tukey's post hoc $P < .05$) shows increased activation (c-FOS) of POMC neurons after either leptin or SC/PF exposure (*main leptin and food type effects*: $F_{1,20} = 19.9$, $P = .0002$ and $F_{1,20} = 12.1$, $P = .002$, respectively; *leptin \times food type interactions*: $F_{1,20} = 2.66$, $P = .12$). C, Levels of NPY mRNA expression are reduced in mice exposed to either leptin and SC/SC or SC/PF (*leptin \times food type interaction*: $F_{1,25} = 7.5$, $P = .01$). Numerical data depicted are mean \pm SEM (n = 6-7/group); *Tukey's post hoc $P < .05$



exogenous leptin concomitantly activates POMC neurons (increased c-FOS expression; Figure 5A,B), and downregulates NPY mRNA expression (Figure 5C) in mice given SC/SC. Moreover, saline- and leptin-treated mice showed similar anorexigenic responses when presented with SC/PF (Figures 4C,D and 5B,C; also see food intake data in Figure 1B), indicating that PF does not immediately disrupt Arc components of the neurocircuitry that regulates feeding behaviour.³⁵⁻³⁷ It deserves pointing out that, since PF is known to induce the release of endogenous leptin from fat tissue,³⁸ it was not surprising to find that exogenous leptin did not potentiate the POMC cell-activating effects of PF per se (Figure 5A,B). The apparent discrepancy between these findings and those showing that PF abrogates the inhibitory effects of leptin on OX neurons (Figure 3B) led us to consider that additional regulatory pathways and mechanisms

may be engaged when hedonic foods are encountered during a state of satiety.

2.4 | Galanin (Gal) restores leptin efficacy during hedonia-driven overeating

Given the unlikelihood that POMC and NPY neurons in the Arc are the loci at which hedonic foods act to override leptin actions (Figure 5B,C), we hypothesized that PF perturbs signal reception and/or transmission in the LHA. This was based on previous *in vitro* work (acute hypothalamic slices) that demonstrated that Gal mediates leptin-induced hyperpolarization of OX neurons²⁵ and the data depicted in Figure 6A that show galanin (Gal) neurons terminating in the vicinity of OX neurons in the LHA, as also described by previous authors.^{14,26}




FIGURE 6 The ability of leptin to suppress hedonic eating is reinstated by intra-LHA injections of a galanin receptor 2 (Gal2R) agonist. A, Galanin-immunoreactive fibres (blue) terminating in the vicinity of activated (c-FOS-immunoreactive, green) orexin A-immunoreactive neurons (red) in the LHA (arrowheads, sites of contact). Tissues analysed were derived from experimental mice used in Figures 1B and 3-5. B, Verification of sites of galanin agonist microinjections; correctly targeted injections of saline (black rectangles), Gal1R (purple triangles) and Gal2R (blue circles) are shown on coronal maps (from Franklin and Paxinos, 2008); LHA outlined (dotted lines) on cresyl violet-stained coronal sections are shown in the lowest panel. C and D, Intra-LHA Gal2R agonist M1145 restores the suppressive effects of ip leptin (3 mg/kg) on PF ingestion. Compared to saline-treated (control) mice, PF intake was significantly reduced in M1145-treated mice after 1 h (in C, $*P < .05$; in D, drug effect: $F_{1,21} = 7.2$, $P = .014$; time effect $F_{3,63} = 142.6$, $P < .0001$; $n = 10-12$ /group). E, Intake of PF over 1 h in 24 h-fasted mice. Data in columns 1 and 2 are from experiment shown in Figure 1B ($n = 23$ /group); columns 3-5 show data obtained in mice co-treated with leptin and either intra-LHA saline, Gal1R (M617; 0.1 nmol/L) or Gal2R (M1145; 0.1 nmol/L) 10 min before food presentation leptin ($F_{4,74} = 2.96$, $P = .025$; $n = 10-12$ /group). Numerical values represent means \pm SEM

To examine the role of Gal in the escape from leptin-induced suppression of PF consumption, leptin-treated mice were given intra-LHA microinjections (Figure 6B) of M617

and M1145 (agonists of the galanin receptor 1 [Gal1R] and galanin receptor 2 [Gal2R], respectively)³⁹ immediately before being presented with PF. The ability of leptin to suppress

(PF) eating was restored after application of the Gal2R agonist M1145 (Figure 6C,D), whereas PF overeating persisted in mice receiving leptin and the Gal1R agonist M617 (Figures 6E and S3A). On the other hand, a low (0.1 nmol/L) (Figure 6C,D), but not high (1 nmol/L) (Figure S3A), dose of the Gal2R agonist “permitted” leptin to exert its inhibitory effects on PF eating (Figure S3A). These observations indicate that blockade of the inhibitory actions of Gal initiate PF overeating, as evidenced by the PF feeding data over 24 h, shown in Figure S3B.

3 | DISCUSSION

Despite considerable advances in the understanding of the neurobiological mechanisms underlying overeating and obesity, the proximal mechanisms that initiate overeating are largely unknown; greater knowledge of the pathways and mechanisms involved can serve to develop preventative strategies against obesity and its associated pathologies. Accordingly, we here examined the hypothesis that disruption of the homeostatic processes that normally signal satiety and terminate eating are disrupted upon initial encounters with palatable foods that have high hedonic value, therefore resulting in their overconsumption.⁵ The present study focused on leptin, a potent satiety factor³⁸ whose role in the regulation of hedonic eating has not been investigated to date. To this end, we developed a paradigm in which fasted, lean mice were injected with leptin before being exposed to either only standard food (SC/SC) or to a choice consisting of a highly palatable (high fat-high sucrose) food (PF) and standard chow (SC/PF).

Two independent experiments showed that systemic leptin injections are more potent at suppressing the consumption of SC in the control condition than PF (SC/PF choice). While chronic over-eating of energy-dense foods is causally linked to leptin resistance,^{2,3,12} these data are the first to demonstrate that a single, brief exposure to hedonic food can override physiological signals of satiety.

Consistent with reports that ghrelin stimulates hedonic eating⁴⁰ and that ghrelin increases the reward salience of high-fat foods in an OX-dependent manner,^{41,42} we observed that ghrelin selectively increases appetite for hedonic food. The fact that leptin, like ghrelin, caused unrestrained consumption of PF indicates the validity of the choice paradigm devised for these studies.

The lateral hypothalamic area (LHA), a brain area that plays a key role in the integration of hedonic and homeostatic information, is responsive to both, ghrelin and leptin.⁴³ The fact that the LHA innervates the VTA, an important driver of motivated behaviour,⁴⁴⁻⁴⁶ also suggests that this hypothalamic nucleus plays a key role in regulating the intake of hedonic foods. In support of the latter view, we found that access to

hedonic food increases neuronal activity in the LHA, as well as other areas responsible for processing reward/hedonic and/or homeostatic signals. Thus, the LHA might indeed represent a nexus at which PF challenges the homeostatic mechanisms that govern food intake. Consistent with reports that hunger^{19,31} and hedonic foods⁴⁷⁻⁵⁰ activate OX-expressing neurons in the LHA, we also observed a rapid induction of pre-pro-OX mRNA transcript levels in the LHA of mice given access to palatable milk after they were already satiated with SC. Our experiments also showed that OX neurons receive inputs that signal the hedonic properties of food: thus, OX expression was further upregulated in mice that were satiated on milk before being presented with a qualitatively different (taste, odour, texture) hedonic stimulus (solid PF comprised of high-fat/high-sucrose). It also deserves mentioning that PF eating was not only accompanied by increased OX neuron activity but also by elevated levels of serum corticosterone (CORT). This catabolic hormone, together with its ability to modulate cognitive and motivational behaviour,^{51,52} was previously shown to be stimulated by OX⁵³ and is likely to amplify the seeking of palatable, energy-dense foods.

The finding that PF defies leptin-mediated signals of satiety and upregulates OX expression (in both, satiated and fasted mice) prompted a detailed examination of the regulation of orexigenic peptide expression in the LHA by leptin. Moreover, while leptin (predictably, see ref. 23) reduced the number of activated OX neurons in mice offered SC/SC, it failed to inhibit OX neuron activity in mice that preferentially ate PF in the SC/PF choice paradigm. The incongruence between the results of the present behavioural observations, the observed insensitivity of OX neurons to leptin in the hedonic eating paradigm used here and reports that leptin inhibits OX neuron activity,^{19,25} prompted us to further investigate the mechanistic basis of PF-triggered non-responsiveness to leptin.

It is known that the hypothalamic actions of leptin depend on the phosphorylation of STAT3, with a parallel increase in the activity of POMC neurons and downregulation of NPY expression.^{2,3,33,34} We found that PF elicits anorexigenic responses in POMC and NPY neurons and, importantly, that PF does not impair the induction of pSTAT3 by leptin per se. Our demonstration that PF ingestion does not impair proximal signalling of leptin, required formulation of an alternative hypothesis to explain how OX neurons might escape from the inhibitory control of leptin during consumption of PF. Accordingly, we chose to examine whether PF perturbs signal reception and/or transmission in the LHA by interfering with the inhibitory control of OX neurons. While it is known that OX neurons do not express leptin receptors, it is worth noting that leptin receptor-expressing galanin (Gal) neurons (some of which co-express neurotensin/GABA) terminate in the vicinity of OX neurons^{14,25,26} (see Figure 6A). Moreover, Gal (but neither neurotensin nor GABA) was

previously demonstrated to mediate leptin-induced hyperpolarization of OX neurons in acute hypothalamic slices.²⁵ Against this background, we considered the possibility that galanin neurons are the locus where PF acts to interfere with the relay of leptin signals to OX neurons, thereby negating physiological restraints on eating. Using pharmacological activation of Gal receptors in the LHA, we found that microinjections of M1145, a specific Gal2R agonist, into the LHA could restore the ability of leptin to suppress PF eating. Thus, the initiation of PF overeating would appear to result from blockade of the inhibitory actions of Gal on neighbouring OX neurons. While this interpretation awaits testing in future studies, our observations provide a basis for understanding the mechanisms that initiate the excessive intake calories and, therefore, the onset of obesity. Notably, the finding that only low doses of M1145 could inhibit PF eating hints at the potential importance of the prevailing Gal-ergic tone. Furthermore, while our results may at first appear to be at odds with previous reports that Gal induces feeding⁵⁴ and food-seeking behaviour,⁵⁵ other work using mice in which leptin receptor (LepRb) expression was constitutively deleted in Gal neurons confirms that the inhibitory actions of leptin are mediated by GalR.²⁴ The precise mechanisms underlying the actions of Gal are unknown. However, previous work suggested that Gal2R are coupled to either excitatory or inhibitory G (Gi) proteins (Webling et al, 2012). Coupling of Gal2R to Gi could result in either hyperpolarization of OX neurons or inhibition of OX neurons through modulation of their presynaptic inputs; the latter mechanism would be consistent with the results of previously-reported experiments.²⁵ When considered in the light of previous publications,^{24,25}

the present results may be interpreted as follows: deficits in the release of Gal from LepRb-expressing galanin neurons in the LHA underpin the disinhibition of OX neurons during hedonic feeding. Unfortunately, technical limitations precluded demonstration that the Gal2 agonist specifically activated OX neurons (following Gal agonist treatments, precision of microinjection sites was verified using methylene blue – an essential step in light of the small size and difficult-to-access location of the LHA). Future studies will be needed to define the origin of the Gal-ergic neurons that innervate the OX neurons that are involved in hedonic overeating.

Recent work shows that leptin reduces the strength of presynaptic excitatory inputs to OX neurons that project to the VTA, an effect blunted in diet-induced obesity.⁴ Given that the LHA is innervated by forebrain structures such as the shell of the nucleus accumbens (NAc) and ventral pallidum (VP)^{43,56} which are activated by PF (Figure 2A), it is plausible that the aforementioned areas increase LHA neuron excitability and dampen competing inhibitory Gal-ergic inputs (from a presently unknown brain area) to OX neurons. At the same time, because OX neurons are known to integrate complex and even counteracting metabolic cues,^{19,43} a role for other competing influences on OX neurons cannot be excluded.

To sum up, our work demonstrates that hedonic eating results from the development of insensitivity to leptin in OX neurons, an effect which can be (at least partly) attributed to a loss of Gal-mediated inhibitory control over OX-ergic neurons in the LHA. This study reveals a potentially new pathway and mechanism that complements existing knowledge of the neural underpinnings of hedonic eating. Clearly,




FIGURE 7 Working model of the circuitry upon which leptin acts to signal energy balance and satiety (Arc nucleus is major component of the blue sphere) during SC feeding. Inhibition of OX neurons in the lateral hypothalamic area (LHA), involving leptin stimulation of Gal secretion from LepRb-bearing Gal neurons in the vicinity of OX neurons (see Figure 5A), is considered a key mechanism in the suppression of feeding. The model postulates that besides positive emotional, cognitive and hedonic drivers, PF overeating results from a disruption of the ability of leptin to influence communication between Gal and OX neurons since activation of Gal2R in the LHA reduces PF consumption (see Figure 5C-E)

feeding behaviour, which is essential for survival, depends on in-built fail-safe mechanisms, rather than simple homeostatic reflexes.⁵⁷ The multiplicity of simultaneously or sequentially-acting regulatory factors makes dissection of the role of individual signalling molecules challenging. Thus, the hypophagic response to just a single hormone (eg leptin) may be small and probabilistic.⁵⁷ In this study, robustness of the data was ascertained by repeating all behavioural experiments on at least two occasions. The model shown in Figure 7 is necessarily simple - it does not include all of the proposed molecular players and switches through which PF may modulate leptin signalling and actions in the hypothalamus.⁵⁸⁻⁶⁰ Rather, it highlights input-output relationships between the LHA and periphery as well as brain areas that receive and process sensory information^{4,43,61}; among these, the central amygdala and VTA, both innervated by OX-ergic neurons in the LHA, contribute to flavour and reward learning.^{4,62-66}

4 | MATERIALS AND METHODS

4.1 | Animals

Three-month old male C57Bl/6N mice (Charles River, Sulzfeld, Germany) were single-housed on arrival and acclimated (2 weeks) to local conditions before use (lights on at ZT0, lights off at ZT12), under standard conditions recommended for housing laboratory mice (temperature $21 \pm 2^\circ\text{C}$, relative humidity $50 + 10\%$), with ad libitum access to standard chow (SC) and water, unless otherwise specified. Experiments were approved by the local ethics board (Government of Upper Bavaria) and complied with Directive 2010/63/EU.

4.2 | Diet composition and sources

Standard chow (SC) was Diet 1320 from Altromin Spezialfutter GmbH & Co. KG (Lage, Germany); it contained 2.84 kcal/g (11% fat, 65% carbohydrate, of which 4.9% was sucrose). Palatable food (PF), purchased from ResearchDiets Inc (Cat. D12451; New Brunswick, NJ, USA), had a calorific content of 4.73 kcal/g (45% kcal from fat, 35% from carbohydrate, of which 17% was sucrose). Low-fat food (LF; Cat. D12450B from Research Diets Inc), contained 4.06 kcal/g (10% energy from fat, 70% from carbohydrate, of which 35% was sucrose); ResearchDiets Inc). The protein contents of the SC, HF and LF were similar (20%, 20% and 24% respectively), but the diets varied in sweetness (LF > HF > SC). Brain activation during the hedonic experience, was assessed in mice with given access to SC and milk (60 mL) of dairy cooking cr me [Molkerei Weihenstephan, Germany] diluted in water to provide 1.64 kcal/g and a final composition of 5% fat, 1.3% carbohydrate, 1% protein).

4.3 | Food choice paradigm

Standard chow (SC) was available throughout, unless otherwise specified. With the exception of one test when mice were presented with milk (1.64 kcal/g; 5% fat) and SC, animals could choose between SC, high-fat/high-sucrose (45% fat) and/or low-fat/high-sucrose (LF, 10% fat) foods that varied in sweetness (see above). The LF option was eventually dropped since mice neglected SC and LF in favour of high-fat food (hereinafter, "palatable food", PF) and comparisons of PF intake in the choice condition (SC/PF) were compared to SC intake when only SC (SC/SC) was provided.

4.4 | Drugs

Ghrelin was injected immediately before, and leptin 30 min before, presentation of test foods. Octanoylated rat ghrelin (Tocris, Bristol, UK) was given at a dose of 0.3 mg/kg, ip, as described in previous studies.⁶⁷⁻⁶⁹ Leptin (LEP-6; Protein Laboratories Rehovot Ltd., Rehovot, Israel) was injected ip at a dose of 3mg/kg.^{62,70,71} The galanin receptor 1 and 2 (Gal1R and Gal2R) agonists (M617 and M1145, respectively, both from Tocris) were injected (intra-LHA) at 0.1 or 1 nmol/L in a volume of 0.5 μL , as described under "Restoration of leptin responsiveness by intra-LHA galanin". Saline (0.9%) served as vehicle and diluent for all drugs injections.

4.5 | Brain activation by acute hedonic food experience

Mice ($n = 24$) were habituated to the presence of two water-containing bottles and had ad libitum access to SC throughout the experiment. At ZT16, 4 groups of mice ($n = 6$) could choose between bottles containing either water/water or water/milk (see above) for 1-24 h; one group received PF for 1 h (after 24 h access to milk. At sacrifice (1 or 24 h), trunk blood was collected for measuring corticosterone by radioimmunoassay (RIA-1364 kit from DRG, Marburg, Germany) and brains were processed to detect *c-fos* mRNA and c-FOS protein (see below). Pilot studies demonstrated greater consumption of milk (high hedonic value ascribable to its high fat and lactose content) vs standard chow (SC); to avoid the confound from differing calorific contents of the food options, the milk was diluted in water to contain 5% fat.

4.6 | Food preference as a function of natural and simulated satiety states

Using a non-randomized treatment/within-subject design (Figure S1A), mice were repeatedly exposed to the choice paradigm (HF, LF and SC over 24 h) or control diet (SC/SC), with a 2 d "wash-out" between exposures starting at ZT3, the resting phase for mice, in order to create a conflict

between homeostatic state and hedonic motivation of the animal. Animals were tested in either satiated or fasted (24 h) states (5 d recovery between fasts). Ghrelin (0.3 mg/kg) or leptin (3 mg/kg) were injected ip immediately (Ghr) or 30 min (leptin) before presentation of SC or SC/PF at ZT3. Ghrelin-injected mice had continuous access to SC and were naturally satiated at the time of injection; leptin was injected to mimic a state of satiety in mice that had been 24 h-fasted (Figure S1A). Food intake was monitored after 1, 3, 6 and 24 h.

Results of the first experiment were subsequently verified by applying a between-subject factorial design to 8 groups of fasted or satiated mice with experiments started at ZT3 ($n = 10$; BW similar between groups) that were exposed to one of the following drug-food combinations: vehicle (saline)-SC, ghrelin (Ghr)-SC or leptin-SC, vehicle-PF, and Ghr-PF or leptin-PF (Figure S1B). As before, test foods were presented to mice after saline or Ghr (0.3 mg/kg) injections (groups previously satiated on SC) or after saline or leptin (3 mg/kg ip) injections (24 h-fasted group). Food consumption was monitored after 1 h, after which the mice were sacrificed (see the next paragraph). Mice were exposed to PF (24 h, 5 day before testing) to avoid novelty confounds. The between-subject experiments with leptin were also repeated with starting point at ZT16 (dark-phase); the results were not different from those obtained at ZT3, (data not shown).

4.7 | Subjugation of leptin activity by hedonia-driven eating: neuroanatomical correlates

Mice from the between-subject design ($n = 56$, see Figure S1B) were sacrificed 1 h after eating either SC or PF. Trunk blood was collected for determination of leptin, insulin and Ghr using multiplex assays (described later). Brains ($n = 7$) were excised and prepared for in situ hybridization histochemistry and/or immunohistochemical analysis of c-fos, orexin, melanin-concentrating hormone (MCH) and neuropeptide Y (NPY) mRNA expression and c-FOS protein expression (for detailed methods, see later). Brains from a subset of mice ($n = 6$) were perfused with 4% *P*-formaldehyde and immunostained for c-FOS and pSTAT3, orexin-A and orexin-B, c-FOS and orexin-B, c-FOS and POMC, or c-FOS, orexin-B and galanin. Free-floating sections from the hypothalamus (every 6th section from Bregma -1.06 to -2.18) were analysed.

4.8 | Restoration of leptin responsiveness by intra-LHA galanin

Mice ($n = 39$) were outfitted with bilateral intra-LHA injection cannulae, as described below. After 2 weeks, mice

were fasted (24 h) and given leptin (3 mg/kg ip) at ZT 3, followed by an intra-LHA microinjection (0.5 μ L) of either 0.9% NaCl (vehicle) or Gal receptor (Gal1R or Gal2R) agonists (0.1 nmol/L each) 20 min later. Following a 10 min recovery period, mice were given the SC/PF choice (intake of each food type was measured for up to 24 h). After a 6 days rest, all procedures were repeated except that mice received intra-LHA saline or a higher dose (1 nmol/L) of the Gal1R or Gal2R agonists.

In preparation for the intra-LHA microinjections of the galanin receptor agonists, mice were anaesthetized with isoflurane, placed in a stereotaxic frame (Stoelting Co.) and subcutaneously injected with analgesics (Metacam, 0.5 mg/kg and atropine sulphate, 0.5 mg/kg; both from Braun Melsungen AG). Bilateral intra-LHA guide cannulas with 2-mm stylets (Plastics One) were implanted using the coordinates A/P: -1.5 , M/L ± 1 from Bregma (Franklin and Paxinos, 2013). After replacement of the stylets with removable dummy injectors (Plastics One), the guide cannulas were fixed to the skull with dental acrylic resin (Paladur; Heraeus Kulzer, Hanau, Germany). Animals were allowed to recover for 2 weeks before experimental manipulations. On the day of experiment, vehicle or drugs (GalR1 and GalR2 agonists) were infused bilaterally via infusion cannulas that targeted the LH at -5.2 mm from Bregma, over 3 min, using a micro-infusion pump. Infusion cannulas were held in place for 3 min after microinjection, removed and then replaced with dummy injectors. After completion of behavioural tests, 0.5 μ L methylene blue was infused via re-inserted injector cannulas. The mice were then anaesthetized and their brains snap-frozen, sectioned on a cryostat (20 μ m) and counterstained with cresyl violet to verify correct placement of the cannulae and site of microinjection by reference to a stereotaxic atlas of the mouse brain.⁷² Cannulae were misplaced in two mice; these animals were excluded from further analysis. The doses of Gal1R (M617^{73,74}) or a Gal2R (M1145^{75,76}) were chosen based on the previous studies.

4.9 | mRNA detection by in situ hybridization histochemistry (ISHH)

After removal from the skull, brains were flash-frozen in isopentane/dry ice and stored at -80°C until sagittal cryosectioning (10 μ m thick; lateral coordinates: 3.25, 2.52, 1.56, 0.96, 0.48, 0.24 mm) or coronal sectioning to detect c-fos mRNA (10 μ m thick; 2.34, 1.70, 0.74, -0.70 , -1.43 , -3.16 mm from Bregma). Sections (between Bregma -1.46 and -1.70 ; 10 μ m) were used to detect prepro-orexin, pro-MCH mRNA and NPY mRNA. The ISHH protocol followed has been previously described.⁷⁷ Briefly, following fixation, acetylation, dehydration and

delipidation, air-dried sections were hybridized with custom-synthesized oligoprobes (Sigma-Aldrich Chemie GmbH, Steinheim, Germany):

c-fos: 5'-GTTGACAGGAGAGCCCATGCTGGAG
AAGGAGTCGGCTGGGGAATG-3'

prepro-
orexin: 5'-AGCAGCGTCACGGCGGCCAGGGAACCT
TTGTAG-3'

pro-MCH: 5'-CAACATGGTCGGTAGACTCTCCACAGCATA
CACCTGAGCATGTCAA-3'

NPY: 5'-GTCCTCTGCTGGCGCGTCTCGCCCGGA
TTGTCCGGCTTGAGGGGTA-3'

Oligoprobes were 3' end-labelled with ³⁵S d-ATP (Perkin Elmer, Waltham, MA, USA). Specificity was controlled by hybridizing alternate sections with a corresponding sense sequence probe. Following an overnight incubation (37°C) and stringent washing, dehydrated sections were exposed to X-ray film (Carestream Biomax MR Film) for 24 h (OX and MCH), 3 days (NPY) or 10 days (c-fos), before developing. Images were captured and the optical densities of radioactive signals were either qualitatively evaluated or quantified, relative to a ¹⁴C standard (range: 0-35 nCi/mg; American Radiolabelled Chemicals Inc), using Image J software (<http://imagej.nih.gov/ij/download/zips/-j149.zip>).

4.10 | Immunohistochemistry (IHC)

Brains were flash-frozen in isopentane/dry ice immediately after sacrifice and stored at -80°C until cryosectioning (10 µm). A freezing microtome was used to cut 25 µm sections from 4% P-formaldehyde (PFA)-fixed brains. In the latter case, brains were initially fixed in situ by intra-cardiac perfusion of pentobarbital-anaesthetized mice with 0.9% saline, followed by PFA; excised brains were subsequently post-fixed in, cryo-preserved (30% sucrose/PBS) and stored at -20°C until sectioned. Unless otherwise specified, phosphate-buffered saline (PBS; 0.01 mol/L, pH 7.4) served as a general diluent and wash solution. Endogenous peroxidases were quenched 0.5% hydrogen peroxide (15 min), 0.2% Triton X-100 (15 min) was used for permeabilization, and non-specific antibody binding was blocked using 2% normal horse serum (20 min). Antigens detected by immunofluorescence were marked with direct-conjugated fluorescent antibodies or biotinylated secondary antibodies from Jackson ImmunoResearch (West Grove, PA). Details of primary antisera used, dilutions and incubation times are given in the Table 1.

Transmitted light and fluorescent microscopic images were captured using an AxioCam MRc 5 digital camera mounted on a Zeiss AxioImager M1 microscope, outfitted with AxioVision 4.6 software (Carl Zeiss). Confocal images were captured with

TABLE 1 Antisera, incubation conditions and detection systems used in immunohistochemistry

Antibodies	Source/Dilution	Incubation/detection
Single targets		
Anti-c-Fos, rb	1:5000; 18 h; Millipore (PC38)	BIOT-dk-anti-rb IgG (1:500; 2h; Jackson (711-65-152), ABC ^a , Ni/DAB (<i>grey-black</i>)
Anti-Orexin A, rb	1:5000; 18h; Phoenix Pharmaceuticals (H-003-30)	FITC-dk-anti-rb IgG (1:500; 2h; Jackson (711-095-152) (<i>green</i>)
Anti-Orexin B, gt	1:2000; 18 h; Santa Cruz Biotechnology (sc-8071)	CY3-dk-anti-gt IgG (1:500; 2h; Jackson (715-165-150) (<i>red</i>)
Anti-pSTAT3, rb	1:500; 48 h, Cell Signaling Technologies (#9131)	BIOT-dk-anti-rb IgG, ABC, biotin-tyramide (BT) amplification ^b , streptavidin (STA) 488 (<i>green</i>)
Multi-targets		
Anti-c-Fos, rb	1:6000; 48 h; Millipore (PC38)	BIOT-dk-anti-rb IgG, ABC, Ni/DAB, silver-gold intensification ^c (<i>black</i>)
Anti-Orexin B, gt	1:3000; 48 h; Santa Cruz Biotechnology (sc-8071)	BIOT-dk-anti-gt IgG (1:500; 2h, RT; Jackson 705-065-147), ABC, DAB (<i>brown</i>)
Anti-c-Fos, gt	1:10 000; 48h; Santa Cruz Biotechnology (sc-52)	BIOT-dk-anti-gt IgG + ABC +BT amplification, STA 488 (<i>green</i>)
Anti-POMC, rb	1:5000; 48 h, Phoenix Pharmaceuticals (H029-30)	CY3-dk-anti-rb IgG (1:500; 2h; Jackson 715-165-152) (<i>red</i>)
Anti-galanin, sheep	1:50 000; 48 h, kindly gifted by Dr I. Merchenthaler, University of Maryland, College Park, MD	AMCA dk-anti-sheep IgG (1:500; 2h; Jackson 713-155-003) (<i>blue</i>)

^aavidin-biotin complex (ABC; Vector Laboratories, Burlingame, CA); 1:1000, 2 h, RT;

^bBiotin-tyramide (BT) amplification (Hopman et al, 1998); secondary streptavidin fluorescent antibody (Alexa 488-streptavidin from Molecular Probes, S-11223 Thermo Fisher; 1:1000, 12 h, RT.

^cLiposits, Setalo and Flerko B (1984).⁷⁸

^dAnti-orexin-Texas Red®-X conjugate prepared using Zenon Tricolor Rabbit IgG Labeling Kit #2, Thermo-Fischer Scientific (Z- 25 370, Zenon®).

an inverted Nikon Eclipse Ti-E microscope, equipped with an AIR confocal system (Nikon, Vienna, Austria). Digital images were processed with Adobe Photoshop CS software (Adobe Systems) and equally adjusted for brightness and contrast. A summary of the antibodies, incubation conditions and detection systems used is provided in Table 1.

4.11 | Capillary electrophoretic detection of STAT3 immunoreactivity

Hypothalami were homogenized in 100 mmol/L Tris-HCl, pH 8 (with 1 mmol/L EDTA, 250 mmol/L NaCl, 25 mmol/L MgCl₂, 10% glycerol, 1% Nonidet and Roche cOmplete Protease Inhibitor Cocktail and phosphatase inhibitor cocktails 2 and 3). After determining protein contents, samples were assayed for pSTAT3, total STAT3 and actin by semi-automatic capillary electrophoresis (2 µg protein/µL per capillary) using a *Wes-Mouse* Masterkit (PS-MK17, ProteinSimple); electrophoretic separation and quantification of concentrations of proteins of interest was performed with a Wes™ (ProteinSimple) apparatus. The following primary antibodies were used: anti-pSTAT3, Tyr705 (1:25, #9145, Cell Signaling Technologies), anti-STAT3 (1:100, #9139, Cell Signaling Technologies) and actin (1:3000 for pSTAT3 and 1:1000 for STAT3, #4967, Cell Signaling Technologies), using anti-rabbit IgG (042-206, ProteinSimple) as secondary antibody. Incubations were carried out at room temperature for 2 h. Total STAT3 and pSTAT3 signals were normalized against actin signals.

4.12 | Multiplex assay of serum leptin, insulin and ghrelin

Serum concentrations of leptin, insulin and ghrelin were measured using Luminex®-based technology Milliplex MAP kit (#MMHMAG-44K-07.Mouse, Merck Chemicals, Am Kronberger, Schwalbach, Germany) and the Bio-Plex system for signal detection (Bio-Rad Laboratories GmbH, Munich, Bayern, Germany). The assay sensitivities for leptin, insulin and ghrelin were 40, 30 and 3 pg/mL, respectively; values below these limits were excluded from the analysis. Assays were performed at the Else Kröner-Fresenius Center for Nutritional Medicine, Technical University Munich, courtesy of Professor Hans Hauner and Manuela Hubersberger.

4.13 | Statistics

Statistical differences were tested using GraphPad Prism 6 software. Student's *t*-tests were used to compare data from pairs of groups. Following 2-way ANOVA, *post hoc* multiple comparisons were made using Tukey's or Sidak's (repeated measures) tests (significance level: $P < .05$).

ACKNOWLEDGEMENTS

We thank Barna László (Budapest), Manuela Hubersberger (Weihenstephan) and Cornelia Flackskamm and Kathrin Hafner (Munich) for excellent technical assistance. This work was partly supported by the EU FP7 Consortium SwitchBox (Contract 259772 to OFXA) and by the National Science Foundation of Hungary (OTKA K101326 to IK).

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

AUTHORS' CONTRIBUTION

EL, AVP and OFXA conceived the studies; EL, MG, AVP and IK planned the experiments; EL, MG, AVP, AP, NCG, MK and OFX performed experiments; EL, AVP, ZL and IK carried out histological analyses; EL, AVP, IK and OFXA analysed and interpreted the data; EL, IK and OFX wrote the manuscript; all authors read and approved the manuscript.

ORCID

Este Leidmaa  <https://orcid.org/0000-0001-9409-2134>

Mary Gazea  <https://orcid.org/0000-0003-1468-8419>

Imre Kallo  <https://orcid.org/0000-0003-3412-4100>

Osborne F. X. Almeida  <https://orcid.org/0000-0001-7331-6928>

REFERENCES

1. Swinburn BA, Sacks G, Hall KD, et al. The global obesity pandemic: shaped by global drivers and local environments. *Lancet (London, England)*. 2011;378(9793):804-814.
2. Myers MG, Leibel RL, Seeley RJ, Schwartz MW. Obesity and leptin resistance: distinguishing cause from effect. *Trends Endocrinol Metab*. 2010;21(11):643-651.
3. Morton GJ, Meek TH, Schwartz MW. Neurobiology of food intake in health and disease. *Nat Rev Neurosci*. 2014;15(6):367-378.
4. Liu J-J, Bello NT, Pang ZP. Presynaptic regulation of leptin in a defined lateral hypothalamus-ventral tegmental area neurocircuitry depends on energy state. *J Neurosci*. 2017;37(49):11854-11866.
5. Berridge KC, Kringelbach ML. Neuroscience of affect: brain mechanisms of pleasure and displeasure. *Curr Opin Neurobiol*. 2013;23(3):294-303.
6. Kojima M, Hosoda H, Date Y, Nakazato M, Matsuo H, Kangawa K. Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature*. 1999;402(6762):656-660.
7. Kojima M, Kangawa K. Ghrelin: structure and function. *Physiol Rev*. 2005;85(2):495-522.
8. Tschöp M, Weyer C, Tataranni PA, Devanarayan V, Ravussin E, Heiman ML. Circulating ghrelin levels are decreased in human obesity. *Diabetes*. 2001;50(4):707-709.

9. Greenman Y, Golani N, Gilad S, Yaron M, Limor R, Stern N. Ghrelin secretion is modulated in a nutrient- and gender-specific manner. *Clin Endocrinol (Oxf)*. 2004;60(3):382-388.
10. Friedman J. 20 years of leptin: leptin at 20: an overview. *J Endocrinol*. 2014;223(1):T1-T8.
11. El-Haschimi K, Pierroz DD, Hileman SM, Bjørbaek C, Flier JS. Two defects contribute to hypothalamic leptin resistance in mice with diet-induced obesity. *J Clin Invest*. 2000;105(12):1827-1832.
12. Münzberg H, Flier JS, Bjørbaek C. Region-specific leptin resistance within the hypothalamus of diet-induced obese mice. *Endocrinology*. 2004;145(11):4880-4889.
13. Elmquist JK, Coppari R, Balthasar N, Ichinose M, Lowell BB. Identifying hypothalamic pathways controlling food intake, body weight, and glucose homeostasis. *J Comp Neurol*. 2005;493(1):63-71.
14. Laque A, Zhang Y, Gettys S, et al. Leptin receptor neurons in the mouse hypothalamus are colocalized with the neuropeptide galanin and mediate anorexigenic leptin action. *Am J Physiol Endocrinol Metab*. 2013;304(9):E999-E1011.
15. López-Ferreras L, Richard JE, Anderberg RH, et al. Ghrelin's control of food reward and body weight in the lateral hypothalamic area is sexually dimorphic. *Physiol Behav*. 2017;176:40-49.
16. Shintani M, Ogawa Y, Ebihara K, et al. Ghrelin, an endogenous growth hormone secretagogue, is a novel orexigenic peptide that antagonizes leptin action through the activation of hypothalamic neuropeptide Y/Y1 receptor pathway. *Diabetes*. 2001;50(2):227-232.
17. Chen HY, Trumbauer ME, Chen AS, et al. Orexigenic action of peripheral ghrelin is mediated by neuropeptide Y and agouti-related protein. *Endocrinology*. 2004;145(6):2607-2612.
18. Diano S, Horvath B, Urbanski HF, Sotonyi P, Horvath TL. Fasting activates the nonhuman primate hypocretin (orexin) system and its postsynaptic targets. *Endocrinology*. 2003;144(9):3774-3778.
19. Yamanaka A, Beuckmann CT, Willie JT, et al. Hypothalamic orexin neurons regulate arousal according to energy balance in mice. *Neuron*. 2003;38(5):701-713.
20. Toshinai K, Date Y, Murakami N, et al. Ghrelin-induced food intake is mediated via the orexin pathway. *Endocrinology*. 2003;144(4):1506-1512.
21. Elias CF, Aschkenasi C, Lee C, et al. Leptin differentially regulates NPY and POMC neurons projecting to the lateral hypothalamic area. *Neuron*. 1999;23(4):775-786.
22. Nair SG, Golden SA, Shaham Y. Differential effects of the hypocretin 1 receptor antagonist SB 334867 on high-fat food self-administration and reinstatement of food seeking in rats. *Br J Pharmacol*. 2009;154(2):406-416.
23. Borgland SL, Chang S-J, Bowers MS, et al. Orexin A/hypocretin-1 selectively promotes motivation for positive reinforcers. *J Neurosci*. 2009;29(36):11215-11225.
24. Laque A, Yu S, Qualls-Creekmore E, et al. Leptin modulates nutrient reward via inhibitory galanin action on orexin neurons. *Mol Metab*. 2015;4(10):706-717.
25. Goforth PB, Leininger GM, Patterson CM, Satin LS, Myers MG. Leptin acts via lateral hypothalamic area neurotensin neurons to inhibit orexin neurons by multiple GABA-independent mechanisms. *J Neurosci*. 2014;34(34):11405-11415.
26. Louis GW, Leininger GM, Rhodes CJ, Myers MG. Direct innervation and modulation of orexin neurons by lateral hypothalamic LepRb neurons. *J Neurosci*. 2010;30(34):11278-11287.
27. Leininger GM, Jo Y-H, Leshan RL, et al. Leptin acts via leptin receptor-expressing lateral hypothalamic neurons to modulate the mesolimbic dopamine system and suppress feeding. *Cell Metab*. 2009;10(2):89-98.
28. Leininger G, Opland D, Jo Y-H, et al. Leptin action via neurotensin neurons controls orexin, the mesolimbic dopamine system and energy balance. *Cell Metab*. 2011;14(3):313-323.
29. Bullitt E. Expression of c-fos-like protein as a marker for neuronal activity following noxious stimulation in the rat. *J Comp Neurol*. 1990;296(4):517-530. <https://doi.org/10.1002/cne.902960402>.
30. Perrin-Terrin A-S, Jeton F, Pichon A, et al. The c-FOS protein immunohistological detection: a useful tool as a marker of central pathways involved in specific physiological responses in vivo and ex vivo. *J Vis Exp*. 2016;110:e53613.
31. Sakurai T, Amemiya A, Ishii M, et al. Orexins and orexin receptors: a family of hypothalamic neuropeptides and G protein-coupled receptors that regulate feeding behavior. *Cell*. 1998;92(4):573-585.
32. López M, Seoane L, del Carmen García M, et al. Leptin regulation of prepro-orexin and orexin receptor mRNA levels in the hypothalamus. *Biochem Biophys Res Commun*. 2000;269(1):41-45.
33. Schwartz MW, Seeley RJ, Campfield LA, Burn P, Baskin DG. Identification of targets of leptin action in rat hypothalamus. *J Clin Invest*. 1996;98(5):1101-1106.
34. Elias CF, Lee C, Kelly J, et al. Leptin activates hypothalamic CART neurons projecting to the spinal cord. *Neuron*. 1998;21(6):1375-1385.
35. Kleinridders A, Schenten D, Könnner AC, et al. MyD88 signaling in the CNS is required for development of fatty acid-induced leptin resistance and diet-induced obesity. *Cell Metab*. 2009;10(4):249-259.
36. Sohn J-W, Elmquist JK, Williams KW. Neuronal circuits that regulate feeding behavior and metabolism. *Trends Neurosci*. 2013;36(9):504-512.
37. Magnan C, Levin BE, Luquet S. Brain lipid sensing and the neural control of energy balance. *Mol Cell Endocrinol*. 2015;418:3-8.
38. Lee M-J, Fried SK. Integration of hormonal and nutrient signals that regulate leptin synthesis and secretion. *Am J Physiol Metab*. 2009;296(6):E1230-E1238.
39. Webbing K, Ruesson J, Bartfai T, Langel Ü. Galanin receptors and ligands. *Front Endocrinol (Lausanne)*. 2012;3:146.
40. Egecioglu E, Jerlhag E, Salomé N, et al. Ghrelin increases intake of rewarding food in rodents. *Addict Biol*. 2010;15(3):304-311.
41. Perello M, Sakata I, Birnbaum S, et al. Ghrelin increases the rewarding value of high-fat diet in an orexin-dependent manner. *Biol Psychiatry*. 2010;67(9):880-886.
42. Cone JJ, McCutcheon JE, Roitman MF. Ghrelin acts as an interface between physiological state and phasic dopamine signaling. *J Neurosci*. 2014;34(14):4905-4913.
43. Berthoud H-R, Münzberg H. The lateral hypothalamus as integrator of metabolic and environmental needs: from electrical self-stimulation to opto-genetics. *Physiol Behav*. 2011;104(1):29-39.
44. Shizgal P, Bielajew C, Corbett D, Skelton R, Yeomans J. Behavioral methods for inferring anatomical linkage between rewarding brain stimulation sites. *J Comp Physiol Psychol*. 1980;94(2):227-237.
45. Nieh E, Matthews G, Allsop S, et al. Decoding neural circuits that control compulsive sucrose seeking. *Cell*. 2015;160(3):528-541.
46. Tyree SM, de Lecea L. Lateral hypothalamic control of the ventral tegmental area: reward evaluation and the driving of motivated behavior. *Front Syst Neurosci*. 2017;11:50.
47. Wortley KE, Chang G-Q, Davydova Z, Leibowitz SF. Peptides that regulate food intake: orexin gene expression is increased during

- states of hypertriglyceridemia. *Am J Physiol Regul Integr Comp Physiol.* 2003;284(6):R1454-R1465.
48. Harris GC, Wimmer M, Aston-Jones G. A role for lateral hypothalamic orexin neurons in reward seeking. *Nature.* 2005;437(7058):556-559.
 49. Valdivia S, Patrone A, Reynaldo M, Perello M. Acute high fat diet consumption activates the mesolimbic circuit and requires orexin signaling in a mouse model. Chowen JA, ed. *PLoS ONE.* 2014;9(1):e87478.
 50. Valdivia S, Cornejo MP, Reynaldo M, De Francesco PN, Perello M. Escalation in high fat intake in a binge eating model differentially engages dopamine neurons of the ventral tegmental area and requires ghrelin signaling. *Psychoneuroendocrinology.* 2015;60:206-216.
 51. Ossewaarde L, Qin S, Van Marle H, van Wingen GA, Fernández G, Hermans EJ. Stress-induced reduction in reward-related prefrontal cortex function. *NeuroImage.* 2011;55(1):345-352.
 52. Graf EN, Wheeler RA, Baker DA, et al. Corticosterone Acts in the Nucleus Accumbens to Enhance Dopamine Signaling and Potentiate Reinstatement of Cocaine Seeking. *J Neurosci.* 2013;33(29):11800-11810.
 53. Bonnavion P, Jackson AC, Carter ME, de Lecea L. Antagonistic interplay between hypocretin and leptin in the lateral hypothalamus regulates stress responses. *Nat Commun.* 2015;6(1):6266.
 54. Leibowitz SF. Regulation and effects of hypothalamic galanin: relation to dietary fat, alcohol ingestion, circulating lipids and energy homeostasis. *Neuropeptides.* 2005;39(3):327-332. <https://doi.org/10.1016/j.npep.2004.12.022>.
 55. Qualls-Creekmore E, Yu S, Francois M, et al. Galanin-Expressing GABA Neurons in the Lateral Hypothalamus Modulate Food Reward and Noncompulsive Locomotion. *J Neurosci.* 2017;37(25):6053-6065.
 56. Root DH, Melendez RI, Zaborszky L, Napier TC. The ventral pallidum: Subregion-specific functional anatomy and roles in motivated behaviors. *Prog Neurobiol.* 2015;130:29-70.
 57. Woods SC, Begg DP. Food for Thought: Revisiting the Complexity of Food Intake. *Cell Metab.* 2015;22(3):348-351.
 58. Koch M, Varela L, Kim JG, et al. Hypothalamic POMC neurons promote cannabinoid-induced feeding. *Nature.* 2015;519(7541):45-50.
 59. Fenselau H, Campbell JN, Verstegen A, et al. A rapidly acting glutamatergic ARC→PVH satiety circuit postsynaptically regulated by α -MSH. *Nat Neurosci.* 2017;20(1):42-51.
 60. Campbell JN, Macosko EZ, Fenselau H, et al. A molecular census of arcuate hypothalamus and median eminence cell types. *Nat Neurosci.* 2017;20(3):484-496.
 61. Castro DC, Berridge KC. Opioid and orexin hedonic hotspots in rat orbitofrontal cortex and insula. *Proc Natl Acad Sci U S A.* 2017;114(43):E9125-E9134.
 62. Hommel JD, Trinko R, Sears RM, et al. Leptin Receptor Signaling in Midbrain Dopamine Neurons Regulates Feeding. *Neuron.* 2006;51(6):801-810.
 63. Domingos AI, Vaynshteyn J, Voss HU, et al. Leptin regulates the reward value of nutrient. *Nat Neurosci.* 2011;14(12):1562-1568.
 64. Castro DC, Cole SL, Berridge KC. Lateral hypothalamus, nucleus accumbens, and ventral pallidum roles in eating and hunger: interactions between homeostatic and reward circuitry. *Front Syst Neurosci.* 2015;9:90.
 65. Risco S, Mediavilla C. Orexin-1 receptor antagonist in central nucleus of the amygdala attenuates the acquisition of flavor-taste preference in rats. *Pharmacol Biochem Behav.* 2014;126:7-12.
 66. Terrill SJ, Hyde KM, Kay KE, et al. Ventral tegmental area orexin 1 receptors promote palatable food intake and oppose post-ingestive negative feedback. *Am J Physiol Integr Comp Physiol.* 2016;311(3):R592-R599.
 67. Wang L, Saint-Pierre DH, Taché Y. Peripheral ghrelin selectively increases Fos expression in neuropeptide Y - synthesizing neurons in mouse hypothalamic arcuate nucleus. *Neurosci Lett.* 2002;325(1):47-51.
 68. Yannielli PC, Molyneux PC, Harrington ME, Golombek DA. Ghrelin effects on the circadian system of mice. *J Neurosci.* 2007;27(11):2890-2895.
 69. Jerlhag E. Systemic administration of ghrelin induces conditioned place preference and stimulates accumbal dopamine. *Addict Biol.* 2008;13(3-4):358-363.
 70. Vaisse C, Halaas JL, Horvath CM, Darnell JE, Stoffel M, Friedman JM. Leptin activation of Stat3 in the hypothalamus of wild-type and ob/ob mice but not db/db mice. *Nat Genet.* 1996;14(1):95-97.
 71. Margetic S, Gazzola C, Pegg G, Hill R. Leptin: a review of its peripheral actions and interactions. *Int J Obes.* 2002;26(11):1407-1433.
 72. Franklin K, Paxinos G. *The Mouse Brain in Stereotaxic Coordinates.* Boston: Elsevier Academic Press; 2008.
 73. Lundström L, Sollenberg U, Brewer A, et al. A Galanin Receptor Subtype 1 Specific Agonist. *Int J Pept Res Ther.* 2005;11(1):17-27.
 74. Blackshear A, Yamamoto M, Anderson BJ, et al. Intracerebroventricular administration of galanin or galanin receptor subtype 1 agonist M617 induces c-Fos activation in central amygdala and dorsomedial hypothalamus. *Peptides.* 2007;28(5):1120-1124.
 75. Runesson J, Saar I, Lundström L, Järv J, Langel U. A novel GalR2-specific peptide agonist. *Neuropeptides.* 2009;43(3):187-192.
 76. Saar I, Runesson J, McNamara I, Järv J, Robinson JK, Langel U. Novel galanin receptor subtype specific ligands in feeding regulation. *Neurochem Int.* 2011;58(6):714-720.
 77. Murgatroyd C, Patchev AV, Wu Y, et al. Dynamic DNA methylation programs persistent adverse effects of early-life stress. *Nat Neurosci.* 2009;12(12):1559-1566.
 78. Liposits Z, Sétáló G, Flerkó B. Application of the silver-gold intensified 3,3'-diaminobenzidine chromogen to the light and electron microscopic detection of the luteinizing hormone-releasing hormone system of the rat brain. *Neuroscience.* 1984;13(2):513-525.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Leidmaa E, Gazea M, Patchev AV, et al. Blunted leptin sensitivity during hedonic overeating can be reinstated by activating galanin 2 receptors (Gal2R) in the lateral hypothalamus. *Acta Physiol.* 2020;228:e13345. <https://doi.org/10.1111/apha.13345>