

**MOLECULAR CHARACTERIZATION AND
BIOINFORMATICS ANALYSIS OF MICRORNA-
221-5P REGULATED BY STANDARDIZED
Polyalthia longifolia (Sonn.) Thwaites LEAF
EXTRACT IN HELA CELL LINES**

SHANMUGAPRIYA

UNIVERSITI SAINS MALAYSIA

2019

**MOLECULAR CHARACTERIZATION AND
BIOINFORMATICS ANALYSIS OF MICRORNA-
221-5P REGULATED BY STANDARDIZED
Polyalthia longifolia (Sonn.) Thwaites LEAF
EXTRACT IN HELA CELL LINES**

by

SHANMUGAPRIYA

**Thesis submitted in fulfilment of the requirements
for the degree of
Doctor of Philosophy**

May 2019

ACKNOWLEDGEMENT

First and foremost, I would like to thank God who granted me all the grace and strength to complete this research to a success. My genuine gratitude is always to my mother, Ms. Manimagalai and my sister, Ms. Subhasree for their endless love, support and encouragement throughout the duration of my Doctor of Philosophy programme. Their moral support and their trust on me have always been the bridge for the success of my research.

Without an expert's guidance, completion of this research would have been an arduous work to be submitted on time. Hence, I would like to express my heartfelt appreciation to my supervisor, Associate Professor Dr. Sasidharan Sreenivasan who has not only given me the opportunity to learn and conduct my research work under his supervision but also for his guidance and encouragement throughout the course of my study. I would also like to thank my co-supervisor, Dr. Nurulhasanah Othman who has always guided me throughout my research. I would like to extend my highest gratitude to both of them for sharing their invaluable knowledge and advice to conduct my Doctor of Philosophy programme to a successful end.

I would also like to acknowledge the financial assistance of Universiti Sains Malaysia (USM) by awarding me the Graduate Assistant Scheme which financially supported me throughout the duration of my PhD degree. Furthermore, I would also like to express my sincere appreciation to Professor Dr. Narazah and Mrs. Nor Fadhila from Advanced Medical & Dental Institute (AMDI), for their technical support and assistance especially in handling flow cytometry. I would also like to pay my gratitude to Mrs. Jamilah from the School of Biological Sciences, USM for

guiding me in the handling of SEM, TEM and epifluorescence microscopy. I would like to extend my acknowledgment for the imperative technical support of laboratory equipments and indispensable workshops offered by Institute for Research in Molecular Medicine (INFORMM).

This research was funded by Bridging Grant from Universiti Sains Malaysia, Pulau Pinang, Malaysia with grant number 304.CIPPM.6316068 and partly supported by the Fundamental Research Grant Scheme (FRGS; Grant No.: 203/CIPPM/6711379) from the Ministry of Education Malaysia, Government of Malaysia, Malaysia.

Last but not least, I would like to pay my bona fide gratitude to all my friends for supporting and encouraging me throughout the completion of my PhD programme. I pray to God that may all the individuals acknowledged above be enthroned with constant love, health and happiness.

SHANMUGAPRIYA

Institute for Research in Molecular Medicine

Universiti Sains Malaysia

May 2019

TABLE OF CONTENT

ACKNOWLEDGEMENT	ii
TABLE OF CONTENTS	iv
LIST OF TABLES	xi
LIST OF PLATES	xii
LIST OF FIGURES	xiii
LIST OF ABBREVIATIONS	xv
ABSTRAK	xx
ABSTRACT	xxii
CHAPTER 1: INTRODUCTION	1
1.1 Overview and rationale of study	1
1.2 <i>Polyalthia longifolia</i> as an important medicinal plant	2
1.3 Objectives	5
1.4 Workflow	6
CHAPTER 2: LITERATURE REVIEW	7
2.1 Cancer	7
2.1.1 Current development in cancer therapy	8
2.2 <i>Polyalthia longifolia</i>	10
2.2.1 Botanical description of <i>P. longifolia</i>	10
2.2.2 Pharmacological activities of <i>P. longifolia</i>	11
2.2.1(a) Anti-microbial activity	11
2.2.1(b) Anti-oxidant activity	12
2.2.1(c) Anti-cancer activity	12
2.3 Plant extraction methods	13
2.3.1 Maceration	13
2.3.2 Soxhlet extraction	15
2.3.3 Ultrasound extraction or sonication extraction	15

2.4	MicroRNA	16
2.4.1	MicroRNA biogenesis	17
2.4.2	MicroRNA and cancer	21
2.4.2(a)	Mechanisms involved in microRNA deregulation in cancer	22
2.4.2(b)	Pathways involved in microRNA regulation in cancer	23
2.4.3	MicroRNA and medicinal plants	24
2.4.4	Validation and Functional Analysis of microRNA	27
2.4.4(a)	Validation of miRNA	27
2.4.4(b)	Functional analysis of miRNA	30
2.5	Bioinformatics analysis of microRNA	31
2.6	Mass spectrometry	34
2.6.1	Ionization	34
2.6.2	Mass analyzer	36
2.6.3	Detection of ions	37
2.7	Mechanisms of cell death	39
2.7.1	Apoptosis	39
2.7.1(a)	Intrinsic Pathway	41
2.7.1(b)	Extrinsic pathway	42
2.7.2	Necrosis	43
CHAPTER 3: METHANOLIC <i>P. longifolia</i> LEAF EXTRACTION AND <i>IN VITRO</i> CYTOTOXICITY TESTING ON HELA CELL LINES		45
3.1	Introduction	45
3.1.1	Objectives	46
3.2	Materials and methods	47
3.2.1	Sample collection	47
3.2.2	Extraction of <i>P. longifolia</i> leaves	47
3.2.3	Cell Lines	48
3.2.4	Revitalisation of frozen cells	48
3.2.5	Subculturing of cells	47

3.2.6	Cell counting	50
3.2.7	<i>In vitro</i> cytotoxicity study of <i>P. longifolia</i> leaf extract	51
3.3	Results	53
3.3.1	Extraction of <i>P. longifolia</i> leaves	53
3.3.2	<i>In vitro</i> cytotoxicity study of <i>P. longifolia</i> leaf extract	55
3.4	Discussion	57
3.4.1	Extraction of <i>P. longifolia</i> leaves	57
3.4.2	<i>In vitro</i> cytotoxicity study of <i>P. longifolia</i> leaf extract	59
3.5	Conclusion	64
 CHAPTER 4: VALIDATION AND FUNCTIONAL ANALYSIS OF MIRNA-221-5P ALONGSIDE WITH <i>P. longifolia</i> LEAF EXTRACT TREATMENT IN HELA CELLS		 65
4.1	Introduction	65
4.1.1	Objectives	66
4.2	Materials and methods	67
4.2.1	Chemicals and reagents	67
4.2.2	Transfection of miRNA mimics and anti-miRNA	67
4.2.3	Quantification of miR-221-5p Expression	68
4.2.3(a)	Total RNA Isolation	68
4.2.3(b)	Reverse Transcription (RT)	69
4.2.3(c)	Real time Polymerase Chain Reaction (qPCR)	71
4.2.4	Functional analysis of miR-221-5p	73
4.2.4(a)	MTT cell viability assay	73
4.2.4(b)	Annexin V/PI assay	73
4.2.4(c)	Caspase-3 assay	75
4.2.4(d)	Bicinchoninate protein assay	76
4.2.5	Statistical analysis	76
4.3	Results	77
4.3.1	Total RNA Isolation	77
4.3.2	Validation and quantification of miRNA	81

4.3.3	MTT cell viability assay	84
4.3.4	Annexin V/Pi assay	88
4.3.5	Caspase 3 assay	91
4.4	Discussion	94
4.4.1	Transfection of miRNA mimics and anti-miRNA	94
4.4.2	Total RNA isolation	96
4.4.3	Taqman RTqPCR	99
4.4.4	MTT cell viability assay	104
4.4.5	Flow cytometric annexin V/PI analysis	107
4.4.6	Caspase-3 analysis	111
4.5	Conclusion	116
CHAPTER 5: <i>IN VITRO</i> ULTRA-MORPHOLOGICAL ASSESSMENT OF MICRORNA OVER-EXPRESSED HELA CELLS IN RELATION WITH <i>P. longifolia</i> LEAF EXTRACT TREATMENT		117
5.1	Introduction	117
5.1.1	Objective	118
5.2	Materials and methods	119
5.2.1	Cell culture and miRNA transfection	119
5.2.2	Scanning Electron Microscopy (SEM)	119
5.2.3	Transmission Electron Microscopy (TEM)	119
5.2.4	Acridine Orange / Propidium Iodide (AO/PI) Staining	120
5.3	Results	121
5.3.1	Scanning Electron Microscopy (SEM)	121
5.3.2	Transmission Electron Microscopy (TEM)	126
5.3.3	Acridine Orange / Propidium Iodide (AO/PI) Staining	129
5.4	Discussion	131
5.4.1	Scanning Electron Microscopy (SEM)	131
5.4.2	Transmission Electron Microscopy (TEM)	134
5.4.3	Acridine Orange / Propidium Iodide (AO/PI) Staining	137
5.5	Conclusion	140

CHAPTER 6: BIOINFORMATICS ANALYSIS OF MIR-221-5P	141
 BY PREDICTION OF GENES AND PROTEIN- PROTEIN INTERACTION NETWORKING FOR MIR-221-5P	
6.1 Introduction	141
6.1.1 Objectives	142
6.2 Materials and methods	143
6.2.1 Identification of mature sequences of miR-221-5p	143
6.2.2 microRNA-gene interaction analysis of miR-221-5p	143
6.2.3 Gene enrichment analysis	144
6.2.4 Protein-Protein Interaction Network (PPIN) analysis	144
6.3 Results	146
6.3.1 Identification of mature sequences of miR-221-5p	146
6.3.2 microRNA-gene interaction analysis of miR-221-5p	150
6.3.3 Gene enrichment analysis	154
6.3.4 Protein-Protein Interaction Network (PPIN) analysis	161
6.4 Discussion	163
6.4.1 microRNA-gene interaction analysis of miR-221-5p	163
6.4.2 Gene enrichment analysis	170
6.4.3 Protein-Protein Interaction Network (PPIN) Analysis	176
6.5 Conclusion	180
CHAPTER 7: PROTEOMICS ANALYSIS OF MIR-221-5P	181
 TARGET PROTEINS ALONGSIDE WITH <i>P.</i> <i>longifolia</i> LEAF EXTRACT TREATMENT IN HELA CELLS	
7.1 Introduction	181
7.1.1 Objectives	182
7.2 Materials and Methods	183
7.2.1 Sample preparation	183
7.2.2 Extraction of total protein using RIPA lysis buffer	183
7.2.3 Acetone precipitation and buffer exchange	183
7.2.4 Determination of protein concentration	184

7.2.5	Buffer preparation and SDS-PAGE	184
7.2.5(a)	10% SDS	184
7.2.5(b)	Resolving buffer (1.5 M Tris-Cl, pH 8.8)	184
7.2.5(c)	Stacking buffer, (0.5 M Tris-Cl, pH 6.8)	185
7.2.5(d)	Ammonium persulfate (APS), 10% (w/v)	185
7.2.5(e)	Loading buffer, 5×	185
7.2.5(f)	SDS-PAGE running buffer	185
7.2.5(g)	Coomassie Brilliant Blue, CBB R250, 0.05% (w/v)	186
7.2.5(h)	Separating gel and stacking gel	186
7.2.5(i)	SDS-PAGE	186
7.2.6	Buffer preparation and In-solution digestion	187
7.2.6(a)	RapiGest	187
7.2.6(b)	100 mM Dithiothreitol (DTT)	187
7.2.6(c)	200 mM Iodoacetamide (IAA)	187
7.2.6(d)	1 µg/µL Trypsin	188
7.2.6(e)	Trypsin Digestion	188
7.2.7	LC-ESI-MS/MS	189
7.2.8	Functional annotation of identified proteins	190
7.3	Results	191
7.3.1	Determination of protein concentration	191
7.3.2	SDS-PAGE	194
7.3.3	Protein identification by LC-ESI-MS/MS	196
7.3.4	Functional annotation of proteins	202
7.4	Discussion	208
7.4.1	Isolation of total protein	208
7.4.2	Protein identification by LC-ESI-MS/MS	211
7.4.3	Functional annotation of proteins identified by LC- ESI-MS/MS	216
7.5	Conclusion	220

CHAPTER 8: GENERAL CONCLUSION AND FUTURE RECOMMENDATIONS	221
8.1 General conclusion	221
8.2 Suggestions for future studies	221
REFERENCES	225
APPENDICES	
LIST OF PUBLICATIONS	

LIST OF TABLES

		Page
Table 3.1	Origin and the source of the cell line used with its complete growth medium requirements	49
Table 4.1	RT Master mix components	70
Table 4.2	Thermocycler set up parameter	72
Table 4.3	PCR master mix components	72
Table 4.4	Applied Biosystems 7900HT Fast Real-Time PCR System set up parameter	72
Table 4.5	Quantification of total RNA	79
Table 6.1	Mature sequences of miR-221-5p	147
Table 6.2	Representative target genes predicted by miRGate bioinformatics tool	153
Table 6.3	GO terms for biological process adapted from DAVID bioinformatics	155
Table 6.4	GO terms for molecular function adapted from DAVID bioinformatics	157
Table 6.5	GO terms for cellular component adapted from DAVID bioinformatics	159
Table 7.1	Examples of proteins identified from untreated HeLa cells by LC-ESI-MS/MS	198
Table 7.2	Examples of proteins identified from <i>P. longifolia</i> leaf extract treated HeLa cells by LC-ESI-MS/MS	199
Table 7.3	Examples of proteins identified from miR-221-5p mimics transfected <i>P. longifolia</i> leaf extract treated HeLa cells by LC-ESI-MS/MS	200
Table 7.4	Examples of proteins identified from anti-miR-221-5p transfected <i>P. longifolia</i> leaf extract treated HeLa cells by LC-ESI-MS/MS	201

LIST OF PLATES

		Page
Plate 3.1	<i>P. longifolia</i> methanolic leaf extraction process	54
Plate 4.1	The absorbance spectrums of RNA	78
Plate 4.2	Agarose gel electrophoresis of RNA extracted	80
Plate 4.3	The 24-well plate used in MTT assay	85
Plate 4.4	Flow cytometric analysis of HeLa cells by Annexin V/PI double staining	89
Plate 5.1	SEM micrographs of untreated HeLa cells.	122
Plate 5.2	SEM micrographs of <i>P. longifolia</i> leaf extract treated HeLa cells	123
Plate 5.3	SEM micrographs of <i>P. longifolia</i> leaf extract treated miR-221-5p mimic transfected HeLa cells	124
Plate 5.4	SEM micrographs of <i>P. longifolia</i> leaf extract treated miR-484 mimic transfected HeLa cells	125
Plate 5.5	TEM micrograph of untreated HeLa cell	127
Plate 5.6	TEM micrograph of <i>P. longifolia</i> leaf extract treated HeLa cell	127
Plate 5.7	TEM micrograph of miR-221-5p mimic transfected <i>P. longifolia</i> leaf extract treated HeLa cell	128
Plate 5.8	TEM micrograph of miR-484 mimic transfected <i>P. longifolia</i> leaf extract treated HeLa cell	128
Plate 5.9	Photomicrograph of AO and PI stained HeLa cells.	130
Plate 7.1	The 96-well micro plate used in BSA assay	192
Plate 7.2	Protein profiles of total protein extracted from HeLa cells by SDS-PAGE	195

LIST OF FIGURES

	Page
Figure 1.1 Workflow of the overall research	6
Figure 2.1 Different types of plant extraction methods	14
Figure 2.2 Overview of miRNA biogenesis process	20
Figure 2.3 The mechanism of Taqman real time qPCR	29
Figure 2.4 Pipeline of bioinformatics analysis	33
Figure 2.5 Components involved in mass spectrometry	38
Figure 2.6 The intrinsic and extrinsic pathway of apoptosis	40
Figure 2.7 Types of cell death	44
Figure 3.1 Cell viability of HeLa cells treated with <i>P. longifolia</i> leaf extract by using MTT assay	56
Figure 4.1 Amplification plot from real time RTqPCR	82
Figure 4.2 Fold change expression from real time RTqPCR	83
Figure 4.3 Cell viability of HeLa cells by using MTT assay	86
Figure 4.4 Percentage of HeLa cells in viable and apoptosis form based on flow cytometric analysis by Annexin V/PI double staining.	90
Figure 4.5 Typical standard curve from the Bicinchoninate Protein assay	92
Figure 4.6 Caspase-3 enzyme activity in HeLa cells treated with <i>P. longifolia</i> leaf extract for 24 hours	93
Figure 6.1 Location of miR-221 in Chromosome X	148
Figure 6.2 Conserved secondary structure of miR-221	149
Figure 6.3 Protein coding genes predicted by different computational algorithms	151

Figure 6.4	Summary of target genes for further analysis	152
Figure 6.5	Gene ontology bar chart adapted from Enrichr bioinformatics represented based on combinatory scores	160
Figure 6.6	Protein-protein interaction networking generated by STRING bioinformatics tool	162
Figure 7.1	A standard curve of protein concentration using Bicinchoninate Protein assay.	193
Figure 7.2	The Venn diagram represents the number of proteins identified by LC-ESI-MS/MS from total proteins extracted from HeLa cells	197
Figure 7.3	PANTHER generated protein classification obtained for the proteins identified from the (A) Untreated HeLa cells and (B) <i>P. longifolia</i> leaf extract treated HeLa cells.	204
Figure 7.4	PANTHER generated protein classification obtained for the proteins identified from the (A) miR-221-5p mimic transfected <i>P. longifolia</i> leaf extract treated HeLa cells and (B) anti-miR-221-5p transfected <i>P. longifolia</i> leaf extract treated HeLa cells	205
Figure 7.5	PANTHER generated gene ontology for biological process obtained for the proteins identified from the (A) Untreated HeLa cells and (B) <i>P. longifolia</i> leaf extract treated HeLa cells.	206
Figure 7.6	PANTHER generated gene ontology for biological process obtained for the proteins identified from the (A) miR-221-5p mimic transfected <i>P. longifolia</i> leaf extract treated HeLa cells and (B) anti-miR-221-5p transfected <i>P. longifolia</i> leaf extract treated HeLa cells	207

LIST OF ABBREVIATIONS

$\Delta\psi_m$	Mitochondrial membrane potential
$\mu\text{g/mL}$	Microgram per milliliter
μM	Micro molar
2D	Two-dimensional
Å	Angstrom
ADCs	Analog-to-digital converters
AIF	Apoptosis inducing factor
ANOVA	Analysis of variance
AO/PI	Acridine orange and propidium iodide
Apaf-1	Apoptotic protease activating factor-1
APCI	Atmospheric-pressure chemical ionization
APL	Acute promyelocytic leukemia
<i>APLP2</i>	Amyloid beta (A4) precursor-likeprotein 2
AP-MS	Affinity purification–mass spectrometry
APS	Ammonium persulfate
AVD	Apoptotic volume decrease
BCA	Bicinchoninate assay
Bcl	B-cell lymphoma
BIND	Biomolecular interaction network database
BMF	B-cell lymphoma 2 modifying factor
bp	Base pair
BSA	Bovine serum albumin
<i>C. elegans</i>	<i>Caenorhabditis elegans</i>
CAD	Caspase-Activated dnase
CARD	Caspase-recruitment domain
CBB	Coomassie brilliant blue
CCDB	Cervical cancer gene database
<i>CDH15</i>	Cadherin-15
CDK	Cyclin-dependent kinase
CDKN1B	Cyclin-dependent kinase inhibitor 1B
cDNA	Complementary DNA

<i>CEACAM1</i>	Carcinoembryonic antigen-related cell adhesion molecule 1
CID	Collision-induced dissociation
cm	Centimeter
<i>CPEB1</i>	Cytoplasmic polyadenylation element binding protein 1
Ct	Cycle threshold
dATP	Deoxyadenosine triphosphate
DAVID	Database for Annotation, Visualization and Integrated Discovery
DGCR8	<i>Di George Syndrome critical region gene 8</i>
DISC	Death inducing signalling complex
DMEM	Dubelcco's minimum essential medium
DMSO	Dimethyl sulfoxide
DNA	Deoxyribonucleic acid
DPPH	2,2-diphenyl-1-picrylhydrazyl
dsDNA	Double-stranded deoxyribonucleic acid
<i>DSG1</i>	Desmoglein 1
DTT	Dithiothreitol
EBV	Epstein-Barr virus
ECD	Electron capture dissociation
ERK	Extracellular signal-regulated kinase
ESI	Electrospray ionization
ETD	Electron-transferdissociation
FAF1	Fas-associated factor 1
FBS	Fetal bovine serum
FDA	Food and Drug Administration
FDR	False discovery rate
FRET	Forster resonance energy transfer
GO	Gene ontology
H ₂ O ₂	Hydrogen peroxide
HCl	Hydrochloric acid
HPRD	Human protein reference database
HPV	Human papilloma virus
IAA	Iodoacetamide
IC ₅₀	Half maximal inhibitory concentration

IEF	Isoelectric focusing
<i>IGFBP5</i>	Insulin-like growth factor binding proteins 5
<i>IL18</i>	Interleukin 18
<i>ITGAV</i>	Integrin alpha-V
<i>ITGB1</i>	Integrin, beta 1
JNK	Jun N-terminal kinases
K ⁺	Potassium
kHz	Kilo hertz
KISS	Kinase substrate sensor
LC-ESI-MS/MS	Liquid chromatography-electrospray ionization-mass spectrometry/ mass spectrometry
LDH	Lactate dehydrogenase
m/z	Mass to charge ratio
<i>MAL</i>	Myelin and lymphocyte protein
MALDI	Matrix-assisted laser desorption/ionization
MAPPIT	Mammalian protein-protein interaction trap
mg/g	Milligram per gram
mg/kg	Milligram per kilogram
mg/mL	Milligram per milliliter
MGB	Minor groove binder
min	Minutes
MINT	Molecular interaction database
miRISC	Mirna-Induced Silencing Complex
miRNA	Microna
MOMP	Mitochondrial outer membrane permeabilization
mRNA	Messenger RNA
<i>MSN</i>	Moesin
MTT	3-(4, 5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide
NCBI	National Center for Biotechnology Information
NCI	National cancer institute
NFQ	Nonfluorescent quencher
nm	Nano meter
nM	Nano molar

°C	Degree celsius
OD	Optical density
<i>P. longifolia</i>	<i>Polyalthia longifolia</i>
PANTHER	Protein Analysis through Evolutionary Relationships
PAZ	Piwi, Arganoate and Zwillie
PBS	Phosphate-buffered saline
PCD	Programmed cell death
PIK3R1	Phosphoinositide-3-kinase regulatory subunit 1
PLA	Proximity ligation assay
Plk1	Polo-like kinase 1
<i>p</i> NA	<i>P</i> nitroanilide
PPIN	Protein-protein interaction networking
pre-miRNA	Precursor mirna
pri-miRNA	Primary mirna
PS	Phospholipid phosphatidylserine
PTEN	Phosphatase and tensin
PTP	Permeability transition pore
PTPC	Permeability transition pore complex
PUMA	P53 upregulated modulator of apoptosis
Q-TOF	Quadrupole Time-of-Flight
<i>RARG</i>	Retinoic acid receptor gamma
RCF	Relative centrifugal force
RIA	Radioimmunoassay
RIPA	Radioimmunoprecipitation assay
RNA	Ribonucleic acid
ROS	Reactive oxygen species
RP	Ribosomal protein
RPM	Revolutions per minute
rRNA	Ribosomal RNA
RT	Reverse transcription
RT	Room temperature
RT-qPCR	Real time quantitative polymerase chain reaction
RVI	Regulatory volume increase

SD	Standard deviation
SDG	Sustainable development goals
SDS	Sodium dodecyl sulfate
SEM	Scanning electron microscopy
SID	Surface-induced dissociation
siRNA	Small interfering RNA
snoRNA	Small nucleolar RNA
TEM	Transmission electron microscopy
<i>THBS2</i>	Thrombospondin-2
TIMP3	Tissue inhibitor of metalloproteinase-3
T _m	Melting temperature
TNF	Tumor necrosis factor
TOF	Time-of-flight
TRAF-6	Tnf receptor-associated factor 6
TRAIL	TNF-related apoptosis-inducing ligand
tRNA	Transfer RNA
UTR	Untranslated region
UV	Ultraviolet
XIAP	X-linked inhibitor of apoptosis
$\Delta\Delta CT$	Double delta cycle threshold

**PENCIRIAN MOLEKUL DAN ANALISIS BIOINFORMATIK MICRORNA-
221-5P YANG DIKAWALATUR OLEH EKSTRAK PIAWAI
DAUN *Polyalthia longifolia* (Sonn.) Thwaites DI DALAM SEL HELA**

ABSTRAK

Polyalthia longifolia (Sonn.) Thwaites adalah sejenis spesies tumbuhan yang kaya dengan nilai-nilai perubatan. Penyelidikan saintifik terkini ekstrak daripada daun *P. longifolia* telah mendedahkan sifat anti kansernya terhadap sel HeLa melalui pengaruh apoptosis yang berdasarkan kaspase dengan mengawal ekspresi miRNA. Walau bagaimanapun, belum ada kajian dilaksanakan untuk melaporkan analisis fungsi mikroRNA terkawal-atur dan tiada bukti saintifik pengesahan pengawalan ekspresi mikroRNA dalam sel HeLa yang dirawat dengan ekstrak daun methanolic *P. longifolia*. Oleh itu, kajian ini dijalankan untuk mengesahkan ekspresi miRNA dalam sel HeLa yang dirawat dengan ekstrak daun *P. longifolia* berbanding dengan sel-sel HeLa kawalan yang tidak dirawat. Kajian ini telah memberi butiran yang khusus tentang analisis fungsian dan analisis proteomik terhadap ekspresi miRNA. Dalam kajian ini, ekstrak daun *P. longifolia* yang segar telah disediakan dan ujian MTT telah dilakukan untuk mengenal pasti nilai IC₅₀ terhadap sel HeLa. Ekspresi miR-221-5p disahkan dengan menjalankan RTqPCR masanya TaqMan yang mengesahkan regulasi penurunan ekspresi miR-221-5p dalam sel HeLa yang telah dirawat dengan ekstrak daun *P. longifolia* berbanding dengan sel yang tidak dirawat. Analisis fungsi miR-221-5p dilakukan dengan pendekatan mengaktifkan dan menyahaktifkan fungsi miRNA tersebut bersama-sama dengan menggunakan ujian sel pertumbuhan MTT, ujian pewarna Annexin-V FITC dan propidium iodida dan diukur melalui aliran sitometri dan ujian kaspase-3. Selain itu, perubahan

morfologi struktur ultra sel HeLa telah dikaji melalui mikroskop elektron transmisi (TEM), imbasan (SEM) dan mikroskop pendarfluor bagi sel yang diwarana dengan pewarna akridin dan propidium iodida (AO/PI). Analisis proteomik tentang protein yang dikawalatur oleh miR-221-5p juga dilakukan dengan dengan penggunaan LC-ESI-MS/MS. Di samping itu, analisis bioinformatik telah dilakukan untuk mengenal pasti sasaran mRNA untuk miR-221-5p melalui alat bioinformatik miRGate dan gen ontology telah dikaji dengan penggabungan DAVID dan Enrichr serta rangkaian interaksi protein-protein dianalisis dengan menggunakan alat bioinformatik STRING. Penurunan ekspresi miR-221-5p telah disahkan dengan jayanya dalam sel HeLa yang dirawat dengan ekstrak daun *P. longifolia*. Selain itu, mekanisme molekul komprehensif miR-221-5p yang dikaji melalui analisis fungsi dan proteomik jelas menunjukkan peranan miR-221-5p dalam induksi apoptotik intrinsik yang bersandarkan kaspase oleh rawatan ekstrak daun *P. longifolia*. Memandangkan pengekspresan mikroRNA mengambil bahagian dalam pengawalan/ patogenesis kanser dan kecekapannya dalam terapeutik gen semasa yang menasarkan kanser sebagai pendekatan rawatan klinikal yang boleh diterima, maka ekstrak daun *P. longifolia* boleh menjadi calon ubat antikanser baru yang baik.

MOLECULAR CHARACTERIZATION AND BIOINFORMATICS
ANALYSIS OF MICRORNA-221-5P REGULATED BY STANDARDIZED
***Polyalthia longifolia* (Sonn.) Thwaites LEAF EXTRACT IN HELA CELL**
LINES

ABSTRACT

Polyalthia longifolia (Sonn.) Thwaites is an exquisite plant species with rich ethnomedicinal values. Recent scientific investigations on *P. longifolia* leaf extract have also revealed its anti-cancer property against HeLa cells through the induction of caspase-dependent apoptosis by regulating microRNA (miRNA) expressions. However, there were no further investigations performed to report the functional analysis of the regulated miRNA, with absolutely no means of scientific evidence of validation of miRNA dysregulation in HeLa cells treated with the methanolic *P. longifolia* leaf extract. Hence, this study was conducted to validate the miRNA expression in methanolic *P. longifolia* leaf extract treated HeLa cells in comparison with untreated HeLa cells with an intricate elucidation of functional and proteomic analysis of miRNA expression. In this study, methanolic *P. longifolia* leaf extract was freshly prepared and MTT assay was performed to identify the IC₅₀ value against HeLa cells. The expression of miR-221-5p was validated by performing Taqman real time RTqPCR which confirmed the down-regulation of miR-221-5p in HeLa cells treated with methanolic *P. longifolia* leaf extract compared to the untreated HeLa cells. The functional analysis of miR-221-5p was conducted through gain-of-function and loss-of-function approach by MTT assay, flow cytometric analysis of Annexin V/ Propidium Iodide assay and caspase-3 assay. Besides, the ultra-structural morphological changes in HeLa cells were investigated through

scanning electron microscopy (SEM), transmission electron microscopy (TEM) and fluorescence microscopy for cells stained with acridine orange and propidium iodide (AO/PI). Proteomic analysis of proteins that regulated by miR-221-5p was carried out by LC-ESI-MS/MS. In addition, bioinformatics analysis was performed to identify the mRNA targets of miR-221-5p by miRGate bioinformatics tool and the gene ontology of the predicted genes was determined by DAVID and Enrichr bioinformatics tool while the protein-protein interaction network was analysed using STRING bioinformatics tool. Conclusively, based on the data obtained from this overall study, expression of miR-221-5p has been successfully validated upon *P. longifolia* leaf extract treatment in HeLa cells with its comprehensive molecular mechanism involved through functional and proteomic analysis which clearly indicated the role of miR-221-5p in the induction of caspase-dependent intrinsic apoptotic pathway of cell death by *P. longifolia* leaf extract treatment. Considering the participation of miRNA expression in cancer regulation/ pathogenesis and its efficiency in current gene therapeutic target for cancer as a clinically admissible treatment approach, *P. longifolia* leaf extract could be a promising novel anticancer drug candidate.

CHAPTER 1: INTRODUCTION

1.1 Overview and rationale of study

Cells as the “building blocks of life” are well organized to balance its proliferation and death to ensure the proper development of healthy tissue and to get rid of the damaged cells, respectively (Mason and Rathmell, 2011). The coordination of the cell proliferation as well as the cell death mechanisms is highly important because an unbalanced process may lead to deadly diseases such as cancer. Cancer occurs when cell growth exceeds cell death due to some alteration in the cell causing them to multiply out of control and invade to other parts of the body. Cancer which is also known as malignancy or malignant neoplasm has been accounted to be one of the leading causes of mortality worldwide. Based on the recently published paper on cancer facts and figures 2019, more than 1.7 million new cancer cases are expected to be diagnosed and about 606, 880 deaths is expected to be recorded in United States of America due to cancer in 2019 (American Cancer Society, 2019).

Cancer can be categorized based on the types of cells it occurs such as the carcinomas (epithelial cells), sarcoma (bone and soft tissues), leukemia (blood cells and bone marrow) and lymphoma (lymphocytes). Among the 100 over types of cancer, cervical cancer has attained an alarming health concern among women globally which needs a compelling prevention. Cervical cancer is the third most prevailing gynecologic cancer in USA which peaks in elderly women, usually in their 50s (Dawkins *et al.*, 2018). In fact, 569, 847 incidences of cervical cancer and 311, 365 deaths due to cervical cancer have been reported by world region (Bray *et al.*, 2018). Recent study estimated fourteen million new cancer cases to occur by

2035 which may consequently lead to challenging preventive medicine structure (Pilleron *et al.*, 2019).

Although there are few general treatments available for cervical cancer like radiotherapy, immunotherapy, chemotherapy and surgery; chemoprevention of cancer through the regulation of miRNA with the utilization of medicinal plants is presently being advanced in cancer therapy field. Since dysregulation of genes involved in the biological processes have been convincingly demonstrated to be associated with cancer, miRNA therapeutic approach is highly trustworthy in cancer treatments (Ji *et al.*, 2017). MiRNAs are small, single stranded, non-coding RNA molecules of 20-22 nucleotides that control the expression of target genes by imperfect pairing to multiple mRNA targets. The incomplete base pairing of miRNAs to their mRNA targets at 3' UTRs causes the degradation of mRNA targets and subsequent down-regulation of gene expression and protein inhibition. The participation of miRNAs in the regulation of gene expression at post transcriptional level and subsequent protein translational repression (Bartel, 2004) clearly substantiate the major role of miRNA in diverse biological processes such as cell death (Xu *et al.*, 2004) and cell proliferation (Cheng *et al.*, 2005). Interestingly, various plants rich in anti-cancer bioactive compounds have been shown to regulate the mammalian miRNA levels (Xie *et al.*, 2016; Gezici and Sekeroglu, 2017).

1.2 *Polyalthia longifolia* as an important medicinal plant

Polyalthia longifolia var. *Angustifolia* Thw. (Annonaceae) is one of the most important medicinal plants which is found throughout Malaysia and widely used in traditional medicine as febrifuge and tonic (The Wealth of India, 1969). The local name of *P. longifolia* is Glodogan tiang. *Polyalthia longifolia* is a small medium-

sized tree with linear-lanceolate leaves, 1 to 1.5 cm broad, occurring in Sri Lanka and now grown in tropical parts of India along road sides and in gardens for their beautiful appearance (Bose *et al.*, 1998). The diterpenes, alkaloids, steroid and miscellaneous lactones were isolated from its bark (The Wealth of India, 1969). The stem bark extracts and isolated compounds were studied for various biological activities like anti-bacterial, cytotoxicity and anti-fungal activity (Goutham *et al.*, 2010). Study conducted by Jothy *et al.* (2012) reported the anti-oxidant activity and hepatoprotective potential of *P. longifolia*. The results of this study revealed that *P. longifolia* leaf extract could protect the liver against paracetamol-induced oxidative damage. They also reported that the observed hepatoprotective activity of *P. longifolia* in their study might be due to its antioxidant activity, resulting from the presence of phenolic compounds in the extracts. Jothy *et al.* (2013) also tested the genotoxic potential of *P. longifolia* leaf against H₂O₂-radical-mediated DNA damage by using plasmid relation, comet, and *Allium cepa* assay. The results of the *in vitro* tests in their study demonstrated that *P. longifolia* leaf was devoid of a significant genotoxic effect under experimental conditions. Their *Allium cepa* assay results showed that, applied in lower concentrations, the methanol extract of *P. longifolia* leaf could be important for maintaining the genetic stability of the organism. An acute oral toxicity study revealed that *P. longifolia* leaf extract was safe after oral administration as a single dose to female albino Wistar rats with up to 5000 mg/kg body weight (Jothy *et al.*, 2013).

The induction of apoptosis by *P. longifolia* treatment has been revealed to cause cell cycle arrest at sub G₀/G₁, G₀/G₁ and G₂/M phases as well as to increase the mitochondria membrane potential depolarization (Vijayarathna *et al.*, 2017a). The proteomic profiling array conducted by Vijayarathna *et al.* (2017a) has also

demonstrated an up-regulation of pro-apoptotic proteins and down-regulation of anti-apoptotic proteins in HeLa cells upon *P. longifolia* treatment. Besides, morphological study through various microscopic approaches also evidently showed the induction of apoptosis in HeLa cells by *P. longifolia* leaf extract treatment based on typical apoptotic cell morphology observed in *P. longifolia* treated cells (Vijayarathna *et al.*, 2017b). The most recent study showed that *P. longifolia* leaf extract induced apoptosis in HeLa cells through the regulation of miRNAs (Vijayarathna *et al.*, 2017c; Vijayarathna, 2017). However, there are no further reports on the relationship between the cytotoxicity of *P. longifolia* leaf extract with the detailed mechanism of apoptosis through miRNA deregulation in HeLa cell lines. There is absolutely no further validation and functional analysis of the deregulated miRNAs which has been previously reported. Hence, the current study was undertaken to validate the role of dysregulated miRNA in inducing the apoptotic cell death in HeLa cells and to identify their targets proteins.

The miRNA deregulation, validation and functional analysis in HeLa cell treated with standardized *P. longifolia* leaf extract was conducted by transfection of miRNA mimic and anti-miR into HeLa cells to over-express and silence the miRNA expression, respectively. Validation of miRNA-221-5p in HeLa cells upon *P. longifolia* treatment was carried out through taqman real time RT-qPCR. Functional analysis of miR-221-5p in relation to *P. longifolia* leaf extract treatment in HeLa cells was carried out through MTT, annexin V/PI and caspase 3 assays. Preliminary study of morphology of HeLa cells upon transfection of miRNA mimic and treatment with methanolic *P. longifolia* leaf extract was accomplished using scanning electron microscopy (SEM), transmission electron microscopy (TEM) and dual-fluorescence nuclear staining method using acridine orange (AO) and

propidium iodide (PI) dyes. The detailed proteomics analysis of miR-221-5p in relation to the *P. longifolia* treatment in HeLa cells was implemented through LC-ESI-MS/MS mass spectrometry systems. Further bioinformatics analysis of miR-221-5p was accomplished with the aid of various bioinformatics tools such as mirgate (<http://mirgate.bioinfo.cnio.es>), The Database for Annotation, Visualization and Integrated Discovery (DAVID) v6.8 (<https://david.ncifcrf.gov/>) and a web-based PPI prediction tool called the STRING resource v10.5 (<http://string-db.org/>). Figure 1.1 shows the workflow of the overall research.

1.3 Objectives

The study was conducted with the objectives:

1. To validate the expression of miR-221-5p in HeLa cells treated with methanolic *P. longifolia* leaf extract treatment in comparison with untreated control HeLa cells.
2. To study the function of miR-221-5p in inducing the apoptotic cell death in HeLa cells treated with standardized *P. longifolia* leaf extract by using miRNA mimic and anti-miR transfection to enhance and knock down the miRNA expression, respectively.
3. To study the morphological changes in miR-221-5p and miR-484 regulated HeLa cells, alongside with *P. longifolia* leaf extract treatment.
4. To investigate the role of miRNAs and their targets in HeLa cells to induce apoptotic cell death after treatment with standardized *P. longifolia* leaf extract using bioinformatics approaches.
5. To identify the target apoptotic related proteins of miR-221-5p in HeLa cells treated with standardized *P. longifolia* leaf extract through proteomic approach.

1.4 Workflow




Figure 1.1: Workflow of the overall research

CHAPTER 2: LITERATURE REVIEW

2.1 Cancer

Genetically programmed cell division and differentiation occur in the process of formation of specific tissues and eventually functional organs. However, intermittently the uncontrollable cell division may give rise to tissue masses called tumors, or neoplasms. A single mass of benign tumor is usually not life threatening since it can be cured completely by surgical removal. However, when the cells of a tumor start to invade and interrupt the surrounding tissues, the tumor is said to be malignant and is identified as cancer which can consequently lead to death due to injury to vital organs, secondary infection, metabolic problems, secondary malignancies, or hemorrhage. The place where cancer begins is known as the original or primary site. A malignant tumor can break away from its original location and invade far-away sites through the lymphatic system, forming new tumors. This process is known as metastasis. The uncontrollable growth of cells may occur in any parts of the body leading to more than 100 types of cancer including lung cancer, breast cancer, cervical cancer, stomach cancer, prostate cancer, bowel cancer and ovarian cancer.

According to the American Cancer Society (2019), risk factors for cancer include genetic factors such as inherited genetic mutations and immune conditions as well as the lifestyle of a person such as tobacco use, alcohol use, diet, and physical activity. Other disposing factors to cancer are certain type of infections such as human papilloma virus (HPV), Epstein-Barr virus (EBV), hepatitis B, hepatitis C and *Helicobacter pylori*. Environmental exposures to diverse range of chemicals, radiations and even overexposure to ultraviolet (UV) light from the sun may also

lead to cancer. Smoking and alcohol intake can be associated with several cancers such as the mouth, oral cavity, pharynx, larynx, esophagus, lung, stomach, pancreas and even colon (Schmidt and Popham, 1981). Besides, viral infection can be related to cancer because of their capability to integrate into the DNA of the human stem cell where it mutates and transforms the cell to be the parent of the malignant clone (Doll and Peto, 1981).

2.1.1 Current development in cancer therapy

There are several types of treatments available for cancer including surgery, radiation therapy, chemotherapy, immunotherapy, gene therapy, hyperthermia, and stem cell transplant. However, these treatments have excruciating side effects that vary from person to person depending on the frequency of treatment, the age of the person and other health conditions. Commonly occurring side effects generated by cancer treatments include anemia, alopecia (hair loss), constipation, edema, fatigue, memory problems, peripheral neuropathy, nausea and vomiting (National Cancer Institute, 2018; Wilkes, 2018). Chemotherapy is one of the popular cancer treatments from the 1960s as the degree of curing cancer elevated at approximately 33% through radical local treatments. Eventually Cancer Chemotherapy National Service Centre was established in the effort of developing methods to screen chemicals using transplantable tumors in rodents (Devita and Chu, 2008).

The evolving knowledge on cancer mechanisms has expedited the expansion of novel anticancer approaches. One of the most extensive conventions is to slow down or to inhibit the prime characteristic of cancer cells that grow uncontrollably. This can be correlated with the elevation of tendency of the cells to go through the process of cell suicide, or apoptosis. This effective route is eventually achieved

through a mechanistic manner where the cytotoxic drugs are designed so as to impede the DNA replication by damaging the DNA of the cancer cells, subsequently inducing apoptosis.

Crude extracts from plant samples have been established to be selectively toxic to cancer cells after passing through various bioassays including *in vitro* and *in vivo* screenings which led to the plant collection program by the United States National Cancer Institute (NCI), followed by screening of plant species for anticancer activity which resulted in a revelation of enormous number of new anticancer agents such as taxanes and camptothecin (Cassady and Douros, 1980; Shoeb, 2006). Plants rich in phytochemicals such as berberine, curcumin, genistein, daidzein, glyceollin, apigenin, quercetin, baicalein, resveratrol, luteolin, matrine, garcinol, silibinin, mangiferin, doxorubicin, and paclitaxel have been evidently demonstrated to exhibit anti-cancer property by participating in biological processes including cell differentiation, proliferation and apoptosis (Sala-Cirtog *et al.*, 2015; Biersack, 2016; Devi *et al.*, 2017). The recent evolution of miRNA investigations and its involvement in regulating biological processes such as cell proliferation, and migration, metastasis, apoptosis, and cell differentiation in numerous malignancies by partial complementary binding to mRNA targets (Sethi *et al.*, 2013; Thakur *et al.*, 2014). This has led to a viewpoint concerning to associate phytochemical-rich medicinal plants to regulate the expression of miRNAs involved in tumorigenesis. Regulation of miRNA expression by medicinal plant extracts has become a promising novel strategy for cancer treatment, exclusively through inducing apoptosis and inhibiting cell proliferation in cancer cells (Gezici and Sekeroglu, 2017).

2.2 *Polyalthia longifolia*

Polyalthia longifolia from Annonaceae family is well known for its sophisticated traditional medicinal values. *Polyalthia longifolia* (Sonn.) Thwaites commonly known as False Ashoka, Buddha Tree, Green champa, Indian mast tree, and Indian Fir tree while its synonyms include *Uvaria longifolia* (Sonn.), *Guatteria longifolia* (Sonn.) Wallich, *Unona longifolia* (Sonn.) (Jothy *et al.*, 2013). *Polyalthia longifolia* indigenously belongs to Sri Lanka and cultivated in Pakistan which is now widely found throughout Malaysia. The classification of *P. longifolia* is as follows:

Kingdom : Plantae
Division : Magnoliophyta
Class : Magnoliopsida
Subclass : Magnoliidae
Order : Magnoliids
Family : Annonaceae
Tribe : Annoneae
Genus : *Polyalthia*
Species : *longifolia*

2.2.1 Botanical description of *P. longifolia*

The evergreen pyramid-like *P. longifolia* trees are known to grow tall upto an altitude of 15 to 20 m. Its dark green glossy leaves are seen to be long and narrow approximately accounting to 7.5-23 by 1.5-3.8 cm (Katkar *et al.*, 2010). The ovate-oblong leaves with wavy margins possess mild fragrance while the mildly pale green flowers with wavy petals are non-fragrant (Lemmens and Bunyapraphatsara, 2003). The oval-shaped ripe fruits are 1.8 to 2 cm long bearing a single smooth and shiny

seed. The *P. longifolia* trees generally bear flowers and fruits during the month of February till June (Wallis, 1985; Yadav and Sardesai, 2000).

2.2.2 Pharmacological activities of *P. longifolia*

The pharmacological properties of *P. longifolia* include anti-microbial, anti-oxidant, anti-cancer, anti-proliferative, radioprotective, anti-inflammatory, and anti-ulcer (Dixit *et al.*, 2014).

2.2.2(a) Anti-microbial activity

Polyalthia longifolia leaves extracted using different solvents have been comparatively tested for anti-microbial activity against microbes such as *Bacillus subtilis*, *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Proteus vulgaris* and *Salmonella typhi* through disc-diffusion method which evidently demonstrated the potential anti-bacterial activity of *P. longifolia* leaf extract (Thenmozhi and Sivaraj, 2010). In another study, methanolic leaf extract of *P. longifolia* was shown to exhibit anti-bacterial activity against *Bacillus subtilis* and *Staphylococcus aureus* reporting diameter of inhibition zones 24 mm and 22.6 mm, respectively (Uzama *et al.*, 2011). In addition, various bioactive components diterpenoids, alkaloids and allantoin isolated from the different parts of *P. longifolia* have also reported to exert significant anti-bacterial and anti-fungal activities against *Aspergillus fumigatus*, *Saccharomyces caulbequence*, *Saccharomyces cerevaceae*, *Candida albicans*, and *Hensila californica* (Rashid *et al.*, 1996; Faizi *et al.*, 2003; Murthy *et al.*, 2005).

2.2.2(b) Anti-oxidant activity

In vitro anti-oxidant activity of ethyl acetate and methanolic *P. longifolia* seed extracts by DPPH, nitric oxide radical, hydroxyl radical, and superoxide radical considering ascorbic acid as standard have been successfully demonstrated by Thonangi and Akula (2018) which was further correlated with total phenolic content assay reporting 114 ± 1.7 and 146.5 ± 2.4 mg/g gallic acid equivalents respectively. Another study also reported the anti-oxidant potential of *P. longifolia* stem bark ethanol extract by analysing the DPPH radicals scavenging effect, ferric ions reduction effect and lipid peroxidation inhibition with IC₅₀ values of 18.14, 155.41 and 73.33 µg/mL, respectively (Manjula *et al.*, 2010). *In vitro* antioxidant activity of methanolic *P. longifolia* leaf extract was shown by inhibition of DPPH radical at IC₅₀ value of 2.721 ± 0.116 mg/mL (Jothy *et al.*, 2012).

2.2.2(c) Anti-cancer activity

Ethanollic *Polyalthia longifolia* leaf extract was evidently shown to exhibit anti-cancer activity by inhibiting cell viability of several human cancers including colon cancer, liver cancer, prostate cancer, neuroblastoma, with a highest anti-cancer effect on colon cancer cells SW-620 accounting an IC₅₀ value 6.1 µg/ml (Verma *et al.*, 2008). Another study reported the anti-cancer activity of *Polyalthia longifolia* stem bark ethanol extract against HeLa and MCF-7 cells with IC₅₀ values of 25.24 and 50.49 µg/mL, respectively (Manjula *et al.*, 2010). Recent study conducted by Vijayarathna *et al.* (2017a) revealed the anti-cancer activity of standardized methanolic *Polyalthia longifolia* leaf extract against HeLa cells through MTT assay and CyQuant assay with an average IC₅₀ value of 22 µg/mL.

2.3 Plant extraction methods

Plants are highly utilized pharmaceutically due to the rich medicinal values of their phytochemicals such as phenolics and flavonoids. Various plant extraction methods are widely practiced in galenical development such as the maceration method (Figure 2.1 A), soxhlet extraction method (Figure 2.1 B) and ultrasound extraction method (Figure 2.1 C).

2.3.1 Maceration

Maceration is a well-established plant extraction method which involves soaking of the coarse powdered plant material in a closed vessel with an appropriate solvent (Jones and Kinghorn, 2006; Handa *et al.*, 2008). The selection of solvent mainly depends on the bioactive compounds of interest from the plant material and thus the solubility of the compounds in the selected solvent is important to be taken into account while choosing an appropriate solvent. In addition, chemical characterization of the solvent and extraction yield is also equally imperative to consider while choosing a solvent for the plant extraction through maceration method. Commonly used solvents for the maceration process include hexane, chloroform, ethyl acetate, methanol or ethanol (Yan *et al.*, 2008). The maceration process is usually carried out at room temperature for at least three days with occasional agitation to allow the solubilisation of phytochemicals from the plant material. The mixture is then filtered and the final marc is pressed out to completely extract out the dissolved bioactive compounds (Pandey *et al.*, 2014).






Figure 2.1: Different types of plant extraction methods

(A) Maceration Method, (B) Soxhlet Method and (C) Ultrasound or Sonication Method

2.3.2 Soxhlet extraction

Soxhlet extraction, also known as the Hot Continuous Extraction was developed by van Soxhlet in 1879 (Soxhlet, 1879). The finely powdered plant material is placed in a “thimble” which is usually attributed from a strong filter paper or cellulose. The sample containing “thimble” is placed in the thimble-holder of the soxhlet extractor while the extraction solvent is filled in the distillation flask at the bottom. When the solvent is heated, the vapourized solvent is condensed and consecutively fills the thimble containing the plant material. When the solvent level rises, a siphon tube aspirates it from the thimble-holder and discharges it back into the distillation flask. This process is continued until the solvent from the siphon tube does not leave residue when evaporated and thus, it is a continuous–discrete plant extraction method (Luque de Castro and Priego-Capote, 2010).

2.3.3 Ultrasound extraction or sonication extraction

This extraction method comprises the utilization of ultrasound with frequencies ranging from 20 kHz to 2000 kHz (Handa *et al.*, 2008). The acoustic effect from the ultrasound causes produces cavitation which in turn increases the permeability of the cell wall promoting the release of phytochemicals from the plant material into the solvent. Although this extraction method is easy and cost-effective, higher ultrasound energy is known to cause undesirable degradation of bioactive compounds.

2.4 MicroRNA

Small endogenous RNA molecules can be classified into several types, including transfer RNA (tRNA), ribosomal RNA (rRNA), small nucleolar RNA (snoRNA), small interfering RNA (siRNA) and micro RNA (miRNA). The endogenous small miRNA molecules which are approximately 20-22 nucleotides long are derived from the double stranded RNA precursor molecules (Ketting, 2010). The breakthrough of miRNA was first discovered in *Caenorhabditis elegans* and the disclosure of small non-coding lin-4 transcript from *C. elegans* which was 22 nucleotides long found to down regulate LIN-14 protein expression via sequence complementary binding to 3' untranslated region (UTR) of lin-14 mRNA (Lee *et al.*, 1993). Since then, miRNA has attained an increasing deliberation and led to detailed investigation of miRNA biogenesis and function in the advancement of molecular biology. Contemporarily, 28645 distinguish hairpin precursor miRNA, expressing 35828 mature miRNA from 223 species have been identified (Griffiths-Jones, 2004; Griffiths-Jones *et al.*, 2006; Griffiths-Jones *et al.*, 2008; Kozomara and Griffiths-Jones, 2011; Kozomara and Griffiths-Jones, 2014) (<http://www.mirbase.org/> accessed March 10, 2018). This arising principal class of regulatory genes have been identified by bioinformatics prediction approaches and validated through several experimental methods. The involvement of miRNA in the negative regulation of gene expression at post transcriptional level and subsequent protein translational repression (Bartel, 2004) clearly substantiate the major role of miRNA in diverse biological processes such as cell death (Xu *et al.*, 2004), cell proliferation (Cheng *et al.*, 2005), cell development (Yoo and Greenwald, 2005), cell differentiation (Naguibneva *et al.*, 2006), stress resistance (Dresios *et al.*, 2005), haematopoiesis (Garzon *et al.*, 2006), fat metabolism (Ambros, 2003; Esau *et al.*, 2006) and insulin secretion (Poy *et al.*,

2004). Hence, the evolution of miRNA has exposed a novel and attractive therapeutic target and diagnostic tool for various diseases including cancer.

2.4.1 MicroRNA biogenesis

In like manner of precursor mRNA synthesis, miRNA are also generated by RNA polymerase II by initially producing a lengthy transcript called the primary miRNA (pri-miRNA) (Bartel, 2004). The pri-miRNA transcripts have been evidently validated to possess 5' cap and poly (A) tail at 3' end as any other typical mRNA (Cai *et al.*, 2004; Lee *et al.*, 2004). Previous studies suggest that the length of pri-miRNA transcript can be approximately 1000 nucleotide (Lee *et al.*, 2003; Cai *et al.*, 2004). Considering the length of pri-miRNA is pretty long with complementary bases within the transcript, it is legitimate to form a partially paired stem-loop structure (Treiber *et al.*, 2012). This structure acts as substrate for RNase III class of enzymes, namely DROSHA and *Di George Syndrome critical region gene 8* (DGCR8) which eventually recognises the hairpin-loop structure of pri-miRNA and catalyzes it into a short precursor miRNA (pre-miRNA) (Denli *et al.*, 2004; Gregory *et al.*, 2004; Landthaler *et al.*, 2004). This first cleavage process is initiated by the binding of the microprocessor complex (complex of DROSHA and DGCR8) to the open-ended part of the stem-looped miRNA and finally the double-stranded cleavage produces a concise hair-pin shaped RNA molecule with a two nucleotide over hang at the 3' end (Han *et al.*, 2004; Han *et al.*, 2006). The double stranded stem-loop structure of pre-miRNA has been identified to be approximately 70-100 bp long (Treiber *et al.*, 2012).

Subsequently, the transportation of pre-miRNA from nucleus to the cytoplasm is mediated by the nuclear export receptor, known as the Exportin 5 (Yi *et*

al., 2003; Lund *et al.*, 2004). Previous studies demonstrated that the Exportin 5 performs its role as nuclear cargo with the aid of RanGTP in which stable complexes of pre-miRNA·Exportin 5·RanGTP are productively exported to cytoplasm down the RanGTP gradient across the nuclear envelope and pre-miRNA and Exportin 5 are dissociated upon the hydrolysis of RanGTP to RanGDP in cytoplasm (Bohnsack *et al.*, 2004). The free Exportin 5 is then returned back to the nucleus to mediate new pre-miRNA exportation.

Instantaneously, the second cleavage in the biogenesis process of miRNA takes place in the cytoplasm by RNase III enzyme called the DICER (Grishok *et al.*, 2001; Ketting *et al.*, 2001). DICER incorporates PAZ (Piwi, Argonaute and Zwiille) domain that binds to the two nucleotide 3' overhang and anchors the pre-miRNA in position while placing the stem loop terminal at the positively charged catalytic domain of the DICER (Macrae *et al.*, 2006; MacRae *et al.*, 2007). This arrangement enables the DICER to act as a molecular ruler, thereby assisting the cleavage to occur efficiently at approximately 65 angstrom (Å) from PAZ domain and cleaves off the loop from the pre-miRNA (MacRae *et al.*, 2007; Ketting, 2010; Park *et al.*, 2011). The subsequent shorter double stranded RNA of about 20-25 nucleotides in length, with two nucleotide 3' overhangs at both terminals is known as miRNA duplex or miRNA/miRNA* (MacFarlane and Murphy, 2010).

miRNA duplex is then loaded into the miRNA-Induced Silencing Complex (miRISC) and releases one of the strand while selectively bound to one strand in order to generate an active complex (Bartel, 2004). The strand which is integrated into the miRISC is termed as the guide strand (miRNA) while the strand which is released and degraded is termed as the passenger strand (miRNA*). The Argonaute protein being the major component of RISC acts as the capital for catalytic process.

The Argonaute protein comprises two essential domains, namely PAZ and PIWI. The PAZ domain has been demonstrated to bind to the backbone of the guide strand (Song *et al.*, 2003; Ma *et al.*, 2004) while the PIWI domain acts as the RNase H which breaks down the passenger strand (Martinez and Tuschl, 2004; Song *et al.*, 2004; Ma *et al.*, 2005). Figure 2.2 shows an overview of miRNA biogenesis process.




Figure 2.2: Overview of miRNA biogenesis process

(Source: Lin and Gregory, 2015)

Abbreviation: RNA polymerase II (Pol II), Primary microRNA (pri-miRNA), *Di George Syndrome critical region gene 8* (DGCR8), Precursor microRNA (pre-miRNA), Exportin 5 (XPO5), miRNA-Induced Silencing Complex (miRISC)

2.4.2 MicroRNA and cancer

Ever since the exploration of miRNA and its' correlation with the widespread biological processes mainly including apoptosis and cell proliferation, the fundamental significance of miRNA in tumorigenesis are strongly postulated. Henceforth, the miRNA-mediated molecular mechanism in cancer biology has unfastened a novel dimension for cancer therapeutic targets as well as cancer biomarkers. The miRNA binds to its target mRNA by partial complementary binding, thus silences the gene expression and represses the post translational activity. The means of function of miRNA via alteration of gene expression and consecutive translational expression, pinpoints that miRNAs can act as tumor suppresser genes or oncogenes depending on their target genes (Kent and Mendell, 2006; Zhang *et al.*, 2007).

For instance, up-regulation of specific miRNA targeting the tumor suppressor genes, eventually, promoting cell growth and cancer initiation acts as an oncogene. On the other hand, up-regulation of specific miRNA targeting genes responsible for oncogenic activities which ultimately lead to cancer inhibition or repression acts as tumor suppressor gene (Shenouda and Alahari, 2009). However, the up-growing investigations on miRNA have uncovered the dual role of miRNA in cancer, in which various evidence supports the concept that a same individual miRNA can act as both oncogene and tumor-suppressor gene depending on the cellular environment (Schetter *et al.*, 2012; Sharma *et al.*, 2014; Ding *et al.*, 2018). Based on the literature, extensive studies have reported the correlation between miRNAs and cancer to date (>38, 000 Pubmed hits as of January 2019).

2.4.2(a) Mechanisms involved in microRNA deregulation in cancer

The dysregulation of miRNAs in cancer occur through numerous overlapping mechanisms including chromosomal abnormalities, transcriptional control alterations, epigenetic modulation and disruption in the miRNA processing machinery (Peng and Croce, 2016). For instance, chromosomal alterations may occur due to amplification of a chromosome site harbouring a specific miRNA, leading to an over-expression of the particular miRNA (Hayashita *et al.*, 2005; Tagawa and Seto, 2005) while deletion of the chromosome site may result in down-regulation of the specific miRNA (Calin *et al.*, 2002; Calin and Croce, 2006).

Other than that, various transcriptional factors have been evidently reported to control the expression of miRNAs such as c-Myc (Chang *et al.*, 2008; Han *et al.*, 2013), p53 (He *et al.*, 2007; Hermeking, 2010), myeloid transcription factors PU.1 and C/EBPs (Fukao *et al.*, 2007) and transcription factors NFI-A and C/EBP α (Fazi *et al.*, 2005). Besides, miRNAs have also been reported to undergo epigenetic changes through CpG methylation (Fazi *et al.*, 2007), DNA methylation with histone acetylation inhibitors (Saito *et al.*, 2006), and hypermethylation (Lujambio *et al.*, 2008; Lehmann *et al.*, 2008).

Finally, dysregulation or mutation of any proteins involved in miRNA biogenesis process as described in section 2.1.1 such as DROSHA (Thomson *et al.*, 2006), DGCR8 (Walz *et al.*, 2015), Dicer (Kumar *et al.*, 2009; Iliou *et al.*, 2014), Argonaute proteins (Zhang *et al.*, 2013; Völler *et al.*, 2013), TRBP (Melo *et al.*, 2009) and Exportin 5 (Melo *et al.*, 2010) which leads to miRNA dysregulation.

2.4.2(b) Pathways involved in microRNA regulation in cancer

The current chemotherapy targeting miRNA is hugely attaining interest due to their important participation in cancer pathway. Numerous miRNAs were also evidently shown to regulate tumor necrosis factor (TNF)-related apoptosis-inducing ligand (TRAIL)-induced apoptosis pathway. Based on the research conducted by Garofalo *et al.* (2008), over-expression of miR-221 and miR-222 was shown to hinder the TRAIL-induced apoptosis in non-small cell lung cancer by suppressing the important apoptotic protein expressions as well as targeting the tumor suppressor p27^{kip}mRNAs. The up-regulation of miR-221 and miR-222 was also demonstrated to be over-expressed, leading to the down-regulation of tumor suppressor p27^{kip} in prostate carcinoma (Galardi *et al.*, 2007) and melanoma (Felicetti *et al.*, 2008).

Besides, the involvement of miR-21 regulation in AKT-dependent pathway was also reported to inhibit apoptosis by directly targeting and eventually repressing the expression of FasL (Sayed *et al.*, 2010). Another example of miRNA to be participating in Fas-mediated apoptotic pathway is miR-24 which was shown to target Fas-binding pro-apoptotic protein, namely Fas-associated factor 1 (FAF1), leading to inhibition of apoptosis in different types of cancer (Qin *et al.*, 2010; Schickel *et al.*, 2010). Another cancer pathway, namely the phosphatase and tensin homologue (PTEN) pathway was also shown to be regulated by the expression of miRNAs. For instance, many miRNAs are reported to target and suppress the expression PTEN which is one of the prominent tumor suppressor genes such as miR-17-5p (Xiao *et al.*, 2008), miR-19305p (Xiao *et al.*, 2008), miR-2127 (Sayed *et al.*, 2010) and miR-221 and miR-222 (Chun-Zhi *et al.*, 2010).

There are various microRNAs which have been reported to regulate the cell cycle regulatory pathway, in which oncogenic microRNAs tend to expedite cell cycle progression while the microRNAs with tumor suppressor effect tend to facilitate cell cycle arrest. Exemplary oncogenic microRNA include miR-106b and miR-17-92 families which have been reported to be over-expressed in various cancers are known to target one of the important inducer of G1 arrest, namely p21 from the Cip/Kip family of CDK inhibitors (Ivanovska *et al.*, 2008; Kim *et al.*, 2009). Other studies have also experimentally validated numerous other miRNAs to target other genes involved in cell cycle which eventually regulate the RAS/RAF/MAPK pathway as well as the p53 pathway (Jansson and Lund, 2012). Furthermore, microRNAs are also very well known to target numerous genes involved in DNA damage response in cancer cells. For instance, miR-421 was reported to be highly over-expressed in neuroblastoma and B-cell lymphoma cell lines and was shown to target the apical damage sensor kinase ATM (Hu *et al.*, 2010).

2.4.3 MicroRNA and medicinal plants

Endogenous microRNAs (miRNAs) are short single-stranded RNA molecules that cause mRNA cleavage or translational repression through partial complementary binding to the 3' untranslated region (UTR) of specific protein coding mRNAs (Ling *et al.*, 2013). This plays important role in regulation of various biological activities, thus leading to the break thorough of miRNA-based therapeutic approaches for several diseases. As medicinal plants are highly utilized in the treatment of diseases for centuries due to its rich bioactive phytochemicals, the understanding of its mechanistic actions in regulating the miRNA expressions has recently drawn an