



***ERGOGENIC EFFECT OF MENKUDU (*Morinda citrifolia* L.) LEAF
EXTRACT ON OBESE SPRAGUE DAWLEY RATS USING METABOLOMIC
APPROACH***

NORDIANA BINTI ABDUL MAJID

FSTM 2019 26



**ERGOGENIC EFFECT OF *MENGKUDU* (*Morinda citrifolia* L.) LEAF
EXTRACT ON OBESE SPRAGUE DAWLEY RATS USING METABOLOMIC
APPROACH**

By

NORDIANA BINTI ABDUL MAJID

**Thesis Submitted to the School of Graduate Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirements for the Degree of Master of
Science**

April 2019

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

ERGOGENIC EFFECT OF *MENKUDU* (*Morinda citrifolia* L.) LEAF EXTRACT ON OBESE SPRAGUE DAWLEY RATS USING METABOLOMIC APPROACH

By

NORDIANA BINTI ABDUL MAJID

April 2019

Chair : Professor Azizah Abdul Hamid, PhD
Faculty : Food Science and Technology

Natural products are getting much acceptance as ergogenic aids, in enhancing physical performance, not only among the athletes but also the general population. Obese person mostly has reduced desire and ability to exercise; resulting in difficulty to reduce weight and fat in the body. Thus, they need to boost their energy production so that they can be more active and healthier. In this study, *Morinda citrifolia* leaf extract (MLE) believed to possess ergogenic property was used to evaluate its effect on an obese animal model by using Forced Swimming Test (FST) as endurance exercise and the changes in metabolic profiles of exercise obese rats after treatment was also identified using ¹H-NMR based metabolomics. The rats were fed with high fat diet (HFD) for 12 weeks for obese development. Once become obese, all the rats undergone endurance exercise every two weeks for 8 weeks together with treatment. Three different dosages of MLE used were 50 mg/kg, 100 mg/kg and 200 mg/kg of body weight (BW) together with two positive Control, 5 mg/kg caffeine and 100 mg/kg green tea. Blood and urine were collected for the metabolomic study. Animal study carried out showed that rats fed 200 mg/kg BW MLE demonstrated the longest endurance capacity of approximately three times as long as that of green tea and caffeine. The rats were also found to have lower lactate level, suggesting that energy metabolism was more effective in these rats. In addition, lactate dehydrogenase enzyme (LDH) activity, muscle injury indicator, was found to be lowest in rats fed the highest MLE level. Interestingly, the same effect was not seen in rats fed either caffeine or green tea, indicating that MLE treatment able to protect rat's muscle. In metabolomic, multivariate analysis including unsupervised and supervised analysis were used to identify the potential

biomarkers. The study showed that feeding the rats at a dose of 200 mg/kg BW MLE altered metabolites present in the serum of exercised obese rat. The PLS-DA score plots showed distinct separation between normal rats with that of green tea and caffeine treatments, and instead, were very similar to that of 200 mg/kg BW MLE suggesting that the 200 MLE feeding was more effective in improving endurance capacity in comparison to that of either synthetic or natural ergogenic substances. Metabolites such as glutamine, glycerol, glycine, acetoacetate, taurine, carnitine, succinate, pyruvate and 2-hydroxybutyrate were found to be higher after MLE treatment suggested that changes in metabolic pathways which included carbohydrate, lipid and energy metabolism. In conclusion, this study reports on the potential ergogenic property of high dose of MLE based on the enhancement swimming capacity, energy metabolism and metabolic perturbation in the exercised obese rats.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains

**KESAN ERGOGENIK DAUN MENGKUDU (*Morinda citrifolia* L.) TERHADAP
TIKUS SPRAGUE DAWLEY OBESE MENGGUNAKAN PENDEKATAN
METABOLOMIK**

Oleh

NORDIANA BINTI ABDUL MAJID

April 2019

Pengerusi : Profesor Azizah Abdul Hamid, PhD
Fakulti : Sains dan Teknologi Makanan

Produk semulajadi semakin diterima sebagai alat bantuan ergogenik, dalam meningkatkan prestasi fizikal, bukan sahaja di kalangan atlet tetapi juga di kalangan umum. Orang gemuk kebiasaannya telah berkurang keinginan dan keupayaan untuk bersenam; menyebabkan kesukaran untuk mengurangkan berat badan dan lemak di dalam badan. Oleh itu, mereka perlu meningkatkan pengeluaran tenaga supaya mereka boleh menjadi lebih aktif dan lebih sihat. Dalam kajian ini, ekstrak daun *Morinda citrifolia* (MLE) dipercayai mempunyai sifat ergogenik digunakan untuk menilai kesannya ke atas model haiwan obes dengan menggunakan Ujian Berenang Paksa (FST) sebagai latihan ketahanan dan perubahan dalam profil metabolik tikus gemuk yg berlatih selepas rawatan juga dikenalpasti menggunakan metabolomik berasaskan $^1\text{H-NMR}$. Tikus diberi makan dengan diet lemak tinggi (HFD) selama 12 minggu untuk menjadikannya obes. Setelah obes, semua tikus menjalani latihan ketahanan setiap dua minggu selama 8 minggu bersama-sama dengan rawatan. Tiga dos MLE berbeza yang digunakan ialah 50 mg / kg, 100 mg / kg dan 200 mg / kg berat badan (BW) bersama-sama dengan dua Kawalan positif, 5 mg / kg kafein dan 100 mg / kg teh hijau. Darah dan air kencing dikumpulkan untuk kajian metabolomik. Kajian haiwan menunjukkan bahawa tikus yang diberi makan 200 mg / kg BW MLE menunjukkan kapasiti ketahanan paling lama kira-kira tiga kali selagi teh hijau dan kafein. Tikus juga didapati mempunyai tahap laktat yang rendah, menunjukkan bahawa metabolisme tenaga lebih berkesan dalam tikus ini. Di samping itu, aktiviti enzim laktat dehidrogenase (LDH), penunjuk kecederaan otot, didapati paling rendah dalam tikus yang diberi peringkat tertinggi MLE. Menariknya, kesan yang sama tidak dilihat pada tikus yang diberi minum kafein atau teh hijau, menunjukkan bahawa rawatan MLE dapat melindungi otot tikus. Dalam analisis metabolomik, multivariate termasuk analisis yang tidak diselia dan diselia digunakan untuk mengenal pasti biomarker berpotensi. Kajian menunjukkan bahawa pada dos 200 mg / kg BW MLE mengubah metabolit yang hadir dalam serum tikus gemuk. Skop

skor PLS-DA menunjukkan pemisahan berbeza antara tikus normal dengan teh hijau dan rawatan kafein, dan sebaliknya, sangat mirip dengan 200 mg / kg BW MLE yang menunjukkan bahawa memberi makan dengan 200 MLE lebih berkesan dalam meningkatkan kapasiti ketahanan berbanding dengan bahan ergogenik sintetik atau semula jadi. Metabolit seperti glutamin, gliserol, glisin, acetonacetate, taurine, carnitine, succinate, piruvat dan 2-hydroxybutyrate didapati lebih tinggi selepas rawatan MLE menunjukkan perubahan dalam metabolisme karbohidrat, lipid dan tenaga. Sebagai kesimpulan, kajian ini melaporkan tentang potensi dos tertinggi MLE sebagai agen ergogenik berdasarkan keupayaan meningkatkan kapasiti berenang, metabolisme tenaga dan perubahan metabolik dalam tikus obes yang dijalankan.



ACKNOWLEDGEMENTS

Alhamdulillah, praised to Allah The Almighty, without His will no one can achieve anything such as my works today. I feel great gratitude for His graciousness and mercifulness in giving me strength and spirit everyday until I finished this research, and salawat to His messenger the beloved prophet, Muhammad s.a.w who bring inspiration in my everyday life.

I would like to express my deep appreciation and gratitude to my supervisor committee chairman, Prof. Dr Azizah Abdul Hamid for her incalculable suggestion, guidance, patience, opportunity and support along my master's journey. I would like to express my gratitude to my other supervisory committee, Prof Dr Nazamid Saari and Assoc Prof Dr. Faridah Abas for their constructive comment, ideas and enthusiastic encouragement for this research work. My sincere gratitude also goes to SGS UPM for their financial support in term of scholarship provided to me during my postgraduate study period.

I would like to extend my thankfulness to all my friends particularly my lab mates that had contributed to my works by sharing ideas, giving comments and supports. Assistance provided by Universiti Putra Malaysia (UPM), Institut Biosains staff was greatly appreciated.

Finally, I would like to thanks to my parents, siblings and my husband for their incalculable support and strength in this journey of pursuing knowledge.

I certify that a Thesis Examination Committee has met on 22 April 2019 to conduct the final examination of Nordiana binti Abdul Majid on her thesis entitled "Ergogenic Effect of *Mengkudu (Morinda citrifolia L.)* Leaf Extract on Obese Sprague Dawley Rats Using Metabolomic Approach" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Hasanah bt Mohd Ghazali, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

Son Radu, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Internal Examiner)

Noriham Abdullah, PhD

Professor
Faculty of Applied Science
Universiti Teknologi MARA
Malaysia
(External Examiner)



RUSLI HAJI ABDULLAH, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 23 July 2019

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Azizah Abdul Hamid, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

Nazamid Saari, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

Faridah Abas, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

ROBIAH BINTI YUNUS, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Nordiana Binti Abdul Majid (GS48694)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____

Name of Chairman of
Supervisory
Committee:

Prof. Dr Azizah Abdul Hamid

Signature: _____

Name of Member of
Supervisory
Committee:

Prof. Dr Nazamid Saari

Signature: _____

Name of Member of
Supervisory
Committee:

Assoc. Prof. Dr. Faridah Abas

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xii
LIST OF FIGURES	xiii
LIST OF SYMBOLS AND ABBREVIATIONS	xv
CHAPTER	
1. INTRODUCTION	1
2. LITERATURE REVIEW	4
2.1 Ergogenic	4
2.1.1 Description of ergogenic	4
2.1.2 Nutritional ergogenic aids	5
2.1.2.1 Green tea	5
2.1.2.2 Coffee	5
2.1.2.3 Tongkat Ali	6
2.1.2.4 Ginseng	7
2.1.2.5 Carbohydrate	8
2.1.2.6 L-carnitine	8
2.2 Obesity	8
2.2.1 Description of obesity	8
2.2.2 Causes of obesity	9
2.2.3 Complication associated with obesity	10
2.2.4 Strategies for weight management	11
2.3 Exercise and Energy production	12
2.3.1 Aerobic pathway for energy production	12
2.3.2 Anaerobic and lactic acid pathway	13
2.4 Fatigue	14
2.4.1 Obesity and fatigue	15
2.4.2 Oxidative stress	15
2.4.3 Lactate and fatigue	16
2.4.4 Lactate dehydrogenase and fatigue	16
2.5 Endurance Exercise Model	16
2.5.1 Weight Loaded Forced swimming test	16
2.5.2 Voluntary wheel running	18
2.5.3 Treadmill running	18
2.6 <i>Morinda citrifolia</i> L.	19
2.6.1 General description	19
2.6.2 Phytochemical composition of <i>M. citrifolia</i> L	20
2.6.3 Pharmacological effect of <i>M. citrifolia</i> L	21
2.7 Metabolomic	23

2.7.1	NMR spectroscopy	24
2.7.2	Samples type	24
2.7.3	Multivariate data analysis	25
3.	MATERIALS AND METHODS	27
3.1	Materials	27
3.2	Methods	27
3.2.1	Preparation of <i>M. citrifolia</i> leaf extract (MLE)	27
3.2.2	Animal Model	28
3.2.3	Obesity induction	28
3.2.4	Treatment with MLE	29
3.2.5	Weight loaded Forced swimming test	29
3.2.6	Blood collection from the rats	31
3.2.7	Biochemical measurements	31
3.2.8	Preparation of urine and serum for ¹ H-NMR analysis	31
3.2.9	¹ H-NMR data processing and statistical analysis	32
4.	RESULT AND DISCUSSION	33
4.1	Percent yield of the extract	33
4.2	Body weight gain in Sprague Dawley rats	33
4.3	Forced swimming test	34
4.4	Biochemicals parameter related to energy metabolism	37
4.5	¹ H-NMR spectra of rat's serum and urine	41
4.6	Multivariate analysis of serum and urine metabolites of Sprague Dawley rats fed HFD or ND for 12 weeks	50
4.7	Identification of potential metabolites for obesity induction in urine and serum	53
4.8	Multivariate data analysis of exercised rat's serum after 8 weeks of treatment	63
4.9	Metabolic changes in obese exercised rat's serum after MLE treatment.	73
5.	CONCLUSION AND RECOMMENDATION	79
5.1	Summary and Conclusions	79
5.2	Recommendation	80
	REFERENCES	81
	APPENDICES	101
	BIODATA OF STUDENT	103
	LIST OF PUBLICATIONS	104

LIST OF TABLES

Table		Page
2.1	Specific Characterization of <i>M. Citrifolia</i> L.	20
2.2	Summary of Pharmacological Effect of <i>M. Citrifolia</i> L.	23
4.1	List of Metabolites in Rat's Serum and Urine	49
4.2	PLS-DA and OPLS-DA Model's Validation for Serum and Urine of Sprague-Dawley Rats Fed High Fat (HFD) or Normal Diet (ND) for 12 Weeks	53
4.3	Relative Quantitative of Discriminating Metabolites of Serum in the Obese and Normal Rats after 12 Weeks Fed HFD using Chenomx NMR Suite	56
4.4	Relative Quantitative of Discriminating Metabolites of Urine in the Obese and Normal Rats after 12 Weeks Fed HFD using Chenomx NMR Suite	60
4.5	Relative Quantification of Significant Discriminating Metabolites Based on the Concentration of 0.1% Of 3-Trimethylsilyl Propionic-2,2,3,3-D4 Acid Sodium Salt (TSP) as Internal Standard and Quantified using Chenomx NMR Suite	72

LIST OF FIGURES

Figure		Page
2.1	Metabolism Pathways of Carbohydrate, Protein and Fat During Exercise	13
2.2	Forced Swimming Test Apparatus for Exercise Model	17
2.3	Voluntary Wheel Running Apparatus for Exercise Model	18
2.4	Treadmill Running Apparatus for Exercise Model	19
3.1	Experimental Design for the Ergogenic Activity of <i>M. Citrifolia</i> Leaf Extract in Exercised HFD Induced Obese Sprague Dawley Rats	30
4.1	Percent Increase in Body Weight of the Rats with Two Different Diet; Normal Diet (ND) or High Fat Diet (HFD)	34
4.2	Effect of MLE Treatment on FST (Second) of Rats at 0 Week (Before the Treatment) and after 8 Weeks (Final Week of Treatment)	35
4.3	Lactate Levels of Exercised Rats after 8 Weeks of MLE Treatment	38
4.4	The LDH Levels on Exercised Rats after 8 Weeks of MLE Treatment	40
4.5a	Representative of ¹ H Nuclear Magnetic Resonance (NMR) of Serum Spectra Collected from Normal Rat Fed Normal Chow and Obese Rat Fed High Fat Diet with Expanded Region Δ 1.00-5.40 ppm	42
4.5b	Typical of ¹ H-NMR Serum Spectra (Δ 0.9-2.1) Collected from the Normal Rat Fed Normal Chow and Obese Rat Fed High Fat Diet	43
4.5c	Typical of ¹ H-NMR Serum Spectra (Δ 3.0-5.4) Collected from the Normal Rat Fed Normal Chow and Obese Rat Fed High Fat Diet	44
4.6a	Typical of ¹ H-NMR Serum Spectra (Δ 3.0-5.4) Collected from the Normal Rat Fed Normal Chow and Obese Rat Fed High Fat Diet	45
4.6b	Typical of ¹ H-NMR Urine Spectra (Δ 0.7-2.9) Collected from the Normal Rats Fed Normal Chow and Obese Rats Fed High Fat Diet	46
4.6c	Typical of ¹ H-NMR Urine Spectra (Δ 3.0-6.0) Collected from the Normal Rat Fed Normal Chow and Obese Rat Fed High Fat Diet	47
4.6d	Typical of ¹ H-NMR Urine Spectra (Δ 6.1-9.5) Collected from the Normal Rat Fed Normal Chow and Obese Rat Fed High Fat Diet	48

4.7	PCA Derived Score Plot obtained using ¹ H-NMR Data for Serum Samples from Sprague Dawley Rats after Fed a High Fat and Normal Diet	51
4.8	PCA Derived Score Plot obtained using ¹ H-NMR Data for Urine Samples from Sprague Dawley Rats after Fed a High Fat and Normal Diet	52
4.9	OPLS-DA Score Plot (A), Loading Plot (B) and S-Plot obtained using ¹ H-NMR Data for Serum Samples from the Normal and Obese Sprague Dawley Rats	55
4.10	OPLS-DA Score Plot (A), Loading Plot (B) and S-Plot obtained using ¹ H-NMR Data for Urine Samples from the Normal and Obese Sprague Dawley Rats	60
4.11	The Representative of ¹ H-NMR Spectra of Serum Sample from Different Groups of Exercised Sprague Dawley Rats after 8 Weeks of Treatment	64
4.12	PLS-DA Score Plot (A) and Loading Plot (B) Model for Nontargeted Approach Comparing the Three-Different Dosages of MLE and Normal Rats after 8 Weeks of Treatment	66
4.13	PLS-DA Model for Nontargeted Approach Comparing the Five Different Groups of Rats; 200 mg/kg BW MLE, 5 mg/kg BW Caffeine, 100 mg/kg Green Tea, Normal and Obese after 8 Weeks of exercise and Treatment	67
4.14	OPLS-DA Score Scatter (A) and Loading Scatter (B) Derived from Rat Serum from Exercised Obese HFD and Exercised Obese 200 mg/kg BW MLE After 8 Weeks of Exercise and Treatment with 200 mg/kg BW MLE	68
4.15	OPLS-DA Score Scatter (A) and Loading Scatter (B) Derived from Rat Serum of Exercised Obese HFD and Exercised Obese 5 mg/kg BW Caffeine after 8 Weeks of Exercise and Treatment with 5 mg/kg BW Caffeine	69
4.16	OPLS-DA Score Scatter (A) and Loading Scatter (B) Derived from Rat Serum of Exercised Obese HFD and Exercised Obese 100 mg/kg BW Green Tea after 8 Weeks of Exercise and Treatment with 100 mg/kg BW Green Tea	70
4.17	VIP Value of the Metabolites. Only the Metabolites with the VIP Value > 0.5 were Chosen	71
4.18	OPLS-DA Trajectory Score Plot obtained using ¹ H-NMR Spectra for Serum Samples from Normal Rats (NDF), Obese Exercised Rats before (200MLEI) and after (200 MLEF) Treated with 200 mg/kg BW MLE	74

4.19 OPLS-DA Score (A) and Loading Column (B) obtained on Serum ¹H-NMR Spectra of the Exercised Obese Rats before and after Treated with 200 mg/kg BW MLE. 75



LIST OF ABBREVIATIONS

^1H NMR	Proton Nuclear Magnetic Resonance Spectroscopy
<i>D</i>	Doublet
<i>Dd</i>	Doublet of doublet
DPPH	Diphenylpicrylhydrazyl
g	Gram
HFD	High fat diet
HPLC	High Pressure Liquid Chromatography
LCMS	Liquid Chromatography Mass Spectrometry
<i>M</i>	Multiplet
MHz	Mega Hertz
mL	Milliliter
MLE	<i>M. citrifolia</i> leaf extract
Mmol	Millimole
ND	Normal diet
°C	Degree in Celsius
OPLS-DA	Orthogonal Partial Least Squares–Discriminant Analysis
PC	Principal Component
PCA	Principal Component Analysis
PLS-DA	Partial Least Squares–Discriminant Analysis
<i>S</i>	Singlet
SIMCA	Soft Independent Modelling of Class Analogy
TPC	Total Phenolic Contents
VIP	variable importance in the projection
Δ	Chemical Shift in ppm

CHAPTER 1

INTRODUCTION

Obesity is one of the most problematic condition worldwide, showing an increment in the risk of morbidity and mortality throughout most countries around the world. Malaysia known as Asian's fattest country, recorded an increase in its obesity rate, with the latest data showing half of its population are obese and overweight (NHMS, 2015). Obesity is related to metabolic perturbation that causes complications which include hypertension, hyperlipidaemia, diabetes mellitus, cancer and cardiovascular diseases. Exercise and reduction of caloric intake are most effective in making obese people to be healthier and have an active lifestyle (Mukherjee, 2003). Despite being the most effective way in weight management, individual's compliance is inconsistent, resulting from low performance and oxidative stress associated fatigue, justifying the need for better alternatives, including herbal sources. This is useful, in particular, for the herbs that possess ergogenic property.

Ergogenic aid is defined as substance that enhances the use of energy such as energy generation, efficiency and control. Athletes consume ergogenic aids to enhance their performances, elevate possibility to win in competitions, and improve their physicals (Silver, 2001; Palou and Bonet, 2007). Ergogenic aids will be beneficial for the obese and non-obese in efficient energy production and potentially in weight management as well as to improve health, have more active lifestyle and free from chronic diseases. There are many types of ergogenic aids such as doping agents (Saugy et al., 2006), amino acid and protein supplements (Kerksick et al., 2006), ginseng (Oliynyk and Oh 2013), caffeine (Beck et al., 2006) and some minerals (Williams, 2005). Creatine supplementation is an example of amino acid supplement that can be obtained naturally from meat. Creatine can act as dietary antioxidant because it can regulate oxidative stress (Sung et al., 2016). Caffeine is also regarded as ergogenic aid because it can enhance endurance by stimulating central nervous system. Some of ergogenic substances such as doping agents (anabolic steroids, growth hormone and other anabolic agents) are illegal and over consumption can lead to serious adverse effects including metabolic diseases, mental problem, cancer, renal and hepatic damage (Bird et al., 2016; Maravelias et al., 2005). Choosing the natural ergogenic aids are becoming more popular and might be a safer way to enhance performance without worrying any dangerous adverse effects. Nowadays, various plants and herbs have been shown to exhibit ergogenic property and this include Astragali radix (Li et al., 2014), capsicum (Lim et al., 1997) and ginseng (Bahrke and Morgan, 2000). Flavonoids content in some of the plants may contribute to the ergogenic property due to their potent antioxidant activity. Hence, study on ergogenic property of some plants are recommended because most plants and herbs possess antioxidant activity.

Morinda citrifolia commonly called 'mengkudu' in Malaysia was discovered more than 2000 years ago (Gerlach, 1996). There are various bioactive compounds that have been found in this plant including anthraquinones, anthraquinones glycosides, lignans, flavonol glycosides, phenylpropanoids, saccharides, triterpenoids and fatty acids (Pawlus and Kinghorn, 2007). Each part of the plant (leaf, bark, stem, fruit and root) has different chemical composition and thus exhibits different biological activity. A study on bioactivities of *M. citrifolia* (Jambocus et al., 2016) revealed that the leaves extracted with 60 – 40% ethanol consisted more bioactive compounds and showed potent antioxidant activity as compared to that using 100% of ethanol. In the same study, the researchers reported the ability of *M. citrifolia* leaves extract in the prevention and treatment of obesity in the obese rats. In other study, *M. citrifolia* leaves water extract has been reported to enhance performance in swimming mice better than *M. citrifolia* fruit extract and green tea extract (Shalan et al., 2016). Despite as anti-obese and exhibit ergogenic property, this plant also has been studied for cancer treatment (Hirazumi et al, 1994), inflammation, diabetes, asthma, hypertension and pain (Solomon, 1999). There are many studies done on *M. citrifolia* activities, but the mechanisms of this plant as ergogenic property in an obese has not been discovered yet. Based on its potent antioxidant activity and ability to repair free radical damages and slow down oxidation process that might lead to oxidative fatigue, *M. citrifolia* leaf is expected to have ergogenic property.

Combination of exercise and consumption of the extract can cause changes in body metabolism that can increase energy production. Nuclear Magnetic Resonance (NMR) is one of the methods for metabolite assessment that involves in measuring the overall metabolic signature of biological samples. Metabolomics is most common available technology in detection, identification, quantification and differentiation of dynamic multivariate metabolic changes of biology system. Urine and blood serum are most widely used biofluid in metabolomics. Alteration in metabolomic profile also can be detected using urinary and serum metabolomics. Non-targeted approach of metabolomic using urine sample could provide new information related to development of food metabolome. Recent study by Hodgson et al., (2013) had identified the effects of drinking a green tea extract on the metabolic profiling of athletes by using ¹NMR-based metabolomic. Results showed an effect of the green tea-based sports drink that is related with energy metabolism of athletes based on the metabolites identified. Hence, this approach will be used to investigate metabolic changes due to physical exercise in an obese rat.

Problem Statements and Significant of the Study

An obese person usually will easily suffer from fatigue upon physical exercises. This is the main problem faced by the obese individuals, when they want to do reduce their weight. Consequently, they are in need of substance that can boost their energy production so that they can be more active and have healthier lifestyle. Pharmaceuticals may help increase energy production but most of them will show many side effects. Hence, the use of a natural product that has ergogenic property with minimal side effect is recommended.

M. citrifolia leaf consisted of abundant of bioactive compounds that give benefits to human health. Previous studies, this plant leaf known to have potential as an anti-obesity property through *in vitro* and *in vivo* studies and its mechanism by using ¹H-NMR approaches. However, there are no reported study on the ergogenic property of the same in high fat induced obese rats. By virtue of its antioxidant activity, this plant is expected to have an ergogenic property. Metabolomics is useful in confirming the effectiveness of the extract by identifying the changes of the metabolites produced by the different groups. Hence, this study hypothesizes that, the leaf extract of *M. citrifolia* exhibit the ergogenic property based on the perturbation of metabolic pathways in exercise high fat diet induced obese rat model.

The present study has been initiated with following objectives:

- 1) To demonstrate the ergogenic effect of *M. citrifolia* leaf extract in the obese rats using Forced Swimming Test.
- 2) To evaluate the effect of exercise on metabolic profile of obese rats using ¹H-NMR based metabolomic.

REFERENCES

- Alessio, H. M., Hagerman, A. E., Fulkerson, B. K., Ambrose, J., Rice, R. E. and Wiley, R. L. (2000). Generation of reactive oxygen species after exhaustive aerobic and isometric exercise. *Medicine and science in sports and exercise*, 32(9), 1576-1581.
- Ali, F., Ismail, A., Esa, N. M. and Pei, C. P. (2015). Transcriptomics expression analysis to unveil the molecular mechanisms underlying the cocoa polyphenol treatment in diet-induced obesity rats. *Genomics*, 105(1), 23-30.
- Allen, D. and Westerblad, H. (2004). Lactic acid—the latest performance-enhancing drug. *Science*, 305(5687), 1112-1113.
- Alpert, M. A. (2001). Obesity cardiomyopathy: pathophysiology and evolution of the clinical syndrome. *The American journal of the medical sciences*, 321(4), 225-236.
- An, L., Shi, Q. and Feng, F. (2015). Metabolomics approach to identify therapeutically potential biomarkers of the Zhi-Zi-Da-Huang decoction effect on the hepatoprotective mechanism. *RSC Advances*, 5(102), 84048-84055.
- Angulo, P. (2002). Nonalcoholic fatty liver disease. *New England Journal of Medicine*, 346(16), 1221-1231.
- Anugweje, K. C. and E. O. Ayalogu. (2014) "Effect of *Morinda citrifolia* Supplementation on the Lactate Dehydrogenase (LDH) levels of Athletes." *New York Science Journal*.
- Anugweje, K. C. and Willey-Abey, B. (2014) "Effect of *Morinda citrifolia* Supplementation on the Lactate levels of Athletes." *New York Science Journal*.
- Bäckhed, F., Ding, H., Wang, T., Hooper, L. V., Koh, G. Y., Nagy, A. and Gordon, J. I. (2004). The gut microbiota as an environmental factor that regulates fat storage. *Proceedings of the national academy of sciences*, 101(44), 15718-15723.
- Bäckhed, F., Manchester, J. K., Semenkovich, C. F. and Gordon, J. I. (2007). Mechanisms underlying the resistance to diet-induced obesity in germ-free mice. *Proceedings of the National Academy of Sciences*, 104(3), 979-984.
- Bahrke, M. S., and Morgan, W. P. (2000). Evaluation of the ergogenic properties of ginseng. *Sports Medicine*, 29(2), 113-133.
- Bailey, T. G., Jones, H., Gregson, W., Atkinson, G., Cable, N. T. and Thijssen, D. H. (2012). Effect of ischemic preconditioning on lactate accumulation and running performance. *Medicine & science in sports & exercise*, 44(11), 2084-2089.

- Balentine, D. A., Wiseman, S. A. and Bouwens, L. C. (1997). The chemistry of tea flavonoids. *Critical Reviews in Food Science & Nutrition*, 37(8), 693-704.
- Balsom, P. D., Gaitanos, G. C., Ekblom, B. and Sjodin, B. (1994). Reduced oxygen availability during high intensity intermittent exercise impairs performance. *Acta Physiologica Scandinavica*, 152(3), 279-285.
- Balsom, P. D., Gaitanos, G. C., Söderlund, K. and Ekblom, B. (1999). High-intensity exercise and muscle glycogen availability in humans. *Acta Physiologica Scandinavica*, 165, 337-346.
- Bangsbo, J., Graham, T. E., Kiens, B. and Saltin, B. (1992). Elevated muscle glycogen and anaerobic energy production during exhaustive exercise in man. *The Journal of Physiology*, 451(1), 205-227.
- Barba, I., de León, G., Martín, E., Cuevas, A., Aguade, S., Candell-Riera, J. and Garcia-Dorado, D. (2008). Nuclear magnetic resonance-based metabolomics predicts exercise-induced ischemia in patients with suspected coronary artery disease. *Magnetic Resonance in Medicine: An Official Journal of the International Society for Magnetic Resonance in Medicine*, 60(1), 27-32.
- Barton, D. L., Soori, G. S., Bauer, B. A., Sloan, J. A., Johnson, P. A., Figueras, C. and Christensen, B. (2010). Pilot study of *Panax quinquefolius* (American ginseng) to improve cancer-related fatigue: a randomized, double-blind, dose-finding evaluation: NCCTG trial N03CA. *Supportive Care in Cancer*, 18(2), 179.
- Bastien, M., Poirier, P., Lemieux, I. and Després, J. P. (2014). Overview of epidemiology and contribution of obesity to cardiovascular disease. *Progress in cardiovascular diseases*, 56(4), 369-381.
- Beck, T. W., Housh, T. J., Schmidt, R. J., Johnson, G. O., Housh, D. J., Coburn, J. W. and Malek, M. H. (2006). The acute effects of a caffeine-containing supplement on strength, muscular endurance, and anaerobic capabilities. *The Journal of Strength & Conditioning Research*, 20(3), 506-510.
- Beneke, R., Leithäuser, R. M. and Ochentel, O. (2011). Blood lactate diagnostics in exercise testing and training. *International journal of sports physiology and performance*, 6(1), 8-24.
- Bray, G. A., Ryan, D. H., Gordon, D., Heidingsfelder, S., Cerise, F. and Wilson, K. (1996). A double-blind randomized placebo-controlled trial of sibutramine. *Obesity Research*, 4(3), 263-270.
- Beckwith-Hall, B. M., Thompson, N. A., Nicholson, J. K., Lindon, J. C. and Holmes, E. (2003). A metabolomic investigation of hepatotoxicity using diffusion-edited ¹H NMR spectroscopy of blood serum. *Analyst*, 128(7), 814-818.

- Bener, A., Yousafzai, M. T., Darwish, S., Al-Hamaq, A. O., Nasralla, E. A. and Abdul-Ghani, M. (2013). Obesity index that better predict metabolic syndrome: body mass index, waist circumference, waist hip ratio, or waist height ratio. *Journal of obesity*, 2013.
- Benotti, P. N. and Forse, R. A. (1995). The role of gastric surgery in the multidisciplinary management of severe obesity. *The American Journal of Surgery*, 169(3), 361-367.
- Bernini, P., Bertini, I., Luchinat, C., Nincheri, P., Staderini, S. and Turano, P. (2011). Standard operating procedures for pre-analytical handling of blood and urine for metabolomic studies and biobanks. *Journal of biomolecular NMR*, 49(3-4), 231-243.
- Billat, V. L., Sirvent, P., Py, G., Koralsztein, J. P. and Mercier, J. (2003). The concept of maximal lactate steady state. *Sports medicine*, 33(6), 407-426.
- Bird, S. R., Goebel, C., Burke, L. M. and Greaves, R. F. (2016). Doping in sport and exercise: anabolic, ergogenic, health and clinical issues. *Annals of clinical biochemistry*, 53(2), 196-221.
- Blue, J. G. and Lombardo, J. A. (1999). Steroids and steroid-like compounds. *Clinics in Sports Medicine*, 18(3), 667-689.
- Bogdanova, O. V., Kanekar, S., D'Anci, K. E. and Renshaw, P. F. (2013). Factors influencing behaviour in the forced swim test. *Physiology & behaviour*, 118, 227-239.
- Bollard, M. E., Stanley, E. G., Lindon, J. C., Nicholson, J. K. and Holmes, E. (2005). NMR-based metabolomic approaches for evaluating physiological influences on biofluid composition. *NMR in Biomedicine: An International Journal Devoted to the Development and Application of Magnetic Resonance In vivo*, 18(3), 143-162.
- Boussouar, F. and Benahmed, M. (2004). Lactate and energy metabolism in male germ cells. *Trends in Endocrinology & Metabolism*, 15(7), 345-350.
- Brancaccio, P., Lippi, G. and Maffulli, N. (2010). Biochemical markers of muscular damage. *Clinical Chemistry and Laboratory Medicine*, 48(6), 757-767.
- Burke, L. M. (2008). Caffeine and sports performance. *Applied Physiology, Nutrition, and Metabolism*, 33(6), 1319-1334.
- Call, J. A., Voelker, K. A., Wolff, A. V., McMillan, R. P., Evans, N. P., Hulver, M. W. and Grange, R. W. (2008). Endurance capacity in maturing mdx mice is markedly enhanced by combined voluntary wheel running and green tea extract. *Journal of applied physiology*, 105(3), 923-932.
- Calvani, R. I. C. C., Miccheli, A., Capuani, G., Miccheli, A. T., Puccetti, C., Delfini, M. and Mingrone, G. (2010). Gut microbiome-derived metabolites

- characterize a peculiar obese urinary metabotype. *International journal of obesity*, 34(6), 1095.
- Carpentier, A., Olbrechts, N., Vieillevoye, S. and Poortmans, J. R. (2015). β -Alanine supplementation slightly enhances repeated plyometric performance after high-intensity training in humans. *Amino acids*, 47(7), 1479-1483.
- Casuso, R. A., Martínez-Amat, A., Martínez-López, E. J., Camiletti-Moirón, D., Porres, J. M. and Aranda, P. (2013). Ergogenic effects of quercetin supplementation in trained rats. *Journal of the International Society of Sports Nutrition*, 10(1), 3.
- Chan, Y. Y., Lim, K. K., Lim, K. H., Teh, C. H., Kee, C. C., Cheong, S. M. and Ahmad, N. A. (2017). Physical activity and overweight/obesity among Malaysian adults: findings from the 2015 National Health and morbidity survey (NHMS). *BMC public health*, 17(1), 733.
- Chan-Blanco Y, Vaillant F, Perez AM, Reynes M, Brillouet JM, Brat P. The noni fruit (*Morinda citrifolia* L.): A review of agricultural research, nutritional and therapeutic properties. *Journal of food composition and analysis*. 2006 Sep 1;19(6-7):645-54.
- Chen, C. K., Muhamad, A. S. and Ooi, F. K. (2012). Herbs in exercise and sports. *Journal of physiological anthropology*, 31(1), 4.
- Chen, C., Shah, Y. M., Morimura, K., Krausz, K. W., Miyazaki, M., Richardson, T. A. and Gonzalez, F. J. (2008). Metabolomics reveals that hepatic stearoyl-CoA desaturase 1 downregulation exacerbates inflammation and acute colitis. *Cell metabolism*, 7(2), 135-147.
- Choi, Y. H., Tapias, E. C., Kim, H. K., Lefeber, A. W., Erkelens, C., Verhoeven, J. T. J. and Verpoorte, R. (2004). Metabolic discrimination of *Catharanthus roseus* leaves infected by phytoplasma using ¹H-NMR spectroscopy and multivariate data analysis. *Plant Physiology*, 135(4), 2398-2410.
- Chung, H. H., Lee, K. S. and Cheng, J. T. (2013). Decrease of obesity by allantoin via imidazoline I1-receptor activation in high fat diet-fed mice. *Evidence-Based Complementary and Alternative Medicine*, 2013.
- Colditz, G. A., Willett, W. C., Rotnitzky, A. and Manson, J. E. (1995). Weight gain as a risk factor for clinical diabetes mellitus in women. *Annals of internal medicine*, 122(7), 481-486.
- Colquitt, J. L., Picot, J., Loveman, E. and Clegg, A. J. (2009). Surgery for obesity. *Cochrane database of systematic reviews*, (2).
- Cori, G. T. and Cori, C. F. (1933). Changes in hexosephosphate, glycogen, and lactic acid during contraction and recovery of mammalian muscle. *Journal of Biological Chemistry*, 99(2), 493-505.

- Coyle, E. F. (1992). Carbohydrate supplementation during exercise. *The Journal of nutrition*, 122(suppl_3), 788-795.
- Coyle, E. F. (1995). Substrate utilization during exercise in active people. *The American journal of clinical nutrition*, 61(4), 968S-979S.
- Craig, S. A. (2004). Betaine in human nutrition. *The American journal of clinical nutrition*, 80(3), 539-549.
- Cupisti, A., Meola, M., D'Alessandro, C., Bernabini, G., Pasquali, E., Carpi, A. and Barsotti, G. (2007). Insulin resistance and low urinary citrate excretion in calcium stone formers. *Biomedicine & Pharmacotherapy*, 61(1), 86-90.
- D'angelo, L., Grimaldi, R., Caravaggi, M., Marcoli, M., Perucca, E., Lecchini, S. and Crema, A. (1986). A double-blind, placebo-controlled clinical study on the effect of a standardized ginseng extract on psychomotor performance in healthy volunteers. *Journal of ethnopharmacology*, 16(1), 15-22.
- Dixon, A. R., McMillen, H. and Etkin, N. L. (1999). Ferment this: the transformation of Noni, a traditional Polynesian medicine (*Morinda citrifolia*, *Rubiaceae*). *Economic botany*, 53(1), 51-68.
- Dona, A. C., Kyriakides, M., Scott, F., Shephard, E. A., Varshavi, D., Veselkov, K. and Everett, J. R. (2016). A guide to the identification of metabolites in NMR-based metabonomic/metabolomics experiments. *Computational and structural biotechnology journal*, 14, 135-153.
- Duggan, G. E., Hittel, D. S., Hughey, C. C., Weljie, A., Vogel, H. J. and Shearer, J. (2011). Differentiating short-and long-term effects of diet in the obese mouse using ¹H-nuclear magnetic resonance metabolomics. *Diabetes, Obesity and Metabolism*, 13(9), 859-862.
- Dulloo, A. G., Duret, C., Rohrer, D., Girardier, L., Mensi, N., Fathi, M. and Vandermander, J. (1999). Efficacy of a green tea extract rich in catechin polyphenols and caffeine in increasing 24-h energy expenditure and fat oxidation in humans—. *The American journal of clinical nutrition*, 70(6), 1040-1045.
- Essig, D., Costill, D. L. and Van Handel, P. J. (1980). Effects of caffeine ingestion on utilization of muscle glycogen and lipid during leg ergometer cycling. *International Journal of Sports Medicine*, 1(02), 86-90.
- Eriksson, L., Byrne, T., Johansson, E., Trygg, J. and Vikström, C. (2013). *Multi- and megavariate data analysis basic principles and applications* (Vol. 1). Umetrics Academy.
- Farooqi, I. S. and O'Rahilly, S. (2006). Genetics of obesity in humans. *Endocrine reviews*, 27(7), 710-718.

- Ferreira, J. C., Rolim, N. P., Bartholomeu, J. B., Gobatto, C. A., Kokubun, E. and Brum, P. C. (2007). Maximal lactate steady state in running mice: effect of exercise training. *Clinical and Experimental Pharmacology and Physiology*, 34(8), 760-765.
- French, S. A., Story, M. and Jeffery, R. W. (2001). Environmental influences on eating and physical activity. *Annual review of public health*, 22(1), 309-335.
- Fukuhara, K., Ohno, A., Ota, Y., Senoo, Y., Maekawa, K., Okuda, H. and Takikawa, O. (2013). NMR-based metabolomics of urine in a mouse model of Alzheimer's disease: identification of oxidative stress biomarkers. *Journal of clinical biochemistry and nutrition*, 52(2), 133-138.
- Gerlach, J. (1996). Native or introduced plant species? *Phelsuma*, 4, 70-74.
- Ghani, Z. D. F. A., Husin, J. M., Ab Rashid, A. H., Shaari, K. and Chik, Z. (2016). Biochemical studies of *Piper betle* L leaf extract on obese treated animal using ¹H-NMR-based metabolomic approach of blood serum samples. *Journal of ethnopharmacology*, 194, 690-697.
- Giovannucci, E. (1995). Insulin and colon cancer. *Cancer Causes & Control*, 6(2), 164-179.
- Gladden, L. B. (2010). Lactate transport and exchange during exercise. *Comprehensive Physiology*, 614-648.
- Gleeson, M., Blannin, A. K., Walsh, N. P., Field, C. N. and Pritchard, J. C. (1998). Effect of exercise-induced muscle damage on the blood lactate response to incremental exercise in humans. *European Journal of Applied Physiology and Occupational Physiology*, 77(3), 292-295.
- Goh, J. and Ladiges, W. (2015). Voluntary wheel running in mice. *Current protocols in mouse biology*, 5(4), 283-290.
- Go, H. K., Rahman, M. M., Kim, G. B., Na, C. S., Song, C. H., Kim, J. S. and Kang, H. S. (2015). Antidiabetic effects of yam (*Dioscorea batatas*) and its active constituent, allantoin, in a rat model of streptozotocin-induced diabetes. *Nutrients*, 7(10), 8532-8544.
- Goldstein, B. (1975). Ginseng: its history, dispersion, and folk tradition. *The American journal of Chinese medicine*, 3(03), 223-234.
- Gooda Sahib, N., Abdul Hamid, A., Saari, N., Abas, F., Pak Dek, M. S. and Rahim, M. (2012). Anti-pancreatic lipase and antioxidant activity of selected tropical herbs. *International Journal of Food Properties*, 15(3), 569-578.
- Grandhi, A., Mujumdar, A. M. and Patwardhan, B. (1994). A comparative pharmacological investigation of Ashwagandha and Ginseng. *Journal of ethnopharmacology*, 44(3), 131-135.

- Kim, H. J., Kim, J. H., Noh, S., Hur, H. J., Sung, M. J., Hwang, J. T. and Yoon, S. H. (2010). Metabolomic analysis of livers and serum from high-fat diet induced obese mice. *Journal of proteome research*, 10(2), 722-731.
- Hermansen, L., Hultman, E. and Saltin, B. (1967). Muscle glycogen during prolonged severe exercise. *Acta Physiologica Scandinavica*, 71(2-3), 129-139.
- Hill, J. O. and Peters, J. C. (1998). Environmental contributions to the obesity epidemic. *Science*, 280(5368), 1371-1374.
- Hiramatsu, T., Imoto, M., Koyano, T. and Umezawa, K. (1993). Induction of normal phenotypes in *ras*-transformed cells by damnacanthol from *Morinda citrifolia*. *Cancer letters*, 73(2-3), 161-166.
- Hirazumi, A., Furusawa, E., Chou, S. C. and Hokama, Y. (1994). Anticancer activity of *Morinda citrifolia* (noni) on intraperitoneally implanted Lewis lung carcinoma in syngeneic mice. In *Proceedings of the Western pharmacology society* (Vol. 37, pp. 145-146).
- Hirvonen, J., Rehunen, S., Rusko, H. and Härkönen, M. (1987). Breakdown of high-energy phosphate compounds and lactate accumulation during short supramaximal exercise. *European journal of applied physiology and occupational physiology*, 56(3), 253-259.
- Hodgson, A. B., Randell, R. K. and Jeukendrup, A. E. (2013). The metabolic and performance effects of caffeine compared to coffee during endurance exercise. *PloS one*, 8(4), e59561.
- Hodgson, A. B., Randell, R. K., Boon, N., Garczarek, U., Mela, D. J., Jeukendrup, A. E. and Jacobs, D. M. (2013). Metabolic response to green tea extract during rest and moderate-intensity exercise. *The Journal of nutritional biochemistry*, 24(1), 325-334.
- Hohl, R., de Oliveira, R. B., Ferraresso, R. L. P., Brenzikofer, R. and Macedo, D. V. (2011). Effect of body weight variation on swimming exercise workload in rats with constant and size-adjusted loads. *Scandinavian Journal of Laboratory Animal Sciences*, 38(3), 145-154.
- Holloszy, J. O. and Coyle, E. F. (1984). Adaptations of skeletal muscle to endurance exercise and their metabolic consequences. *Journal of applied physiology*, 56(4), 831-838.
- Hopkinson, Z. E., Sattar, N., Fleming, R. and Greer, I. A. (1998). Polycystic ovarian syndrome: the metabolic syndrome comes to gynaecology. *BMJ*, 317, 329-32.
- Hortemo, K. H., Munkvik, M., Lunde, P. K. and Sejersted, O. M. (2013). Multiple causes of fatigue during shortening contractions in rat slow twitch skeletal muscle. *PloS one*, 8(8), e71700.

- Hou, Z., Lambert, J. D., Chin, K. V. and Yang, C. S. (2004). Effects of tea polyphenols on signal transduction pathways related to cancer chemoprevention. *Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis*, 555(1-2), 3-19.
- Hu, S. Y. (1976). The genus *Panax* (ginseng) in Chinese medicine. *Economic Botany*, 30(1), 11-28.
- Huang, W. C., Chiu, W. C., Chuang, H. L., Tang, D. W., Lee, Z. M., Wei, L. and Huang, C. C. (2015). Effect of curcumin supplementation on physiological fatigue and physical performance in mice. *Nutrients*, 7(2), 905-921.
- Huszar, D., Lynch, C. A., Fairchild-Huntress, V., Dunmore, J. H., Fang, Q., Berkemeier, L. R. and Smith, F. J. (1997). Targeted disruption of the melanocortin-4 receptor results in obesity in mice. *Cell*, 88(1), 131-141.
- Huvenne, H. and Dubern, B. (2014). Monogenic Forms of Obesity. In *Molecular Mechanisms Underpinning the Development of Obesity* (pp. 9-21). Springer, Cham.
- Iwashita, S., Williams, P., Jabbour, K., Ueda, T., Kobayashi, H., Baier, S. and Flakoll, P. J. (2005). Impact of glutamine supplementation on glucose homeostasis during and after exercise. *Journal of applied physiology*, 99(5), 1858-1865.
- Jabbour, G. and Iancu, H. D. (2017). Acute and chronic exercises: Effect on lipid metabolisms in obese individuals. *Science & Sports*, 32(6), 321-326.
- Jadhav, R. and Puchchakayala, G. (2012). Hypoglycemic and antidiabetic activity of flavonoids: boswellic acid, ellagic acid, quercetin, rutin on streptozotocin-nicotinamide induced type 2 diabetic rats. *Group*, 1, 100g.
- Jain, R. K. and Srivastava, S. D. (1992). Two new anthraquinones in the roots of *Morinda citrifolia*. *Proceedings-National Academy of Sciences India Section A*, 11-11.
- Jambocus, N. G. S., Saari, N., Ismail, A., Khatib, A., Mahomoodally, M. F. and Hamid, A. A. (2016). An investigation into the antiobesity effects of *Morinda citrifolia* L. leaf extract in high fat diet induced obese rats using a H NMR metabolomics approach. *J Diabetes Res* [http://dx. doi. org/10.1155/2016/2391592](http://dx.doi.org/10.1155/2016/2391592).
- Jansson, P. A., Larsson, A., Smith, U. and Lönnroth, P. (1994). Lactate release from the subcutaneous tissue in lean and obese men. *The Journal of clinical investigation*, 93(1), 240-246.
- Jiwajinda, S., Santisopasri, V., Murakami, A., Hirai, N. and Ohigashi, H. (2001). Quassinoids from *Eurycoma longifolia* as plant growth inhibitors. *Phytochemistry*, 58(6), 959-962.

- Jung, J. Y., Lee, H. S., Kang, D. G., Kim, N. S., Cha, M. H., Bang, O. S. and Hwang, G. S. (2011). ¹H-NMR-based metabolomics study of cerebral infarction. *Stroke*, STROKEAHA-110.
- Jung, K., Kim, I. H. and Han, D. (2004). Effect of medicinal plant extracts on forced swimming capacity in mice. *Journal of ethnopharmacology*, 93(1), 75-81.
- Kaaks, R. (1996). Nutrition, hormones, and breast cancer: is insulin the missing link? *Cancer Causes & Control*, 7(6), 605-625.
- Kaaks, R., Lukanova, A. and Kurzer, M. S. (2002). Obesity, endogenous hormones, and endometrial cancer risk: a synthetic review. *Cancer Epidemiology and Prevention Biomarkers*, 11(12), 1531-1543.
- Kamiya, K., Tanaka, Y., Endang, H., Umar, M. and Satake, T. (2004). Chemical constituents of *Morinda citrifolia* fruits inhibit copper-induced low-density lipoprotein oxidation. *Journal of agricultural and food chemistry*, 52(19), 5843-5848.
- Kannel, W. B., Cupples, L. A., Ramaswami, R., Stokes III, J., Kreger, B. E. and Higgins, M. (1991). Regional obesity and risk of cardiovascular disease; the Framingham Study. *Journal of clinical epidemiology*, 44(2), 183-190.
- Katz, A. T. and Sahlin, K. (1988). Regulation of lactic acid production during exercise. *Journal of applied physiology*, 65(2), 509-518.
- Keisler, B. D. and Armsey, T. D. (2006). Caffeine as an ergogenic aid. *Current sports medicine reports*, 5(4), 215-219.
- Kemppainen, J., Fujimoto, T., Kalliokoski, K. K., Viljanen, T., Nuutila, P. and Knuuti, J. (2002). Myocardial and skeletal muscle glucose uptake during exercise in humans. *The Journal of physiology*, 542(2), 403-412.
- Kennedy, G., Spence, V. A., McLaren, M., Hill, A., Underwood, C. and Belch, J. J. (2005). Oxidative stress levels are raised in chronic fatigue syndrome and are associated with clinical symptoms. *Free radical biology and medicine*, 39(5), 584-589.
- Kerksick, C. M., Rasmussen, C. J., Lancaster, S. L., Magu, B., Smith, P., Melton, C. and Kreider, R. B. (2006). The effects of protein and amino acid supplementation on performance and training adaptations during ten weeks of resistance training. *The Journal of Strength & Conditioning Research*, 20(3), 643-653.
- Keys, A., Fidanza, F., Karvonen, M. J., Kimura, N. and Taylor, H. L. (1972). Indices of relative weight and obesity. *Journal of chronic diseases*, 25(6-7), 329-343.
- Khanam, Z., Wen, C. S. and Bhat, I. U. H. (2015). Phytochemical screening and antimicrobial activity of root and stem extracts of wild *Eurycoma*

- longifolia* Jack (Tongkat Ali). *Journal of King Saud University-Science*, 27(1), 23-30.
- Khambalia, A. Z. and Seen, L. S. (2010). Trends in overweight and obese adults in Malaysia (1996–2009): a systematic review. *Obesity reviews*, 11(6), 403-412.
- Kiew, O. K., Singh, R., Sirisinghe, R. G., Ang, B. and Jamalullail, S. (2001). Effects of a herbal ergogenic drink on cycling performance in young cyclists. *Malays J Nutr*, 7, 33-40.
- Kiew, O. F., Singh, R., Sirisinghe, R. G., Suen, A. B., and Jamalullail, S. M. S. (2003). Effects of a herbal drink on cycling endurance performance. *The Malaysian journal of medical sciences: MJMS*, 10(1), 78.
- Kim, S. H., Yang, S. O., Kim, H. S., Kim, Y., Park, T. and Choi, H. K. (2009). ¹H-nuclear magnetic resonance spectroscopy-based metabolic assessment in a rat model of obesity induced by a high-fat diet. *Analytical and bioanalytical chemistry*, 395(4), 1117-1124.
- Kimball, S. R. and Jefferson, L. S. (2001). Regulation of protein synthesis by branched-chain amino acids. *Current Opinion in Clinical Nutrition & Metabolic Care*, 4(1), 39-43.
- Kregel, K. C., Allen, D. L., Booth, F. W., Fleshner, M. R., Henriksen, E. J., Musch, T. I. and Sheriff, D. D. (2006). Resource book for the design of animal exercise protocols. *American Physiological Society*, 152.
- Krishnaiah, D., Nithyanandam, R. and Sarbatly, R. (2012). Phytochemical Constituents and Activities of *Morinda citrifolia* L. In *Phytochemicals-A Global Perspective of Their Role in Nutrition and Health*. InTech.
- Kulakowski, E. C. and Maturo, J. (1984). Hypoglycemic properties of taurine: not mediated by enhanced insulin release. *Biochemical pharmacology*, 33(18), 2835-2838.
- Kume, S., Yamato, M., Tamura, Y., Jin, G., Nakano, M., Miyashige, Y. and Yamano, E. (2015). Potential biomarkers of fatigue identified by plasma metabolome analysis in rats. *PLoS one*, 10(3), e0120106.
- Kurien, B. T., Everds, N. E. and Scofield, R. H. (2004). Experimental animal urine collection: a review. *Laboratory animals*, 38(4), 333-361.
- Lännergren, J., & Westerblad, HA. (1991). Force decline due to fatigue and intracellular acidification in isolated fibres from mouse skeletal muscle. *The Journal of Physiology*, 434(1), 307-322.
- Leal, E. C. P., Lopes-Martins, R. Á. B., Frigo, L., De Marchi, T., Rossi, R. P., De Godoi, V. and de Valls Corsetti, F. (2010). Effects of low-level laser therapy (LLLT) in the development of exercise-induced skeletal muscle fatigue and changes in biochemical markers related to postexercise recovery. *Journal of orthopaedic & sports physical therapy*, 40(8), 524-532.

- Lee, S. P., Mar, G. Y. and Ng, L. T. (2009). Effects of tocotrienol-rich fraction on exercise endurance capacity and oxidative stress in forced swimming rats. *European journal of applied physiology*, 107(5), 587-595.
- Leibson, C. L., Williamson, D. F., Melton, L. J., Palumbo, P. J., Smith, S. A., Ransom, J. E. and Narayan, K. V. (2001). Temporal trends in BMI among adults with diabetes. *Diabetes care*, 24(9), 1584-1589.
- Lever, M. and Slow, S. (2010). The clinical significance of betaine, an osmolyte with a key role in methyl group metabolism. *Clinical biochemistry*, 43(9), 732-744.
- Llorach, R., Garrido, I., Monagas, M., Urpi-Sarda, M., Tulipani, S., Bartolome, B. and Andres-Lacueva, C. (2010). Metabolomics study of human urinary metabolome modifications after intake of almond (*Prunus dulcis* (Mill.) DA Webb) skin polyphenols. *Journal of proteome research*, 9(11), 5859-5867.
- Li Li, Z. Y., He, P., Sun, H. F., Qin, X. M. and Du, G. H. (2014). ¹H NMR based metabolomic study of the antifatigue effect of *Astragali Radix*. *Molecular BioSystems*, 10(11), 3022-3030.
- Lieberman, H. R. (2001). The effects of ginseng, ephedrine, and caffeine on cognitive performance, mood and energy. *Nutrition reviews*, 59(4), 91-102.
- Liland, K. H. (2011). Multivariate methods in metabolomics—from pre-processing to dimension reduction and statistical analysis. *TrAC Trends in Analytical Chemistry*, 30(6), 827-841.
- Lim, K., Yoshioka, M., Kikuzato, S., Kiyonaga, A., Tanaka, H., Shindo, M. and Suzuki, M. (1997). Dietary red pepper ingestion increases carbohydrate oxidation at rest and during exercise in runners. *Medicine and Science in Sports and Exercise*, 29(3), 355-361.
- Lim, W., Hong, S., Nelesen, R. and Dimsdale, J. E. (2005). The association of obesity, cytokine levels, and depressive symptoms with diverse measures of fatigue in healthy subjects. *Archives of internal medicine*, 165(8), 910-915.
- Liu, J. M., Haroun-Bouhedja, FE. and Boison-Vidal, C. (2000). Analysis of the *in vitro* inhibition of mammary adenocarcinoma cell adhesion by sulphated polysaccharides. *Anticancer research*, 20(5/A), 3265-3272.
- Liu, L., Wu, X., Zhang, B., Yang, W., Li, D., Dong, Y. and Chen, Q. (2017). Protective effects of tea polyphenols on exhaustive exercise-induced fatigue, inflammation and tissue damage. *Food & nutrition research*, 61(1), 1333390.
- Liu, G., Yang, G., Fang, T., Cai, Y., Wu, C., Wang, J. and Chen, X. (2014). NMR-based metabolomic studies reveal changes in biochemical profile of

urine and plasma from rats fed with sweet potato fiber or sweet potato residue. *RSC advances*, 4(45), 23749-23758.

Lv, L., Chen, H., Ho, C. T. and Sang, S. (2011). Chemical components of the roots of Noni (*Morinda citrifolia*) and their cytotoxic effects. *Fitoterapia*, 82(4), 704-708.

Ma, G. D., Chiu, C. H., Hsu, Y. J., Hou, C. W., Chen, Y. M. and Huang, C. C. (2017). Changbai Mountain ginseng (*Panax ginseng* CA Mey) extract supplementation improves exercise performance and energy utilization and decreases fatigue-associated parameters in mice. *Molecules*, 22(2), 237.

Malaguti, M., Angeloni, C. and Hrelia, S. (2013). Polyphenols in exercise performance and prevention of exercise-induced muscle damage. *Oxidative medicine and cellular longevity*, 2013.

Maravelias, C., Dona, A., Stefanidou, M. and Spiliopoulou, C. (2005). Adverse effects of anabolic steroids in athletes: a constant threat. *Toxicology letters*, 158(3), 167-175.

Marconi, C., Sassi, G., Carpinelli, A. and Cerretelli, P. (1985). Effects of L-carnitine loading on the aerobic and anaerobic performance of endurance athletes. *European journal of applied physiology and occupational physiology*, 54(2), 131-135.

Maughan, R. J., King, D. S. and Lea, T. (2004). Dietary supplements. *Journal of sports sciences*, 22(1), 95-113.

McArdle, W. D., Katch, F. I. and Katch, V. L. (2006). *Essentials of exercise physiology*. Lippincott Williams & Wilkins.

Melzer, K. (2011). Carbohydrate and fat utilization during rest and physical activity. *e-SPEN, the European e-Journal of Clinical Nutrition and Metabolism*, 6(2), e45-e52.

Mena, P., Maynar, M., Gutierrez, J. M., Maynar, J., Timon, J. and Campillo, J. E. (1991). Erythrocyte free radical scavenger enzymes in bicycle professional racers. Adaptation to training. *International journal of sports medicine*, 12(06), 563-566.

Miccheli, A., Marini, F., Capuani, G., Micheli, A. T., Delfini, M., Di Cocco, M. E. and Spataro, A. (2009). The influence of a sports drink on the postexercise metabolism of elite athletes as investigated by NMR-based metabolomics. *Journal of the American College of Nutrition*, 28(5), 553-564.

Mohtar, M., Shaari, K., Ali, N. A. M. and Ali, A. M. (1998). Antimicrobial activity of selected Malaysian plants against micro-organisms related to skin infections. *Journal of Tropical Forest Products*, 4, 199-206.

Mu, C., Yang, Y., Luo, Z. and Zhu, W. (2015). Metabolomic analysis reveals distinct profiles in the plasma and urine of rats fed a high-protein diet. *Amino acids*, 47(6), 1225-1238.

- Muhamad, A. S., Keong, C. C., Kiew, O. F., Abdullah, M. R. and Lam, C. K. (2010). Effects of *Eurycoma longifolia* Jack supplementation on recreational athletes' endurance running capacity and physiological responses in the heat. *Int J Appl Sports Sci*, 22(2), 1-19.
- Mukherjee, M. (2003). Human digestive and metabolic lipases—a brief review. *Journal of Molecular Catalysis B: Enzymatic*, 22(5-6), 369-376.
- Mustaffa, F., Indurkar, J., Ali, N. I. M., Hanapi, A., Shah, M., Ismail, S. and Mansor, S. M. (2011). A review of Malaysian medicinal plants with potential antidiabetic activity. *Journal of Pharmacy Research*, 4(11), 4217-4224.
- Murase, T., Haramizu, S., Shimotoyodome, A., Tokimitsu, I. and Hase, T. (2006). Green tea extract improves running endurance in mice by stimulating lipid utilization during exercise. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 290(6), R1550-R1556.
- Murase, T., Haramizu, S., Shimotoyodome, A., Nagasawa, A. and Tokimitsu, I. (2005). Green tea extract improves endurance capacity and increases muscle lipid oxidation in mice. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 288(3), R708-R715.
- Nayak, B. S., Marshall, J. R., Isitor, G. and Adogwa, A. (2011). Hypoglycemic and hepatoprotective activity of fermented fruit juice of *Morinda citrifolia* (Noni) in diabetic rats. *Evidence-Based Complementary and Alternative Medicine*, 2011.
- Nayak, B. S., Sandiford, S. and Maxwell, A. (2009). Evaluation of the wound-healing activity of ethanolic extract of *Morinda citrifolia* L. leaf. *Evidence-based complementary and alternative medicine*, 6(3), 351-356.
- Neel, J. V. (1962). Diabetes mellitus: a "thrifty" genotype rendered detrimental by "progress"? *American journal of human genetics*, 14(4), 353.
- Newcomer, B. R., Larson-Meyer, D. E., Hunter, G. R. and Weinsier, R. L. (2001). Skeletal muscle metabolism in overweight and post-overweight women: an isometric exercise study using ³¹P magnetic resonance spectroscopy. *International journal of obesity*, 25(9), 1309.
- Nieman, D. C. (2003). Current perspective on exercise immunology. *Current sports medicine reports*, 2(5), 239-242.
- Nualsanit, T., Rojanapanthu, P., Gritsanapan, W., Lee, S. H., Lawson, D. and Baek, S. J. (2012). Damnacanthol, a noni component, exhibits antitumorigenic activity in human colorectal cancer cells. *The Journal of nutritional biochemistry*, 23(8), 915-923.
- Oh, J. K., Shin, Y. O., Yoon, J. H., Kim, S. H., Shin, H. C. and Hwang, H. J. (2010). Effect of supplementation with *Ecklonia cava* polyphenol on

- endurance performance of college students. *International journal of sport nutrition and exercise metabolism*, 20(1), 72-79.
- Oliynyk, S. and Oh, S. (2013). Actoprotective effect of ginseng: improving mental and physical performance. *Journal of ginseng research*, 37(2), 144.
- Osman, W. N. W., & Mohamed, S. (2018). Standardized *Morinda citrifolia* L. and *Morinda elliptica* L. leaf extracts alleviated fatigue by improving glycogen storage and lipid/carbohydrate metabolism. *Phytotherapy Research*, 32(10), 2078-2085.
- Pandareesh, M. D. and Anand, T. (2013). Ergogenic effect of dietary L-carnitine and fat supplementation against exercise induced physical fatigue in Wistar rats. *Journal of physiology and biochemistry*, 69(4), 799-809.
- Pawlus, A. D. and Kinghorn, A. D. (2007). Review of the ethnobotany, chemistry, biological activity and safety of the botanical dietary supplement *Morinda citrifolia* (noni). *Journal of Pharmacy and pharmacology*, 59(12), 1587-1609.
- Palou, A. and Bonet, M. L. (2007). Controlling lipogenesis and thermogenesis and the use of ergogenic aids for weight control. *Novel food ingredients for weight control*, 58.
- Palu, A. K., Kim, A. H., West, B. J., Deng, S., Jensen, J. and White, L. (2008). The effects of *Morinda citrifolia* L (noni) on the immune system: its molecular mechanisms of action. *Journal of Ethnopharmacology*, 115(3), 502-506.
- Philp, A., Macdonald, A. L. and Watt, P. W. (2005). Lactate—a signal coordinating cell and systemic function. *Journal of Experimental Biology*, 208(24), 4561-4575.
- Pillard, F., Van Wymelbeke, V., Garrigue, E., Moro, C., Crampes, F., Guillard, J. C. and Brondel, L. (2010). Lipid oxidation in overweight men after exercise and food intake. *Metabolism*, 59(2), 267-274.
- Porsolt, R. D., Anton, G., Blavet, N. and Jalfre, M. (1978). Behavioural despair in rats: a new model sensitive to antidepressant treatments. *European journal of pharmacology*, 47(4), 379-391.
- Powers, S. K. and Jackson, M. J. (2008). Exercise-induced oxidative stress: cellular mechanisms and impact on muscle force production. *Physiological reviews*, 88(4), 1243-1276.
- Powers, S. K., Talbert, E. E. and Adhihetty, P. J. (2011). Reactive oxygen and nitrogen species as intracellular signals in skeletal muscle. *The Journal of physiology*, 589(9), 2129-2138.
- Radak, Z., Chung, H. Y., Koltai, E., Taylor, A. W. and Goto, S. (2008). Exercise, oxidative stress and hormesis. *Ageing research reviews*, 7(1), 34-42.
- Rahman, H. A., Sahib, N. G., Saari, N., Abas, F., Ismail, A., Mumtaz, M. W. and Hamid, A. A. (2017). Anti-obesity effect of ethanolic extract from

- Cosmos caudatus* Kunth leaf in lean rats fed a high fat diet. *BMC complementary and alternative medicine*, 17(1), 122.
- Rains, T. M., Agarwal, S. and Maki, K. C. (2011). Antiobesity effects of green tea catechins: a mechanistic review. *The Journal of nutritional biochemistry*, 22(1), 1-7.
- Rao, U. M. and Subramanian, S. (2009). Biochemical evaluation of antihyperglycemic and antioxidative effects of *Morinda citrifolia* fruit extract studied in streptozotocin-induced diabetic rats. *Medicinal chemistry research*, 18(6), 433-446.
- Rebouche, C. J. and Paulson, D. J. (1986). Carnitine metabolism and function in humans. *Annual review of nutrition*, 6(1), 41-66.
- Rezzi, S., Ramadan, Z., Martin, F. P. J., Fay, L. B., Van Bladeren, P., Lindon, J. C. and Kochhar, S. (2007). Human metabolic phenotypes link directly to specific dietary preferences in healthy individuals. *Journal of proteome research*, 6(11), 4469-4477.
- Fitts, R. H. (1994). Cellular mechanisms of muscle fatigue. *Physiological reviews*, 74(1), 49-94.
- Robergs, R. A., Ghiasvand, F. and Parker, D. (2004). Biochemistry of exercise-induced metabolic acidosis. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 287(3), R502-R516.
- Ross, I. A. (2001). Medical plants of the world. Chemical constituents, traditional and modern medical uses. New Jersey: *Humana Press*.
- Rosenbaum, M., Knight, R. and Leibel, R. L. (2015). The gut microbiota in human energy homeostasis and obesity. *Trends in Endocrinology & Metabolism*, 26(9), 493-501.
- Saito, S. T., Gosmann, G., Pungartnik, C. and Brendel, M. (2009). Green tea extract-patents and diversity of uses. *Recent patents on food, nutrition & agriculture*, 1(3), 203-215.
- Sajak, A. A. B., Mediani, A., Ismail, A. and Abas, F. (2017). Metabolite Variation in Lean and Obese Streptozotocin (STZ)-Induced Diabetic Rats via ¹H NMR-Based Metabolomics Approach. *Applied biochemistry and biotechnology*, 182(2), 653-668.
- Sahlin, K. (1986). Muscle fatigue and lactic acid accumulation. *Acta physiologica Scandinavica. Supplementum*, 556, 83-91.
- Salazar, D. E. and Corcoran, G. B. (1988). Predicting creatinine clearance and renal drug clearance in obese patients from estimated fat-free body mass. *The American journal of medicine*, 84(6), 1053-1060.
- Saludes, J. P., Garson, M. J., Franzblau, S. G. and Aguinaldo, A. M. (2002). Antitubercular constituents from the hexane fraction of *Morinda citrifolia* Linn (Rubiaceae). *Phytotherapy Research: An International*

Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives, 16(7), 683-685.

Saludes JP, Garson MJ. And Franzblau SG. (2005): Chemical and antitubercular studies of *Morinda citrifolia* Linn. (noni) fruits. *Sci Augustin* (Manila) 2:42-64.

Sang, S., Cheng, X., Zhu, N., Stark, R. E., Badmaev, V., Ghai, G. and Ho, C. T. (2001). Flavonol glycosides and novel iridoid glycoside from the leaves of *Morinda citrifolia*. *Journal of agricultural and food chemistry*, 49(9), 4478-4481.

Satoh, H. and Sperelakis, N. (1998). Review of some actions of taurine on ion channels of cardiac muscle cells and others. *General Pharmacology: The Vascular System*, 30(4), 451-463.

Satoh, H. (1994). Cardioprotective actions of taurine against intracellular and extracellular calcium-induced effects. In *Taurine in Health and Disease* (pp. 181-196). Springer, Boston, MA.

Saugy, M., Robinson, N., Saudan, C., Baume, N., Avois, L. and Mangin, P. (2006). Human growth hormone doping in sport. *British journal of sports medicine*, 40(suppl 1), i35-i39.

Serkova, N. J., Jackman, M., Brown, J. L., Liu, T., Hirose, R., Roberts, J. P. and Niemann, C. U. (2006). Metabolic profiling of livers and blood from obese Zucker rats. *Journal of hepatology*, 44(5), 956-962.

Shalan, N. A. A. M., Mustapha, N. M. and Mohamed, S. (2016). *Morinda citrifolia* leaf enhanced performance by improving angiogenesis, mitochondrial biogenesis, antioxidant, anti-inflammatory & stress responses. *Food chemistry*, 212, 443-452. Shetty AK, Rashmi R, Rajan MGR, Sambaiah K, Salimath PV: Antidiabetic influence of quercetin in streptozotocin-induced diabetic rats. *Nutr Res* 2004, 24:373-381.

Shearer, J., Duggan, G., Weljie, A., Hittel, D. S., Wasserman, D. H. and Vogel, H. J. (2008). Metabolomic profiling of dietary-induced insulin resistance in the high fat-fed C57BL/6J mouse. *Diabetes, Obesity and Metabolism*, 10(10), 950-958.

Shovic, A. C. and Whistler, W. A. (2001). Food sources of provitamin A and vitamin C in the American Pacific. *Trop. Sci*, 41, 199-202.

Sickmann, H. M., Waagepetersen, H. S., Schousboe, A., Benie, A. J. and Bouman, S. D. (2010). Obesity and type 2 diabetes in rats are associated with altered brain glycogen and amino-acid homeostasis. *Journal of Cerebral Blood Flow & Metabolism*, 30(8), 1527-1537.

Singh, J. and Tiwari, R. D. (1976). Flavone glycosides from the flowers of *Morinda citrifolia*. *Journal*.

- Silver, M. D. (2001). Use of ergogenic aids by athletes. *JAAOS-Journal of the American Academy of Orthopaedic Surgeons*, 9(1), 61-70.
- Soga, T., Sugimoto, M., Honma, M., Mori, M., Igarashi, K., Kashikura, K. and Otsuka, M. (2011). Serum metabolomics reveals γ -glutamyl dipeptides as biomarkers for discrimination among different forms of liver disease. *Journal of hepatology*, 55(4), 896-905.
- Solomon, N. and Udall, C. (1999). *The Noni Phenomenon*. Direct Source Publishing.
- Song, X., Wang, J., Wang, P., Tian, N., Yang, M. and Kong, L. (2013). ^1H NMR-based metabolomics approach to evaluate the effect of Xue-Fu-Zhu-Yu decoction on hyperlipidemia rats induced by high-fat diet. *Journal of pharmaceutical and biomedical analysis*, 78, 202-210.
- Sousa, M., Teixeira, V. H. and Soares, J. (2014). Dietary strategies to recover from exercise-induced muscle damage. *International journal of food sciences and nutrition*, 65(2), 151-163.
- Souto, G., Donapetry, C., Calvino, J. and Adeva, M. M. (2011). Metabolic acidosis-induced insulin resistance and cardiovascular risk. *Metabolic Syndrome and Related Disorders*, 9(4), 247-253.
- Speakman, J. R. (2008). Thrifty genes for obesity, an attractive but flawed idea, and an alternative perspective: the 'drifty gene' hypothesis. *International journal of obesity*, 32(11), 1611.
- Spradley, B. D., Crowley, K. R., Tai, C. Y., Kendall, K. L., Fukuda, D. H., Esposito, E. N. and Moon, J. R. (2012). Ingesting a pre-workout supplement containing caffeine, B-vitamins, amino acids, creatine, and beta-alanine before exercise delays fatigue while improving reaction time and muscular endurance. *Nutrition & metabolism*, 9(1), 28.
- Spriet, L. L. and Watt, M. J. (2003). Regulatory mechanisms in the interaction between carbohydrate and lipid oxidation during exercise. *Acta physiologica scandinavica*, 178(4), 443-452.
- Stoner, G. D., Wang, L. S., Seguin, C., Rocha, C., Stoner, K., Chiu, S. and Kinghorn, A. D. (2010). Multiple berry types prevent N-nitrosomethylbenzylamine-induced oesophageal cancer in rats. *Pharmaceutical research*, 27(6), 1138-1145.
- Stunkard, A. J. (1996). Current views on obesity. *The American journal of medicine*, 100(2), 230-236.
- Sung, D. J., Kim, S., Kim, J., An, H. S. and So, W. Y. (2016). Role of L-carnitine in sports performance: Focus on ergogenic aid and antioxidant. *Science & Sports*, 31(4), 177-188.
- Suzuki, T., Morita, M., Kobayashi, Y. and Kamimura, A. (2016). Oral L-citrulline supplementation enhances cycling time trial performance in healthy trained men: Double-blind randomized placebo-controlled 2-way

- crossover study. *Journal of the International Society of Sports Nutrition*, 13(1), 6.
- Swamy, M. S., Sivanna, N., Tamatam, A. and Khanum, F. (2011). Effect of poly phenols in enhancing the swimming capacity of rats. *Functional Foods in Health and disease*, 1(11), 482-491.
- Szumilak, D., Sułowicz, W. and Walatek, B. (1998). Rhabdomyolysis: clinical features, causes, complications and treatment. *Przegląd lekarski*, 55(5), 274-279.
- Tang, T., Muneta, T., Ju, Y. J., Nimura, A., Miyazaki, K., Masuda, H. and Sekiya, I. (2008). Serum keratan sulfate transiently increases in the early stage of osteoarthritis during strenuous running of rats: protective effect of intraarticular hyaluronan injection. *Arthritis research & therapy*, 10(1), R13.
- Teng, Q. (2013). NMR-Based metabolomics. In *Structural Biology* (pp. 311-392). Springer, Boston, MA.
- Tiwari, R. D. and Singh, J. (1977). Structural study of the anthraquinone glycoside from the flowers of *Morinda citrifolia*. *Journal*.
- Tiziani, S., Lopes, V. and Günther, U. L. (2009). Early stage diagnosis of oral cancer using ¹H NMR-based metabolomics. *Neoplasia*, 11(3), 269-276.
- Tsushima, Y., Nishizawa, H., Tochino, Y., Nakatsuji, H., Sekimoto, R., Nagao, H. and Tamura, M. (2013). Uric acid secretion from adipose tissue and its increase in obesity. *Journal of Biological Chemistry*, jbc-M113.
- Unger, R., Kreeger, L. and Christoffel, K. K. (1990). Childhood obesity: medical and familial correlates and age of onset. *Clinical pediatrics*, 29(7), 368-373.
- Van Gaal, L. F., Mertens, I. L. and Christophe, E. (2006). Mechanisms linking obesity with cardiovascular disease. *Nature*, 444(7121), 875.
- Wanchai, A., Armer, J. M. and Stewart, B. R. (2010). Complementary and alternative medicine use among women with breast cancer: a systematic review. *Clinical journal of oncology nursing*, 14(4).
- Wang, J., Qin, X., Chen, Z., Ju, Z., He, W., Tan, Y. and Liu, Y. (2016). Two new anthraquinones with antiviral activities from the barks of *Morinda citrifolia* (Noni). *Phytochemistry Letters*, 15, 13-15.
- Wang, L. C. and Lee, T. F. (1998). Effect of ginseng saponins on exercise performance in non-trained rats. *Planta medica*, 64(02), 130-133.
- Wang, M., Kikuzaki, H., Csiszar, K., Boyd, C. D., Maunakea, A., Fong, S. F. and Ho, C. T. (1999). Novel trisaccharide fatty acid ester identified from the fruits of *Morinda citrifolia* (Noni). *Journal of agricultural and food chemistry*, 47(12), 4880-4882.
- Wang, M. Y., West, B. J., Jensen, C. J., Nowicki, D., Su, C., Palu, A. K. and Anderson, G. (2002). *Morinda citrifolia* (Noni): a literature review and

- recent advances in Noni research. *Acta Pharmacologica Sinica*, 23(12), 1127-1141.
- Wang, S. Y., Huang, W. C., Liu, C. C., Wang, M. F., Ho, C. S., Huang, W. P. and Huang, C. C. (2012). Pumpkin (*Cucurbita moschata*) fruit extract improves physical fatigue and exercise performance in mice. *Molecules*, 17(10), 11864-11876.
- Watford, M. (2008). Glutamine metabolism and function in relation to proline synthesis and the safety of glutamine and proline supplementation. *The Journal of nutrition*, 138(10), 2003S-2007S.
- Weiderpass, E., Partanen, T., Kaaks, R., Vainio, H., Porta, M., Kauppinen, T. and Malats, N. (1998). Occurrence, trends and environmental etiology of pancreatic cancer. *Scandinavian journal of work, environment & health*, 165-174.
- Wheelock, Å. M. and Wheelock, C. E. (2013). Trials and tribulations of 'omics data analysis: assessing quality of SIMCA-based multivariate models using examples from pulmonary medicine. *Molecular bioSystems*, 9(11), 2589-2596.
- Williams, M. H. (2005). Dietary supplements and sports performance: minerals. *Journal of the International Society of Sports Nutrition*, 2(1), 43.
- World Health Organization. Obesity and overweight. 2016. Google Scholar. 2016 Sep.
- World Health Organization. (2011). Waist circumference and waist-hip ratio: report of a WHO expert consultation, Geneva, 8-11 December 2008.
- Won, E. Y., Yoon, M. K., Kim, S. W., Jung, Y., Bae, H. W., Lee, D. and Chi, S. W. (2013). Gender-specific metabolomic profiling of obesity in leptin-deficient ob/ob mice by ¹H NMR spectroscopy. *PloS One*, 8(10), e75998.
- Woolf, K., Bidwell, W. K. and Carlson, A. G. (2008). The effect of caffeine as an ergogenic aid in anaerobic exercise. *International journal of sport nutrition and exercise metabolism*, 18(4), 412-429.
- Wu, R. E., Huang, W. C., Liao, C. C., Chang, Y. K., Kan, N. W. and Huang, C. C. (2013). Resveratrol protects against physical fatigue and improves exercise performance in mice. *Molecules*, 18(4), 4689-4702.
- Wu, G. (2009). Amino acids: metabolism, functions, and nutrition. *Amino acids*, 37(1), 1-17.
- Xie, B., Waters, M. J. and Schirra, H. J. (2012). Investigating potential mechanisms of obesity by metabolomics. *BioMed Research International*, 2012.

- Xu, J., Liu, C., Cai, S., Dong, J., Li, X., Feng, J. and Chen, Z. (2013). Metabolomic profilings of urine and serum from high fat-fed rats via ^1H NMR spectroscopy and pattern recognition. *Applied biochemistry and biotechnology*, 169(4), 1250-1261.
- Yang, Q., Jin, W., Lv, X., Dai, P., Ao, Y., Wu, M. and Yu, L. (2016). Effects of *macamides* on endurance capacity and anti-fatigue property in prolonged swimming mice. *Pharmaceutical biology*, 54(5), 827-834.
- Yazdi, F. T., Clee, S. M. and Meyre, D. (2015). Obesity genetics in mouse and human: back and forth, and back again. *PeerJ*, 3, e856.
- Younos, C., Rolland, A., Fleurentin, J., Lanhers, M. C., Misslin, R. and Mortier, F. (1990). Analgesic and behavioural effects of *Morinda citrifolia*. *Planta Medica*, 56(05), 430-434.
- Zin, Z. M., Abdul-Hamid, A. and Osman, A. (2002). Antioxidative activity of extracts from Mengkudu (*Morinda citrifolia* L.) root, fruit and leaf. *Food Chemistry*, 78(2), 227-231.
- Zin, Z. M., Hamid, A. A., Osman, A. and Saari, N. (2006). Antioxidative activities of chromatographic fractions obtained from root, fruit and leaf of Mengkudu (*Morinda citrifolia* L.). *Food Chemistry*, 94(2), 169-178.
- Zanoli, P., Zavatti, M., Montanari, C. and Baraldi, M. (2009). Influence of *Eurycoma longifolia* on the copulatory activity of sexually sluggish and impotent male rats. *Journal of ethnopharmacology*, 126(2), 308-313.
- Ziemba, A. W., Chmura, J., Kaciuba-Uscilko, H., Nazar, K., Wisnik, P. and Gawronski, W. (1999). Ginseng treatment improves psychomotor performance at rest and during graded exercise in young athletes. *International journal of sport nutrition*, 9(4), 371-377.