



Unraveling the ecotoxicity of deep eutectic solvents using the mixture toxicity theory



Inês P.E. Macário ^{a,1}, Fátima Jesus ^{a,1}, Joana L. Pereira ^{a,*}, Sónia P.M. Ventura ^b, Ana M.M. Gonçalves ^{a,c}, João A.P. Coutinho ^b, Fernando J.M. Gonçalves ^a

^a Department of Biology & CESAM, University of Aveiro, 3810-193 Aveiro, Portugal

^b CICECO - Aveiro Institute of Materials & Department of Chemistry, University of Aveiro, 3810-193 Aveiro, Portugal

^c MARE, Faculty of Science and Technology, University of Coimbra, 3004-517 Coimbra, Portugal

HIGHLIGHTS

- Mixture toxicity theory successfully assesses the ecotoxicity of [N_{xxxx}]Cl-based DES.
- DES toxicity followed the same trend as salts: [N₁₁₁₁]Cl < [N₂₂₂₂]Cl < [N₃₃₃₃]Cl.
- IA model, with specific deviations, adjusted better than CA in 5 out of 6 DES.
- Antagonism was found for [N₁₁₁₁]Cl-based DES and synergism for [N₃₃₃₃]Cl-based DES.
- DES toxicity cannot be predicted based solely on the toxicity of HBA and HBD.

ARTICLE INFO

Article history:

Received 27 June 2018

Received in revised form

28 August 2018

Accepted 29 August 2018

Available online 30 August 2018

Handling Editor: Willie Peijnenburg

Keywords:

Quaternary ammonium salts

Microtox

Concentration addition

Independent action

Synergism

Antagonism

ABSTRACT

The interest on deep eutectic solvents (DES) has been increasing. However, the ecotoxicological profile of DES is scarcely known. Also, despite previous studies showed that DES components dissociate in water, none assessed DES toxicity using the classical and adequate models for mixture toxicity prediction - concentration addition (CA) and independent action (IA). This study evaluates the ecotoxicological profile of DES based on [N₁₁₁₁]Cl, [N₂₂₂₂]Cl and [N₃₃₃₃]Cl as hydrogen bond acceptors (HBA) combined with hydrogen-bond donors (HBD) vis. ethylene glycol and 1-propanol, through the Microtox[®] Acute Toxicity Test. CA and IA with deviations describing synergism/antagonism, dose-ratio and dose-level effects were fitted to the toxicity data. Neither the starting materials nor DES were found hazardous to *Aliivibrio fischeri*, in this specific case agreeing with the claimed “green character” of DES. Among the starting materials, ethylene glycol was the least toxic, whereas [N₃₃₃₃]Cl was the most toxic (30 min-EC₅₀ = 96.49 g L⁻¹ and 0.5456 g L⁻¹, respectively). DES toxicity followed the same trend as observed for the salts: [N₁₁₁₁]Cl-based DES < [N₂₂₂₂]Cl-based DES < [N₃₃₃₃]Cl-based DES. The IA model, with specific deviations, adjusted better in 5 out of 6 DES. Antagonism was observed for [N₁₁₁₁]Cl-based DES, and synergism for [N₃₃₃₃]Cl-based DES and for 1-propanol:[N₂₂₂₂]Cl. The application of the mixture toxicity models represents a breakthrough in the problematic of assessing the toxicity of the countless number of DES that can be created with the same starting materials, since they provide the expected toxicity of any virtual combination between HBA and HBD.

© 2018 Elsevier Ltd. All rights reserved.

1. Introduction

Over the past decades, the design of environmentally friendlier solvents has gained increased attention, driven by global awareness

of environmental pollution. Deep eutectic solvents (DES) arose as a new type of tailor-made and “greener” solvents. They are a mixture of two or three components – starting materials (e.g., quaternary ammonium salts, amides, organic acids, polyalcohols), acting as hydrogen bond acceptors (HBA) and hydrogen bond donors (HBD) to form an eutectic mixture based on hydrogen bonding interactions that holds a melting point much lower than either single component (Dai et al., 2013; Ruß and König, 2012; Zhang et al.,

* Corresponding author.

E-mail address: jpereira@ua.pt (J.L. Pereira).

¹ Both authors contributed equally.

2012). DES are simple to prepare and do not need purification, have low-cost production due to the low cost of starting materials, and have been showing good biocompatibility (Hayyan et al., 2013b; Jhong et al., 2009; Singh et al., 2012).

The DES tailor-made character makes them appealing for several applications within chemistry (Radošević et al., 2015), and pharmaceutical/biomedical (Mbous et al., 2017a) fields. Following their production and use, they can reach aquatic systems, causing toxicity to aquatic biota, hence the recent interest on the toxicity profile of DES. (Eco)toxicity of several [Chol]Cl and phosphonium-based DES was assessed through a variety of organisms (Cardellini et al., 2015; de Moraes et al., 2015; Hayyan et al., 2015, 2013a; 2013b; Huang et al., 2014; Juneidi et al., 2016; Radošević et al., 2015; Wen et al., 2015; Zhao et al., 2015), as well as through several cell lines (Hayyan et al., 2016, 2015; Mbous et al., 2017b; Paiva et al., 2014; Radošević et al., 2015).

The assumption of DES low toxicity derives from the low toxicity of their individual components. However, DES components may interact (Hayyan et al., 2013b) depending on their nature and proportion within a DES. According to Hayyan et al. (2013a) and de Moraes et al. (2015) the behavior of DES should be closer to that of a single compound, rather than a mixture. However, in presence of water, the assumption that DES structure remains undissociated has been challenged, since water competes for hydrogen bonding (Dai et al., 2015; Hammond et al., 2017; Shah and Mjalli, 2014; Zhekenov et al., 2017). As diluted by more than 50% in water, DES components indeed dissociate (Dai et al., 2015; Hammond et al., 2017), due to both the disruption of the hydrogen bonding network and to the solvation of the single components by water (Passos et al., 2016). The prevalence of synergistic effects between the starting materials was already suggested (Hayyan et al., 2015, 2013a), but still not quantified. As most (eco)toxicity studies test DES as aqueous solutions comprising two (or more) starting materials, it makes sense to analyse and predict DES toxicity using classical models describing mixtures behavior – concentration addition (CA) and independent action (IA). These models (see material and methods for details) are commonly used to predict the toxicity of a mixture based on the toxicities of its individual components (Backhaus and Faust, 2012).

So far, no rules could be established to predict the (eco)toxicity of DES, which prevents an efficient assessment (mandatory before marketing authorization in most countries) of the countless variations that can be produced for these compounds. The toxic responses seem to depend on DES composition, concentration, and organism tested. This study aims to contribute to the evaluation of the ecotoxicological profile of non-typical DES (e.g. not based on cholinium chloride), using the principles of mixture toxicity theory. The salts tetramethylammonium chloride ([N₁₁₁₁]Cl), tetraethylammonium chloride ([N₂₂₂₂]Cl) and tetrapropylammonium chloride ([N₃₃₃₃]Cl) were used as HBA, whereas ethylene glycol and 1-propanol were used as HBD in the preparation of DES. The marine bacteria *Aliivibrio fischeri* was used as the biological model, and the inhibition of its luminescence was used as the target endpoint through the Microtox[®] Acute Toxicity Test, a standard, sensitive and widely used ecotoxicological test (Johnson, 2005).

2. Material and methods

2.1. Preparation of DES

The following chemical compounds were used for the preparation of DES: ethylene glycol (99.5% purity) and 1-propanol (99.5% purity), as HBDs; [N₁₁₁₁]Cl (97% purity), [N₂₂₂₂]Cl (98% purity) and [N₃₃₃₃]Cl (98% purity), as HBAs, at 4 M ratios (HBD:HBA - 1:2, 1:1, 2:1 and 4:1). All chemicals were purchased from Sigma-Aldrich,

except 1-propanol, which was purchased from Merck. For the preparation of DES, HBDs and HBAs were added gravimetrically to a closed vial and heated in a heat block with constant agitation with a known volume of water. As a transparent liquid formed, the mixture was cooled down to room temperature. Water content of the starting materials was determined by Karl Fischer titration, and was taken into account in the calculation of EC_x values.

2.2. Microtox[®] Toxicity Test assay

DES were tested using the Microtox[®] Toxicity Test (see Johnson, 2005), through the inhibition of the luminescence in the marine bacteria *A. fischeri*. Briefly, the bacteria were exposed to a range of diluted solutions (from 0 to 81.9%) of each DES or each starting material prepared in ultra-pure water. The concentrations of these stocks were adjusted to capture full breadth concentration-response curves in the tests. These standard dilutions were made following the manufacturer protocol with the supplied diluent reagent (2% NaCl solution in water), after a first step of osmotic pressure adjustment of the samples to 2% NaCl using the proprietary osmotic adjusting solution; this ensures that the test is run at optimal osmotic conditions for the bacteria, i.e. 2% NaCl (Johnson, 2005). Measurements of the luminescent output of the bacteria were recorded and compared with the light output of a control sample (bacteria in blank diluent) after 5, 15, and 30 min of exposure. These data were used to estimate the concentrations eliciting 50%, 20% and 10% luminescence inhibition (EC₅₀, EC₂₀ and EC₁₀, respectively) and the corresponding 95% confidence intervals through a non-linear regression, using the least-squares method to fit the data to the logistic equation. Despite calculated directly based on tested mass-per-volume concentrations, these EC_x benchmarks estimated for starting materials and DES, are potentially useful information to include in risk assessment databases as required by different regulatory frameworks (e.g. European Union REACH Directive EC 1907/2006).

2.3. Mixtures toxicity assessment and modelling

The data were analysed on the basis of the mixture toxicity theory, which stems from the application of the baseline CA and IA models.

The CA model relies on the assumption that the components of a mixture have the same site of action (bind to the same biological receptor) in organisms. Thus, each component acts as dilutions of each other (Altenburger et al., 2000). This means that the effect can be predicted by replacing one component totally or in part by the equieffective amount of another component. For a mixture of *n* components, the CA concept can be mathematically expressed as:

$$\sum_{i=1}^n \frac{c_i}{ECx_i} = 1 \quad (1)$$

where *c_i* denotes the concentration (or dose) of the *i*th component which causes *x*% of total effect and EC_{*x*}*i* represents the concentration of that substance causing *x*% of effect if applied singly (Backhaus and Faust, 2012). Each fraction *c_i*/EC_{*x*}*i* is termed a “toxic unit” (TU) and represents the concentration of the *i*th component in the mixture scaled for its relative potency (Backhaus and Faust, 2012).

The model IA assumes that mixture components act on different sites of action of an organism (Altenburger et al., 2000). Thus, the effect of a component is independent of the other. For a mixture of *n* components, the expected effect can be predicted according to the joint probability of statistically independent events following the

equation:

$$E(C_{mix}) = 1 - \prod_{i=1}^n (1 - E(C_i)) \quad (2)$$

where $E(C_{mix})$ denotes the expected effect of the mixture (scaled 0–1); $E(C_i)$ denotes the effect of component i if applied singly, and C_i is the concentration of that component in the mixture (Altenburger et al., 2000).

The application of both models is limited by the assumption that all mixture components affect the same endpoint (e.g. mortality, growth, reproduction) and that there are no interactions between the mixture components, concerning uptake, distribution or metabolization. By characterizing the deviation of experimental data to these reference models, one can assess interactions between mixture components and define corresponding synergic and antagonistic behavior. This was done in the present study as follows.

The experimental responses to all combinations between starting materials within each DES (different molar ratios between starting materials were pooled) along with the responses (bioluminescence inhibition) to each corresponding starting material dosed singly were used to feed mixtures response surface modeling for each DES. First, inhibitory experimental records were fitted to the nonlinear decay model representing toxicological concentration-response curves (Barata et al., 2006). Significant fitting was always achieved, with model accuracy being assessed through adjusted coefficient of determination (r^2) and residual distribution (Quinn and Keough, 2002). The significance of regressions and their regression coefficients were determined by the F-test of overall significance and the t -test, respectively. Then, the joint action within each DES was assessed by fitting the experimental data into the reference mixture models of CA and IA, as well as to CA and IA models added the deviation functions allowing the assessment of synergic/antagonist effect, dose-level and dose-ratio dependent effects; such effects are denoted by two parameters defining the functional form of the deviation pattern, a and b , as described in Jonker et al. (2005). Mixture toxicity modelling and analysis was run in a customised MS[®] Excel[®] spreadsheet (ToxCalcMix, version 1.0, last rev. 19/03/2018; ©Prof. António J.A. Nogueira). Contour plots showing the interactive effects between HBA and HBD within each DES towards *A. fischeri* were created to facilitate interpretation. Several statistic parameters were determined to allow comparison of the model-deviation fit (Jonker et al., 2005; Motulsky and Christopoulos, 2004; Quinn and Keough, 2002): r^2 (coefficient of determination), RMSD (root-mean-square deviation), SSE (sum of squared errors), AIC (Akaike's information criterion) and the p -value for the F-test. RMSD provides a measure of the prediction precision through the differences between predicted and observed values. Both for RMSD and SSE, lower values stand for better fit. AIC allows the comparison of the adequacy of multiple models, with a lower AIC value denoting higher likelihood of the model. The p (F-test) stands for the p -value of the F-test applied to the errors of each model-deviation, which tests the null hypothesis that the errors of respective deviation are equal or higher than the errors of other deviations – we applied the F-test to compare the errors of each deviation with the errors of the baseline model and other deviations with less fitting parameters. P -values ≤ 0.05 mean that the respective model-deviation allows a better adjustment to the experimental data.

3. Results and discussion

The interest on DES has increased in the last years, boosted by

their advantages over other solvents (Vieira et al., 2018). This interest has been accompanied by an increasing number of studies on their toxicity. However, the majority of these studies used the salt [Chol]Cl as HBA, and considered each DES as a single compound, which does not dissociate, regardless the presence of water. Following a different approach, in this study we consider the DES aqueous solutions as a mixture, and aimed to evaluate the applicability of the mixture theory models to the understanding and prediction of the toxicity of DES based on non conventional HBA.

It has been argued that DES can exert their toxicity to bacteria by disrupting the cell wall (Wen et al., 2015). The cell wall in gram negative bacteria such as *A. fischeri* possesses molecules of peptidoglycan outside of the cytoplasmic membrane, which are responsible for the rigidity and strength of the structure (Nelson and Cox, 2012). Since DES were tested as aqueous solutions, they are expected to be dissociated (Dai et al., 2015; Hammond et al., 2017; Passos et al., 2016; Zhekenov et al., 2017). In this context, the ammonium cation could behave similarly to the cholinium cation addressed in other studies, by interacting with the polysaccharides or the peptide chains through hydrogen bonding or electrostatic interactions, leading to distortion or disruption of the cell wall (Wen et al., 2015). Moreover, the saltwater test media of the Microtox[®] test, rich in chloride, can play an important role in the results, since chloride can also interact with the DES cations, reducing their permeability through the cell walls and, therefore, their toxicity (Latała et al., 2005).

The median effective concentrations (EC_{50}) as estimated following 5, 15 and 30 min of exposure of *A. fischeri* to each HBA, HBD and DES are presented in Table 1. There was a general agreement in the benchmarks estimated for different exposure periods; despite this, very slight fluctuations were observed through time with little meaning since they lead to overlapping confidence intervals (thus rendering the benchmarks unlikely to be different between exposure periods). Because a longer exposure period (30 min) is likely to better ensure that the whole bacteria inoculum in each treatment contacts and/or internalizes the waterborne chemicals, further comparative discussion regarding EC_{50} benchmarks will be based on the results retrieved following 30 min of exposure. For a complete characterization of these starting materials and eutectic mixtures, all calculated EC_x values after 30 min of exposure are presented as supplementary material (Table S1).

According to the United Nations Globally Harmonized System of Classification and Labelling Chemicals (GHS) (UN, 2011) none of the starting materials, neither DES, could be considered hazardous to the aquatic environment, as their 30 min- EC_{50} values are above 100 mg L^{-1} . Such a low toxicity agrees with the claimed “green character” of DES. Among the starting materials, ethylene glycol was the least toxic (30 min- $EC_{50} = 96.49 \text{ g L}^{-1}$), whereas [N₃₃₃₃]Cl was the most toxic (30 min- $EC_{50} = 0.5456 \text{ g L}^{-1}$). Regarding the toxicity of HBDs, the lower toxicity of ethylene glycol compared to 1-propanol reflects its lower lipophilicity, represented by its lower log P (logarithm function of 1-octanol/water partition coefficient; log $P = -1.36$ and 0.25 respectively for ethylene glycol and 1-propanol; Chemspider, assessed on 19 January 2018). Previous studies also found lower toxicity of ethylene glycol compared to other HBDs such as glycerine, triethylene glycol, urea, acetamide and glycerol (Hayyan et al., 2013b; Wen et al., 2015). Concerning the toxicity of the tested HBAs, it increased in the following order: [N₁₁₁₁]Cl < [N₂₂₂₂]Cl < [N₃₃₃₃]Cl. This variation agrees with the side-chain effect defining increased toxicity along with the alkyl side chain elongation (Egorova and Ananikov, 2014; Matzke et al., 2010). This effect could be attributed to the increased lipophilicity caused by the elongation of the alkyl chain, with consequent higher reactivity with the biological membranes and embedded proteins (Stolte et al., 2007). It was found valid both for DES with ethylene

Table 1

Median effective concentration (EC₅₀) in g L⁻¹ and the respective 95% confidence intervals within brackets, obtained after 5, 15 and 30 min of exposure the marine bacteria *A. fischeri* (Microtox® toxicity test) to different DES and their starting materials.

	Chemical compounds	5 min - EC ₅₀ (95% CI), g L ⁻¹	15 min - EC ₅₀ (95% CI), g L ⁻¹	30 min - EC ₅₀ (95% CI), g L ⁻¹
Starting materials	[N ₁₁₁₁]Cl	23.79 (19.37–28.21)	20.88 (16.64–25.13)	22.69 (17.48–27.90)
	[N ₂₂₂₂]Cl	17.78 (12.94–22.62)	18.45 (12.482–24.421)	19.24 (13.27–25.21)
	[N ₃₃₃₃]Cl	0.3800 (0.3454–0.4146)	0.4706 (0.4224–0.5187)	0.5456 (0.4870–0.6040)
	1-propanol	8.327 (6.951–9.703)	9158.993 (7.788–10.53)	9.052 (7.475–10.63)
DES	Ethylene glycol	77.12 (52.16–102.08)	87.75 (62.74–112.8)	96.49 (69.96–123.0)
	1-propanol:[N ₁₁₁₁]Cl 1:1	20.46 (17.14–23.78)	19.93 (17.62–22.23)	20.87 (16.92–24.81)
	1-propanol:[N ₁₁₁₁]Cl 2:1	15.43 (11.77–19.10)	15.42 (11.74–19.10)	16.15 (12.10–20.19)
	1-propanol:[N ₁₁₁₁]Cl 4:1	15.68 (12.87–18.49)	16.43 (14.71–18.14)	15.36 (14.00–16.71)
	1-propanol:[N ₂₂₂₂]Cl 1:2	19.53 (16.58–33.47)	20.71 (17.51–23.92)	22.26 (19.15–25.36)
	1-propanol:[N ₂₂₂₂]Cl 1:1	17.20 (11.82–22.95)	17.59 (12.45–22.73)	18.09 (13.44–22.74)
	1-propanol:[N ₂₂₂₂]Cl 2:1	12.51 (10.66–14.37)	13.86 (11.61–16.12)	15.55 (12.80–18.30)
	1-propanol:[N ₂₂₂₂]Cl 4:1	7.821 (6.525–9.117)	8.571 (7.673–9.470)	9.500 (8.555–10.44)
	1-propanol:[N ₃₃₃₃]Cl 1:2	0.8187 (0.6186–1.019)	1.467 (1.139–1.795)	1.555 (1.181–1.929)
	1-propanol:[N ₃₃₃₃]Cl 1:1	10.94 (7.172–14.70)	6.106 (3.756–8.455)	4.981 (2.862–7.099)
	1-propanol:[N ₃₃₃₃]Cl 2:1	1.415 (1.207–1.622)	1.700 (1.490–1.910)	1.845 (1.618–2.072)
	1-propanol:[N ₃₃₃₃]Cl 4:1	0.8254 (0.6444–1.006)	0.9896 (0.8050–1.174)	1.120 (0.9055–1.334)
	Ethylene glycol:[N ₁₁₁₁]Cl 1:2	31.73 (28.97–34.49)	29.01 (26.38–31.63)	30.20 (20.59–33.34)
	Ethylene glycol:[N ₁₁₁₁]Cl 1:1	55.72 (50.02–61.41)	52.11 (46.79–57.43)	53.99 (47.88–60.09)
	Ethylene glycol:[N ₁₁₁₁]Cl 2:1	42.35 (37.62–47.07)	42.41 (38.81–45.99)	49.25 (46.47–52.03)
	Ethylene glycol:[N ₁₁₁₁]Cl 4:1	52.78 (35.46–70.10)	54.52 (37.55–71.49)	65.62 (47.36–83.88)
	Ethylene glycol:[N ₂₂₂₂]Cl 1:2	18.20 (14.66–21.74)	18.98 (15.36–22.60)	18.98 (15.27–22.60)
	Ethylene glycol:[N ₂₂₂₂]Cl 1:1	23.91 (21.54–26.29)	24.50 (21.19–27.81)	23.94 (20.52–27.36)
	Ethylene glycol:[N ₂₂₂₂]Cl 2:1	21.52 (14.21–28.82)	9.069 (8.882–9.257)	18.61 (13.75–23.48)
	Ethylene glycol:[N ₂₂₂₂]Cl 4:1	35.31 (27.59–43.03)	37.18 (27.65–46.72)	36.39 (27.79–44.99)
	Ethylene glycol:[N ₃₃₃₃]Cl 1:2	0.5362 (0.5080–0.5644)	0.7786 (0.7100–0.8472)	0.9713 (0.7336–1.209)
	Ethylene glycol:[N ₃₃₃₃]Cl 1:1	2.520 (2.417–2.624)	3.074 (2.949–3.199)	3.665 (3.441–3.888)
	Ethylene glycol:[N ₃₃₃₃]Cl 2:1	0.7399 (0.6639–0.8138)	0.9087 (0.8042–1.013)	0.9453 (0.8259–1.064)
	Ethylene glycol:[N ₃₃₃₃]Cl 4:1	1.052 (0.9616–1.143)	1.273 (1.163–1.384)	1.285 (1.142–1.429)

glycol, as with 1-propanol, and highlights the central role of HBA in DES toxicity.

The effects of the binary mixtures HBD:HBA to *A. fischeri* are depicted in Fig. 1 through 2D isobolographic diagrams representing response surfaces. In addition, estimated parameters for the fitting of experimental data to the reference mixture models and the corresponding deviation functions are shown in Table 2. The interpretation of such parameters to appraise the behavior of each mixture (i.e. each DES) was performed based on Jonker et al. (2005).

Regarding the toxicity of the DES composed of 1-propanol:[N₁₁₁₁]Cl, only the CA model incorporating dose-level deviation showed significant fitting through the F-test ($p = 0.039$); also, this deviation has high r^2 values and low SSE and RMSD values (Table 2) and, thus, can be considered as the model that best describes the experimental toxicity data. The positive value of “a” indicates antagonism at low dose level and synergism at high dose level. Graphically, the antagonism is evident through the convex shape of the isobologram (Fig. 1A). The theoretical shift between antagonism and synergism occurs at higher dose level than the EC₅₀ value, as given by the “b_{DL}” value (0.322, Table 2), and shown by the approximation to linearity of the isobole denoting 80% luminescence inhibition compared to the control (i.e. the 20%-luminescence isobole in Fig. 1A) while that denoting 40% of luminescence isobole still presents the typical convex shape defining antagonism. The linearity of the isobole for 20% luminescence (approximating to a baseline CA behavior where chemicals do not interact) is indicative that the full inversion into a synergic behavior of the mixture is unlikely to occur in practice or that its occurrence may occur at very high effect levels (higher than 80% luminescence inhibition compared to the control) that may not be biologically significant as the predictions may confirm beyond 100% luminescence inhibition.

All the deviation functions from IA and CA fitted significantly better than the baseline models in the case of ethylene glycol:[N₁₁₁₁]Cl, with dose-level and dose-ratio dependence bearing

higher error than S/A in both cases (P-value from the F test <0.05; Table 2). This supports the assumption that this mixture behaves according to the S/A model. This is the simpler among the deviation-models explaining the data better than the corresponding baseline (CA or IA) and explaining the data at least as well as DL or DR, this reasoning preventing overfitting effects and the hampering of predictive accuracy (‘Principle of Parsimony’; Burnham and Anderson, 2002). Concerning whether CA-S/A or IA-S/A fit better, given that IA-S/A model shows higher r^2 and lower SSE, RMSD and AIC values compared to CA-S/A, one should reasonably assume that IA-S/A is the model best describing this mixture's behavior. The positive value of “a” indicates antagonism, which agrees with the slightly convex shape of the isobologram (Fig. 1B). Antagonism is more evident at low dose level, whereas a tendency for baseline IA behavior or slight synergism is observed at high dose level, as suggested by the “a” value in the DL deviation and retrievable from the isobologram in Fig. 1B. Antagonism is mainly caused by the salt, as given by the positive value of b_{HBA}, whereas synergism is mainly caused by ethylene glycol, as given by the negative value of b_{HBD}.

For 1-propanol:[N₂₂₂₂]Cl, a good fitting to the experimental data was achieved by the IA model, with all the deviation functions adjusting better than the baseline model (Table 2). However, the F-test values show that neither the DL deviation nor the DR deviation fit better than the S/A deviation, which suggests that the model IA with S/A deviation is the most appropriate to describe the experimental data and hence the toxic behavior of the mixture towards *A. fischeri*. The negative “a” value of the S/A deviation indicates synergism, which agrees with the slightly concave shape of the response curve (Fig. 1C).

Similarly to 1-propanol:[N₂₂₂₂]Cl, the CA model and derived deviation models did not fit significantly to the experimental data, and a good fitting of the experimental data to the IA model with all deviations was achieved for ethylene glycol:[N₂₂₂₂]Cl. Indeed, the p-values of the F-test suggest that all the deviations to IA adjust




Fig. 1. Isobolograms illustrating 2D response surfaces, according to the model that best fitted the experimental data, identified at the top (see Table 2 for the fitting parameters) – CA (Concentration Addition) or IA (Independent Action) added S/A (Synergism/Antagonism), DR (Dose Ratio) or DL (Dose Level) deviation functions. The isoboles represent equi-effective levels, the effect being bioluminescence of the bacteria *A. fischeri* relative to control levels.

better than the baseline model. Then, although IA-DR is the most complex among the three deviation models (thus failing to fully comply with the best practices for regression modelling; Burnham and Anderson, 2002), integrative analysis of the statistics suggests that this is the most adequate to predict the behavior of the present mixture. IA-DR explains the data better than S/A ($P < 0.05$ for the direct comparison of the errors; Table 2) but this is not the case while comparing with IA-DL ($P > 0.05$; Table 2). However, IA-DR bears a relevant increase in r^2 , SSE, RMSD and AIC compared to IA-DL, strongly suggesting that the DR deviation indeed fits better to the data. The value of “a” indicates antagonism, which is mainly

due to ethylene glycol (“b_{HBD}”), as observed in the isobologram (Fig. 1D); also, synergism is observed, being mainly caused by the salt [N₂₂₂₂]Cl, as given by the “b_{HBA}” value. Because the DL deviation also reflects very well the mixture behavior, it is worth remarking that synergism mostly occurs at low dose levels while antagonism may establish at high dose level (negative value of the corresponding “a” parameter of -5.064).

Concerning the DES based in 1-propanol:[N₃₃₃₃]Cl, both the CA and IA models with dose-ratio deviations adjusted significantly to the observed toxicity. However, given the lower SSE values associated to IA with DR deviation, it is reasonable to assume that IA-DR

Table 2

Statistical and estimated parameters for the fitting of the experimental data (retrieved while testing DES toxicity to *A. fischeri*) to CA and IA, as well as deviations from these reference models, denoting dose-level dependence (DL), synergism/antagonism (S/A) and dose-ratio (DR) dependence. The models best describing each DES behavior are highlighted in grey. For the purpose, we generally followed the 'Principle of Parsimony' (Burnham and Anderson, 2002) by rationally selecting the simplest model (i.e. that with lower number of added parameters) that significantly fitted the dataset with equal and/or lower uncertainty.

	CA				IA				
	Baseline	S/A	DL	DR	Baseline	S/A	DL	DR	
1-propanol : [N ₁₁₁₁]Cl	r ²	0.926	0.963	0.970	0.964	0.957	0.969	0.971	0.969
	RMSD	-	8.335	7.686	8.656	-	7.653	7.626	7.960
	SSE (df)	3644.292 (21)	1806.420	1476.823	1798.281	2115.481 (21)	1522.746	1453.730	1520.648
	AIC	183.257	132.821	131.249	140.840	133.571	128.209	130.823	136.312
	P (F-test)	-	vs CA: 0.075	vs CA: 0.039; vs S/A: 0.373	vs CA: 0.111; vs S/A: 0.589; vs DL: 0.296	-	vs IA: 0.267	vs IA: 0.271; vs S/A: 0.506	vs IA: 0.349; vs S/A: 0.592; vs DL: 0.415
	a, b	-	a=3.151	a=6.167; b _{DL} =0.322	a=1.941; b ₁ =1.950; b ₂ =-0.009	-	a=1.175	a=2.140; b _{DL} =0.834	a=0.729; b ₁ =0.691; b ₂ =-0.037
1-propanol : [N ₂₂₂₂]Cl	r ²	0.942	0.942	0.948	0.962	0.898	0.958	0.971	0.972
	RMSD	-	10.590	10.232	8.887	-	9.031	7.653	7.583
	SSE (df)	3854.525 (29)	3813.406	3454.843	2527.494	6695.802 (29)	2772.774	1932.601	1840.151
	AIC	179.356	182.045	181.869	174.454	198.686	170.891	161.537	163.346
	P (F-test)	-	vs CA: 0.527	vs CA: 0.461; vs S/A: 0.437	vs CA: 0.205; vs S/A: 0.195; vs DL: 0.757	-	vs IA: 0.014	vs IA: 0.001; vs S/A: 0.201	vs IA: 0.001; vs S/A: 0.196; vs DL: 0.510
	a, b	-	a=-0.496	a=-2.091; b _{DL} =0.629	a=-0.817; b ₁ =-11.784; b ₂ =10.967;2	-	a=-2.761	a=-3.980; b _{DL} =1.129	a=-1.935; b ₁ =-6.408; b ₂ =4.473
1-propanol : [N ₃₃₃₃]Cl	r ²	0.675	0.726	0.727	0.905	0.683	0.728	0.733	0.909
	RMSD	-	19.396	19.656	11.752	-	19.318	19.433	11.517
	SSE (df)	15601.085 (30)	13166.800	13136.099	4557.887	15232.321 (30)	13061.693	12840.183	4377.147
	AIC	233.286	230.203	233.351	198.708	232.425	229.914	232.531	197.251
	P (F-test)	-	vs CA: 0.358	vs CA: 0.392; vs S/A: 0.535	vs CA: 0.002; vs S/A: 0.006; vs DL: 0.995	-	vs IA: 0.374	vs IA: 0.393; vs S/A: 0.520	vs IA: 0.001; vs S/A: 0.005; vs DL: 0.996
	a, b	-	a=20.915	a=19.124; b _{DL} =-0.0273	a=-819.580; b ₁ =-1792.308; b ₂ =972.731	-	a=11.494	a=1.517; b _{DL} =-9.364	a=-468.005; b ₁ =-1022.536; b ₂ =554.532
ethylene glycol : [N ₁₁₁₁]Cl	r ²	0.935	0.976	0.981	0.982	0.948	0.981	0.981	0.985
	RMSD	-	6.353	5.814	5.754	-	5.688	5.744	5.233
	SSE (df)	3886.956 (30)	1412.460	1149.468	1092.750	3066.528 (30)	1132.290	1121.839	903.835
	AIC	183.257	149.838	145.653	147.294	174.722	141.879	144.777	140.461
	P (F-test)	-	vs CA: 0.005	vs CA: 0.001; vs S/A: 0.327	vs CA: 0.001; vs S/A: 0.316; vs DL: 0.514	-	vs IA: 0.006	vs IA: 0.007; vs S/A: 0.528	vs IA: 0.002; vs S/A: 0.345; vs DL: 0.679
	a, b	-	a=4.171	a=6.668; b _{DL} =0.192	a=0.519; b ₁ =-7.587; b ₂ =8.106	-	a=2.436	a=2.972; b _{DL} =0.276	a=0.338; b ₁ =-4.134; b ₂ =4.472
ethylene glycol : [N ₂₂₂₂]Cl	r ²	0.846	0.888	0.892	0.914	0.836	0.921	0.926	0.964
	RMSD	-	13.632	13.578	12.329	-	11.419	11.282	8.011
	SSE (df)	8945.590 (30)	6503.988	6267.991	5016.368	9522.147 (30)	4563.918	4327.335	2117.875
	AIC	213.264	204.813	206.714	202.159	215.512	192.060	193.376	171.116
	P (F-test)	-	vs CA: 0.223	vs CA: 0.221; vs S/A: 0.499	vs CA: 0.103; vs S/A: 0.313; vs DL: 0.685	-	vs IA: 0.031	vs IA: 0.028; vs S/A: 0.482	vs IA: 0.000; vs S/A: 0.036; vs DL: 0.959
	a, b	-	a=-6.552	a=-8.027; b _{DL} =-0.249	a=5.794; b ₁ =34.438; b ₂ =-28.644	-	a=-4.387	a=-5.064; b _{DL} =0.824	a=2.231; b ₁ =18.345; b ₂ =-16.114
ethylene glycol : [N ₃₃₃₃]Cl	r ²	0.783	0.797	0.799	0.866	0.783	0.798	0.803	0.867
	RMSD	-	15.992	16.183	13.404	-	15.987	16.015	13.343
	SSE (df)	9590.928 (30)	8951.095	8903.891	5928.778	9579.092 (30)	8945.239	8720.788	5875.060
	AIC	215.772	216.310	219.352	208.175	215.727	216.286	218.604	207.847
	P (F-test)	-	vs CA: 0.463	vs CA: 0.495; vs S/A: 0.523	vs CA: 0.157; vs S/A: 0.188; vs DL: 0.830	-	vs IA: 0.464	vs IA: 0.474; vs S/A: 0.511	vs IA: 0.152; vs S/A: 0.183; vs DL: 0.823
	a, b	-	a=79.854	a=65.891; b _{DL} =-0.088	a=-31766.216; b ₁ =-64252.457 b ₂ =32485.229	-	a=46.668	a=0.161; b _{DL} =-475.226	a=-18459.990; b ₁ =-37336.398; b ₂ =18876.080

r²: coefficient of determination; RMSD: root-mean-square deviation; SSE (df): sum of squared errors of prediction (degrees of freedom between parenthesis); AIC: Akaike's information criterion; P (F-test) refers to the p-value of the F test that tests the null hypothesis that the errors of the respective deviation model are equal and the alternative hypothesis that they are higher than the errors of other deviations with less fitting parameters; a, b₁, b₂, b_{DL}: estimated parameters, where 1 and 2 stand for the HBD and HBA, respectively.

is the model best describing the mixture toxic behavior. The heavily negative value of “a” (−468.005) indicates strong synergism, which is consistent with the pronounced concave shape of the isobologram (Fig. 1E). The heavily positive value of “b_{HBA}” may indicate antagonism caused by the salt, which is observed at concentrations higher than 0.8 g L^{−1}, approximately. On the other hand, the heavily negative value of “b_{HBD}” confirms the synergistic interaction between DES components as visible in the isobologram, where the toxicity of the mixture is mainly caused by ethylene glycol (Table 2, Fig. 1E).

Regarding ethylene glycol:[N₃₃₃₃]Cl, the F-test to the errors did not highlight any deviation model as better describing the experimental data compared to the baseline model ($P > 0.05$; Table 2). However, the higher r^2 , supported by lower SSE and RMSD values suggest that the experimental data fit better to the CA or IA with DR deviation than to the baseline models. Despite the short differences between the statistical parameters, the IA model with DR deviation fits slightly better than its CA counterpart. Similarly to 1-propanol:[N₃₃₃₃]Cl, the “b_{HBA}” and “b_{HBD}” values for ethylene glycol:[N₃₃₃₃]Cl suggest antagonism caused by the salt, which is not depicted in the isobologram, as well as synergism mainly caused by the HBD is pictured by the strongly concave shape of the isoboles (Fig. 1F).

In summary, the toxicity of the tested DES was well described by the classical models of mixture toxicity, with the IA model with specific deviations being the one that adjusted better to the experimental data for 5 out of 6 DES investigated. The interactions (synergism and antagonism) found between DES components shows that DES toxicity cannot be predicted based solely on the toxicity of the starting materials. Moreover, this approach highlighted the role of both the HBD and HBA on DES toxicity, corroborating results of a previous study assessing DES toxicity to a bacterium (Wen et al., 2015). The HBA markedly affected DES toxicity, as translated by the higher toxicity of [N₃₃₃₃]Cl-based DES compared to that of [N₁₁₁₁]Cl- and [N₂₂₂₂]Cl-based DES, this being related to decreased hydrophilicity of the HBA (de Moraes et al., 2015). The pronounced synergism found for both [N₃₃₃₃]Cl-based DES, but not for other DES, and the antagonism found for both [N₁₁₁₁]Cl-based DES are also noteworthy in this context. A relevant role of HBD on DES toxicity has also been suggested in previous studies, namely concerning their cytotoxicity (Hayyan et al., 2016; Radošević et al., 2015; Wen et al., 2015) and toxicity to fungi and fish (Juneidi et al., 2016). Here, we found that ethylene glycol-based DES were equally or less toxic than 1-propanol-based DES (Table 1), which agrees with the differing toxicity of both these HBD isolated.

The knowledge of the type of interaction between the DES components (HBA and HBD) is of great importance for the toxicological profile of DES. The synergism is of particular concern, as the toxicity of the mixture (DES aqueous solution) is greater than the toxicity of each of its components separately, thus potentially more environmentally hazardous. In these cases, the toxicity of DES solutions would be underestimated if predicted based only on the toxicity of their components. In the present study, synergism was found for [N₃₃₃₃]Cl-based DES and also for ethylene glycol:[N₂₂₂₂]Cl. The toxicity of [N₃₃₃₃]Cl-based DES raises special concern, not only because of the strong synergism between their components (Fig. 1E and F), but also because of their high overall toxicity compared to the other DES (Table 1). However, it should be kept in mind that the response of different organisms to the same DES may vary (Wen et al., 2015). In other words, the fact that synergism (or antagonism) occurred between the components of a DES for a certain species does not mean that the same deviation will occur for other species. For this reason, from an environmentally precautionary perspective, studies covering several functional levels (e.g. Santos et al., 2015 for cholinium-based ILs) are essential to gain a

comprehensive knowledge about the effects of these solvents on the aquatic ecosystems, thus improving the knowledge of their ecotoxicological profile and ultimately, leading to a more realistic environmental risk assessment of such chemicals.

4. Conclusions

Considering the tailor-made character of DES, the number of DES that can be created with the same starting materials is potentially countless, this making their toxicity assessment a daunting task. The appropriateness of mixture toxicity models to describe DES toxicity, as demonstrated for the first time in the present study, configures a breakthrough in this problematic. By feasibly modelling response surfaces of each DES (covering all theoretical combinations, i.e. molar ratios, between a given HBA and HBD) based on the experimental assessment of rationally selected ratios, one gains a broad overview of its toxicity profile, properly appraising interactive effects (synergism/antagonism; dose-level or dose-ratio dependence). Such predictive tools can be valuable to retrieve theoretical EC_x values that may hold as environmental benchmarks useful in the environmental risk assessment of new DES. Additionally, these tools provide the possibility of *a priori* selecting, among DES that fit the desired properties regarding functionality for a given application, those bearing lower environmental hazardous potential. In practice, the advantages of such a possibility reflect in two important arenas. First, regulation worldwide increasingly accounts to the environmental safety of newly developed compounds, specifically requesting the ecotoxicological assessment of chemicals (e.g. REACH Directive in Europe; EC, 2006), eventually comprising read-across (or data gap filling) frameworks (see e.g. ECHA, 2015; Williams et al., 2011 for an overview on this regulatory approach) for designer compounds such as DES. Secondly, important savings are likely to occur through preventing investment in the development of DES deriving from given HBD-HBA that are likely to be more environmentally hazardous, hence more hardly complying with mandatory regulation for the registration and marketing of new compounds.

Acknowledgments

Thanks are due for the financial support to CESAM - Centre for Environmental and Marine Studies, University of Aveiro, Portugal (UID/AMB/50017 - POCI-01-0145-FEDER-007638), to FCT - Portuguese Foundation for Science and Technology/MCTES - Ministry of Science, Technology and Higher Education, Portugal through national funds (PIDDAC), and the co-funding by the FEDER, within the PT2020 Partnership Agreement and Compete 2020. This work was also developed within the scope of the project CICECO-Aveiro Institute of Materials, University of Aveiro, Portugal, POCI-01-0145-FEDER-007679 (FCT Ref. UID/CTM/50011/2013), financed by national funds through the FCT/MEC - Ministry of Education and Science, Portugal and when appropriate co-financed by FEDER under the PT2020 Partnership Agreement. The authors also acknowledge the support by the FCT through the project PTDC/ATP-EAM/5331/2014, the doctoral grant of IPE Macário (SFRH/BD/123850/2016) and post-doctoral grants of AMM Gonçalves and JL Pereira (SFRH/BPD/97210/2013, SFRH/BPD/101971/2014), respectively. SPM Ventura acknowledges FCT for the contract IF/00402/2015. We are also grateful to Prof. António JA Nogueira for giving us permission to use the ToxCalcMix spreadsheet.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.chemosphere.2018.08.153>.

References

- Altenburger, R., Backhaus, T., Boedeker, W., Faust, M., Scholze, M., Grimme, L.H., 2000. Predictability of the toxicity of multiple chemical mixtures to *Vibrio fischeri*: mixtures composed of similarly acting chemicals. *Environ. Toxicol. Chem.* 19, 2341–2347. <https://doi.org/10.1002/etc.5620190926>.
- Backhaus, T., Faust, M., 2012. Predictive environmental risk assessment of chemical mixtures: a conceptual framework. *Environ. Sci. Technol.* 46, 2564–2573. <https://doi.org/10.1021/es2034125>.
- Barata, C., Baird, D.J., Nogueira, A.J.A., Soares, A.M.V.M., Riva, M.C., 2006. Toxicity of binary mixtures of metals and pyrethroid insecticides to *Daphnia magna* Straus. Implications for multi-substance risks assessment. *Aquat. Toxicol.* 78, 1–14. <https://doi.org/10.1016/j.aquatox.2006.01.013>.
- Burnham, K.P., Anderson, D.R., 2002. *Model Selection and Multimodel Inference. A Practical Information-theoretic Approach*, second ed. Springer-Verlag New York, Inc., p. 488.
- Cardellini, F., Germani, R., Cardinali, G., Corte, L., Roscini, L., Spreti, N., Tieccoab, M., 2015. Room temperature deep eutectic solvents of (1S)-(+)-10-camphorsulfonic acid and sulfobetaines: hydrogen bond-based mixtures with low ionicity and structure-dependent toxicity. *RSC Adv.* 40, 31772–31786.
- Dai, Y., van Spronsen, J., Witkamp, G.J., Verpoorte, R., Choi, Y.H., 2013. Natural deep eutectic solvents as new potential media for green technology. *Anal. Chim. Acta* 766, 61–68. <https://doi.org/10.1016/j.aca.2012.12.019>.
- Dai, Y., Witkamp, G.J., Verpoorte, R., Choi, Y.H., 2015. Tailoring properties of natural deep eutectic solvents with water to facilitate their applications. *Food Chem.* 187, 14–19. <https://doi.org/10.1016/j.foodchem.2015.03.123>.
- de Moraes, P., Gonçalves, F., Coutinho, J.A.P., Ventura, S.P.M., 2015. Ecotoxicity of cholinium-based deep eutectic solvents. *ACS Sustain. Chem. Eng.* 3, 3398–3404. <https://doi.org/10.1021/acsschemeng.5b01124>.
- EC, 2006. Regulation (EC) 1907/2006 of the European Parliament and of the Council of 18 December 2006-REACH. Official Journal of the European Union. European Commission, 396:0001:0849:(EN: PDF). <https://eur-lex.europa.eu/LexUriServ/LexUriServ.do?uri=14OJ:L:200606>.
- ECHA, 2015. Read-across Assessment Framework (RAAF). European Chemical Agency, Helsinki, Finland. <https://doi.org/10.2823/546436>.
- Egorova, K.S., Ananikov, V.P., 2014. Toxicity of ionic liquids: eco(cyto)activity as complicated, but unavoidable parameter for task-specific optimization. *ChemSusChem* 7, 336–360. <https://doi.org/10.1002/cssc.201300459>.
- Hammond, O.S., Bowron, D.T., Edler, K.J., 2017. The effect of water upon deep eutectic solvent nanostructure: an unusual transition from ionic mixture to aqueous solution. *Angew. Chem. Int. Ed.* 56, 9782–9785. <https://doi.org/10.1002/anie.201702486>.
- Hayyan, M., Hashim, M.A., Al-Saadi, M.A., Hayyan, A., AlNashef, I.M., Mirghani, M.E.S., 2013a. Assessment of cytotoxicity and toxicity for phosphonium-based deep eutectic solvents. *Chemosphere* 93, 455–459. <https://doi.org/10.1016/j.chemosphere.2013.05.013>.
- Hayyan, M., Hashim, M.A., Hayyan, A., Al-Saadi, M.A., AlNashef, I.M., Mirghani, M.E.S., Saheed, O.K., 2013b. Are deep eutectic solvents benign or toxic? *Chemosphere* 90, 2193–2195. <https://doi.org/10.1016/j.chemosphere.2012.11.004>.
- Hayyan, M., Looi, C.Y., Hayyan, A., Wong, W.F., Hashim, M.A., 2015. *In Vitro* and *In Vivo* toxicity profiling of ammonium-based deep eutectic solvents. *PLoS One* 10, 1–18. <https://doi.org/10.1371/journal.pone.0117934>.
- Hayyan, M., Mbous, Y.P., Looi, C.Y., Wong, W.F., Hayyan, A., Salleh, Z., Mohd-Ali, O., 2016. Natural deep eutectic solvents: cytotoxic profile. *SpringerPlus* 5, 913. <https://doi.org/10.1186/s40064-016-2575-9>.
- Huang, Z.L., Wu, B.P., Wen, Q., Yang, T.X., Yang, Z., 2014. Deep eutectic solvents can be viable enzyme activators and stabilizers. *J. Chem. Technol. Biotechnol.* 89, 1975–1981. <https://doi.org/10.1002/jctb.4285>.
- Jhong, H.R., Wong, D.S.H., Wan, C.C., Wang, Y.Y., Wei, T.C., 2009. A novel deep eutectic solvent-based ionic liquid used as electrolyte for dye-sensitized solar cells. *Electrochem. Commun.* 11, 209–211. <https://doi.org/10.1016/j.elecom.2008.11.001>.
- Johnson, B., 2005. Microtox® Acute toxicity test. In: *Small-scale Freshwater Toxicity Investigations*, pp. 69–105. https://doi.org/10.1007/1-4020-3120-3_2.
- Jonker, M.J., Svendsen, C., Bedaux, J.J.M., Bongers, M., Kammenga, J.E., 2005. Significance testing of synergistic/antagonistic, dose level-dependent, or dose ratio-dependent effects in mixture dose–response analysis. *Environ. Toxicol. Chem.* 24, 2701. <https://doi.org/10.1897/04-431R.1>.
- Juneidi, I., Hayyan, M., Mohd Ali, O., 2016. Toxicity profile of choline chloride-based deep eutectic solvents for fungi and *Cyprinus carpio* fish. *Environ. Sci. Pollut. Res.* 23, 7648–7659. <https://doi.org/10.1007/s11356-015-6003-4>.
- Latala, A., Stepnowski, P., Ne, dzi, M., Mrozik, W., 2005. Marine toxicity assessment of imidazolium ionic liquids: Acute effects on the Baltic algae *Oocystis submarina* and *Cyclotella meneghiniana*. *Aquat. Toxicol.* 73, 91–98. <https://doi.org/10.1016/j.aquatox.2005.03.008>.
- Matzke, M., Arning, J., Ranke, J., Jastorff, B., Stolte, S., 2010. Design of inherently safer ionic liquids: toxicology and biodegradation. In: *Handbook of Green Chemistry*, pp. 233–298.
- Mbous, Y.P., Hayyan, M., Hayyan, A., Wong, W.F., Hashim, M.A., Looi, C.Y., 2017a. Applications of deep eutectic solvents in biotechnology and bioengineering—promises and challenges. *Biotechnol. Adv.* 35, 105–134. <https://doi.org/10.1016/j.biotechadv.2016.11.006>.
- Mbous, Y.P., Hayyan, M., Wong, W.F., Looi, C.Y., Hashim, M.A., 2017b. Unraveling the cytotoxicity and metabolic pathways of binary natural deep eutectic solvent systems. *Sci. Rep.* 7, 41257. <https://doi.org/10.1038/srep41257>.
- Motulsky, H., Christopoulos, A., 2004. *Fitting Models to Biological Data Using Linear and Nonlinear Regression. A Practical Guide to Curve Fitting*. GraphPad Software Inc., San Diego CA.
- Nelson, D.L., Cox, M.M., 2012. *Lehninger - Principles of Biochemistry*, sixth ed. Freeman, W. H. & Company.
- Paiva, A., Craveiro, R., Aroso, I., Martins, M., Reis, R.L., Duarte, A.R.C., 2014. Natural deep eutectic solvents – solvents for the 21st century. *ACS Sustain. Chem. Eng.* 2, 1063–1071.
- Passos, H., Tavares, D.J.P., Ferreira, A.M., Freire, M.G., Coutinho, J.A.P., 2016. Are aqueous biphasic systems composed of deep eutectic solvents ternary or quaternary systems? *ACS Sustain. Chem. Eng.* 4, 2881–2886. <https://doi.org/10.1021/acsschemeng.6b00485>.
- Quinn, G.P., Keough, M.J., 2002. *Experimental Design and Data Analysis for Biologists*. Cambridge University Press, Cambridge, UK.
- Radošević, K., Cvjetko Bubalo, M., Gaurina Srček, V., Grgas, D., Landeka Dragičević, T., Redovniković, R.I., 2015. Evaluation of toxicity and biodegradability of choline chloride based deep eutectic solvents. *Ecotoxicol. Environ. Saf.* 112, 46–53. <https://doi.org/10.1016/j.ecoenv.2014.09.034>.
- Ruß, C., König, B., 2012. Low melting mixtures in organic synthesis – an alternative to ionic liquids? *Green Chem.* 14, 2969–2982. <https://doi.org/10.1039/b000000x>.
- Santos, J.I., Gonçalves, A.M.M., Pereira, J.L., Figueiredo, B.F.H.T., Silva, F.A.E., Coutinho, J.A.P., Ventura, S.P.M., Gonçalves, F., 2015. Environmental safety of olinium-based ionic liquids: assessing structure-ecotoxicity relationships. *Green Chem.* 17, 4657–4668. <https://doi.org/10.1039/c5gc01129a>.
- Shah, D., Mjalli, F.S., 2014. Effect of water on the thermo-physical properties of Reline: an experimental and molecular simulation based approach. *Phys. Chem. Chem. Phys.* 16, 23900–23907. <https://doi.org/10.1039/C4CP02600D>.
- Singh, B.S., Lobo, H.R., Shankarling, G.S., 2012. Choline chloride based eutectic solvents: magical catalytic system for carbon – carbon bond formation in the rapid synthesis of β -hydroxy functionalized derivatives. *Catal. Commun.* 24, 70–74. <https://doi.org/10.1016/j.catcom.2012.03.021>.
- Stolte, S., Matzke, M., Jürgen, A., Bösch, A., Pitner, W.-R., Welz-Biermann, U., Jastorff, B., Ranke, J., 2007. Effects of different head groups and functionalised side chains on the aquatic toxicity of ionic liquids. *Green Chem.* 9, 1170–1179. <https://doi.org/10.1039/b711119c>.
- UN, U.N., 2011. *Globally Harmonized System of Classification and Labelling of Chemicals (GHS)*, fourth rev. ed. United Nations, New York and Geneva. <https://doi.org/10.1265/jjh.65.5>.
- Vieira, V., Prieto, M.A., Barros, L., Coutinho, J.A.P., Ferreira, I.C.F.R., Ferreira, O., 2018. Enhanced extraction of phenolic compounds using choline chloride based deep eutectic solvents from *Juglans regia* L. *Ind. Crop. Prod.* 115, 261–271. <https://doi.org/10.1016/j.indcrop.2018.02.029>.
- Wen, Q., Chen, J.X., Tang, Y.L., Wang, J., Yang, Z., 2015. Assessing the toxicity and biodegradability of deep eutectic solvents. *Chemosphere* 132, 63–69. <https://doi.org/10.1016/j.chemosphere.2015.02.061>.
- Williams, E.S., Berninger, J.P., Brooks, B.W., 2011. Application of chemical toxicity distributions to ecotoxicology data requirements under REACH. *Environ. Toxicol. Chem.* 30, 1943–1954. <https://doi.org/10.1002/etc.583>.
- Zhang, Q.H., Vigier, K.D., Royer, S., Jerome, F., 2012. Deep eutectic solvents: syntheses, properties and applications. *Chem. Soc. Rev.* 41, 7108–7146. <https://doi.org/10.1039/C2cs35178a>.
- Zhao, B.Y., Xu, P., Yang, F.X., Wu, H., Zong, M.H., Lou, W.Y., 2015. Biocompatible deep eutectic solvents based on choline chloride: characterization and application to the extraction of rutin from *Sophora japonica*. *ACS Sustain. Chem. Eng.* 3, 2746–2755. <https://doi.org/10.1021/acssuschemeng.5b00619>.
- Zhekenov, T., Toksanbayev, N., Kazakbayeva, Z., Shah, D., Mjalli, F., 2017. Formation of type III Deep Eutectic Solvents and effect of water on their intermolecular interactions. *Fluid Phase Equil.* 441, 43–48. <https://doi.org/10.1016/j.fluid.2017.01.022>.