


Victoria V. Ruzhenkova,\* Victor A. Ruzhenkov, Natalia K. Rzhetskaya, Alevtina V. Boeva, Julia N. Gomelyak. (2020). Sociophobia in Medical Students (Prevalence and Comorbidity). *International Journal of Early Childhood Special Education (INT-JECSE)*, 12(1): 574-581.

DOI: 10.9756/INT-JECSE/V12I1.201039

Received: 28.02.2020 Accepted: 15.05.2020

 Victoria V. Ruzhenkova<sup>1\*</sup>  
 Victor A. Ruzhenkov<sup>2</sup>  
 Natalia K. Rzhetskaya<sup>3</sup>  
 Alevtina V. Boeva<sup>4</sup>  
 Julia N. Gomelyak<sup>5</sup>

## Sociophobia in Medical Students (Prevalence and Comorbidity)

### Abstract

*Social phobias are one of the most common anxiety-phobic disorders. This study aimed to verify the prevalence of social phobias and comorbid disorders among medical students to develop recommendations for the prevention. A sample of 724 Russian students of 1-6 years of education in the medical institute was examined using the psychometric, clinical, psychopathological and nonparametric statistics methods. It was established that the symptoms of social phobia occur in 33.7% of medical students, more often in females (37.1%) than in male (24.1%). Social phobias have a negative influence on students' attitude to the medical profession and form doubt in their choice. It is advisable to provide psychological consultations for students with social phobias. It is possible that a person-centered psychotherapy, aimed at eliminating the symptoms of social phobia and changing the negative attitude towards the medical profession may be needed.*

**Keywords:** Anxiety. Attitude to the Profession. Comorbidity. Depression. Stress.

### Introduction

Epidemiological studies indicate that social phobias (SF), which have the status of independent mental disorder in ICD-10 and DSM-V, are one of the most common anxiety and phobic disorders (Rowa and Antony 2005; Thai et al. 2019). They manifest a constant and excessive fear of one or more social or service situations (Cremers and Roelofs 2016; Vargas-Hernández, 2016). Most often, symptoms of SF are manifested in the cognitive, somatic and behavioral spheres (Albuquerque and Deshauer 2002; Rendueles 2017).

In the development of SF genetic, family, environmental factors and growth play the role of

(Hudson and Rapee 2000), as well as personal characteristics of patients (Savoia et al. 2010; Nikfarjam et al. 2017; Wersebe et al. 2018; Bicer, et al, 2018).

Phobic disorders are registered in 10%-12.7% of adolescents of 16–17 years old (which is important for high school students and students due to the period of active social activity.) and women showed a higher level of social phobia than men (Olivares-Olivaresa et al. 2016). SF has the third highest prevalence in primary care after depression and GAD and occurs in 7% of general practice patients (Essau et al. 2018) According to the literature (Mohammadi et al. 2019; Shen et al. 2020) only 72% of students can be called "completely

Victoria V. Ruzhenkova<sup>1\*</sup>, Medical Institute, Belgorod State University, Belgorod, Russia  
Victor A. Ruzhenkov<sup>2</sup>, Medical Institute, Belgorod State University, Belgorod, Russia  
Natalia K. Rzhetskaya<sup>3</sup>, Medical Institute, Belgorod State University, Belgorod, Russia  
Alevtina V. Boeva<sup>4</sup>, Medical Institute, Belgorod State University, Belgorod, Russia  
Julia N. Gomelyak<sup>5</sup>, Medical Institute, Belgorod State University, Belgorod, Russia

healthy", the rest have chronic diseases and various kinds of deviations in the state of health. Sociophobia in first-year medical students was observed in 28% of cases, and in 19.3% in pedagogical (Ruzhenkova et al. 2018a) and was more common among foreign students compared to local (Ruzhenkova 2018). The greatest number of social phobia cases was in 1st-year foreign medical students (45,2%) with reduction to 14.9% in the fifth year (Ruzhenkova et al. 2018b).

Social phobias limit opportunities for feedback and can lead to the negative self-perceptions (Henderson et al. 2014), interfere with social adaptation and career growth, leaving the patient socially isolated in some cases (Stein and Vythilingum 2015). SP is highly comorbid with other anxiety disorders, depression, alcohol abuse (Jensen et al. 2013) and pathological depersonalization (Michal et al. 2013; Husky et al. 2018; Rani et al. 2018).

### Objectives

This study aimed to verify the prevalence of social phobias and comorbid disorders among medical students to develop recommendations for the prevention.

### Material and Methods

During 2016-2018 a sample of 724 Russian-speaking students of 1-6 years of education in the medical institute was examined. Students were at the age of 17-26 ( $20.5 \pm 1.8$ ) years.

Sample included 537 (74.2%) female and 187 (25.8%) male.

The main research methods were medico-sociological (an author's questionnaire containing socio-demographic information, information about attitude to study and future professions), psychometric (SPIN-test for social phobia, GAD-7 for verification of generalized anxiety disorder, DASS-21 test to study the severity of depression, anxiety and stress). A clinical interview was conducted with students who showed significant symptoms of social phobia, anxiety, depression, and stress. The study design was approved by the Ethical Committee of the Medical Institute of the Belgorod State National Research University, Protocol №6 dated 21.01.2016

Statistical processing was carried out using non-parametric statistics (the Mann-Whitney test for comparing two independent groups, the  $\chi^2$  criterion with the Yates correction for the contingency tables 2x2, the Spearman rank correlation coefficient) (Yatham et al. 2018; Canals et al. 2019).

The SPIN test revealed various symptoms of social phobia in 244 (33.7%) students (Table 1). Symptoms of social phobia more often ( $\chi^2=9.907$ ;  $p=0.0025$ ;  $OR=1.858$  95%;  $CI=1.25-2.76$ ) were observed in female – 199 (37.1%) cases, than male – 45 (24.1%) cases. The odds ratio indicates that the probability of detecting social phobia in females is almost 2 times higher than in males.

**Table 1.**  
*Social phobia by severity (SPIN-test)*


№	The severity of social phobia	Male (n=187)		Female (n=537)		TOTAL (n=724)	
		n	%	n	%	n	%
1	None	142	75.9	338	62.9	480	66.3
2	Mild	24	12.8	111	20.7	135	18.6
3	Moderate	14	7.5	60	11.2	74	10.2
4	Severe	5	2.7	19	3.5	24	3.3
5	Very Severe	2	1.1	9	1.7	11	1.5
TOTAL		187	100.0	537	100.0	724	100.0

At the same time, a mild degree of social phobia, which did not affect social adaptation, was typical for 18.6% of the students: 20.7% female and 12.8% male (the differences are not statistically significant). Clinically significant

### Results

The GAD-7 test revealed symptoms of generalized anxiety in 401 (55.4%) students (Figure 1).

severity of social phobia (with difficulty and violation in varying degrees of social adaptation) was observed in 15% of cases: 16.4% female and 11.3% male (the differences are not statistically significant).



**Fig. 1.**  
Generalized anxiety by severity

The ratio of social phobia with signs of generalized anxiety disorder (GAD) is presented in Table 2.

**Table 2.**

*The ratio of social phobia with generalized anxiety disorder (The calculation of the proportion of GAD was carried out in rows in relation to social phobia)*


GAD-7	None		Mild		Medium		Severe		TOTAL	
	n	%	n	%	n	%	n	%	n	%
None	254	52.9	161	33.5	42	8.8	23	4.8	480	66.3
Mild	47	34.8	57	42.2	19	14.1	12	8.9	135	18.6
Moderate	19	25.7	31	41.9	18	24.3	6	8.1	74	10.2
Severe	3	12.5	11	45.8	6	25.0	4	16.7	24	3.3
Very Severe	–	–	6	54.5	3	27.3	2	18.2	11	1.5
TOTAL	323	44.6	266	36.7	88	12.3	47	6.5	724	100.0

As can be seen from Table 2, in general a sociophobia (revealed in 244 cases) were combined with generalized anxiety in 175 (71.7%) cases: 43% were mild, 18.9% were moderate, and 9.8% were severe. At the same time, with mild expressed sociophobia, generalized anxiety occurred in 65.2%, with a moderate in 74.3%, with severe – in 87.5% and with an extremely severe – in 100% of cases. In cases of lack of social phobia, generalized anxiety was characteristic of 47.1% (mostly easily expressed - 33.5%) of the examined medical students. The differences in the proportion of generalized anxiety in the presence and absence of social phobia are statistically significant ( $\chi^2=38.75$ ;  $p=0.0005$ ; OR =2.851 95%; CI=2.02-4.03). The odds ratio indicates that the probability of having generalized anxiety with varying severity of

social phobia is almost 3 times higher than in the absence of social phobia. Moderate and severe generalized anxiety was observed in 70 (21.6%) students with social phobia.

Correlation analysis revealed a weak significant correlation dependence of social phobia and generalized anxiety ( $r = 0.343$ ,  $p = 0.000$ ).

The DASS-21 test revealed depression of varying severity in 220 (30.4%) students. The distribution of depression by severity is presented in Figure 2.



**Fig. 2.**  
Depression by severity

As can be seen from the diagram, a severe and extremely severe depression was observed

in 7.5% of students, moderate - in 10.6% and mild - in 12.3%.

The ratio of social phobia with depression (DASS-21 test) is presented in Table 3.

**Table 3.**

The ratio of social phobia with depression (Calculation of the proportion of depression was carried out in rows in relation to social phobia)

Depression	None		Mild		Moderate		Severe		Very Severe		TOTAL	
	n	%	n	%	n	%	n	%	n	%	n	%
None	391	81.5	51	10.6	25	5.2	9	1.9	4	0.8	480	66.3
Mild	4	54.8	20	14.8	26	19.3	10	7.4	5	3.7	135	18.6
Moderate	32	43.2	12	16.2	18	24.3	8	10.8	4	5.4	74	10.2
Severe	6	25.0	6	25.0	5	20.8	1	4.2	6	25.0	24	3.3
Very Severe	1	9.1	–	–	3	27.3	4	36.4	3	27.3	11	1.5
TOTAL	504	69.6	89	12.3	77	10.6	32	4.4	22	3.0	724	100


In 131 (53.7%) cases (out of 244), social phobias were combined with varying degrees of depression: in 15.6% were mild, in 21.3% – moderate, and in 16.8% – severe and extremely severe. The quantity of moderate, severe and extremely severe depression was 71%. At the same time, depression was recorded in 45.1% with mild social phobia, in 56.8% – with moderate, in 75% – with severe and 90.9% – with extremely severe.

In the absence of social phobia, depression occurred in 89 (18.5%) students (more than half of them - 57.3% in mild severity). The differences in the share of depression in the

presence and absence of social phobia are statistically significant ( $\chi^2=92.815$ ;  $p=0.0005$ ; OR =5.094 95%, CI =3.6-7.3). The odds ratio indicates that the probability of having depression with varying degrees of social phobia is 5 times higher than without it.

Correlation analysis revealed the average level of correlation between social phobia and depression ( $r = 0.560$ ,  $p = 0.000$ ).

Anxiety of varying severity (DASS-21 test) was detected in 243 (33.6%) students. The distribution of anxiety by severity is presented in Figure 3.



**Fig. 3.**  
Anxiety by severity

As can be seen from chart 3, 18.4% of students had a clinical level (moderate, severe and extremely severe) of anxiety. Correlation analysis revealed the average correlation

between generalized anxiety (GAD-7) and anxiety according to the DASS-21 test ( $r=0.517$ ,  $p=0.000$ ). The ratio of social phobia with anxiety (test DASS-21) is presented in Table 4.

**Table 4.**


*The ratio of social phobia with anxiety (The calculation of the proportion of anxiety was carried out in rows in relation to social phobia)*

Anxiety	None		Mild		Moderate		Severe		Very Severe		TOTAL	
	n	%	n	%	n	%	n	%	n	%	n	%
None	374	77.9	57	11.9	24	5.0	9	1.9	16	3.3	480	66.3
Mild	67	49.6	33	24.4	18	13.3	8	5.9	9	6.7	135	18.6
Moderate	30	40.5	16	21.6	14	18.9	5	6.8	9	12.2	74	10.2
Severe	8	33.3	4	16.7	3	12.5	3	12.5	6	25.0	24	3.3
Very Severe	2	18.2	–	–	2	18.2	2	18.2	5	45.5	11	1.5
TOTAL	481	66.4	110	15.2	61	8.4	27	3.7	45	6.2	724	100

It was found that in total, out of 244 people with social phobia, 137 (56.1%) had an anxiety. Anxiety was recorded in 50.4% of cases with mild social phobia, in 59.5% – with moderate, in 66.7% – with severe and in 81.8% – with extremely severe. In addition, in 106 (22.1%) cases, anxiety was encountered among students with a lack of social phobia, in half of them (11.9%), it was mild. The differences in the frequency of occurrence of anxiety in students with and without social phobia are statistically significant ( $\chi^2=82.661$   $p=0.0005$ ;  $OR=4.518$  95%;  $CI=3.2-6.4$ ). The odds ratio indicates that the probability of having anxiety with varying degrees of social phobia is 4.5 times higher than without it. Correlation analysis revealed a weak correlation between social phobia and anxiety according to the DASS-21 test ( $r = 0.496$ ,  $p = 0.000$ ).

Significant differences in the severity of anxiety in GAD-7 and DASS-21 in the total population of the subjects (+ 21.8% in GAD-7) and in students with sociophobia (+ 15.6% in GAD-7) are associated with some overdiagnoses of anxiety by the GAD test -7 and underdiagnosis of DASS-21, which is determined by the structure of the questions: in the GAD-7 test, all questions concern only the psychic sphere, and in DASS-21, 4 out of 7 questions concern somato-vegetative anxiety manifestations.

Stress of varying severity (DASS-21 test) was detected in 254 (35.1%) students. The distribution of stress by severity is presented in Figure 4. It shows that the clinical level of stress (moderate, severe and extremely severe) was found in 23.5% of students — in fact, one out of four.



**Fig. 4.**  
*The stress by severity*

The ratio of social phobia with stress (DASS-21 test) is presented in Table 5.

**Table 5.**  
*The ratio of social phobia with stress (Calculation of the share of stress was carried out in rows in relation to social phobia)*

Stress	None		Mild		Moderate		Severe		Very Severe		TOTAL	
	n	%	n	%	n	%	n	%	n	%	n	%
None	364	75.8	44	9.2	44	9.2	19	4.0	9	1.9	480	66.3
Mild	70	51.8	18	13.3	18	13.3	20	14.8	9	6.7	135	18.6
Moderate	29	39.2	17	30.0	17	30.0	7	9.5	4	5.4	74	10.2
Severe	7	29.2	6	25.0	3	12.5	7	29.2	1	4.2	24	3.3
Very Severe	–	–	1	9.1	1	9.1	5	45.5	4	36.4	11	1.5
TOTAL	470	64.9	86	11.9	83	11.5	58	8.0	27	3.7	724	100

It was found that in stress was in 138 (56.6%) of students with social phobia: in 48.1% with mild, 60.8% – with moderate, 70.8% – with severe and 100% with extremely severe.

For students without signs of social phobia, stress was observed in 24.2% of cases ( $\chi^2=73.106$ ;  $p=0.0005$ ;  $OR=4.086$  95%;  $CI =2.9-5.8$ ). The odds ratio indicates that the probability of having a stress in presence of varying degrees of social phobia is 4 times higher than without it.

In total, of all the students which was examined, 566 (78.4%) reported on their adherence to the medical profession, and if there was a choice, they would enroll at the Medical Institute again, while significant number of students – 156 (21.6%) – every fifth, would choose a different profession. There were 70

(28.7%) among those with social phobia, and 86 (17.9%) in the group without social phobia. The differences are statistically significant ( $\chi^2=10.476$ ;  $p=0.002$ ;  $OR=1.843$  95%;  $CI =1.3-2.7$ ). The odds ratio indicates that the probability of disappointment in the medical profession is almost 2 times higher among those who have a social phobia than among students without it.

The study of the correlation between the severity of social phobia and conviction in the correctness of the choice of the medical profession as well as the attractiveness of the medical activity revealed a significant weak inverse relationship: respectively,  $r=-0.229$  and  $r=-0.216$  with  $p=0.000000$ . Thus, sociophobia has a negative impact on the attitude of students to the medical profession and form doubt on the

correctness of its choice. Moreover, the probability of disappointment in the medical profession is almost 2 times higher among students with social phobia than without it.

## Conclusion

Thus, the study showed that the symptoms of sociophobia occur in 33.7% of undergraduate medical students: it is statistically significantly more often in female (37.1%) than male (24.1%). At the same time, a clinically significant degree of manifestation of social phobia (with difficulty and violation in varying degrees of social adaptation) was observed in 15% of cases, equally often among males and females. Comorbidity of social phobia with GAD was 71.7%, with depression - 53.7%, with anxiety - 56.1% and with stress - 35.1%. In students with no signs of social phobia, GAD occurred in 47.1% of cases, depression - in 18.5%, anxiety - in 22.1%, and stress in 24.2% of cases.

Significant correlations revealed between the social phobia severity level and GAD, depression, anxiety and stress severity levels.

The probability of having GAD is almost 3 times, depression is 5 times, anxiety is 4.5 times and stress is 4 times higher among students with social phobia, than in cases without it. In addition, it has been established that social phobia has a negative influence on the students' attitude to the medical profession and forms doubt on the correctness of their choice. In addition, the probability of disappointment in the medical profession is almost 2 times higher among students with social phobia, than without it.

This fact should be taken into account conducting vocational guidance work and such students should be advised to consult a medical psychologist.

## Recommendations

It is possible that a person-centered psychotherapy, aimed at eliminating the symptoms of social phobia and changing the negative attitude towards the medical profession may be needed, such psychotherapy methods can be investigated in further studies. This study was conducted among medical students, further works can investigate different groups under different conditions to develop more comprehensive recommendations for the prevention of social phobia.

## References

- Albuquerque J, Deshauer D 2002. Social Anxiety Disorder: A Syndrome with Many Faces. *The Canadian Journal of CME*, 6: 87-99.
- Canals J, Voltas N, Hernández-Martínez C, Cosi S, Arijá V 2019. Prevalence of DSM-5 anxiety disorders, comorbidity, and persistence of symptoms in Spanish early adolescents. *European child & adolescent psychiatry*, 28(1): 131-43.
- Cremers HR, Roelofs K 2016. Social anxiety disorder: a critical overview of neurocognitive research. *Wiley Interdisciplinary Reviews: Cognitive Science*, 7(4): 218-232.
- Essau CA, Lewinsohn PM, Lim JX, Moon-ho RH, Rohde P 2018. Incidence, recurrence and comorbidity of anxiety disorders in four major developmental stages. *Journal of affective disorders*, 228: 248-53.
- Henderson L, Gilbert P, Zimbardo P 2014. *Shyness, social anxiety, and social phobia. In Social Anxiety*. Cambridge, Massachusetts: Academic Press.
- Hudson JL, Rapee RM 2000. The Origins of social phobia. *Behavior Modification*, 24(1): 102-129.
- Husky MM, Mazure CM, Kovess-Masfety V 2018. Gender differences in psychiatric and medical comorbidity with post-traumatic stress disorder. *Comprehensive psychiatry*, 84: 75-81.
- Jensen V, Hougaard E, Fishman DB 2013. Sara, a social phobia client with sudden change after exposure exercises in an intensive cognitive-behavior group therapy: a case-based analysis of mechanisms of change. *Pragmatic Case Studies in Psychotherapy*, 9(3): 275-336.
- Michal M, Heidenreich T, Engelbach U, Lenz C, Overbeck G, Beutel M, Grabhorn R 2013. Depersonalization, social phobia and shame. *Psychotherapie, Psychosomatik, Medizinische Psychologie*, 56: 383-389.
- Mohammadi MR, Alavi SS, Ahmadi N, Khaleghi A, Kamali K, Ahmadi A, Hooshyari Z, Jaberghaderi N, Nazaribadie M, Farshidfar Z, Kaviani N 2019. The prevalence, comorbidity and socio-demographic factors of depressive disorder among Iranian children and adolescents: To identify the main predictors of depression. *Journal of affective disorders*, 247: 1-0.
- Nikfarjam M, Heidari-Soureshjani S, Khoshdel A, Asmand P, Ganji F 2017. Comparison of spiritual well-being and social health among the students attending group and individual religious rites. *World Family Medicine Journal: Incorporating the Middle East Journal of Family Medicine*, 99(5480): 1-6.
- Olivares-Olivaresa PJ, Olivaresa J, Maciàb D, Maciàc A, Montesinosd L 2016. Community versus Clinical Cognitive-Behavioral Intervention in Young-Adult Spanish

- Population with Generalized Social Phobia. *Terapia Psicológica*, 34(1): 23-30.
- Rani R, Bala N, Garg PD, Bansal A 2018. A Study of Anxiety Disorders in Terms of Severity and Prevalence in Patients of Major Depressive Disorder in a Tertiary Care Centre. *International Journal of Innovative Research in Medical Science*, 3(11): 2311-27.
- Rendueles C 2017. *Sociophobia: Political change in the digital utopia*. New York, USA: Columbia University Press.
- Rowa K, Antony MM 2005. Psychological treatments for social phobia. *Canadian Journal of Psychiatry*, 50(6): 308-316.
- Ruzhenkova VV 2018. Academical stress as a risk factor for the formation of addictive behavior, anxiety and depressive disorders in foreign medical students. *Research Result: Medicine and Pharmacy series*, 4(2): 55-68.
- Ruzhenkova VV, Ruzhenkov VA, Morozova EN, Lukyantseva IS 2018. Affective disorders associated with educational stress among foreign medical students (Dynamics over a 6-year period of study). *Indo American journal of pharmaceutical sciences*, 5(7): 7021-7027.
- Ruzhenkova VV, Tarabaeva VB, Ruzhenkov VA, Lukyantseva IS 2018. Medical and psychological characteristics of the 1st – year students of Medical and Pedagogical Institutes and their features of educational adaptation. *Drug Invention Today*, 10(3): 3240-3246.
- Savoia MG, de Barros Neto TP, Vianna AM, Bernik M 2010. Evaluation of personality traits in social phobia patients. *Revista de Psiquiatria Clinica*, 37(2): 57-9.
- Shen Y, Zhang Y, Chan BS, Meng F, Yang T, Luo X, Huang C 2020. Association of ADHD symptoms, depression and suicidal behaviors with anxiety in Chinese medical college students. *BMC psychiatry*, 20: 1-9.
- Stein DJ, Vythilingum B 2015. *Anxiety Disorders and Gender*. Cham, Switzerland: Springer Publication.
- Thai J, Zhao XD, Huang L, Zhong Y, Peng H, Koran J 2019. The Positive Association Between Empathy and Self-Esteem in Chinese Medical Students: A Multi-Institutional Study. *Frontiers in psychology*, 10: 1921-37.
- Wersebe H, Lieb R, Meyer AH, Mische M, Mikoteit T, Imboden C, Hoyer J, Bader K, Hatzinger M, Gloster AT 2018. Well-being in major depression and social phobia with and without comorbidity. *International Journal of Clinical and Health Psychology*, 18(3): 201-8.
- Yatham S, Sivathasan S, Yoon R, da Silva TL, Ravindran AV 2018. Depression, anxiety, and post-traumatic stress disorder among youth in low and middle income countries: a review of prevalence and treatment interventions. *Asian journal of psychiatry*, 38: 78-91.
- Bicer, A., Perihan, C., & Lee, Y. (2018). The Impact of Writing Practices on Students' Mathematical Attainment. *International Electronic Journal of Mathematics Education*, 13(3), 305-313. <https://doi.org/10.12973/iejme/3922>.
- Vargas-Hernández, J. G. (2016). THE QUESTION OF CHANGING THE CONCEPT, ROLE AND FUNCTIONS OF STATE. *Humanities & Social Sciences Reviews*, 4(1), 08-19. <https://doi.org/10.18510/hssr.2016.412>