



A Nutrigenetic approach to examine the relationship between vitamin B12 status and cardio-metabolic traits in multiple ethnic groups – findings from the GeNulne Collaboration

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Vimalleswaran, K. S. and Surendran, S. (2021) A Nutrigenetic approach to examine the relationship between vitamin B12 status and cardio-metabolic traits in multiple ethnic groups – findings from the GeNulne Collaboration. *Nutrition Bulletin*. ISSN 1471-9827 doi: <https://doi.org/10.1111/nbu.12494>
Available at <https://centaur.reading.ac.uk/97035/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1111/nbu.12494>

Publisher: Wiley-Blackwell

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

A nutrigenetic approach to examine the relationship between vitamin B12 status and cardio-metabolic traits in multiple ethnic groups: Findings from the GeNulne Collaboration

S. Surendran^{*†}  and K. S. Vimalaswaran^{*‡} 

^{*}Hugh Sinclair Unit of Human Nutrition, Department of Food and Nutritional Sciences, University of Reading, Reading, UK;

[†]Faculty of Health and Medical Sciences, School of Biosciences and Medicine, University of Surrey, Guildford, UK;

[‡]Institute for Food, Nutrition and Health (IFNH), University of Reading, Reading, UK

Abstract

Low vitamin B12 concentration has been shown to be a risk factor for cardio-metabolic traits in numerous observational studies; however, the relationship has remained inconsistent. It is possible that certain genotypes jointly contribute to cardio-metabolic diseases and vitamin B12 deficiency, and these may be modulated by dietary factors. The main objective of this article is to summarise the findings from the *GeNulne* (Gene-Nutrient Interactions) Collaboration on the effect of gene–nutrient interactions on vitamin B12 concentrations and cardio-metabolic disease risk factors in population-based studies from different ethnic groups. Interactions between vitamin B12-related single-nucleotide polymorphisms (SNPs) and protein energy intake (%) on waist circumference ($P_{\text{interaction}} = 0.002$) and body fat percentage ($P_{\text{interaction}} = 0.034$) were observed in Sri Lankan and Indonesian populations, respectively. In the study in Brazilian adolescents, the metabolic and vitamin B12-related SNPs showed a significant interaction with dietary carbohydrate and protein intakes on oxidised low-density lipoprotein cholesterol and homocysteine concentrations, respectively. In the Asian Indian population, an association between obesity-related SNPs and vitamin B12 concentrations ($P = 0.018$) was observed. In summary, these studies in multiple ethnic groups show that the association between genetically low vitamin B12 concentrations and cardio-metabolic traits may be modified by dietary intake. Further studies utilising larger sample sizes are needed to confirm or refute our findings.

Keywords: GRS, metabolic traits, nutrigenetics, obesity, SNP, vitamin B12 pathway

Correspondence: Prof. Karani Santhanakrishnan Vimalaswaran, Professor of Nutrigenetics & Nutrigenomics, Hugh Sinclair Unit of Human Nutrition, University of Reading, Reading, RG6 6DZ, UK.

E-mail: v.karani@reading.ac.uk

Dr. Shelini Surendran, Teaching Fellow, Biosciences Foundation, School of Biosciences and Medicine, University of Surrey, Guildford, Surrey, GU2 7XH, UK.

E-mail: s.surendran@surrey.ac.uk

Introduction

Vitamin B12 is an essential water soluble micronutrient, which participates as a cofactor for the synthesis of DNA, fatty acids and myelin (Smith *et al.* 2018). It has long been known that vegans, lacto-ovo vegetarians and older individuals are at risk of vitamin B12

deficiency (Green *et al.* 2017). Vitamin B12 deficiency can also be caused by having inadequate amounts of intrinsic factor (IF), gastric atrophy, intestinal disease, gastric surgery, bacterial overgrowth in the small intestine, alcohol consumption, a tapeworm infection, drug–nutrient interactions and some genetic defects (Institute of Medicine 2000; Allen 2008; O’Leary & Samman 2010). Additionally, alarmingly high prevalence rates of low plasma vitamin B12 status have been recognised to exist in certain regions of the world including the Indian subcontinent, Mexico, Central and South America and selected areas in Africa (Stabler & Allen 2004).

Vitamin B12 functions as a coenzyme of several key enzymes in one-carbon metabolism, which in turn are essential for nucleotide synthesis and methylation reactions (Selhub 2002). The two active forms of vitamin B12, methylcobalamin and 5-deoxyadenosylcobalamin, are essential coenzymes for the homeostasis of methionine and methylmalonic acid (MMA), respectively (Koury & Ponka 2004; Sobczyńska-Malefora *et al.* 2014). Observational studies have shown that low vitamin B12 concentrations are accompanied by a wide range of chronic diseases and conditions, including obesity, insulin dysregulation and adverse cardio-metabolic outcomes (Baltaci *et al.* 2013; Mahalle *et al.* 2013; Adaikalakoteswari *et al.* 2014; Narang *et al.* 2016; Chakraborty *et al.* 2018; Guarnizo-Poma *et al.* 2018; Jayashri *et al.* 2018). The aetiology between low vitamin B12 concentrations and metabolic phenotypes is still under debate and could be the result of several mechanisms. Vitamin B12 deficiency could possibly lead to adipocyte dysfunction by modulating lipid metabolism and enhancing cellular inflammation (Kumar *et al.* 2013). Several studies have suggested that vitamin B12 deficiency is more prevalent in individuals who are obese as result of either low dietary vitamin B12 consumption (Thomas-Valdés *et al.* 2017; Sun *et al.* 2019) or increased vitamin B12 demands secondary to an increased surface area of the body (Wall 1998; Pinhas-Hamiel *et al.* 2006).

Genetic studies have implicated several gene loci associated with circulating vitamin B12 (Surendran *et al.* 2018). To date, only three Mendelian randomisation studies have explored the relationship between a genetically determined decrease in serum vitamin B12 concentrations on cardio-metabolic traits such as body mass index (BMI) (Allin *et al.* 2017), blood pressure (Husemoen *et al.* 2016) and cardio-metabolic risk (Moen *et al.* 2018). However, these studies were unable to support the causal role of decreased serum vitamin B12 concentrations with these

cardio-metabolic traits. Further to this, dietary factors play an important role in the development of vitamin B12 deficiency, but this relationship may differ across countries, due to the variation in food consumed worldwide (Vimalleswaran 2017). Studies have shown that the intrauterine imbalance of vitamin B12 and folate can affect DNA methylation and ‘programme’ the offspring to develop metabolic disorders later in life (Yajnik & Deshmukh 2012) providing evidence for interactions between genes and nutrients in the development of cardio-metabolic disease, including type 2 diabetes, hypertension, obesity and cardiovascular disease (CVD). The aim of this article is to provide an overview of the findings from the four studies in the gene–nutrient interactions (*GeNuIne*) Collaboration that used a genetic approach to explore the relationship between cardio-metabolic disease risk factors and vitamin B12 status and to investigate whether these relationships were modified by macronutrient intake (carbohydrate, fat and protein intake) (Vimalleswaran 2017).

Importance of studying gene–diet interactions in different genetic groups

It has been established that previous genetic studies looking at vitamin B12 status in healthy adults, especially large-scale ones, have been unable to capture the level of diversity which exists worldwide, as they are mainly based on individuals of European ancestry (Surendran *et al.* 2018). The under-representation of diverse ethnic groups hampers our full understanding of the genetic architecture of vitamin B12 concentrations (Sirugo *et al.* 2019). Furthermore, the limited genetic data on non-Caucasian populations in relation to genetic susceptibility to vitamin B12 deficiency can also impede our ability to translate genetic research into clinical care and will exacerbate health inequalities (Sirugo *et al.* 2019). Given that vitamin B12 status can also be determined by environmental factors, it is also important to explore gene–diet interactions in different ethnic groups. It is important to note that different ethnic groups respond differently to specific dietary interventions (Vimalleswaran 2017). Therefore, using estimates of genetic risk for vitamin B12 deficiency from European-based studies in non-Europeans may result in an inaccurate assessment of risk of vitamin B12 deficiency and could result in an inappropriate environmental intervention (dietary or physical activity) in under-studied populations. To address all these issues, the *GeNuIne* Collaboration (Vimalleswaran 2017) was initiated to investigate the effect of

gene–lifestyle interactions on cardio-metabolic disease risk factors using population-based studies from various ethnic groups in lower-middle income countries (LMICs): Brazil (Surendran & Vimalaswaran 2019), Sri Lanka (Surendran *et al.* 2019b), India (Surendran *et al.* 2019c) and Indonesia (Surendran *et al.* 2019a). Another objective of the *GeNuIne* Collaboration was to look at gene–lifestyle interactions on vitamin B12 concentrations (Matusheski *et al.* 2021). A summary of the general characteristics of the populations and the SNPs investigated are provided in Tables 1 and 2.

Impact of genes and diet on homocysteine, vitamin B12, folate and lipids in a Brazilian adolescent population

Epidemiological studies have suggested that increased homocysteine concentrations may contribute to up to 25% of coronary events (Stanger *et al.* 2004) and have been shown to inversely correlate with B-complex vitamins, such as folate and vitamin B12 (Surendran & Vimalaswaran 2019). While B-complex vitamins have a role in the reduction of homocysteine

Table 1 Single-nucleotide polymorphisms (SNPs) under investigation: A comparison of studies in the Brazilian, Sri Lankan, Indian and Indonesian populations

	Brazil (n = 113) ^a	Sri Lanka (n = 109)	India study (n = 548) ^c	Indonesia study (n = 117) ^b
B12-related SNPs analysed	<ul style="list-style-type: none"> Fucosyltransferase [<i>FUT2</i>]-rs602662 Transcobalamin 2 [<i>TCN2</i>]-rs1801198 5-methyltetrahydrofolate-homocysteine methyltransferase or methionine synthase [<i>MTR</i>]-rs1805087 5-methyltetrahydrofolate-homocysteine methyltransferase reductase or methionine synthase reductase [<i>MTRR</i>]-rs1801394 Betaine-homocysteine S-methyltransferase [<i>BHMT</i>]-rs3797546 and rs492842 methylenetetrahydrofolate reductase [<i>MTHFR</i>]-rs1801131 and rs1801133 	<ul style="list-style-type: none"> <i>MTHFR</i>-rs1801133 Carbamoyl-phosphate synthase 1 [<i>CPS1</i>]-rs1047891 Cubulin [<i>CUBN</i>]-rs1801222 CD320 molecule [<i>CD320</i>]-rs2336573 <i>TCN2</i>-rs1131603 Citrate lyase beta like [<i>CLYBL</i>]-rs41281112 <i>FUT2</i>-rs602662 Transcobalamin 1 [<i>TCN1</i>]-rs34324219 Fucosyltransferase 6 [<i>FUT6</i>]-rs778805 Methylmalonyl-CoA mutase [<i>MUT</i>]-rs1141321 	Not applicable	<ul style="list-style-type: none"> <i>MTHFR</i>-rs1801133 <i>CPS1</i>-rs1047891 <i>CUBN</i>-rs1801222 <i>CD320</i>-rs2336573 <i>TCN2</i>-rs1131603 <i>FUT2</i>-rs602662 <i>TCN1</i>-rs34324219 <i>FUT6</i>-rs778805 <i>MUT</i>-rs1141321
Metabolic disease SNPs analysed	Catechol-o-methyl transferase [<i>COMT</i>]-rs4680 and rs4633	<ul style="list-style-type: none"> Fat mass and obesity-associated [<i>FTO</i>]-rs9939609 and rs8050136 Melanocortin 4 Receptor [<i>MC4R</i>]-rs17782313 and rs2229616 Transcription factor 7-like 2 [<i>TCF7L2</i>]-rs12255372 and rs7903146 Potassium voltage-gated channel subfamily J member 11 [<i>KCNJ11</i>]-rs5219 Calpain 10 [<i>CAPN10</i>]-rs3792267, rs2975760 and rs5030952 	<i>FTO</i> - rs8050136 and rs2388405	<ul style="list-style-type: none"> <i>FTO</i>-rs9939609 and rs8050136 <i>MC4R</i>-rs17782313 and rs2229616 <i>TCF7L2</i>-rs12255372 and rs7903146 <i>KCNJ11</i>-rs5219 <i>CAPN10</i>-rs3792267 and rs5030952

^aBrazilian adolescents who were overweight/obese and/or were previously diagnosed with dyslipidaemia, but not with cardiovascular disease, were included

^bOnly women were included for the Indonesian study

^cCase-control study for type 2 diabetes in men and women

Table 2 General characteristics, dietary intake and biochemical levels: A comparison of studies in the Brazilian, Sri Lankan, Indian and Indonesian populations

	Brazil (n = 113) ^a	Sri Lanka study (n = 109)	India study (n = 548) ^c	Indonesia study (n = 117) ^b
Study design	Cross-sectional	Cross-sectional	Case-control	Cross-sectional
General characteristics				
Age (years)	14 ± 2	38 ± 7	49.39 ± 11.45	40 ± 10
BMI (kg/m ²)	24.0 ± 4.9	24.6 ± 4.1	26.75 ± 5.04	25.1 ± 4.2
Waist-to-hip ratio	N/A	0.92 ± 0.11	0.90 ± 0.09	N/A
Fat (%)	N/A	27.25 ± 7.37	N/A	35.70 ± 7.00
Physical activity levels (%)				
Low	31	72.5	82.0	39.3
Medium	69	19.3	16.2	49.6
High		8.3	1.8	11.1
Dietary intake				
Total energy (Kcal/day)	2522 ± 586	2098 ± 456	2597 ± 773	1774 ± 609
Dietary carbohydrate (%)	47.7 ± 20.6	69.6 ± 8.8	64.3 ± 6.3	54.1 ± 9.4
Dietary protein (%)	17.0 ± 8.4	11.3 ± 2.3	11.3 ± 1.1	16.9 ± 3.3
Dietary fat (%)	25.4 ± 13.2	21.9 ± 5.3	23.8 ± 4.7	28.9 ± 8.0
Total fibre (g)	N/A	16.8 ± 8.2	32.2 ± 11.3	8.8 ± 4.5
Clinical parameter				
24 hours ambulatory systolic BP (mm Hg)	N/A	120 ± 15	129 ± 20	113 ± 9
24 hours ambulatory diastolic BP (mm Hg)	N/A	75 ± 16	80 ± 12	77 ± 6
Triacylglycerol (mg/dl)	94.05 ± 54.16	144.19 ± 86.81	146.54 ± 116.88	97.67 ± 42.80
HDL (mg/dl)	46.29 ± 11.79	42.56 ± 8.24	40.89 ± 8.85	58.99 ± 10.20
LDL (mg/dl)	90.28 ± 21.00	134.03 ± 28.60	114.37 ± 35.56	127.77 ± 39.17
VLDL (mg/dl)	18.85 ± 10.82	28.84 ± 17.36	N/A	N/A
Oxidised-LDL (U/l)	6.42 ± 13.69	N/A	N/A	N/A
Fasting plasma glucose (mg/dl)	N/A	86 ± 13	116 ± 49	92 ± 20
Fasting serum insulin (μU/ml)	N/A	9.9 ± 7.2	9.23 ± 6.25	32 959 ± 26 327
Glycated haemoglobin (%)	N/A	5.4 ± 0.5	6.5 ± 1.7	N/A
B-vitamin biomarker status				
Vitamin B12 (pg/ml)	520 ± 232	516 ± 180	417 ± 255	591 ± 579
Homocysteine (μmol/l)	7.04 ± 2.99	N/A	13.67 ± 8.09	N/A
Folic acid (ng/ml)	11.02 ± 3.27	N/A	8.59 ± 5.81	N/A

BP, blood pressure; HDL, high density lipoprotein; LDL, low density lipoprotein; VLDL, very low density lipoprotein.

^aBrazilian adolescents who were overweight/obese and/or were previously diagnosed with dyslipidaemia, but not with cardiovascular disease, were included

^bOnly women were included for the Indonesian study

^cCase-control study for type 2 diabetes in men and women

concentrations, the effect of these vitamins on cardiovascular health remains ambiguous (Ntaios *et al.* 2009). Some studies have suggested that high folate and vitamin B12 status are associated with a lower risk of coronary heart disease (Voutilainen *et al.* 2001; Ishihara *et al.* 2008). Common variants in genes of the one-carbon metabolism pathway have been reported to influence concentrations of homocysteine, folate, vitamin B12 and lipids (Hazra *et al.* 2009). The aim of this study was to examine the association of ten SNPs involved in the one-carbon metabolism pathway with vitamin B12, folic acid, homocysteine and blood lipids, and to investigate whether environmental factors (dietary factors and physical activity levels)

modified the association of the SNPs in 113 Brazilian adolescents (aged 10–19 years) with cardiovascular risk (overweight or obese and/or previously diagnosed dyslipidaemia, but not with CVD).

Since several SNPs were analysed in all the studies, correction for multiple testing was applied. No gene–lifestyle interactions were observed on vitamin B12 concentrations. This study provided evidence for interactions between the metabolic *COMT* SNP rs4680 and carbohydrate intake on ox-LDL concentrations and the B12-associated *FUT2* SNP rs602662 and protein intake on homocysteine concentrations. This is the first study to provide novel gene–diet interactions at the *COMT* and *FUT2* gene loci, on ox-LDL and

homocysteine concentrations; hence, we have no other studies to compare our findings with. Given that ox-LDL and homocysteine are well-known independent risk factors for cardiovascular disease (Shenoy *et al.* 2014; Papageorgiou & Tousoulis 2016), these findings may have significant public health implications.

In addition to the genetic component of metabolic traits and markers of one-carbon metabolites, physical inactivity could be an important contributor that could interact with an individual's genetic predisposition. Our results showed that the *MTHFR* SNP rs1801131 showed a significant interaction with physical activity on folic acid concentrations. Folate plays a critical role in the methylation pathway and is involved in the methylation of DNA, creatine and acetylcholine, all of which are important for physical activity. It is important to note that physical activity levels may interact with folate metabolism by increasing intestinal folate absorption or stimulating the methionine synthase enzyme due to an increased metabolic demand and the associated possible increase in turnover of methylated molecules required for exercise (Kim *et al.* 2016). It is important that in the future, replications of this finding are made, preferably in a larger independent cohort with adequate statistical power utilising more direct measures of physical activity.

Impact of genes and diet on vitamin B12 concentrations and cardio-metabolic diseases in an Asian Sri Lankan population (GOOD study)

South Asians exhibit a unique phenotype collectively known as the 'South Asian Phenotype' which consists of higher levels of total and visceral fat, higher waist circumference and an increased susceptibility to type 2 diabetes (Mohan & Deepa 2006; Surendran *et al.* 2019b). Currently, over 34.4% of the Sri Lankan adult population are diagnosed as being overweight or obese (Katulanda *et al.* 2010). Although there is a strong genetic component to developing the 'South Asian Phenotype', consuming an unhealthy diet and leading a sedentary lifestyle can further contribute to this phenotype (Anjana *et al.* 2014; Vimaleswaran *et al.* 2016). Several studies have suggested that obesity is related to many micronutrient deficiencies including vitamin B12 (Astrup & Bügel 2010; Damms-Machado *et al.* 2012; Chakraborty *et al.* 2018). This is the first study to investigate the relationship between metabolic traits and vitamin B12 status using a gene-based approach in 109 Sinhalese

adults [*Genetics Of Obesity and Diabetes (GOOD)* study; 61 men and 48 women, aged 25–50 years], and to investigate whether these relationships were modified by lifestyle factors (dietary factors/physical activity levels).

The unweighted genetic risk score (GRS) was calculated for each participant based on the sum of risk allele counts across each SNP which predicted a lower vitamin B12 status (B12-GRS). A second, unweighted GRS was created using allele markers previously reported to be associated with metabolic disease traits (metabolic-GRS). A numerical value of 0, 1 or 2 was assigned to each SNP, which indicates the number of risk alleles on that SNP. The scores were then calculated by adding the number of risk alleles across each SNP. The risk allele score was then divided by the median score into those carrying ≤ 9 risk alleles versus those with ≥ 10 risk alleles for the B12 GRS and ≤ 8 risk alleles versus those with ≥ 9 risk alleles for the metabolic-GRS.

The findings from this study demonstrated that a genetically lowered vitamin B12 concentration may be associated with central obesity (waist circumference) in the presence of a dietary influence (protein energy intake %). Additionally, the metabolic-GRS interacted with carbohydrate energy intake (%) to influence waist-to-hip ratio levels, where individuals carrying more than 9 risk alleles had a higher waist-to-hip ratio among those in the highest tertile of carbohydrate energy percentage compared to those in the lowest tertile (mean \pm SD = 78.00 \pm 7.90%; Fig. 1).




Figure 1 Interaction between the metabolic-genetic risk score and carbohydrate energy intake (%) on waist-to-hip ratio ($P_{\text{interaction}} = 0.015$). Among those who consumed a high carbohydrate diet, individuals who carried nine or more risk alleles had significantly higher waist-to-hip ratios compared to individuals carrying eight or less risk alleles ($P = 0.035$)

Further studies are warranted to investigate the mechanisms underlying the effect of these GRSs on metabolic traits. Given that the daily consumption of protein is low and carbohydrate intake is high in Sri Lankan adults (Jayawardena *et al.* 2014), these findings, if confirmed in randomised control trials, may have significant public health implications in terms of revising dietary guidelines for this population, which could prevent central obesity and its related CVD-related outcomes.

Impact of genes and diet on vitamin B12 concentrations and metabolic diseases in an Asian Indian population (CURES study)

Although vitamin B12 deficiency has been linked to obesity (Pinhas-Hamiel *et al.* 2006; MacFarlane *et al.* 2011; Baltaci *et al.* 2013) and diabetes (Yajnik *et al.* 2008; Krishnaveni *et al.* 2009; Knight *et al.* 2015), no study to date has tested the genetic link between metabolic traits and vitamin B12 status in an Asian Indian population (Surendran *et al.* 2019c). Hence, this study examined whether dietary intake and physical activity levels modified associations between a GRS using two previously studied *FTO* SNPs (rs8050136 and rs2388405) and vitamin B12 concentrations and metabolic disease-related outcomes in 548 Asian Indians (220 normal glucose-tolerant individuals, 152 individuals with prediabetes and 176 individuals with type 2 diabetes). In this study, participants were randomly recruited from the *Chennai Urban Rural Epidemiology Study (CURES)*, a cross-sectional case-control epidemiological study conducted on a representative population of Chennai in Southern India (Deepa *et al.* 2003).

To date, the *FTO* (fat mass and obesity associated) gene has been the strongest obesity risk loci in several populations (Vimalaswaran & Loos 2010). Thus, the *FTO* gene was considered as a suitable candidate to establish the genetic link between obesity-related traits and vitamin B12 concentrations and to determine whether this relationship was modified by lifestyle factors. We found no significant interactions between the *FTO*-GRS and lifestyle factors (diet and physical activity levels) in this study. However, a novel finding of this study indicated the potential association between the *FTO*-GRS and vitamin B12 concentrations, after adjustment for potential confounders. Carriers of more than one risk allele for the *FTO*-GRS had lower vitamin B12 concentrations (mean \pm SD: 355 \pm 189 pg/ml), compared to individuals carrying zero risk alleles (mean \pm SD: 410 \pm 202 pg/ml). The mechanism

explaining the differences in vitamin B12 concentrations in the *FTO*-GRS could potentially be due to the *FTO* genotypes modulating gut microbiota and inducing metabolic inflammation, consequently impairing B12 absorption (Caesar *et al.* 2015; Chakraborty *et al.* 2018).

Given that low vitamin B12 concentrations in Asian Indians are common (Yajnik *et al.* 2006; Sivaprasad *et al.* 2016) and that 28–44% of Asians carry at least one copy of the *FTO* risk allele (Li *et al.* 2012), this study highlights the importance of considering obesity as a risk factor for vitamin B12 deficiency. This has implications for the possible targeting of relevant obesity prevention strategies, by focusing on increasing vitamin B12 intakes in obese or overweight individuals. Following such advice could substantially reduce vitamin B12 deficiency among Asian Indians.

Impact of genes and diet on vitamin B12 concentrations and metabolic diseases in an Indonesian women population (Minangkabau community: MINANG study)

Optimal vitamin B12 status is essential for women to maintain adequate maternal health and to avoid fetal developmental complications (Krishnaveni *et al.* 2009; Dror & Allen 2012; Surendran *et al.* 2019a). Additionally, low vitamin B12 concentrations have shown negative correlations with BMI in healthy women (Baltaci *et al.* 2017). Currently, the prevalence of low vitamin B12 status in healthy Indonesian women is unknown and studies on the relationship between low vitamin B12 status and obesity-related traits have yielded conflicting results (Wiebe *et al.* 2018). The Minangkabau population from Indonesia is of particular interest, as it is the largest matrilineal society in the world. Women in this society have relatively greater authority within the family compared to men and have a major role in dictating food choices and maintaining the welfare of the family (Stark 2013). In this population, GRSs based on nine B12-related SNPs and nine metabolic disease-related SNPs were constructed.

In the Indonesian study, a novel interaction between the B12-GRS and dietary fibre intake (g) on glycated haemoglobin was observed ($P_{\text{interaction}} = 0.042$; Fig. 2). Individuals who consumed a low-fibre diet (4.90 \pm 1.00 g/day) and carried ≥ 9 risk alleles for vitamin B12 deficiency had significantly higher HbA1C concentrations compared to those carrying ≤ 8 risk alleles.




Figure 2 Interaction between the B12-genetic risk score and dietary fibre intake (g) on log HbA1C (ng/ml; $P_{\text{interaction}} = 0.042$). Among those who consumed a low-fibre diet, individuals who carried 9 or more risk alleles had significantly higher concentrations of log HbA1C compared to individuals carrying eight or less risk alleles ($P = 0.025$)

Furthermore, interactions were also seen between the B12-GRS and protein (energy %) on log transformed body fat percentage.

Previous studies have shown that dietary fibre consumption is low in the Indonesian population, and thus, this finding, if confirmed in randomised controlled trials, could be the basis of a consumer education campaign centred around encouraging fibre intake to reduce HbA1C concentrations with those with genetic risk, which could improve glycaemic control in this population.

Ethnic-specific differences

The observations described in these studies confirm previous findings on population-based differences in vitamin B12 concentrations (McLean *et al.* 2008). The two South Asian populations demonstrated lower vitamin B12 concentrations than the Indonesian women population [Indian (417 ± 255 pg/ml) and Sri Lankan populations (516 ± 180 pg/ml) vs. Indonesian women (591 ± 579 pg/ml)]. Despite the Brazilian adolescent population exhibiting signs of cardiovascular risk, intermediate vitamin B12 concentrations were observed in this population (520 ± 232 pg/ml). It is difficult to generalise these findings meaningfully, as some of the populations included cohorts at risk of disease and furthermore the sample size of the population was limited (Table 2).

Several genetic loci have supported the presence of ethnic differences for metabolic traits and vitamin B12

status within the populations studied in this review. The *FTO* rs8050136 genotype is an example of genetic heterogeneity according to race. The minor allele 'A' of the SNP rs8050136 was present in 13% of Indian participants versus 23% of Indonesian participants and although India and Sri Lanka are geographically close, it was even more frequent in the Sinhalese population (34%).

The *FUT2* SNP rs602662 is one of the most commonly studied variants related to vitamin B12 status. The rs602662 SNP is an example of a B12 SNP that has demonstrated ethnic specificity, for example within the Indonesian population the minor allele frequency was extremely low (0.03%) in comparison to the Sri Lankan (31%) and Brazilian populations (41%). Although no genotyping errors were identified, the SNP rs602662 did not reach Hardy–Weinberg equilibrium (HWE), when using the chi-square test, within the Indonesian and Sri Lankan populations. It is possible that HWE was not reached in these populations due to the small sample size and the possibility of interbreeding (especially as consanguineous marriages are common in these populations).

How these findings could be translated into effective personalised and public health strategies

Obesity and perhaps vitamin B12 deficiency are important modifiable risk factors for a variety of chronic diseases. Obesity and vitamin B12 status are known to be associated, but the direction of the association and whether it is a causal relationship has been uncertain. Many genetic variants have been identified for vitamin B12 deficiency and metabolic diseases through candidate gene and genome-wide approaches (Bradfield *et al.* 2012; Willer *et al.* 2013; Surendran *et al.* 2018). However, it is possible that these genetic variants may not cause diseases without exposure to certain dietary factors (Rhee *et al.* 2012). Although this study has shown that low vitamin B12 concentrations are associated with metabolic diseases through a dietary interaction, the molecular and pathophysiological mechanisms remain unknown. It is important that mechanistic studies are carried out to determine how genetically low vitamin B12 concentrations interact with dietary factors to influence adipose tissue metabolism or how epigenetic mechanisms contribute to the development of metabolic diseases. Current literature also suggests that the genetic profile of an individual may shape the microbiome of the host, and an altered gut flora has been associated with vitamin B12 deficiency (Hall *et al.*




Figure 3 The main study findings of the *GeNuIne* Collaboration

2017; Surendran *et al.* 2018). This possibility requires investigation, given the known link between an altered distribution in gut microbiota and obesity. Hence, the combined application of nutrigenetics, nutrigenomics [dietary effects on gene expression (Ordovas *et al.* 2018)] and metabolomics (dietary effects on metabolite changes) is needed to clarify the gene–diet interactions identified in this project.

However, before the findings from these studies can be translated into personalised advice and/or public health strategies, they require confirmation from larger observational studies which include individuals from different ethnic groups and randomised controlled trials. It is difficult to truly isolate the macronutrient accountable for any nutrigenetic effects in observational studies, especially for fat and carbohydrates, as one macronutrient usually compensates the other (Sarzynski & Bouchard 2013). Therefore, dietary intervention studies are preferable as they can manipulate dietary intake more specifically. Dietary intervention studies also have the potential to avoid the measurement bias found within food frequency questionnaires and can demonstrate causal relationships between SNPs, diet and metabolic traits. Future studies should also measure intakes of specific dietary micronutrients to enable more detailed investigations of gene–diet interactions to be carried out. More studies that robustly measure both body composition (*e.g.* dual-energy X-ray absorptiometry, magnetic resonance

imaging and/or computed tomography scans) and body size, and which measure vitamin B12 status using markers such as homocysteine, MMA and holotranscobalamin to investigate the effects of adiposity on vitamin B12 status across all body sizes are also needed (Wiebe *et al.* 2018).

Conclusion

In conclusion, the *GeNuIne* Collaboration identified novel interactions between vitamin B12-related genes and dietary factors (in particular protein intake and fibre intake) on cardio-metabolic traits within Brazilian, Sri Lankan and Indonesian populations. In addition, within an Indian population, individuals who were genetically predisposed to obesity had lower vitamin B12 concentrations, without influence from dietary factors (Fig. 3). Further understanding of the role of these gene–diet interactions at the molecular level is needed before diets can be tailored according to each ethnic subgroup.

Acknowledgements

Dr Karani S Vimalaswaran acknowledges support from the British Nutrition Foundation through the Drummond Pump Priming Award and thanks the support from the authors of the *GeNuIne* Collaboration (Vimalaswaran 2020).

Conflict of interest

The authors have no conflict of interest.

References

- Adaikalakoteswari A, Jayashri R, Sukumar N *et al.* (2014) Vitamin B12 deficiency is associated with adverse lipid profile in Europeans and Indians with type 2 diabetes. *Cardiovascular Diabetology* **13**: 1–7.
- Allen, LH, 2008. Causes of vitamin B12 and folate deficiency. *Food and Nutrition Bulletin* **29**(2_suppl1), pp. S20–S34.
- Allin KH, Friedrich N, Pietzner M *et al.* (2017) Genetic determinants of serum vitamin B12 and their relation to body mass index. *European Journal of Epidemiology* **32**: 125–34.
- Anjana RM, Pradeepa R, Das AK *et al.* (2014) Physical activity and inactivity patterns in India—results from the ICMR-INDIAB study (Phase-1)[ICMR-INDIAB-5]. *International Journal of Behavioral Nutrition and Physical Activity* **11**: 1–11.
- Astrup A & Bügel S (2010) Micronutrient deficiency in the aetiology of obesity. *International Journal of Obesity* **34**: 947–8.
- Baltaci D, Deler MH, Turker Y *et al.* (2017) Evaluation of serum vitamin B12 level and related nutritional status among apparently healthy obese female individuals. *Nigerian Journal of Clinical Practice* **20**: 99–105.
- Baltaci D, Kutlucan A, Turker Y *et al.* (2013) Association of vitamin B12 with obesity, overweight, insulin resistance and metabolic syndrome, and body fat composition; primary care-based study. *Medicinski Glasnik* **10**: 203–10.
- Bradfield JP, Taal HR, Timpson NJ *et al.* (2012) A genome-wide association meta-analysis identifies new childhood obesity loci. *Nature Genetics* **44**: 526.
- Caesar R, Tremaroli V, Kovatcheva-Datchary P *et al.* (2015) Cross-talk between gut microbiota and dietary lipids aggravates WAT inflammation through TLR signaling. *Cell Metabolism* **22**: 658–68.
- Chakraborty S, Chopra M, Mani K *et al.* (2018) Prevalence of vitamin B12 deficiency in healthy Indian school-going adolescents from rural and urban localities and its relationship with various anthropometric indices: a cross-sectional study. *Journal of Human Nutrition and Dietetics* **31**: 513–22.
- Damms-Machado A, Weser G & Bischoff SC (2012) Micronutrient deficiency in obese subjects undergoing low calorie diet. *Nutrition Journal* **11**: 1–10.
- Deepa M, Pradeepa R, Rema M *et al.* (2003) The Chennai Urban Rural Epidemiology Study (CURES)—study design and methodology (urban component)(CURES-I). *The Journal of the Association of Physicians of India* **51**: 863–70.
- Dror DK & Allen LH (2012) Interventions with vitamins B6, B12 and C in pregnancy. *Paediatric and Perinatal Epidemiology* **26**: 55–74.
- Green R, Allen LH, Bjørke-Monsen AL *et al.* (2017) Vitamin B 12 deficiency. *Nature Reviews Disease Primers* **3**: 1–20.
- Guarnizo-Poma M, Urrunaga-Pastor D, Montero-Suyo C *et al.* (2018) Association between serum vitamin B12 levels and metabolic syndrome in a euthyroid population. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews* **12**: 943–8.
- Hall AB, Tolonen AC & Xavier RJ (2017) Human genetic variation and the gut microbiome in disease. *Nature Reviews Genetics* **18**: 690–9.
- Hazra A, Kraft P, Lazarus R *et al.* (2009) Genome-wide significant predictors of metabolites in the one-carbon metabolism pathway. *Human Molecular Genetics* **18**: 4677–87.
- Husemoen LLN, Skaaby T, Thuesen BH *et al.* (2016) Mendelian randomisation study of the associations of vitamin B12 and folate genetic risk scores with blood pressure and fasting serum lipid levels in three Danish population-based studies. *European Journal of Clinical Nutrition* **70**: 613–9.
- Institute of Medicine (US) Standing Committee on the Scientific Evaluation of Dietary Reference Intakes and its Panel on Folate, Other B Vitamins, and Choline (2000) *Dietary Reference Intakes for Thiamin, Riboflavin, Niacin, Vitamin B6, Folate, Vitamin B12, Pantothenic Acid, Biotin, and Choline*. National Academies Press: Washington.
- Ishihara J, Iso H, Inoue M *et al.* (2008) Intake of folate, vitamin B6 and vitamin B12 and the risk of CHD: the Japan Public Health Center-Based Prospective Study Cohort I. *Journal of the American College of Nutrition* **27**: 127–36.
- Jayashri R, Venkatesan U, Rohan M *et al.* (2018) Prevalence of vitamin B 12 deficiency in South Indians with different grades of glucose tolerance. *Acta Diabetologica* **55**: 1283–93.
- Jayawardena R, Thennakoon S, Byrne N *et al.* (2014) Energy and nutrient intakes among Sri Lankan adults. *International Archives of Medicine* **7**: 1–11.
- Katulanda P, Jayawardena MAR, Sheriff MHR *et al.* (2010) Prevalence of overweight and obesity in Sri Lankan adults. *Obesity Reviews* **11**: 751–6.
- Kim YN, Hwang JH & Cho YO (2016) The effects of exercise training and acute exercise duration on plasma folate and vitamin B12. *Nutrition Research and Practice* **10**: 161.
- Knight BA, Shields BM, Brook A *et al.* (2015) Lower circulating B12 is associated with higher obesity and insulin resistance during pregnancy in a non-diabetic white British population. *PLoS One* **10**: e0135268.
- Koury MJ & Ponka P (2004) New insights into erythropoiesis: the roles of folate, vitamin B12, and iron. *Annual Review of Nutrition* **24**: 105–31.
- Krishnaveni GV, Hill JC, Veena SR *et al.* (2009) Low plasma vitamin B 12 in pregnancy is associated with gestational 'diabetes' and later diabetes. *Diabetologia* **52**: 2350–8.
- Kumar KA, Lalitha A, Pavithra D *et al.* (2013) Maternal dietary folate and/or vitamin B12 restrictions alter body composition (adiposity) and lipid metabolism in Wistar rat offspring. *The Journal of Nutritional Biochemistry* **24**: 25–31.
- Li H, Kilpeläinen TO, Liu C *et al.* (2012) Association of genetic variation in FTO with risk of obesity and type 2 diabetes with data from 96,551 East and South Asians. *Diabetologia* **55**: 981–95.
- MacFarlane AJ, Greene-Finestone LS & Shi Y (2011) Vitamin B-12 and homocysteine status in a folate-replete population: results from the Canadian Health Measures Survey. *The American Journal of Clinical Nutrition* **94**: 1079–87.
- Mahalle N, Kulkarni MV, Garg MK *et al.* (2013) Vitamin B12 deficiency and hyperhomocysteinemia as correlates of cardiovascular risk factors in Indian subjects with coronary artery disease. *Journal of Cardiology* **61**: 289–94.
- Matusheski N, Caffrey A, Christensen L *et al.* (2021) Diets, nutrients, genes and the microbiome: recent advances in personalised

- nutrition. *British Journal of Nutrition* 125: 1–9. <https://doi.org/10.1017/S0007114521000374>
- Mclean E, De Benoist B & Allen LH (2008) Review of the magnitude of folate and vitamin B12 deficiencies worldwide. *Food and Nutrition Bulletin* 29: S38–51.
- Moen GH, Qvigstad E, Birkeland KI *et al.* (2018) Are serum concentrations of vitamin B-12 causally related to cardiometabolic risk factors and disease? A Mendelian randomization study. *The American Journal of Clinical Nutrition* 108: 398–404.
- Mohan V & Deepa R (2006) Adipocytokines and the expanding 'Asian Indian Phenotype'. *The Journal of the Association of Physicians of India* 54: 685–6.
- Narang M, Singh M & Dange S (2016) Serum homocysteine, vitamin B12 and folic acid levels in patients with metabolic syndrome. *The Journal of the Association of Physicians of India* 64: 22–6.
- Ntaios G, Savopoulos C, Grekas D *et al.* (2009) The controversial role of B-vitamins in cardiovascular risk: an update. *Archives of Cardiovascular Diseases* 102: 847–54.
- O'Leary F & Samman S (2010) Vitamin B12 in health and disease. *Nutrients* 2: 299–316.
- Ordovas JM, Ferguson LR, Tai ES *et al.* (2018) Personalised nutrition and health. *British Medical Journal* 361: 1–6.
- Papageorgiou N & Tousoulis D (2016) Oxidized-LDL immunization for the treatment of atherosclerosis: how far are we? *International Journal of Cardiology* 222: 93–4.
- Pinhas-Hamiel O, Doron-Panush N, Reichman B *et al.* (2006) Obese children and adolescents: a risk group for low vitamin B12 concentration. *Archives of Pediatrics & Adolescent Medicine* 160: 933–6.
- Rhee KE, Phelan S & McCaffery J (2012) Early determinants of obesity: genetic, epigenetic, and in utero influences. *International Journal of Pediatrics* 2012: 1–9.
- Sarzynski MA & Bouchard C (2013) The challenging chase for nutrigenetic predictors of metabolic responses to dietary interventions. *Diabetes Care* 36: 3379–81.
- Selhub J (2002) Folate, vitamin B12 and vitamin B6 and one carbon metabolism. *The Journal of Nutrition, Health & Aging* 6: 39–42.
- Shenoy V, Mehendale V, Prabhu K *et al.* (2014) Correlation of serum homocysteine levels with the severity of coronary artery disease. *Indian Journal of Clinical Biochemistry* 29: 339–44.
- Sirugo G, Williams SM & Tishkoff SA (2019) The missing diversity in human genetic studies. *Cell* 177: 26–31.
- Sivaprasad M, Shalini T, Balakrishna N *et al.* (2016) Status of vitamin B12 and folate among the urban adult population in South India. *Annals of Nutrition and Metabolism* 68: 94–102.
- Smith AD, Warren MJ & Refsum H (2018) Vitamin B12. *Advances in Food and Nutrition Research* 83: 215–79.
- Sobczyńska-Malefora A, Gorska R, Pelisser M *et al.* (2014) An audit of holotranscobalamin ("Active" B12) and methylmalonic acid assays for the assessment of vitamin B12 status: application in a mixed patient population. *Clinical biochemistry* 47: 82–6.
- Stabler SP & Allen RH (2004) Vitamin B12 deficiency as a worldwide problem. *Annual Review of Nutrition* 24: 299–326.
- Stanger O, Herrmann W, Pietrzik K *et al.* (2004) Clinical use and rational management of homocysteine, folic acid, and B vitamins in cardiovascular and thrombotic diseases. *Zeitschrift für Kardiologie* 93: 439–53.
- Stark A (2013) The matrilineal system of the Minangkabau and its persistence throughout history: a structural perspective. *Southeast Asia: A Multidisciplinary Journal* 13: 1–13.
- Sun Y, Sun M, Liu B *et al.* (2019) Inverse association between serum vitamin B12 concentration and obesity among adults in the United States. *Frontiers in Endocrinology* 10: 414.
- Surendran S, Adaikalakoteswari A, Saravanan P *et al.* (2018) An update on vitamin B12-related gene polymorphisms and B12 status. *Genes & Nutrition* 13: 1–35.
- Surendran S, Aji AS, Ariyasra U *et al.* (2019a) A nutrigenetic approach for investigating the relationship between vitamin B12 status and metabolic traits in Indonesian women. *Journal of Diabetes & Metabolic Disorders* 18: 389–99.
- Surendran S, Alsulami S, Lankeshwara R *et al.* (2019b) A genetic approach to examine the relationship between vitamin B12 status and metabolic traits in a South Asian population. *International Journal of Diabetes in Developing Countries* 40: 21–31.
- Surendran S, Jayashri R, Drysdale L *et al.* (2019c) Evidence for the association between FTO gene variants and vitamin B12 concentrations in an Asian Indian population. *Genes & Nutrition* 14: 1–9.
- Surendran S, Morais, CC, Abdalla, DSP, Shatwan, IA, Lovegrove, JA, Cominetti, C, Vimalaswaran, VK & Horst, MA. (2019) The influence of one-carbon metabolism gene polymorphisms and gene-environment interactions on homocysteine, Vitamin B12, folate and lipids in a Brazilian adolescent population. *Journal of Diabetology* 10: 110–22.
- Thomas-Valdés S, Tostes MDGV, Anunciação PC *et al.* (2017) Association between vitamin deficiency and metabolic disorders related to obesity. *Critical Reviews in Food Science and Nutrition* 57: 3332–43.
- Vimalaswaran KS (2017) Gene–nutrient interactions on metabolic diseases: findings from the GeNuIne Collaboration. *Nutrition Bulletin* 42: 80–6.
- Vimalaswaran KS (2020) A nutrigenetics approach to study the impact of genetic and lifestyle factors on cardiometabolic traits in various ethnic groups: findings from the GeNuIne Collaboration. *The Proceedings of the Nutrition Society* 79: 194–204.
- Vimalaswaran KS, Bodhini D, Lakshmi Priya N *et al.* (2016) Interaction between FTO gene variants and lifestyle factors on metabolic traits in an Asian Indian population. *Nutrition & Metabolism* 13: 1–10.
- Vimalaswaran KS & Loos RJ (2010) Progress in the genetics of common obesity and type 2 diabetes. *Expert Reviews in Molecular Medicine* 12: e7.
- Voutilainen S, Rissanen TH, Virtanen J *et al.* (2001) Low dietary folate intake is associated with an excess incidence of acute coronary events: The Kuopio Ischemic Heart Disease Risk Factor Study. *Circulation* 103: 2674–80.
- Wall C (1998) *Food and Nutrition Guidelines for Healthy Adolescents*. Ministry of Health: Wellington, New Zealand.
- Wiebe N, Field CJ & Tonelli M (2018) A systematic review of the vitamin B12, folate and homocysteine triad across body mass index. *Obesity Reviews*. 19: 1608–18
- Willer CJ, Schmidt EM, Sengupta S *et al.* (2013) Discovery and refinement of loci associated with lipid levels. *Nature Genetics* 45: 1274.

Yajnik CS & Deshmukh US (2012) Fetal programming: maternal nutrition and role of one-carbon metabolism. *Reviews in Endocrine and Metabolic Disorders* 13: 121–7.

Yajnik C, Deshpande SS, Lubree HG *et al.* (2006) Vitamin B12 deficiency and hyperhomocysteinemia in rural and urban Indians. *The Journal of the Association of Physicians of India* 54: 82.

Yajnik CS, Deshpande SS, Jackson AA *et al.* (2008) Vitamin B 12 and folate concentrations during pregnancy and insulin resistance in the offspring: the Pune Maternal Nutrition Study. *Diabetologia* 51: 29–38.