



# Nicotinic acetylcholine receptors: Diversity and physiological importance for neurodegenerative disorders and development of organophosphate antidotes

ANTONIO ZANDONA  
MAJA KATALINIĆ

Institute for Medical Research and Occupational Health, POB 291, HR-10001 Zagreb, Croatia

**\*Corresponding author:**

Antonio Zandona  
azandona@imi.hr

**Keywords:** cholinergic; subunits; nAChR; nicotine; AChE, Alzheimer's; Parkinson's

**Abbreviations:**

ACh	– acetylcholine
AChE	– acetylcholinesterase
AChR	– acetylcholine receptor
CHRN	– nicotinic acetylcholine receptor gene
COMT	– catechol- <i>O</i> -methyltransferase
DA	– dopamine
GABA	– gamma-aminobutyric acid
IP3/DAG	– inositol trisphosphate/diacylglycerol
L-DOPA	– L-3,4-dihydroxyphenylalanin (levodopa)
mAChR	– muscarinic acetylcholine receptor
MAO-B	– monoamine oxidase B
NA	– noradrenaline
nAChR	– nicotinic acetylcholine receptor
NMDA	– N-methyl-D-aspartate
OP	– organophosphorus compound
VTA	– ventral tegmental area

Received March 20, 2020  
Revised June 30, 2020  
Accepted July 7, 2020

## Abstract

*The communication between the nervous and other systems in the organism is carried out by the transmission of nerve impulses. Diverse neurotransmitters are released into the synaptic cleft and bind to the specific receptors at the neighbouring cell to transmit the signal further. One of such receptors are nicotinic acetylcholine receptors (nAChR), integrated membrane proteins (ligand-gated ion channels) activated by the binding of a neurotransmitter acetylcholine. nAChR's main characteristic is their diversity, as they consist of five of the same or mutually different subunits, which contribute to the specific receptors properties and biological activity. During the assembly of a pentameric protein structure, various combinations of subunits are linked together. After the discovery of nAChR's involvement in various diseases, they became an important therapeutic target, for example in the treatment of neurodegenerative diseases (Alzheimer's and Parkinson's) and in the treatment of organophosphorus compound poisoning. This paper presents an overview of current knowledge on nicotinic receptors and an accompanying discussion on diseases, poisonings, potential drugs and treatments is given.*

## INTRODUCTION

By studying nicotine at the turn of the 20<sup>th</sup> century, the concept of *receptive substance* was set up, from which the idea of a receptor was developed (1). A crucial event in nicotinic receptor studies was the chemical identification of the first neurotransmitter receptor. Successful identification allowed for further metabolic, pharmacological and biochemical research (2). Nicotinic acetylcholine receptors (nAChRs) are proteins integrated into membranes and belong to the superfamily of ligand-gated ion-channels (3, 4). Since they are formed by a specific combination of five subunits, there are different subtypes of nAChRs that mediate different physiological functions (5). Nicotinic receptors are expressed in the peripheral nervous system (PNS), where they mediate the transmission of impulses between nerve and muscle cells; in the central nervous system (CNS) and in other types of cells (keratinocytes, epithelial cells, macrophages, *etc.*; (5)). After understanding that the subtypes of nAChR contributed to the neuropathology of many diseases, there arose an interest in the development of therapeutic nicotine agonists and other drugs for specific subtypes and the use of cloned nAChR subunits as possible therapeutic agents (1). nAChRs have be-

come an important therapeutic targets for the treatment of neurodegenerative diseases, primarily Alzheimer's and Parkinson's (5), but also in the treatment of poisoning with highly toxic organophosphorus compounds (6).

### Historical overview

After studying nerve-muscle preparations, John Newport Langley found that muscle tissue possesses something that mixed with nicotine and *curare* (reversible inhibitor of nAChR) receives stimulus and transfers it to other cells (7). In the 1960s, David Nachmansohn and other scientists identified the receptor for acetylcholine (ACh) in the electric eel (*Electrophorus electricus*) electric organ by radioactive ligands, which was unusually rich in nicotinic synapses (8). In addition, Chen-Yuan Lee with snake venom,  $\alpha$ -bungarotoxin ( $\alpha$ -Bgt), specifically inhibited *in vivo* nerve-muscle transmission without interaction with acetylcholinesterase (AChE, EC 3.1.1.7, an enzyme that hydrolyses the ACh neurotransmitter in synapses). It has been shown that nAChR is a hydrophobic protein of high molecular weight and can be physically separated from AChE, and studies then confirmed that AChE and nAChR are different protein entities (9). In 1972, significant influence on the investigation of nicotinic acetylcholine receptors was the isolation of the new generation of nAChR from homogenates of marbled electric ray (*Torpedo marmorata*) (10). In 1973, the purified protein nAChR was observed under a microscope and rings-like structures (with a hydrophilic core linked to compact clusters) were detected. Such a compact cluster was made up of several subunits (11). A group of scientists in 1988 was able to generate 3D crystals of *Torpedo* nAChR for the first time (12, 13). Soon thereafter, the physiology and a variety of nicotinic receptors expression in the vertebrates was determined, including five different subunits that could be assembled in different ratios (14). Also, in the 1990s, nicotinic receptors and nicotine were associated with various diseases, such as schizophrenia (15), Alzheimer's (16), Parkinson's disease (17) and Tourette's syndrome (18) and research took a turn toward finding drugs and effective therapies for nAChR-related states. In association with various diseases, the effect of organophosphorus compounds (OP) on the cholinergic system was also observed (6, 19). More precisely, the symptoms of OP poisoning are caused by the irreversible inhibition of the enzyme AChE, which leads to the accumulation of

ACh and extensive receptor stimulation (20, 21). In addition to the inhibition of AChE, several organophosphorus compounds directly affect the receptors of the cholinergic system and modulate the level of their receptor expression (22, 23). OPs thus also affect nicotine receptors, block the opening of the ion channel and desensitize the receptors (24–27).

### Acetylcholine receptors and nerve effects

In general, acetylcholine receptors (AChR) are integrated membrane proteins, which are activated by neurotransmitter ACh binding. There are two types of such receptors: ionotropic nicotinic receptors (nAChRs), which exist in two forms (nervous and muscular), and five types of metabotropic muscarinic receptors (mAChR) M1 to M5, which can be classified into two traditional pharmacological groups: M1 and M2 receptors. The M1 group receptors (M1 and M3) interact via the IP3/DAG system (by activating phospholipase C), inactivate  $K^+$  channels, while M2 (M2, M4 and M5) receptors regulate adenyl cyclase (intracellular cAMP concentration) and activate  $K^+$  channels (28, 29). Like other transmembrane receptors, AChRs are classified according to their pharmacology or affinity and sensitivity to different compounds (Figure 1). Although all acetylcholine receptors, by definition, are sensitive to acetylcholine, they also react to other xenobiotics. For example, mAChRs are sensitive to muscarine (30, 31).

nAChRs are ionic channels for  $Na^+$ ,  $K^+$  and  $Ca^{2+}$  and are particularly sensitive to nicotine. They are found in the central and peripheral nervous system, muscle and many other tissues (32). In the nerve-muscle synapse, they are the primary receptors for communication with the muscles and controlling muscle contraction. In this way, the peripheral nervous system transmits outgoing signals from presynaptic to postsynaptic cells within the sympathetic and parasympathetic nervous system. In the immune system, nAChRs regulate inflammatory processes and participate in signalling different intracellular pathways (1, 33, 34).

Full activation of a postsynaptic nAChR occurs when two molecules of the neurotransmitter ACh are bound to it. The cationic channel of the receptor is opened because of the electrochemical gradients through which  $Na^+$  and




Fig. 1. Structures: (A) acetylcholine, (B) nicotine and (C) muscarine

K<sup>+</sup> flow. Every second a nerve vesicle releases neurotransmitters into the synapse. In addition to the acetylcholine receptors, there are many AChEs in the synapses, whose task is to hydrolyse the ACh, thus terminating the signalling (35). The long-term binding of ACh to the receptor results in desensitization and thus inactivation of the ion channel. For example, irreversible AChE inhibitors lead to the accumulation of ACh in a synapse (36, 37), and one molecule of ACh can stimulate a few additional openings of ion channels. This may also have its advantages; like in diseases where the number of nAChR is reduced and causes muscular weakness, AChE-inhibiting drugs enable signal reactivation with the same neurotransmitter molecule, thereby improving signalling and replacing the lack of receptors (38, 39).

### Types, localization and receptor structure


It has already been mentioned that nAChRs are ligand-gated ion channels forming pores on a cell plasma membrane mediating fast signal transmission through synapses. They are involved in a wide range of physiological processes and are divided into two groups: nerve and muscle (40). Muscle nAChRs are localized in the nerve-muscle synapses, where the electrical nerve impulse signals contraction to the muscle cells and is responsible for muscle activity. Many different types of nerve nAChR are localized in synapses on postsynaptic nerve cells, as in the CNS, where they are involved in cognitive functions, learning, memory, *etc.* nAChR activation causes cation movement through the ion channel pore with the calcium ions which affect the release of neurotransmitters (1, 41).

**Table 1.** nAChR subunits subfamilies based on similarities in protein sequence (42)

Subfamily	Subtype	Type	Subunits
I		Neuronal	$\alpha 9, \alpha 10$
II		Neuronal	$\alpha 7, \alpha 8$
III	1	Neuronal	$\alpha 2, \alpha 3, \alpha 4, \alpha 6$
	2	Neuronal	$\beta 2, \beta 4$
	3	Neuronal	$\beta 3, \alpha 5$
IV		Muscle	$\alpha 1, \beta 1, \delta, \gamma, \epsilon$

Studies revealed the existence of 5 different types of nAChR subunits with small molecular weight differences. Thus, nAChR consists of five subunit types: alpha ( $\alpha 1$ - $\alpha 10$ ), beta ( $\beta 2$ - $\beta 5$ ), gamma ( $\gamma$ ) delta ( $\delta$ ) and epsilon ( $\epsilon$ ); found in various combinations. The nAChR subunits have been grouped into 4 subfamilies (I-IV) based on similarities in protein sequence, and subfamily III has been further divided into 3 subtypes (Table 1) (42). The most widely expressed nAChR subtypes are  $\alpha 4, \alpha 7$  and  $\beta 2$  (43). Nerve nAChRs are assembled just from alpha and beta units, while muscle nAChR from all five subunit types (41, 44).

Nicotinic receptors are always pentamers (Figure 2), with subunits disposed symmetrically in a circle around the central receptor channel pore. The receptors always contain two or more  $\alpha$ -subunits, which are crucial for ACh binding (45, 46). The ACh binding site is formed



**Figure 2.** The basic structure of nicotinic acetylcholine receptors. (A) Torpedo marmorata nAChR structure (model from PDB code 2BG9; 49) with extracellular (1), transmembrane (2) and intracellular (3) domains, (B) the receptor is formed from 5 subunits around the central pore with Cys-loop (4) in  $\alpha$  subunit.

when dimers of  $\alpha$ -subunits and adjacent subunits associate (47, 48).

Nerve nAChRs are divided into two main groups based on their sensitivity to the snake venom  $\alpha$ -bungarotoxin.  $\alpha$ -Bungarotoxin-sensitive receptors are homomers ( $\alpha7$  and  $\alpha9$ ), while  $\alpha$ -bungarotoxin-non-susceptible receptors are heteromeric (other subunits types). Most nAChRs are asymmetric heteromers in which neither ACh binding sites nor subunit-subunit interactions are identical (48, 49).

After years of experiments, the nicotinic receptor localization and structure, which contains the extracellular synaptic domain, four transmembrane  $\alpha$ -helix and the intracellular cytoplasmic domain, have been proposed (Figure 2). Further analysis revealed the existence of so-called Cys-loops in nAChR structure (2, 47). nAChR is also a glycosylated protein and has N-linked glycosylation sites which vary depending on the subunits. A cys-cys pair is required for agonist binding and its presence denotes the existence of an  $\alpha$ -subunit (41, 47). A Cys-loop is characteristic for this type of receptor and in mammals these cysteines are most often separated with 13 conserved amino acids. Other receptors that belong to the same receptor superfamily as nAChR include: 5-hydroxytryptamine (5-HT<sub>3</sub>),  $\gamma$ -aminobutyric acid type A (GABA<sub>A</sub>) and GABA<sub>C</sub>, glycine receptors, glutamate and histidine receptors (50).

### The variety of nicotine subunits

The diversity of nAChR subunits is the main feature of the specific properties and functions of mature receptors. Receptor pentamers can be assembled from various

combinations of  $\alpha$ ,  $\beta$  and other structural subunits (38, 41). Nowadays, 16 genes encoding for nAChR subunits are known, named *CHRNA1–10* for  $\alpha$  subunits, *CHRNB1–4* for  $\beta$  subunits, *CHRNA3*, *CHRNA4*, *CHRNA5*, *CHRNA6*, *CHRNA7*, *CHRNA8*, *CHRNA9*, *CHRNA10*, *CHRNA11*, *CHRNA12*, *CHRNA13*, *CHRNA14*, *CHRNA15*, *CHRNA16* for  $\gamma$ ,  $\delta$  and  $\epsilon$ , respectively (1, 52, 53). Different nAChR gene details are presented in Table 1. Mutations in *CHRN* genes (causing loss or gain of nAChR function, increased or decreased sensitivity to ACh or desensibilisation of the nAChR) can cause phenotypes with different manifestations such as congenital myasthenic syndrome, nocturnal frontal lobe epilepsy and multiple pterygiums (54–56). The genes *CHRNA3/CHRNA5/CHRNB4* or *CHRNB3/CHRNA6* are in gene clusters on chromosomes 15q25 or 8p11, respectively; while other *CHRN* genes are separately located on chromosomes 1, 2, 4, 8, 11, 15, 17 or 20 as shown in Table 2.

In some brain areas, diverse specific subtype subunits participate in order to form an nAChR of high affinity for substrates. In the *basal ganglia*, including the *ventral tegmental area* (VTA) and *substantia nigra*, the  $\alpha6$  and  $\beta3$  nAChR subunits were included in  $\alpha4\beta2$  nAChR complexes to create a high affinity receptor for ACh (Table 3). Currently, these are the only areas in the brain where  $\alpha6$  and  $\beta3$  subunits are co-expressed alongside the  $\alpha4$  subunit, which is very important for the appearance and therapy of Parkinson's disease (57). The autonomic nervous system is characterized by an abundant expression of  $\alpha3$  and  $\beta4$  nAChR, while expression of  $\alpha4$  and  $\beta2$  subunits dominates the central nervous system. In example, coordinated expression of essential subunits is strongly regulated in the brain during the development of the organism and during various injuries (58).

**Table 2.** nAChR genes: subunit receptor name, gene names, chromosomal location, gene length (kb), gene cluster (if applicable), and number of aminoacids (AA) in expressed protein (1, 52)

Subunit receptor	Gene name	Chromosome	Gene (kb)	Gene cluster	Protein (AA)
$\alpha1$	<i>CHRNA1</i>	2q31.1	16.64		482
$\alpha2$	<i>CHRNA2</i>	8p21.2	18.51		529
$\alpha3$	<i>CHRNA3</i>	15q25.1	28.24	<i>CHRA3-A5-B4</i>	622
$\alpha4$	<i>CHRNA4</i>	20q13.33	14.75		627
$\alpha5$	<i>CHRNA5</i>	15q25.1	29.71	<i>CHRA3-A5-B4</i>	515
$\alpha6$	<i>CHRNA6</i>	8p11.21	15.93	<i>CHRNB3-A6</i>	494
$\alpha7$	<i>CHRNA7</i>	15q13.2	142.25		534
$\alpha9$	<i>CHRNA9</i>	4p14	19.63		479
$\alpha10$	<i>CHRNA10</i>	11p15.4	5.8		450
$\beta1$	<i>CHRNB1</i>	17p13.1	12.65		501
$\beta2$	<i>CHRNB2</i>	1q21.3	12.25		502
$\beta3$	<i>CHRNB3</i>	8p11.21	39.99	<i>CHRNB3-A6</i>	458
$\beta4$	<i>CHRNB4</i>	15q25.1	17.48	<i>CHRA3-A5-B4</i>	498
$\delta$	<i>CHRNA11</i>	2q37.1	10.48		517
$\gamma$	<i>CHRNA12</i>	2q37.1	6.6		517
$\epsilon$	<i>CHRNA13</i>	17p13.2	5.3		496

**Table 3.** The presence of different  $\alpha$  and  $\beta$  subunits in certain parts of the brain (\*specific subtype; - not determined) (62).

Localization	Type nAChR	Involved in specific neurotransmitter release
Amygdala	$\alpha 4\beta 2, \alpha 7$	–
Cerebellum	$\alpha 4\beta 2, \alpha 7, \alpha 3\beta 3^*, \alpha 3\beta 4^*$	Glu
Cortex	$\alpha 4\beta 2, \alpha 4\alpha 5\beta 2, \alpha 7$	Glu, ACh, DA
Hedial habenula	$\alpha 4\beta 2, \alpha 3\beta 2^*, \alpha 7, \alpha 3\beta 3\beta 4, \alpha 3\beta 4^*$	–
Hippocampus	$\alpha 4\beta 2, \alpha 4\alpha 5\beta 2, \alpha 3\beta 4, \alpha 7$	Glu, ACh, DA
Hypothalamus	$\alpha 4\beta 2, \alpha 7$	NA
Interpeduncular nucleus	$\alpha 4\beta 2^*, \alpha 7, \alpha 3\beta 3\beta 4, \alpha 3\beta 4, \alpha 2\beta 2^*$	ACh, GABA, NA
Locus coeruleus	$\alpha 3\beta 4, \alpha 6\beta 2\beta 3^*$	–
Olfactor bulb	$\alpha 4\beta 2, \alpha 7$	Glu, DA
Pineal gland	$\alpha 3\beta 4, \alpha 7$	–
Rapche nuclei	$\alpha 4\beta 2$	–
Spinal cord	$\alpha 4\beta 2, \alpha 7, \alpha 3\beta 2^*$	–
Striatum	$\alpha 4\beta 2, \alpha 4\alpha 5\beta 2, \alpha 6\beta 2\beta 3, \alpha 6\alpha 4\beta 2\beta 3$	Glu, DA
Substantia nigra, VTA	$\alpha 4\beta 2, \alpha 4\alpha 5\beta 2, \alpha 7, \alpha 3\beta 4^*, \alpha 6\beta 2\beta 3$	–
Thalamus	$\alpha 4\beta 2, \alpha 4\alpha 5\beta 2$	DA, GABA

$\alpha 7$  are the most researched subunits of all known nicotinic subunits types. Receptors assembled of  $\alpha 7$  subunits have a high  $\text{Ca}^{2+}:\text{Na}^+$  bandwidth ratio (59). As a result, the opening of the  $\alpha 7$  nAChR channel may cause several  $\text{Ca}^{2+}$ -dependent mechanisms, including the activation of secondary signal pathways (60). Co-expression and assembly, i.e. binding of the  $\alpha 7$  nAChR subunit to other subunits, affects receptor ionic permeability, so nAChR consisting of  $\alpha 7$  and  $\beta 2$  subunits has pharmacological properties different from those homozygous  $\alpha 7$  nAChR (61).

Pentameric structures, unlike those with pair subunits structures (e.g., tetramers), require multiple mechanisms for overcoming problems related to receptor assembly. Coordination of regulatory mechanisms between transcription and receptors assembly, which respond to external changes, are a common biological feature of nicotinic receptors. The first level of regulation of the region-specific expression of nAChR is transcriptional control of subunit expression.

Posttranslational modifications are crucial for the regulation of mature receptor functions, their expression and localization. There are several checkpoints in the cell to ensure proper assembly. Experiments with the chimeric recombinant subunits showed that the 23-amino acid region (from G23 to D46) contains the residues needed for proper  $\alpha 7$  subunit assembly to the homopentamer receptors and a Cys-loop is needed for proper domain folding (63). Another significant control point for nAChR is the endoplasmic reticulum where degradation of proteins is an important part of regulating receptor concentration. In total, 80% of the synthesized subunits are improperly assembled and never leave the endoplasmic

reticulum, where they are degraded (64). Reduction of the degradation of precursor subunits in the endoplasmic reticulum results in increased concentration of nAChR on the cell membrane (65). Also, continuous nicotine exposure increases the concentration of nAChR on the surface of the cell by reduction of the degradation of the intracellular pool of receptors (66). Another important posttranslational regulation is the modification achieved by N-glycosylation, where more sites in the  $\text{NH}_2$ -terminal domain of the nAChR subunit are glycosylated. It has been found that muscular nAChR glycosylation is not required for receptor assembly or the formation and functioning of Cys-loops (67). Still, proper glycosylation is necessary for their subsequent incorporation into the plasma membrane (68, 69). In addition, palmitoylation and phosphorylation are also mechanisms that regulate the expression, receptor function and binding efficacy of different nAChR subunits. Palmitoylation is a reversible posttranslational modification where the palmitate covalently binds to Cys residues in the endoplasmic reticulum (70, 71). As opposed to that, in phosphorylation, phosphate binds to specific residues within the cytoplasmic domain (72). Besides, as mentioned nicotinic receptors also possess the ability of increasing upregulation when exposed to high nicotine concentrations; still, some subtypes have a lower level of expression (73).

### Insight into the function of nAChR

Activation of mammalian nerve nAChR promotes the opening of an unselected cationic channel leading to  $\text{Na}^+$  flow, depolarization of the membrane and activation of  $\text{Ca}^{2+}$  channels (49).  $\text{Ca}^{2+}$  flow through nAChR or voltage-gated channels is essential for nicotine modulation, synaptic plasticity, nerve viability, differentiation and migra-

tion. In the nervous system expressing specific nAChR subtypes, nicotine mediates activation of several  $\text{Ca}^{2+}$  independent kinases including PI3K, protein kinase C (PKC), protein kinase A (PKA) and extracellular signal-regulatory kinase (ERK) (74, 75). Also these kinases activate a large number of transcription factors by downstream regulation, such as cAMP-binding protein (CREB) that further activates transcription factor 2 (ATF-2) (76) and the ETS-like transcription factor (Elk-1), and transcript signal transducer (STAT3) (77).

nAChR are present at essential regulatory sites and lead to multiple outcomes in sensitivity to exogenous stimuli or participation in neurodegeneration and inflammation. The presence of expressed nAChR in adipose tissue has provided additional research into the effects of nicotine on the body. It has shown that average smokers are skinnier and more likely to develop metabolic syndrome such as type II diabetes.

Also, patients with ulcerative colitis who stopped consuming tobacco developed more serious progression of the disease, *i.e.* nicotine had a protective effect. By contrast, patients with Crohn's disease improve disease severity if they smoke (78).

However, small concentrations of nicotine protect cells from damage induced by  $\beta$ -amyloids, inflammation, alcohol, hypoxia, but mechanism of action is unknown (40, 79, 80). High nicotine concentrations increase the number of nAChRs, due to desensitization instead of enhanced receptor function. Nicotine effects on nAChR are complex at molecular level and it is difficult to separate their consequences at system level (81).


### Neurodegenerative disorders

The involvement of nicotinic receptors in many diseases, autoimmune responses and epilepsy, stimulated the development of nAChR subtypes of specific ligands as possible drugs (80). Reduced function and/or nAChR

expression in the brain is associated with the pathophysiology of for example Alzheimer's disease (AD) or Parkinson's disease (PD), severe, incurable, degenerative brain disorders (40). Since the expression of the receptors cannot be induced easily, drug design for the treatment of neurodegenerative disorders is based on improving existing receptors' stimulation. One of the approaches is the use of nicotinic agonists or AChE inhibitors that are investigated to prevent present neurotransmitter acetylcholine from being degraded rapidly, which in turns enables activation of nAChR and the signal transmission to the next postsynaptic cell (82).

AD progression (Figure 3) leads to dementia, memory loss, thinking impairment, and changes in behaviour and personality. AD and other forms of dementia affect about 47 million people worldwide, in Croatia about 86,000 (83). The histopathology of the disease is well-known and is accompanied by a loss of cholinergic transmission, deposition of extracellular amyloid  $\beta$ -peptides ( $A\beta$ ) in plaques, and hyperphosphorylation of  $\tau$ -protein leading to excessive formation of neurofibrillary tangles (84, 85). Also, receptor loss in Alzheimer's disease is much higher than in a normal aging brain (86, 87). Specifically, the degree of loss of  $\alpha 4\beta 2$  nAChR and  $\alpha 7$ -expressed expression are well correlated with the progressive range of cognitive decline in patients with mild to moderate AD (88). Two effects are important for cognitive improvement of AD, improvement of synaptic transfer and reduction of neurodegeneration (89). Interestingly,  $\alpha 4$  nAChR expression decreases by 80% in AD (90).

The simplest therapy to nAChR loss is the long-term use of nicotine (91) and in human clinical trials; nicotine alleviated AD symptoms (92). Additionally, nicotine has shown beneficial effects on learning, memory, attention, and cognitive functions in patients with AD. Now, the focus of the development of AD therapy is on specific nAChR agonists (Table 4), which should have low affinity for muscle nAChR to reduce unwanted muscle stimu-



**Figure 3.** A sketch of the difference in the brain structure of the healthy brain and in patients with Alzheimer's disease (mild or severe) accompanied by severe cortical shrinkage (according to (99) and (100)).

lation. Two types of receptors,  $\alpha 7$  and  $\alpha 4\beta 2$ , have become a target for AD treatment, since these receptors are expressed in the brain regions for memory and information processing. A crucial step in AD treatment could be the introduction of positive allosteric modulators (PAMs) to ensure multiple modulation of receptors to suppress disease (93), as well as a combination of drugs based on synergistic mechanisms. For example, a combination of galantamine that would inhibit AChE and modulate nAChRs, curcumine that would inhibit  $\beta$ -secretase and  $\tau$ -phosphorylation and rivastigmine to reversibly inhibit AChE; or a combination of tacrine, curcumine and huperzine A for the same reason (94). Five medications are currently used to treat the cognitive problems of AD: four are acetylcholinesterase inhibitors (tacrine (withdrawn due to toxicity (95)), rivastigmine, galantamine and donepezil) and the other (memantine) is an NMDA receptor antagonist. The way to increase the function of nAChR in the brain without nicotine is to sensitize receptors using allosteric potentiating ligands (APLs) that include drugs such as physostigmine and galantamine, one of the approved drugs for the treatment of AD (89). Second, non-competitive agonists (NCAs) have agonistic activity and bind to a site close but still further than the ACh binding site on  $\alpha$ -subunits of nAChR (96). Although NCA encourages the opening of nAChR channels in many nerve and non-nerve cells, the likelihood of opening channels with such compounds is very low. Namely, galantamine acts as a nicotine NCA but not as a nicotine APL (97). Subunit  $\alpha 7$  is the only nAChR subunit in mammals with positively charged residues within the  $\alpha 118$ -140 segment that is assumed to be an APL binding site (98).

Along with AD, Parkinson's disease (PD) is one of the most common neurodegenerative diseases in the world. It appears in all ethnic groups, in both sexes, although men are affected twice as much than women. The occurrence of illness increases with age and affects 1% of the population over 60. Symptoms usually begin to appear for the first time between the age of 50 and 65, but the disease may occur early in life, childhood or adolescence, albeit rarely. There are around 7-10 million people in the world suffering from Parkinson's disease, while in Croatia the number of registered people with the disease is 20,000 (122). Disease is caused by the degeneration of dopaminergic nerves of *substantia nigra* with symptoms of muscle stiffness, bradykinesia and blocking movement, loss of reflexes and tremors. Movement disorders are caused by a lack of dopamine-producing nerve cells, since dopamine is important in movement control. Current treatment for PD includes therapy with dopamine precursor L-DOPA (Figure 4), but neither does it stop the progression of the disease nor does its effectiveness last for a long time (123). nAChR-specific nicotine-based agonists are another direction in the development of therapy. One of the synthesized agonists, SIB-1765F, showed high affinity for  $\alpha 4\beta 2$ , and was as equally effective as nicotine in stimulating


**Table 4.** Various known nicotinic agonists, antagonists and allosteric modulators

Name	Action	Specific target	Refs.
18-Methoxycoronaridine (18-MC)	antagonist	$\alpha 3\beta 4$	(101)
A-582941	agonist	$\alpha 7$	(102)
ABBF	agonist	$\alpha 7$	(103)
ABT-089 <sup>a</sup>	agonist	$\alpha 4\beta 2$	(104)
ABT126 <sup>a</sup>	agonist	$\alpha 7$	(104)
ABT-418 <sup>a</sup>	agonist	$\alpha 4\beta 2$ , $\alpha 3\beta 4$	(104)
AR-R 17779	agonist	$\alpha 7$	(105)
Atropine <sup>b</sup>	antagonist	–	(106)
AVL-3288 <sup>a</sup>	allosteric modulator	$\alpha 7$	(107)
Benthiactzine <sup>b</sup>	antagonist	$\alpha 7$ , $\alpha 4\beta 4$ , $\alpha 4\beta 2$	(108)
CCMI	allosteric modulator	$\alpha 7$	(104)
Dihydro- $\beta$ -erythroidine (Dh $\beta$ E)	antagonist	$\alpha 7$ , $\alpha 3\beta 4$ , $\alpha 4\beta 2$	(109)
Dizocilpine (MK-801)	antagonist	$\alpha 7$	(110)
DMPP	agonist	–	(111)
<i>d</i> -Tubocurarine (dTC)	antagonist	$\alpha 4\beta 2$ , $\alpha 7$	(112)
Encenicline <sup>a</sup> (EVP-6124)	agonist	$\alpha 7$	(113)
GTS-21	agonist	$\alpha 7$	(104)
Ibogaine	antagonist	$\alpha 3\beta 4$	(114)
Ispronidine	agonist	$\alpha 4\beta 2$	(104)
MB266 <sup>b</sup>	antagonist	–	(106)
Memantine	antagonist	$\alpha 7$	(115)
Metylylcaconitine (MLA)	antagonist	$\alpha 7$	(116)
NS-1738	allosteric modulator	$\alpha 7$	(111)
PNU-120596	allosteric modulator	$\alpha 7$	(111)
PNU-282987	agonist	$\alpha 7$	(117)
QNB <sup>b</sup>	antagonist	–	(106)
Sazetidine	agonist	$\alpha 4\beta 2$	(110)
SEN123333	agonist	$\alpha 7$	(118)
SSR-180711	agonist	$\alpha 7$	(119)
Tropisetron	agonist	$\alpha 7$	(120)
Vinblastine	antagonist	$\alpha 3\beta 4$	(114)
$\alpha$ -bungarotoxin	antagonist	$\alpha 7$	(121)

<sup>a</sup>in clinical trial phases (104, 107, 113), <sup>b</sup>for organophosphorus poisoning treatment (106, 108)

dopamine release. This agonist in combination with varenicline can attenuate brain overstimulation (124).

Other drugs that cause symptomatic relief and act by raising dopamine levels (L-DOPA, MAO-BI, COMT inhibitors), stimulating dopamine receptors (DA agonists), inhibiting cholinergic effects (anticholinergics) or inhibit-



**Figure 4.** L-DOPA precursor structure and decarboxylation in dopamine in the organism (123).

ing glutaminergic NMDA receptors (amantadines). Thus, in such a system, L-DOPA replaces dopamine; COMT inhibitors maintain L-DOPA, which will metabolize and synthesize dopamine within the nerves, which can then be released into the synapse, MAO-B inhibitors serve to preserve existing dopamine and DA agonists mimic dopamine and bind to dopamine binding sites, as it is necessary to increase the dopamine concentration in PD patients. Interestingly, the  $\alpha 6$  subtype was discovered in the *substantia nigra* and has become attractive as a pharmacological target for Parkinson's disease therapy (125). Epidemiological studies have shown that heavy smokers are less likely to have PD but also a whole spectrum of physiological changes are present in the function and expression of nAChR and desensitization (126). However, improvement of dopamine release was observed on rodents exposed to nicotine and degeneration of DA nerves was prevented (57, 127). The assumption is that nicotine mediates nerve protection because it prevents cell death, reduces symptoms, binds to nAChR and promotes expression of dopamine receptors in *substantia nigra* (128). The exact mechanism is not known yet, although nicotine effects are well explored.

Along with physiological changes, some studies suggest that DA and cholinergic systems operate in a dynamic balance, whose disturbance causes a variety of neurological disorders such as Parkinson's disease. DA-deficiency triggers a reduction in the efficacy of AChR, which result in increase in ACh release (129).

On the other side, an increased kinurenic acid (KYNA), the metabolite of tryptophan that is primarily produced in the brain and releases by astrocytes, follows neurodegenerative disorders and decreased activity/expression of nAChR. Acting as an endogenous activity regulator of  $\alpha 7$  nAChR, KYNA can modulate synaptic transmission, synaptic plasticity, nerve endurance, and nervous connectivity in various brain regions (130).

### Organophosphorus compounds poisonings and nAChRs


Organophosphorus compounds (OP) are a group of highly toxic xenobiotics used as pesticides, lubricants and oils for engines in the industry and as warfare nerve agents. The mechanism of their toxicity is the irreversible

inhibition of the AChE, which participates in the regulation of nerve impulse transfer by hydrolysing ACh (Figure 5). By chemical structure, OP compounds are esters, anhydrides or halides obtained by complete substitution of phosphorous, phosphonic and phosphinic acids. The base consists of a five-valent phosphorus atom coordinated by covalently bound oxygen or sulfur, and two different substituents (alkyl, aryl, alkoxy, alkylthio, aryloxy, mono- or dialkylamino) and the leaving group (fluorine, cyanide, etc.) (131).

Moshnin and Clermont synthesized the first OP compound, tetraethylpyrophosphate (TEPP). After some years, the potential of TEPP was recognized and it began being used as a pesticide. However, OP compounds showed to not be selective to species. During the Second World War, the toxic effects of OP compounds encouraged research and their use as chemical nerve agents. Soon, two series of nerve agents were developed: the G-series (tabun, sarin, soman and cyclosarin), and V-series (like VX, from *Venom, Victory or Viscous*) (37, 132). In the 1990s, the production, stockpiling and use of chemical weapons was prohibited by Organisation for the Prohibition of Chemical Weapons (OPCW), and existing stocks were ordered to be destroyed. Since then, the use of nerve agents for research purposes has been strictly regulated. Still, the use of nerve agents has been recorded several times in recent years by terroristic groups attacking civilians (133).

As mentioned before in the text, when OP inhibits AChE in the nervous system, the neurotransmitter acetylcholine (ACh) accumulates in the synaptic cleft and over-stimulates cholinergic receptors, affecting the appearance of some specific symptoms. The effects of excessive activation of receptors are muscular weakness, cramps and tremor, hypertension and tachycardia, with additional effects such as headaches, memory disorders, wakefulness, confusion, loss of reflexes, anxiety, insomnia and respiratory depression and paralysis (134, 135). The main cause of the death of poisoned patients is the failure of the respiratory system. It is caused by paralysis of respiratory muscle, bronchoconstriction in combination with increased secretion and depression of the respiratory centre (136). Moreover, death can occur within minutes of exposure to nerve agents. Since OPs pass through the blood-brain barrier, they also cause a number of other undesir-





**Figure 5.** Synthesis of ACh in the synapse and association with receptors (according to (136))

able effects due to the action on receptors involved in the transmission of the impulse. Therapy for acute exposure to organophosphorus compounds is a combination of several different antidotes. Atropin is used as an antimuscarinic, ACh antagonist, which reduces the effects of ACh on muscarinic receptors. Compounds known as oximes are used as reactivators of inhibited AChE, which after reactivation reduces the concentration of accumulated acetylcholine in synaptic cleft. The third part of the therapy is administration of benzodiazepine to reduce convulsions and thus prevent brain damage (137). However, none of these drugs is effective enough and the effects of accumulated ACh overstimulating nAChRs persist.

There is ongoing research for selective nAChR inhibitors that could serve in therapy of OP poisonings. Scientists have tried different already available nicotinic antagonist drugs and ligands like  $\delta$ -tubocurarine (138). Furthermore, ketamine and similar drugs also have specific anticholinergic inhibitory activity on  $\alpha 7$ -nicotinic receptors and potentiated testing of a new group of compounds as drug candidates for clinical trials (139). In addition, procyclidine (an anticholinergic drug principally used for the treatment of drug-induced parkinsonism) has been shown to be an antimuscarin, antinicotinic and anti-NMDA receptor drug (140). Although many drugs are available to test from, the research in this field of nicotine antagonists application is still ongoing and none of the candidates has been introduced into medical practice of treating OP poisoning yet.

### Perspectives for future

nAChR isolation paved the way for the design of new pharmacologically active compounds. As they co-operate with other receptors during different physiological processes, it is important to consider them in a wider picture as well. More specifically, the relationship, linkage and communication between different nAChR subtypes and other receptors should be investigated in the future research. In addition, we need to consider that understanding the role that nAChR plays in regulating immune system under normal physiological conditions. Furthermore, the development of ligands that selectively enhance the activity of a particular nAChR subtype is crucial for the future design of drugs for the treatment of, for example, neurodegenerative diseases and OP poisonings. At the same time, the application of such drugs should be as simple, effective and fast as possible to improve existing therapies. In this search, novel *in silico*, *in vitro* and *in vivo* models available for studies of potential agonists and antagonist of nicotinic receptors, could provide an essential platform.

**Acknowledgments:** this review was supported by the Croatian Science Foundation project UIP-2017-05-7260 to M.K. We are grateful to Makso Herman for language editing. Graphics from Servier Medical ART: SMART (<https://smart.servier.com/>) were used for the figures.

### REFERENCES

- ALBUQUERQUE EX, PEREIRA EFR, ALKONDON M, ROGERS SW 2009 Mammalian Nicotinic Acetylcholine Receptors: From Structure to Function. *Physiol Rev* 89 (1): 73–120. <https://doi.org/10.1152/physrev.00015.2008>
- CHANGEUX JP 2012 The Nicotinic Acetylcholine Receptor: The Founding Father of the Pentameric Ligand-gated Ion Channel Superfamily. *J Biol Chem* 287: 40207–40215. <https://doi.org/10.1074/jbc.R112.407668>
- STROUD RM, MCCARTHY MP, SHUSTER M 1990 Nicotinic acetylcholine receptor superfamily of ligand-gated ion channels. *Biochemistry* 29 (50): 11009–23. <https://doi.org/10.1021/bi00502a001>
- GRUTTER T, DE CARVALHO LP, DUFRESNE V, TALY A, EDELSTEIN SJ, CHANGEUX JP 2007 Molecular tuning of fast gating in pentameric ligand-gated ion channels. *Proc Natl Acad Sci USA* 102 (50): 18207–18212. <https://doi.org/10.1073/pnas.0509024102>
- KALAMIDA D, POULAS K, AVRAMOPOULOU V, FOSTIERI E, LAGOUMINTZIS G, LAZARIDIS K, SIDERI A, ZOURIDAKIS M, TZARTOS SJ 2007 Muscle and neuronal nicotinic acetylcholine receptors. Structure, function and pathogenicity. *FEBS Journal* 274: 3799–3845. <https://doi.org/10.1111/j.1742-4658.2007.05935.x>
- KING AM, AARON CK 2014 Organophosphate and carbamate poisoning. *Emerg Med Clin North Am* 33 (1): 133–51. <https://doi.org/10.1016/j.emc.2014.09.010>
- LANGLEY JN 1905 On the reaction of cells and of nerve-endings to certain poisons, chiefly as regards the reaction of striated muscle to nicotine and to curare. *J Physiol* 33: 374–413. <https://doi.org/10.1113/jphysiol.1905.sp001128>

8. NACHMANSOHN D 1961 Chemical factors controlling nerve activity. *Science* 134 (3494): 1962–8. <https://doi.org/10.1126/science.134.3494.1962>
9. CHANG CC, LEE CY 1963 Isolation of neurotoxins from the venom of *Bungarus multicinctus* and their modes of neuromuscular blocking action. *Arch Int Pharmacodyn Ther* 144: 241–257.
10. COHEN JB, WEBER M, HUCHET M, CHANGEUX JP 1972 Purification from *Torpedo marmorata* electric tissue of membrane fragments particularly rich in cholinergic receptor protein. *FEBS Lett* 26 (1): 43–7. [https://doi.org/10.1016/0014-5793\(72\)80538-6](https://doi.org/10.1016/0014-5793(72)80538-6)
11. CARTAUD J, BENEDETTI EL, COHEN JB, MEUNIER JC, CHANGEUX JP 1973 Presence of a lattice structure in membrane fragments rich in nicotinic receptor protein from the electric organ of *Torpedo marmorata*. *FEBS Lett* 33: 109–13. [https://doi.org/10.1016/0014-5793\(73\)80171-1](https://doi.org/10.1016/0014-5793(73)80171-1)
12. HERTLING-JAWEED S, BANDININ G, MULLER-FAHROW A, DOMMES V, HUCHO F 1988 Rapid preparation of the nicotinic acetylcholine receptor for crystallization in detergent solution. *FEBS Lett* 241: 29–32. [https://doi.org/10.1016/0014-5793\(88\)81024-x](https://doi.org/10.1016/0014-5793(88)81024-x)
13. PAAS Y, CARTAUD J, RECOUVREUR M, GRAILHE R, DUFRESNE V, PEBAY-PEYROULA E, LANDAU EM, CHANGEUX JP 2003 Electron microscopic evidence for nucleation and growth of 3D acetylcholine receptor microcrystals in structured lipid-detergent matrices. *Proc Natl Acad Sci USA* 100: 11309–11314. <https://doi.org/10.1073/pnas.1834451100>
14. MCGEHEE DS, HEATH MJ, GELBER S, DEVAY P, ROLE LW 1995 Nicotine enhancement of fast excitatory synaptic transmission in CNS by presynaptic receptors. *Science* 269: 1692–1696. <https://doi.org/10.1126/science.7569895>
15. FREEDMAN R, COON H, MYLES-WORSLEY M, ORR-URTREGER A, OLINCEY A 1997 Linkage of a neurophysiological deficit in schizophrenia to a chromosome 15 locus. *Proc Natl Acad Sci USA* 94: 587–592. <https://doi.org/10.1073/pnas.94.2.587>
16. WHITEHOUSE P, MATINO A, MARCUS K, ZWEIG R, SINGER H, PRICE D, KELLAR K 1988 Reduction in acetylcholine and nicotine binding in several degenerative diseases. *Arch Neurol* 45: 722–724. <https://doi.org/10.1001/archneur.1988.00520310028012>
17. PERRY E, MORRIS C, COURT J, CHENG A, FAIRBAIRN A, MCKEITH I, IRVING D, BROWN A, PERRY R 1995 Alteration in nicotine binding sites in Parkinson's disease, Lewy Body Dementia and Alzheimer's disease, possible index of early neuropathology. *Neuroscience* 64: 385–395. [https://doi.org/10.1016/0306-4522\(94\)00410-7](https://doi.org/10.1016/0306-4522(94)00410-7)
18. SILVERA A, SHYTLER R, PHILIP M, SANBERG P 1995 Transdermal nicotine in Tourette's syndrome. In: Clarke P, Quick M, Adelskofer F, Thurau K (ed) *Effects of Nicotine on Biological Systems II*. Birkhauser, Basel. <https://doi.org/10.1007/978-3-0348-7445-8>
19. SMULDERS CJGM, BUETERS TJH, VAILATI S, VAN KLEEF RGD, VIJVERBERG HPM 2004 Block of neuronal nicotinic acetylcholine receptors by organophosphate insecticides. *Toxicol Sci* 82: 545–554. <https://doi.org/10.1093/toxsci/kfh269>
20. O'MALLEY M 1997 Clinical evaluation of pesticide exposure and poisonings. *Lancet* 349: 1161–1166. <https://doi.org/10.1093/toxsci/kfh269>
21. SOLBERG Y, BELKIN M 1997 The role of excitotoxicity in organophosphorous nerve agents central poisoning. *Trends Pharmacol Sci* 18: 183–185. [https://doi.org/10.1016/s0165-6147\(97\)89540-5](https://doi.org/10.1016/s0165-6147(97)89540-5)
22. ALUIGI MG, ANGELINI C, FALUGI C, FOSSAR, GENEVER P, GALLUS L, LAYER PG, PRESTIPINO G, RAKONCZAY Z, SGRO M, THIELECKE H, TROMBINO S 2005 Interaction between organophosphate compounds and cholinergic functions during development. *Chem Biol Interact* 157–158: 305–16. <https://doi.org/10.1016/j.cbi.2005.10.037>
23. KATZ EJ, CORTES VI, ELDEFRAWI ME, ELDEFRAWI AT 1997 Chlorpyrifos, parathion, and their oxons bind to and desensitize a nicotinic acetylcholine receptor: Relevance to their toxicities. *Toxicol Appl Pharmacol* 146: 227–236. <https://doi.org/10.1006/taap.1997.8201>
24. BAKRY NM, EL-RASHIDY AH, ELDEFRAWI AT, ELDEFRAWI ME 1988 Direct actions of organophosphate anticholinesterases on nicotinic and muscarinic acetylcholine receptors. *J Biochem Toxicol* 3: 235–259. <https://doi.org/10.1002/jbt.2570030404>
25. ELDEFRAWI ME, SCHWEIZER G, BAKRY NM, VALDES JJ 1988 Desensitization of the nicotinic acetylcholine receptor by diisopropylfluorophosphate. *J Biochem Toxicol* 3: 21–32. <https://doi.org/10.1002/jbt.2570030104>
26. RAO KS, ARACAVALA Y, RICKETT DL, ALBUQUERQUE EX 1987 Noncompetitive blockade of the nicotinic acetylcholine receptor-ion channel complex by an irreversible cholinesterase inhibitor. *J Pharmacol Exp Ther* 240: 337–344.
27. TATTERSALL JE 1990 Effects of organophosphorus anticholinesterases on nicotinic receptor ion channels at adult mouse muscle endplates. *Br J Pharmacol* 101: 349–357. <https://doi.org/10.1111/j.1476-5381.1990.tb12713.x>
28. CAULFIELD MP, BIRDSALL NJM 1998 International Union of Pharmacology. XVII. Classification of Muscarinic Acetylcholine Receptors. *Pharmacol Rev* 50 (2): 279–290.
29. ZHANG H, LIU Y, XU J, ZHANG F, LIANG Z, DU X, ZHANG H 2013 Membrane microdomain determines the specificity of 3 receptor-mediated modulation of Kv7/M potassium currents. *Neuroscience* 254: 70–9. <https://doi.org/10.1016/j.neuroscience.2013.08.064>
30. HOLT JC, JORDAN PM, LYSAKOWSKI A, SHAH A, BARSZ K, CONTINI D 2017 Muscarinic acetylcholine receptors and M-currents underlie efferent-mediated slow excitation in calyx-bearing vestibular afferents. *J Neurosci* 37 (7): 1873–1887. <https://doi.org/10.1523/JNEUROSCI.2322-16.2017>
31. MALLOY CA, RITTER K, ROBINSON J, ENGLISH C, COOPER RL 2016 Pharmacological identification of cholinergic receptor subtypes on *Drosophila melanogaster* larval heart. *J Comp Physiol B* 186 (1): 45–57. <https://doi.org/10.1007/s00360-015-0934-4>
32. DI PINTO M, DI BARI R, MARTIN-ALVAREZ R, SPERDUTI S, SERRANO-ACEDO S, GATTA V, TATA AM, MENGOD G 2018 Comparative study of the expression of cholinergic system components in the CNS of experimental autoimmune encephalomyelitis mice: acute vs remitting phase. *Eur J Neurosci* 48 (5): 2165–2181. <https://doi.org/10.1111/ejn.14125>
33. WANG DW, ZHOU RB, YAO YM 2009 Role of cholinergic anti-inflammatory pathway in regulating host response and its interventional strategy for inflammatory diseases. *Chin J Traumatol* 12 (6): 355–64. <https://doi.org/10.3760/cma.j.issn.1008-1275.2009.06.007>
34. KAWASHIMA K, FUJII T, MORIWAKI Y, MISAWA H, HORIGUCHI K 2012 Reconciling neuronally and nonneuronally derived acetylcholine in the regulation of immune function. *Ann N Y Acad Sci* 1261: 7–17. <https://doi.org/10.1111/j.1749-6632.2012.06516.x>
35. LORKE DE, PETROIANU GA 2018 Reversible cholinesterase inhibitors as pretreatment for exposure to organophosphates. A review. *J Appl Toxicol* 39 (1): 101–116. <https://doi.org/10.1002/jat.3662>
36. COSTA LG 2006 Current issue in organophosphate toxicology. *Clin Chim Acta* 366: 1–13. <https://doi.org/10.1016/j.cca.2005.10.008>
37. BAJGAR J, FUSEK J, KASSA J, KUČA K 2015 Global impact of chemical warfare agents used before and after 1945. In: Gupta RC (ed) *Handbook of Toxicology of Chemical Warfare Agents*, 2<sup>nd</sup> ed. Academic Press, p 17–25. <https://doi.org/10.1016/C2013-0-15402-5>
38. UNWIN N 2013 Nicotinic acetylcholine receptor and the structural basis of neuromuscular transmission: insights from Torpedo

- postsynaptic membranes. *Q Rev Biophys* 46 (4): 283–322. <https://doi.org/10.1017/S0033583513000061>
39. FERREIRA-VIEIRA TH, GUIMARAES IM, SILVA FR, RIBEIRO FM 2016 Alzheimer's disease: Targeting the Cholinergic System. *Curr Neuropharmacol* 14 (1): 101–15. <https://doi.org/10.2174/1570159x13666150716165726>
  40. BUCKINGHAM SD, JONES AK, BROWN LA, SATTELE DB 2009 Nicotinic acetylcholine receptor signalling: roles in Alzheimer's disease and amyloid neuroprotection. *Pharmacol Rev* 61 (1): 39–61. <https://doi.org/10.1124/pr.108.000562>
  41. PAPKE RL 2014 Merging old and new perspectives on nicotinic acetylcholine receptors. *Biochem Pharmacol* 89 (1): 1–11. <https://doi.org/10.1016/j.bcp.2014.01.029>
  42. EXLEY R, CRAGG SJ 2008 Presynaptic nicotinic receptors: a dynamic and diverse cholinergic filter of striatal dopamine neurotransmission. *Br J Pharmacol* 153: S283–S297. <https://doi.org/10.1038/sj.bjp.0707510>
  43. WU J, ISHIKAWA M, ZHANG J, HASHIMOTO K 2010 Brain imaging of nicotinic receptors in Alzheimer's disease. *Int J Alzheimers Dis* 2010: 548913. <https://doi.org/10.4061/2010/548913>
  44. GOTTI C, CLEMENTI F, FORNARI A, GAIMARRI A, GUIDUCCI S, MANFREDI I, MORETTI M, PEDRAZZI P, PUCCI L, ZOLI M 2009 Structural and functional diversity of native brain neuronal nicotinic receptors. *Biochem Pharmacol* 78 (7): 703–11. <https://doi.org/10.1016/j.bcp.2009.05.024>
  45. PEREIRA EF, REINHARDT-MAELICKE S, SCHRATTENHOLZ A, MAELICKE A, ALBUQUERQUE EX 1993 Identification and functional characterization of a new agonist site on nicotinic acetylcholine receptors of cultured hippocampal neurons. *J Pharmacol Exp Ther* 265: 1474–1491.
  46. SCHRATTENHOLZ A, COBAN T, SCHRÖDER B, OKONJO KO, KUHLMANN J, PEREIRA EFR, ALBUQUERQUE EX, MAELICKE A 1993 Biochemical characterization of a novel channel-activating site on nicotinic acetylcholine receptors. *J Recept Res* 13: 393–412.
  47. DANI JA 2015 Neuronal Nicotinic Acetylcholine Receptor Structure and Function and Response to Nicotine. *Int Rev Neurobiol* 124: 3–19. <https://doi.org/10.3109/10799899309073669>
  48. ZOLI M, PUCCI S, VILELLA A, GOTTI C 2018 Neuronal and extraneuronal nicotinic acetylcholine receptors. *Curr Neuropharmacol* 16 (4): 338–349. <https://doi.org/10.2174/1570159X15666170912110450>
  49. MCKAY BE, PLACZEK AN, DANI JA 2007 Regulation of synaptic transmission and plasticity by neuronal nicotinic acetylcholine receptors. *Biochem Pharmacol* 74 (8): 1120–33. <https://doi.org/10.1016/j.bcp.2007.07.001>
  50. KARLIN A 2002 Emerging structure of the nicotinic acetylcholine receptors. *Nat Rev Neurosci* 3: 102–114. <https://doi.org/10.1038/nrn731>
  51. UNWIN N 2005 Refined structure of the nicotinic acetylcholine receptor at 4Å resolution. *J Mol Biol* 346 (4): 967–89. <https://doi.org/10.1016/j.jmb.2004.12.031>
  52. GREENBAUM L, LERER B 2009 Differential contribution of genetic variation in multiple brain nicotinic cholinergic receptors to nicotine dependence: recent progress and emerging open questions. *Mol Psychiatry* 14 (10):912–45. <https://doi.org/10.1038/mp.2009.59>
  53. HURST R, ROLLEMA H, BERTRAND D Nicotinic acetylcholine receptors: from basic science to therapeutics. 2013 *Pharmacol Ther* 137 (1): 22–54. <https://doi.org/10.1016/j.pharmthera.2012.08.012>
  54. ENGEL AG, OHNO K, MILONE M, WANG HL, NAKANO S, BOUZAT C, PRUITT JN 2<sup>ND</sup>, HUTCHINSON DO, BRENGMAN JM, BREN N, SIEB JP, SINE SM 1996 New mutations in acetylcholine receptor subunit genes reveal heterogeneity in the slow-channel congenital myasthenic syndrome. *Hum Mol Genet* 5: 1217–1227. <https://doi.org/10.1093/hmg/5.9.1217>
  55. ARIDON P, MARINI C, DI RESTA C, BRILLI E, DE FUSCO M, POLITI F, PARRINI E, MANFREDI I, PISANO T, PRUNA D, CURIA G, CIANCHETTI C, PASQUALETTI M, BECCHETTI A, GUERRINI R, CASARI G 2006 Increased sensitivity of the neuronal nicotinic receptor alpha 2 subunit causes familial epilepsy with nocturnal wandering and ictal fear. *Am J Hum Genet* 79: 342–350. <https://doi.org/10.1086/506459>
  56. MICHALKA, STRICKER S, BECKER J, RUPPS R, PANTZAR T, MIERTUS J, BOTTA G, NARETTO VG, JANETZKI C, YAQOUB N, OTT CE, SEELOW D, WIECZOREK D, FIEBIG B, WIRTH B, HOOPMANN M, WALTHER M, KÖRBER F, BLANKENBURG M, MUNDLOS S, HELLER R, HOFFMANN K 2008 Acetylcholine receptor pathway mutations explain various fetal akinesia deformation sequence disorders. *Am J Hum Genet* 82: 464–476. <https://doi.org/10.1016/j.ajhg.2007.11.006>
  57. QUIK M, BORDIA T, O'LEARY K 2007 Nicotinic receptors as CNS targets for Parkinson's disease. *Biochem Pharmacol* 74: 1224–1234. <https://doi.org/10.1016/j.bcp.2007.06.01>
  58. KELSO ML, WEHNER JM, COLLINS AC, SCHEFF SW, PAULY JR 2006 The pathophysiology of traumatic brain injury in  $\alpha 7$  nicotinic cholinergic receptor knockout mice. *Brain Res* 1083: 204–210. <https://doi.org/10.1016/j.brainres.2006.01.127>
  59. BROIDE RS, LESLIE FM 1999 The  $\alpha 7$  nicotinic acetylcholine receptor in neuronal plasticity. *Mol Neurobiol* 20: 1–16. <https://doi.org/10.1007/BF02741361>
  60. MORLEY BJ, HAPPE HK 2000 Cholinergic receptors: dual roles in transduction and plasticity. *Hear Res* 147: 104–112. [https://doi.org/10.1016/s0378-5955\(00\)00124-6](https://doi.org/10.1016/s0378-5955(00)00124-6)
  61. KHIROUG SS, HARKNESS PC, LAMB PW, SUDWEEKS SN, KHIROUG L, MILLAR NS, YAKEL JL 2002 Rat nicotinic ACh receptor  $\alpha 7$  and  $\beta 2$  subunits co-assemble to form functional heteromeric nicotinic receptor channels. *J Physiol* 540: 425–434. <https://doi.org/10.1113/jphysiol.2001.013847>
  62. GOTTI C, ZOLI M, CLEMENTI F 2006 Brain nicotinic acetylcholine receptors: native subtypes and their relevance. *Trends Pharmacol Sci* 27 (9): 482–91. <https://doi.org/10.1016/j.tips.2006.07.004>
  63. DUNCKLEY T, WU J, ZHAO L, LUKAS RJ 2003 Mutational analysis of roles for extracellular cysteine residues in the assembly and function of human  $\alpha 7$ -nicotinic acetylcholine receptors. *Biochemistry* 42: 870–876. <https://doi.org/10.1021/bi020586x>
  64. WANAMAKER CP, CHRISTIANSON JC, GREEN WN 2003 Regulation of nicotinic acetylcholine receptor assembly. *Ann NY Acad Sci* 998: 66–80. <https://doi.org/10.1196/annals.1254.009>
  65. CELIE PH, VAN ROSSUM-FIKKERT SE, VAN DIJK WJ, BREJC K, SMIT AB, SIXMA TK 2004 Nicotine and carbamylcholine binding to nicotinic acetylcholine receptors as studied in AChBP crystal structures. *Neuron* 41: 907–914. [https://doi.org/10.1016/s0896-6273\(04\)00115-1](https://doi.org/10.1016/s0896-6273(04)00115-1)
  66. PENG X, GERZANICH V, ANAND R, WHITING PJ, LINDSTROM J 1994 Nicotine-induced increase in neuronal nicotinic receptors results from a decrease in the rate of receptor turnover. *Mol Pharmacol* 46: 523–530.
  67. GREEN WN, WANAMAKER CP 1997 The role of the cysteine loop in acetylcholine receptor assembly. *J Biol Chem* 272: 20945–20953. <https://doi.org/10.1074/jbc.272.33.20945>
  68. GEHLE VM, WALCOTT EC, NISHIZAKI T, SUMIKAWA K 1997 N-glycosylation at the conserved sites ensures the expression of properly folded functional ACh receptors. *Brain Res* 45: 219–229. [https://doi.org/10.1016/s0169-328x\(96\)00256-2](https://doi.org/10.1016/s0169-328x(96)00256-2)
  69. daCOSTA CJB, KAISER DE, BAENZIGER JE 2005 Role of glycosylation and membrane environment in nicotinic acetylcholine receptor stability. *Biophys J* 88 (3): 1755–64. <https://doi.org/10.1529/biophysj.104.052944>
  70. ROTH AF, WAN J, BAILEY AO, SUN B, KUCHAR JA, GREEN WN, PHINNEY BS, YATES JR III, DAVIS NG 2006 Global

- analysis of protein palmitoylation in yeast. *Cell* 125: 1003–1013. <https://doi.org/10.1016/j.cell.2006.03.042>
71. ALEXANDER JK, GOVIND AP, DRISDEL RC, BLANTON MP, VALLEJO Y, LAM TT, GREEN WN 2010 Palmitoylation of nicotinic acetylcholine receptors. *J Mol Neurosci* 40 (1–2): 12–20. <https://doi.org/10.1007/s12031-009-9246-z>
  72. POLLOCK, VV, PASTOOR T, KATNIK C, CUEVAS J, WECKER L 2009 Cyclic AMP-dependent protein kinase A and protein kinase C phosphorylate  $\alpha 4\beta 2$  nicotinic receptor subunits at distinct stages of receptor formation and maturation. *Neuroscience* 158: 1311–1325. <https://doi.org/10.1016/j.neuroscience.2008.11.032>
  73. LAI A, PARAMESWARAN N, KHWAJA M, WHITEAKER P, LINDSTROM JM, FAN H, MCINTOSH JM, GRADY SR, QUIK M 2005 Long-term nicotine treatment decreases striatal  $\alpha 6^*$  nicotinic acetylcholine receptor sites and function in mice. *Mol Pharmacol* 67: 1639–1647. <https://doi.org/10.1124/mol.104.006429>
  74. DAJAS-BAILADOR FA, HEIMALA K, WONNACOTT S 2003 The allosteric potentiation of nicotinic acetylcholine receptors by galantamine is transduced into cellular responses in neurons:  $Ca^{2+}$  signals and neurotransmitter release. *Mol Pharmacol* 64: 1217–1226. <https://doi.org/10.1124/mol.64.5.1217>
  75. FENSTER CP, WHITWORTH TL, SHEFFIELD EB, QUICK MW, LESTER RA 1999 Upregulation of surface  $\alpha 4\beta 2$  nicotinic receptors is initiated by receptor desensitization after chronic exposure to nicotine. *J Neurosci* 19: 4804–4814. <https://doi.org/10.1523/JNEUROSCI.19-12-04804.1999>
  76. HIREMAGALUR B, SABBAN EL 1995 Nicotine elicits changes in expression of adrenal catecholamine biosynthetic enzymes, neuropeptide Y and immediate early genes by injection but not continuous administration. *Brain Res* 32: 109–115. [https://doi.org/10.1016/0169-328x\(95\)00068-4](https://doi.org/10.1016/0169-328x(95)00068-4)
  77. DE JONGE WJ, VAN DER ZANDEN EP, THE FO, BIJLSMA MF, VAN WESTERLOO DJ, BENNINK RJ, BERTHOUD HR, UEMATSU S, AKIRA S, VAN DEN WIJNGAARD RM, BOECKXSTAENS GE. Stimulation of the vagus nerve attenuates macrophage activation by activating the Jak2-STAT3 signaling pathway. *Nat Immunol* 6: 844–851. <https://doi.org/10.1038/nri1229>
  78. RUBIN DT, HANAUER SB 2000 Smoking and inflammatory bowel disease. *Eur J Gastroenterol Hepatol* 12: 855–862. <https://doi.org/10.1097/00042737-200012080-00004>
  79. PARK HJ, LEE PH, AH NYW, CHOI YJ, LEE G, LEE DY, CHUNG ES, JIN BK 2007 Neuroprotective effect of nicotine on dopaminergic neurons by anti-inflammatory action. *Eur J Neurosci* 26: 79–89. <https://doi.org/10.1111/j.1460-9568.2007.05636.x>
  80. BROWN D, RAMLOCHANSINGH C, MANAYE KF, TIZABI Y 2013 Nicotine promotes survival of cells expressing amyloid precursor protein and presenilin: implication for Alzheimer's disease. *Neurosci Lett* 535: 57–61. <https://doi.org/10.1016/j.neulet.2012.12.046>
  81. SABBAGH MN, LUKAS RJ, SPARKS DL, REID RT 2002 The nicotinic acetylcholine receptor, smoking, and Alzheimer's disease. *J Alzheimers Dis* 4: 317–325. <https://doi.org/10.3233/jad-2002-4407>
  82. POSADAS I, LÓPEZ-HERNÁNDEZ B, CEÑA V. 2013 Nicotinic receptors in neurodegeneration. *Curr Neuropharmacol* 11 (3): 298–314. <https://doi.org/10.2174/1570159X11311030005>
  83. TAYLOR ME, CLOSE JCT 2018 Dementia. *Handb Clin Neurol* 159: 303–321. <https://doi.org/10.1016/B978-0-444-63916-5.00019-7>
  84. KAR S, SLOWIKOWSKI SP, WESTAWAY D, MOUNT HT 2004 Interactions between beta-amyloid and central cholinergic neurons: implications for Alzheimer's disease. *J Psychiatry Neurosci* 29 (6): 427–41.
  85. ŠIMIĆ G, BABIĆ LEKO M, WRAY S, HARRINGTON C, DELALLE I, JOVANOVIĆ MILOŠEVIĆ N, BAŽADONA D, BUÉE L, DE SILVA R, DI GIOVANNI G, WISCHIK C, HOF PR. Tau protein hyperphosphorylation and aggregation in Alzheimer's disease and other tauopathies, and possible neuroprotective strategies. *Biomolecules* 6 (1): 6. <https://doi.org/10.3390/biom6010006>
  86. LOMBARDO S, MASKOS U 2014 Role of the nicotinic acetylcholine receptor in Alzheimer's disease pathology and treatment. *Neuropharmacology* 96 (Pt B): 255–62. <https://doi.org/10.1016/j.neuropharm.2014.11.018>
  87. SHEN J, WU J 2015 Nicotinic Cholinergic Mechanisms in Alzheimer's Disease. *Int Rev Neurobiol* 124: 275–92. <https://doi.org/10.1016/bs.irn.2015.08.002>
  88. AULD DS, KORNECOOK TJ, BASTIANETTO S, QUIRION R 2002 Alzheimer's disease and the basal forebrain cholinergic system: relations to  $\beta$ -amyloid peptides, cognition, treatment strategies. *Prog Neurobiol* 68: 209–245. [https://doi.org/10.1016/s0301-0082\(02\)00079-5](https://doi.org/10.1016/s0301-0082(02)00079-5)
  89. MAELICKE A, SCHRATTENHOLZ A, SAMOCHOCKI M, RADINA M, ALBUQUERQUE EX 2000 Allosterically potentiating ligands of nicotinic receptors as a treatment strategy for Alzheimer's disease. *Behav Brain Res* 113 (1–2): 199–206. [https://doi.org/10.1016/s0166-4328\(00\)00214-x](https://doi.org/10.1016/s0166-4328(00)00214-x)
  90. MARTIN-RUIZ CM, COURT JA, MOLNAR E, LEE M, GOTTI C, MAMALAKI A, TSOULOUFIS T, TZARTOS S, BAL-LARD C, PERRY RH, PERRY EK 1999  $\alpha 4$  but not  $\alpha 3$  and  $\alpha 7$  nicotinic acetylcholine receptor subunits are lost from the temporal cortex in Alzheimer's disease. *J Neurochem* 73: 1635–1640. <https://doi.org/10.1046/j.1471-4159.1999.0731635.x>
  91. FUJII S, SUMIKAWA K 2001 Acute and chronic nicotine exposure reverse age-related declines in the induction of long-term potentiation in the rat hippocampus. *Brain Res* 894: 347–353. [https://doi.org/10.1016/s0006-8993\(01\)02057-1](https://doi.org/10.1016/s0006-8993(01)02057-1)
  92. CROWE M, ANDEL R, PEDERSEN NL, JOHANSSON B, GATZ M 2003 Does participation in leisure activities lead to reduced risk of Alzheimer's disease? A prospective study of Swedish twins. *J Gerontol B Psychol Sci Soc Sci* 58: 249–255. <https://doi.org/10.1093/geronb/58.5.p249>
  93. HAYDAR SN, DUNLOP J 2010 Neuronal Nicotinic Acetylcholine Receptors – Targets for the Development of Drugs to Treat Cognitive Impairment Associated with Schizophrenia and Alzheimer's Disease. *Curr Top Med Chem* 10: 140–152. <https://doi.org/10.2174/156802610790410983>
  94. SAHOO AK, DANDAPAT J, DASH UC, KANHAR S 2018 Features and outcomes of drugs for combination therapy as multi-targets strategy to combat Alzheimer's disease. *J Ethnopharmacol* 215: 42–73. <https://doi.org/10.1016/j.jep.2017.12.015>
  95. WATKINS PB, ZIMMERMAN HJ, KNAPP MJ, GRACON SI, LEWIS KW 1994 Hepatotoxic effects of tacrine administration in patients with Alzheimer's disease. *JAMA* 271: 992–998. <https://doi.org/10.1001/jama.1994.03510370044030>
  96. PAPKE RL, TROCMÉ-THIBIERGE C, GUENDISCH D, AL RUBAIY SA, BLOOM SA 2011 Electrophysiological perspectives on the therapeutic use of nicotinic acetylcholine receptor partial agonists. *J Pharmacol Exp Ther* 337 (2): 367–79. <https://doi.org/10.1124/jpet.110.177485>
  97. AKK G, STEINBACH, JH 2005 Galantamine activates muscarinic nicotinic acetylcholine receptors without binding to the acetylcholine binding site. *J Neurosci* 25: 1992–2001. <https://doi.org/10.1523/JNEUROSCI.4985-04.2005>
  98. TARIOT PN, SCHNEIDER L, KATZ IR, MINTZER JE, STREET J, COPENHAVER M, WILLIAMS-HUGHES C 2006 Quetiapine treatment of psychosis associated with dementia: a double-blind, randomized, placebo controlled clinical trial. *Am J Geriatr Psychiatry* 14: 767–776. <https://doi.org/10.1097/01.JGP.0000196628.12010.35>
  99. JOHNSON KA, FOX NC, SPERLING RA, KLUNK WE 2012 Brain imaging in Alzheimer disease. *Cold Spring Harb Perspect Med* 2 (4): a006213. <https://doi.org/10.1101/cshperspect.a006213>

100. LEDIG C, SCHUH A, GUERRERO R, HECKEMANN RA, RUECKERT D 2018 Structural brain imaging in Alzheimer's disease and mild cognitive impairment: biomarker analysis and shared morphometry database. *Sci Rep* 8 (1): 11258. <https://doi.org/10.1038/s41598-018-29295-9>
101. GLICK SD, SELL EM, MCCALLUM SE, MAISONNEUVE IM 2011 Brain regions mediating  $\alpha 3\beta 4$  nicotinic antagonist effects of 18-MC on nicotine self-administration. *Eur J Pharmacol* 669 (1–3): 71–75. <https://doi.org/10.1016/j.ejphar.2011.08.001>
102. TIETJE KR, ANDERSON DJ, BITNER RS, BLOMME EA, BRACKEMEYER PJ, BRIGGS CA, BROWMAN KE, BURY D, CURZON P, DRESCHER KU, FROST JM, FRYER RM, FOX GB, GRONLIEN JH, HÅKERUD M, GUBBINS EJ, HALM S, HARRIS R, HELFRICH RJ, KOHLHAAS KL, LAW D, MALYSZ J, MARSH KC, MARTIN RL, MEYER MD, MOLESKY AL, NIKKEL AL, OTTE S, PAN L, PUTTFARCKEN PS, RADEK RJ, ROBB HM, SPIES E, THORIN-HAGENE K, WARING JF, WEEN H, XU H, GOPALAKRISHNAN M, BUNNELLE WH 2008 Preclinical characterization of A-582941: a novel  $\alpha 7$  neuronal nicotinic receptor agonist with broad spectrum cognition-enhancing properties. *CNS Neurosci Ther* 14 (1): 65–82. <https://doi.org/10.1111/j.1527-3458.2008.00037.x>
103. BOESS FG, DE VRY J, ERB C, FLESSNER T, HENDRIX M, LUTHLE J, METHFESSEL C, RIEDL B, SCHNIZLER K, VAN DER STAAY F-J, VAN KAMPEN M, WIESE WB, KOENIG G 2007 The novel  $\alpha 7$  nicotinic acetylcholine receptor agonist N-[(3R)-1-azabicyclo[2.2.2]oct-3-yl]-7-[2-(methoxyphenyl)-1-benzofuran-2-carboxamide improves working and recognition memory in rodents. *J Pharmacol Exp Ther* 321 (2): 716–725. <https://doi.org/10.1124/jpet.106.118976>
104. BERTRAND D, TERRY AV Jr 2018 The wonderland of neuronal nicotinic acetylcholine receptors. *Biochem Pharmacol* 151: 214–225. <https://doi.org/10.1016/j.bcp.2017.12.008>
105. VAN KAMPEN M, SELBACH K, SCHNEIDER R, SCHIEGEL E, BOESS F, SCHREIBER R 2004 AR-R 17779 improves social recognition in rats by activation of nicotinic  $\alpha 7$  receptors. *Psychopharmacology (Berl)* 172 (4): 375–383. <https://doi.org/10.1007/s00213-003-1668-7>
106. TIMPERLEY CM, BIRD M, GORE SJ, LINDSAY CD, RICE H, TATTERSALL JEH, WHITMORE CL, GREEN AC 2020 3-Quinuclidinyl- $\alpha$ -methoxydiphenylacetate: A multi-targeted ligand with antimuscarinic and antinicotinic effects designed for the treatment of anticholinesterase poisoning. *Toxicol Lett* 325: 67–76. <https://doi.org/10.1016/j.toxlet.2020.01.027>
107. GEE KW, OLINCY A, KANNER R, JOHNSON L, HOGENKAMP D, HARRIS J, TRAN M, EDMONDS SA, SAUER W, YOSHIMURA R, JOHNSTONE T, FREEDMAN R 2017 First in human trial of a type I positive allosteric modulator of  $\alpha 7$ -nicotinic acetylcholine receptors: Pharmacokinetics, safety, and evidence for neurocognitive effect of AVL-3288. *J Psychopharmacol* 31 (4): 434–441. <https://doi.org/10.1177/0269881117691590>
108. LUO W, GEX, CUI W, WANG H 2018 Targeting cholinesterase inhibitor poisoning with a novel blocker against both nicotinic and muscarinic receptors. *Neurotox Res* 18 (2): 173–186. <https://doi.org/10.1007/s12640-009-9119-9>
109. CANNADY R, WEIR R, WEE B, GOTSCHLICH E, KOLIA N, LAU E, BROTHERTON J, LEVIN ED 2009 Nicotinic antagonist effects in the mediodorsal thalamic nucleus: regional heterogeneity of nicotinic receptor involvement in cognitive function. *Biochem Pharmacol* 78 (7): 788–794. <https://doi.org/10.1016/j.bcp.2009.05.021>
110. REZVANI AH, CAULEY M, SEXTON H, XIAO Y, BROWN ML, PAIGE MA, MCDOWELL BE, KELLAR KJ, LEVIN ED 2011 Sazetidine-A, a selective  $\alpha 4\beta 2$  nicotinic acetylcholine receptor ligand: effects on dizocilpine and scopolamine-induced attentional impairments in female Sprague-Dawley rats. *Psychopharmacology (Berl)* 215 (4): 620–630. <https://doi.org/10.1007/s00213-010-2161-8>
111. FERREIRA-VIEIRA TH, GUIMARAES IM, SILVA FR, RIBEIRO FM 2016 Alzheimer's disease: targeting the cholinergic system. *Curr Neuropharmacol* 14 (1): 101–115. <https://doi.org/10.2174/1570159x13666150716165726>
112. FELIX R, LEVIN ED 1997 Nicotinic antagonist administration into the ventral hippocampus and spatial working memory in rats. *Neuroscience* 81 (4): 1009–1017. [https://doi.org/10.1016/s0306-4522\(97\)00224-8](https://doi.org/10.1016/s0306-4522(97)00224-8)
113. PRICKAERTS J, VAN GOETHEM NP, CHESWORTH R, SHAPIRO G, BOESS FG, METHFESSEL C, RENEERKENS OAH, FLOOD DG, HILT D, GAWRYL M, BERTRAND S, BERTRAND D, KÖNIG G 2012 EVP-6124, a novel and selective  $\alpha 7$  nicotinic acetylcholine receptor partial agonist, improves memory performance by potentiating the acetylcholine response of  $\alpha 7$  nicotinic acetylcholine receptors. *Neuropharmacology* 62 (2): 1099–1110. <https://doi.org/10.1016/j.neuropharm.2011.10.024>
114. ARIAS HR, FEUERBACH D, TARGOWSKA-DUDA KM, JOZWIAK K 2010 Catharanthine alkaloids are noncompetitive antagonists of muscle-type nicotinic acetylcholine receptors. *Neurochem Int* 57(2): 153–161. <https://doi.org/10.1016/j.neuint.2010.05.007>
115. ROGAWSKI MA, WENK GL 2003 The neuropharmacological basis for the use of memantine in the treatment of Alzheimer's disease. *CNS Drug Rev* 9 (3): 275–308. <https://doi.org/10.1111/j.1527-3458.2003.tb00254.x>
116. CANNADY R, WEIR R, WEE B, GOTSCHLICH E, KOLIA N, LAU E, BROTHERTON J, LEVIN ED 2009 Nicotinic antagonist effects in the mediodorsal thalamic nucleus: regional heterogeneity of nicotinic receptor involvement in cognitive function. *Biochem Pharmacol* 78 (7): 788–794. <https://doi.org/10.1016/j.bcp.2009.05.021>
117. HAJÓS M, HURST RS, HOFFMANN WE, KRAUSE M, WALL TM, HIGDON NR, GROPPI VE 2005 The selective  $\alpha 7$  nicotinic acetylcholine receptor agonist PNU-282987 [N-[(3R)-1-Azabicyclo[2.2.2]oct-3-yl]-4-chlorobenzamide hydrochloride] enhances GABAergic synaptic activity in brain slices and restores auditory gating deficits in anesthetized rats. *J Pharmacol Exp Ther* 312 (3): 1213–1222. <https://doi.org/10.1124/jpet.104.076968>
118. RONCARATI R, SCALI C, COMERY TA, GRAUER SM, ASCHMI S, BOTHMANN H, JOW B, KOWAL D, GIANFRIDDO M, KELLEY C, ZANELLI U, GHIRON C, HAYDAR S, DUNLOP J, TERSTAPPEN GC 2009 Pro-cognitive and neuro-protective activity of a novel  $\alpha 7$  nicotinic acetylcholine receptor agonist for treatment of neurodegenerative and cognitive disorders. *J Pharmacol Exp Ther* 329 (2): 459–468. <https://doi.org/10.1124/jpet.108.150094>
119. PICCHAT P, BERGIS OE, TERRANOVA JP, URANI A, DUARTEC, SANTUCCI V, GUEUDET C, VOLTZ C, STEINBERG R, STEMMELIN J, OURY-DONAT F, AVENET P, GRIEBEL G, SCATTON B 2007 SSR180711, a novel selective  $\alpha 7$  nicotinic receptor partial agonist: (II) efficacy in experimental models predictive of activity against cognitive symptoms of schizophrenia. *Neuropsychopharmacology* 32 (1): 17–34. <https://doi.org/10.1038/sj.npp.1301188>
120. CALLAHAN PM, BERTRAND D, BERTRAND S, PLAGENHOEF MR, TERRY AV Jr 2017 Tropisetron sensitizes  $\alpha 7$  containing nicotinic receptors to low levels of acetylcholine *in vitro* and improves memory-related task performance in young and aged animals. *Neuropharmacology* 117: 422–433. <https://doi.org/10.1016/j.neuropharm.2017.02.025>
121. MOORE MA, MCCARTHY MP 1995 Snake venom toxins, unlike smaller antagonists, appear to stabilize a resting state conformation of the nicotinic acetylcholine receptor. *Biochim Biophys Acta* 1235 (2): 336–342. [https://doi.org/10.1016/0005-2736\(95\)80022-8](https://doi.org/10.1016/0005-2736(95)80022-8)
122. STAUDT MD, DI SEBASTIANO AR, XU H, JOG M, SCHMID S, FOSTER P, HEBB MO 2016 Advances in Neuro-

- trophic Factor and Cell-Based Therapies for Parkinson's Disease: A Mini-Review. *Gerontology* 62 (3): 371–380. <https://doi.org/10.1159/000438701>
123. WHITFIELD AC, MOORE BT, DANIELS RN 2014 Classics in chemical neuroscience: levodopa. *ACS Chem Neurosci* 5 (12): 1192–7. <https://doi.org/10.1021/cn5001759>
124. SPILLER K, XI ZX, LI X, ASHBY CR JR, CALLAHAN PM, TEHIMA A, GARDNER EL 2009 Varenicline attenuates nicotine-enhanced brain-stimulation reward by activation of alpha4beta2 nicotinic receptors in rats. *Neuropharmacology* 57 (1): 60–6. <https://doi.org/10.1016/j.neuropharm.2009.04.006>
125. Haddad F, Sawalha M, Khawaja Y, Najjar A, Karaman R 2017 Dopamine and levodopa prodrugs for the treatment of Parkinson's disease. *Molecules* 23 (1): pii: E40. <https://doi.org/10.3390/molecules23010040>
126. COLLINS AC, MARKS MJ 1991 Progress towards the development of animal models of smoking-related behaviors. *J Addict Dis* 10: 109–126. [https://doi.org/10.1300/J069v10n01\\_08](https://doi.org/10.1300/J069v10n01_08)
127. QUIK M 2004 Smoking, nicotine and Parkinson's disease. *Trends Neurosci* 27: 561–568. <https://doi.org/10.1016/j.tins.2004.06.008>
128. RYAN RE, ROSS SA, DRAGO J, LOIACONO RE 2001 Dose-related neuroprotective effects of chronic nicotine in 6-hydroxydopamine treated rats, and loss of neuro-protection in  $\alpha 4$  nicotinic receptor subunit knockout mice. *Br J Pharmacol* 132: 1650–1656. <https://doi.org/10.1038/sj.bjp.0703989>
129. LESTER DB, ROGERS TD, BLAHA CD 2010 Acetylcholine-dopamine interactions in the pathophysiology and treatment of CNS disorders. *CNS Neurosci Ther* 16 (3): 137–162. <https://doi.org/10.1111/j.1755-5949.2010.00142.x>
130. KISS C, CERESOLI-BORRONI G, GUIDETTI P, ZIELKE CL, ZIELKE HR, SCHWARCZ R 2003 Kynurenate production by cultured human astrocytes. *J Neural Transm* 110: 1–14. <https://doi.org/10.1007/s00702-002-0770-z>
131. BOSAK A 2006 Organofosforni spojevi: klasifikacija i reakcije s enzimima. *Arh Hig Rada Toksikol* 57: 445–57.
132. VALE JA, MARRS TC OBE, MAYNARD RL CBE 2018 Novichok: a murderous nerve agent attack in the UK. *Clin Toxicol (Phila)* 14: 1–5. <https://doi.org/10.1080/15563650.2018.1469759>
133. VOICU V, BAJGAR J, MEDVEDOVICI A, RADULESCU FS, MIRON DS 2010 Pharmacokinetics and pharmacodynamics of some oximes and associated therapeutic consequences: A critical review. *J Appl Toxicol* 30 (8): 719–29. <https://doi.org/10.1002/jat.1561>
134. MYHRER T, PÅL A 2015 Prophylactic and therapeutic measures in nerve agent poisoning. In: Gupta RC (ed) *Handbook of Toxicology of Chemical Warfare Agents*, 2<sup>nd</sup> ed. Academic Press, p 989–1001. <https://doi.org/10.1016/C2013-0-15402-5>
135. JOHN H, BALZUWEIT F, KEHE K, WOREK F, THIERMANN H 2015 Toxicokinetic aspects of nerve agents and vesicants. In: Gupta RC (ed) *Handbook of Toxicology of Chemical Warfare Agents*, 2<sup>nd</sup> ed. Academic Press, p 817–856. <https://doi.org/10.1016/C2013-0-15402-5>
136. GUPTA RC 2015 Introduction. In: Gupta RC (ed) *Handbook of Toxicology of Chemical Warfare Agents*, 2<sup>nd</sup> ed. Academic Press, p 3–5. <https://doi.org/10.1016/C2013-0-15402-5>
137. WESTFALL TC, WESTFALL DP 2006 Neurotransmission: The autonomic and somatic motor nervous systems. In: Brunton LL, Lazo JS, Parker KL (ed) *Goodman and Gilman's the Pharmacological Basis of Therapeutics*, 11<sup>th</sup> ed. New York: McGraw-Hill, p 137–182. <https://doi.org/10.1021/jm058286b>
138. PEREIRA DM, FERRERES F, OLIVEIRA JM, GASPAR L, FARIA J, VALENTÃO P, SOTTOMAYOR M, ANDRADE PB 2010 Pharmacological effects of *Catharanthus roseus* root alkaloids in acetylcholinesterase inhibition and cholinergic neurotransmission. *Phytomedicine*. 17 (8–9): 646–52. <https://doi.org/10.1016/j.phymed.2009.10.008>
139. MOADDEL R, ABDRAKHMANOVA G, KOZAK J, JOZWIAK K, TOLL L, JIMENEZ L, ROSENBERG A, TRAN T, XIAO Y, ZARATE CA, WAINER IW 2013 Sub-anesthetic concentrations of (*R,S*)-ketamine metabolites inhibit acetylcholine-evoked currents in  $\alpha 7$  nicotinic acetylcholine receptors. *Eur J Pharmacol* 698 (1–3): 228–34. <https://doi.org/10.1016/j.ejphar.2012.11.023>
140. KIM YB, CHEON KC, HU GH 2002 Effects of combinational prophylactics composed of physostigmine and procyclidine on soman-induced lethality, seizures and brain injuries. *Env Toxicol Pharmacol* 11: 15–21. [https://doi.org/10.1016/S1382-6689\(01\)00096-5](https://doi.org/10.1016/S1382-6689(01)00096-5)