



Hot topic

Human *Brucella melitensis* infections in southern Vietnam

Brucellosis is a collective term for infections caused by small Gram-negative coccobacilli belonging to genus *Brucella*. This genus incorporates the well-described animal pathogens *Brucella melitensis*, *Brucella abortus*, *Brucella ovis*, *Brucella suis*, and *Brucella canis*, which are associated with disease in goats, cattle, sheep, pigs, and dogs, respectively. *Brucella* are facultative intracellular pathogens, and are sequestered by monocytes and macrophages, spreading throughout the body to the liver, spleen, lymph nodes, and bone marrow [1]. These pathogens are synonymous with an aggressive disease syndrome in animals causing abortion, stillbirth, and the delivery of weak offspring. The organisms replicate to high concentrations in the affected tissues and are transmitted through contact with the placenta, foetus, foetal fluids, and vaginal discharge. Notably, goats can shed *B. melitensis* in vaginal discharge for up to 3 months after abortion and organisms can be shed in milk for the lifetime of an infected animal [2].

Many *Brucella* species have zoonotic potential and can be transmitted from animals to humans. Brucellosis in humans is typically contracted by contact with infected animals or through the ingestion of animal products prepared from infected animals. In symptomatic cases, disease presentation is highly variable and may arise rapidly or progressively. Classically, brucellosis in humans is a sub-acute, non-specific febrile disease, characterized by high temperatures, headaches, malaise, night sweats, and body aches [3]. Some individuals recover quickly, while others develop more persistent, long-term complications including arthritis, spondylitis, endocarditis, dermatitis, and chronic fatigue, and neurological complications [3]. The disease is treated using antimicrobials; however, relapses are common, even after apparent bacteriological cure.

Many middle-high income countries employ successful control programmes to reduce brucellosis in animals and humans. However, such control programmes or surveillance infrastructures are less common in low-middle income countries (LMICs). Consequently, animal brucellosis is endemic in parts of Asia, the Middle East, East and North Africa, Latin America, and some southern and eastern European countries, and poses a potential risk to human health in these locations [4]. The complex variety of symptoms in humans makes the disease difficult to diagnose and it is likely that it is underreported. Vietnam is an LMIC in Southeast Asia where brucellosis is not a commonly reported cause of febrile disease [5], and no cases have been reported since three cases of *B. abortus* in 1962 [6].

The Hospital for Tropical Diseases (HTD) in Ho Chi Minh City (HCMC), serves as a primary and secondary facility for the surrounding local population and a tertiary referral centre for

infectious diseases for southern Vietnam. Non-specific febrile disease is a common reason for admission to HTD [7]. Blood culture is performed routinely on patients in whom an infection is suspected on the basis of fever (>38°C) or clinical evidence of sepsis. For adult patients, 8–15 mL of venous blood is drawn and inoculated into BACTECplus aerobic bottles (Becton Dickinson, USA), prior to incubation at 37°C in a BACTEC9050 analyzer.

From 14 June 2016 to 18 January 2017, 10 febrile patients attending HTD had a positive blood culture containing Gram-negative coccobacilli. These organisms were sub-cultured onto sheep chocolate blood agar and subjected to biochemical identification and antimicrobial susceptibility testing. The organisms stained red using a modified cold Ziehl-Neelsen stain, and were identified as *Brucella* spp. on a VITEK2 system (BioMerieux, France). All organisms were susceptible to amikacin, ciprofloxacin, gentamycin, doxycycline, imipenem, rifampicin, and trimethoprim sulphate. Nucleic acid was extracted from organisms and subjected to Bruce-ladder multiplex PCR to identify the infecting species [8]; all produced an identical collection of amplicons indicative of *B. melitensis*. We next performed MLVA-16 (Multiple Locus VNTR Analysis) on the 10 *Brucella* isolates [9], comprising three panels, panel 1 (eight minisatellite loci), panel 2A (three microsatellite loci), and panel 2B (five microsatellite loci). Panel 1 allows clustering the different *Brucella* species while panels 2A and 2B provide finer resolution characterization. The MLVA profiles of the Vietnamese isolates were compared with a global collection of various *Brucella* species (Fig. 1(a)). The organisms were all confirmed as *B. melitensis* and produced independent VNTR profiles falling into four subgroups, clustering with organisms originating from Southern Europe, the Middle East, and China.

The presumptive diagnoses of the brucellosis patients prior to bacterial culture and identification were sepsis (patients 1 and 2), non-specific viral infection (3), dengue (4), tuberculosis (5 and 7), and non-specific inflammatory disease (6); patients 8, 9, and 10 were correctly diagnosed with brucellosis after laboratory diagnosis of the first seven patients. These infections were additionally confirmed using Rose Bengal agglutination with titres in plasma ranging from 1/4 to 1/256. Differential blood counts were largely unremarkable but almost all patients had elevated AST, ALT, and GGT; several patients had elevated CRP and procalcitonin indicating systemic inflammatory response. On review of the medical histories, all patients had reported exposure to goats prior to the febrile episodes; eight kept goats, two had consumed goat meat, and one was a veterinarian who had been vaccinating goats. The 10 cases originated from four provinces, with the primary cases occurring in Binh Phuoc and Tay Ninh near the Cambodian border,




Fig. 1. Locations and MLVA genotypes of 10 human *Brucella melitensis* cases in southern Vietnam. (a) Circular tree representing the global population structure of the genus *Brucella* using Multiple Locus VNTR Analysis-16 (MLVA-16) analysis. Nine different *Brucella* species (labelled) are shown and associated with main branches; *B. melitensis* forms the majority of the tree. The 10 Vietnamese isolates are on black branches within the *B. melitensis* population. (b) Map of southern Vietnam showing the locations of provinces where the 10 cases of brucellosis originated between June 2016 and January 2017. The solid circle shows the location of the Hospital for Tropical Diseases in Ho Chi Minh City.

and later detection in the south (Ho Chi Minh City and Long An) (Fig. 1(b)). All patients, apart from patient 5 (ceftriaxone 2 g/day for 7 days) received doxycycline (200 mg/day for 6 weeks) with gentamycin (240 mg/day for 7 days). All patients recovered without relapse with the exception of patient 5, who was treated with imipenem (2 g/day), gentamycin (240 mg/day) for 7 days and doxycycline (200 mg/day) for 6 weeks. Patient 5 made a complete recovery without additional relapse.

B. melitensis is a known zoonotic pathogen that can cause an invasive febrile disease in humans exposed to infected animals [10]. Diagnosing brucellosis in humans is complicated by its non-specific presentation and may not be included in a differential diagnosis. The lack of a confirmatory diagnosis with appropriate antimicrobial therapy can lead to lasting physical effects through chronic infections. Persistent intracellular infection is associated with chronic arthritis, endocarditis, and neurobrucellosis. Here we have reported the first identified cases of human brucellosis caused by *B. melitensis* in Vietnam. The organisms were susceptible to the advocated antimicrobial agents, again highlighting the importance of pathogen isolation and identification. By performing MLVA genotyping, we found that the organisms fell into four different subgroups, suggesting that these organisms circulate widely in goats in southern Vietnam and were not part of an isolated outbreak. Therefore, we suggest that patients with non-specific febrile disease in Vietnam and comparable locations in Southeast Asia reporting contact with goats, sheep, and cattle receive a blood culture and a Rose-Bengal test. If the diagnosis is strongly suspected, empirical doxycycline can be given.

Vietnam is a hotspot for zoonotic infections and brucellosis is an important disease globally, stemming from the circulation of undiagnosed sick animals coming into contact with humans. Given the distribution of these cases and the collective exposure to goats, these findings suggest that these organisms are circulating widely in the goat population, which predicts that human cases may become increasingly common. We recommend that sick animals with a suspicion of brucellosis and their owners be screened in the identified provinces to assess the magnitude of the problem. This type of screening requires interaction between the

Departments of Preventative Medicine and Animal Health. Our work indicates the importance of zoonotic infections in Vietnam and highlights the need for sustained surveillance in human and animal populations.

Transparency declaration

This study was supported by a strategic award from the Wellcome Trust of Great Britain (WT/093724). S. Baker is a Sir Henry Dale Fellow, jointly funded by the Wellcome Trust and the Royal Society (100087/Z/12/Z). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

- [1] de Figueiredo P, Ficht TA, Rice-Ficht A, Rossetti CA, Adams LG. Pathogenesis and immunobiology of brucellosis. *Am J Pathol* 2015;185:1505–17.
- [2] Tittarelli M, Di Ventura M, De Massis F, Scacchia M, Giovannini A, Nannini D, et al. The persistence of *Brucella melitensis* in experimentally infected ewes through three reproductive cycles. *J Vet Med Ser B* 2005;52:403–9.
- [3] Jia B, Zhang F, Lu Y, Zhang W, Li J, Zhang Y, et al. The clinical features of 590 patients with brucellosis in Xinjiang, China with the emphasis on the treatment of complications. *PLoS Negl Trop Dis* 2017;11, e0005577.
- [4] Rubach MP, Halliday JEB, Cleaveland S, Crump JA. Brucellosis in low-income and middle-income countries. *Curr Opin Infect Dis* 2013;26:404–12. <http://dx.doi.org/10.1097/QCO.0b013e3283638104>.
- [5] Nga TTT, de Vries PJ, Abdoel TH, Smits HL. Brucellosis is not a major cause of febrile illness in patients at public health care facilities in Binh Thuan Province. *Vietnam J Infect* 2006;53:12–5.
- [6] Spinu I, Vasilescu T, Dang-Tran-Dung. The presence of human and animal brucellosis in North Vietnam. *Microbiol Parazitol Si Epidemiol* 1962;7: 459–69.
- [7] Nga TVT, Parry CM, Le T, Lan NPH, Diep TS, Campbell JI, et al. The decline of typhoid and the rise of non-typhoid salmonellae and fungal infections in a changing HIV landscape: bloodstream infection trends over 15 years in southern Vietnam. *Trans R Soc Trop Med Hyg* 2012;106:26–34.
- [8] Lopez-Goni I, Garcia-Yoldi D, Marin CM, de Miguel MJ, Munoz PM, Blasco JM, et al. Evaluation of a multiplex PCR assay (Bruce-ladder) for molecular typing of all *Brucella* species, including the vaccine strains. *J Clin Microbiol* 2008;46: 3484–7.
- [9] Maquart M, Le Fleche P, Foster G, Tryland M, Ramisse F, Djonje B, et al. MLVA-16 typing of 295 marine mammal *Brucella* isolates from different animal and

geographic origins identifies 7 major groups within *Brucella ceti* and *Brucella pinnipedialis*. *BMC Microbiol* 2009;9:145.

- [10] Moreno E, Moriyon I. *Brucella melitensis*: a nasty bug with hidden credentials for virulence. *Proc Natl Acad Sci USA* 2002;99:1–3.

J.I. Campbell

The Hospital for Tropical Diseases, Wellcome Trust Major Overseas Programme, Oxford University Clinical Research Unit, Ho Chi Minh City, Viet Nam

Centre for Tropical Medicine and Global Health, Nuffield Department of Clinical Medicine, Oxford University, Oxford, UK

N.P.H. Lan

The Hospital for Tropical Diseases, Wellcome Trust Major Overseas Programme, Oxford University Clinical Research Unit, Ho Chi Minh City, Viet Nam

P.M. Phuong, L.B. Chau

The Hospital for Tropical Diseases, Ho Chi Minh City, Viet Nam

Trung Pham Duc

The Hospital for Tropical Diseases, Wellcome Trust Major Overseas Programme, Oxford University Clinical Research Unit, Ho Chi Minh City, Viet Nam

C. Guzmán-Verri, N. Ruiz-Villalobos
Universidad Nacional, Heredia, Costa Rica

T.P.T. Minh

The Hospital for Tropical Diseases, Wellcome Trust Major Overseas Programme, Oxford University Clinical Research Unit, Ho Chi Minh City, Viet Nam

P.M. Muñoz Álvaro

Unidad de Producción y Sanidad Animal del Instituto Agroalimentario de Aragón, Universidad de Zaragoza, Spain

E. Moreno

Universidad Nacional, Heredia, Costa Rica

G.E. Thwaites, M.A. Rabaa

The Hospital for Tropical Diseases, Wellcome Trust Major Overseas Programme, Oxford University Clinical Research Unit, Ho Chi Minh City, Viet Nam

Centre for Tropical Medicine and Global Health, Nuffield Department of Clinical Medicine, Oxford University, Oxford, UK

N.V.V. Chau

The Hospital for Tropical Diseases, Ho Chi Minh City, Viet Nam

S. Baker*

The Hospital for Tropical Diseases, Wellcome Trust Major Overseas Programme, Oxford University Clinical Research Unit, Ho Chi Minh City, Viet Nam

Centre for Tropical Medicine and Global Health, Nuffield Department of Clinical Medicine, Oxford University, Oxford, UK

The Department of Medicine, Cambridge University, Cambridge, UK

* Corresponding author. Professor S. Baker, Oxford University Clinical Research Unit, 764 Vo Van Kiet, Ward 1, District 5, Ho Chi Minh, Viet Nam.

E-mail address: sbaker@oucru.org (S. Baker).

19 May 2017

Available online 29 June 2017

Editor: G. Greub