

Search for Gravitational Waves from High-Mass-Ratio Compact-Binary Mergers of Stellar Mass and Subsolar Mass Black Holes

Alexander H. Nitz^{*} and Yi-Fan Wang (王一帆)

Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut), D-30167 Hannover, Germany and Leibniz Universität Hannover, D-30167 Hannover, Germany



(Received 8 July 2020; accepted 11 December 2020; published 12 January 2021)

We present the first search for gravitational waves from the coalescence of stellar mass and subsolar mass black holes with masses between $20\text{--}100 M_{\odot}$ and $0.01\text{--}1 M_{\odot}$ ($10\text{--}10^3 M_J$), respectively. The observation of a single subsolar mass black hole would establish the existence of primordial black holes and a possible component of dark matter. We search the ~ 164 day of public LIGO data from 2015–2017 when LIGO-Hanford and LIGO-Livingston were simultaneously observing. We find no significant candidate gravitational-wave signals. Using this nondetection, we place a 90% upper limit on the rate of $30\text{--}0.01 M_{\odot}$ and $30\text{--}0.1 M_{\odot}$ mergers at $< 1.2 \times 10^6$ and $< 1.6 \times 10^4 \text{ Gpc}^{-3} \text{ yr}^{-1}$, respectively. If we consider binary formation through direct gravitational-wave braking, this kind of merger would be exceedingly rare if only the lighter black hole were primordial in origin ($< 10^{-4} \text{ Gpc}^{-3} \text{ yr}^{-1}$). If both black holes are primordial in origin, we constrain the contribution of $1(0.1)M_{\odot}$ black holes to dark matter to $< 0.3(3)\%$.

DOI: [10.1103/PhysRevLett.126.021103](https://doi.org/10.1103/PhysRevLett.126.021103)

Introduction.—The first gravitational wave observation from the merger of black holes was detected on September 14, 2015 [1]. Over a dozen binary black hole (BBH) mergers [2–8] have since been reported along with two binary neutron star mergers [9,10] by Advanced LIGO [11] and Virgo [12]. Recently, two compact binary coalescences with unequal masses have been reported [13,14]; the mass ratios are ~ 3 and ~ 9 , respectively.

The nature of dark matter remains a mystery given null results from direct searches using particle experiments (see, e.g., Refs. [15,16] and recent notable exception in Ref. [17]). The observation of BBH mergers has sparked renewed interest in primordial black holes (PBHs) as a contributor to dark matter [18–26]. However, the merger of stellar-mass PBHs may be difficult to separate from standard stellar formation channels. Black holes may form through standard stellar evolution between $2\text{--}50 M_{\odot}$ [27–32]. Furthermore, gravitational-wave observation alone is not always able to determine if a component of a binary is either a neutron star or black hole [33,34]. Although the observation of coincident gamma-ray bursts or kilonovae, such as in the case of GW170817 [9,35–38], can confirm the presence of nuclear matter. In contrast, there is no

known model which can produce subsolar mass black holes by conventional formation mechanisms; the observation of a single subsolar mass black hole would provide strong evidence for PBHs.

There are a variety of constraints for the contributing fraction of PBHs to dark matter (see Refs. [39,40] and references therein). Gravitational-wave astronomy provides a unique window; notably, a direct search for subsolar mass black holes constrained the mass range $0.2\text{--}2 M_{\odot}$ for near equal-mass sources [41,42] and the nondetection of a gravitational-wave astrophysical background by LIGO and Virgo has constrained PBHs with $0.01\text{--}100 M_{\odot}$ [43]. Recently, tight constraints from the NANOGrav pulsar timing array [44] are given by Ref. [45] for $0.001\text{--}1 M_{\odot}$ black holes based on the nondetection of gravitational waves induced by scalar perturbations during the expected PBH formation epoch.

So far, all observations of gravitational waves from BBH mergers were identified by searches targeting stellar-mass sources, i.e., those with component masses $1 - \mathcal{O}(100)M_{\odot}$. Targeted searches for subsolar mass binaries with component masses between $0.2\text{--}2 M_{\odot}$ have null results [41,42]. We report a search for subsolar mass black holes in an unexplored region of parameter space: the merger of $0.01\text{--}1 M_{\odot}$ subsolar mass black holes with $20\text{--}100 M_{\odot}$ stellar-mass black holes. We summarize the region we search in comparison to past analyses in Fig. 1. We find no statistically significant candidates and place the first constraints from gravitational-wave observation on the merger rate of these sources.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Open access publication funded by the Max Planck Society.




FIG. 1. The primary and secondary masses of the sources searched by our analysis (blue), 2-OGC/GWTC-1 (green) [3,8], and the subsolar mass LVC search (orange) [41].

Search.—We analyze the public LIGO data from 2015–2017, which contains ~ 164 day of joint LIGO-Hanford and LIGO-Livingston observation time [46,47]. Virgo was observing for the final month of this period, but had limited range in comparison to the LIGO instruments. We use the open-source PyCBC-based search pipeline [48,49] configured similarly to the analysis of Ref. [3] to analyze the LIGO data, identify potential candidates, apply tests of each candidate’s signal consistency [50,51], rank each candidate, and finally assess each candidate’s statistical significance [52–54]. The statistical significance of any candidate is assessed by comparing to the empirically estimated rate of false alarms. This rate is estimated by creating numerous fictitious analyses analyzed in an identical manner to the search, but where time shifts between the data of the two LIGO observatories are applied. The time shift of each background analysis is greater than the light-travel-time between the two LIGO observatories, which ensures astrophysical signals are not in coincidence. The average sensitive distance of our analysis at a false alarm rate of 1 per 100 years is shown in Fig. 2.

Matched filtering is used to extract the signal-to-noise from data for a given template waveform [48,55]. Each template corresponds to the gravitational-wave signal of a single source type. To search for sources with varied component masses, a discrete bank of template waveforms is required. We use a brute-force stochastic method [56] to find the nearly 9 million templates required by our analysis. To reduce computational cost, we search at most the final 60 seconds of each gravitational waveform. For the lightest sources, this implies we analyze the data starting at a higher gravitational-wave frequency than for the heaviest sources, where we analyze the data starting from 20 Hz.

To model the gravitational-wave signal, we use EOBNRv2, a model based on an effective one-body Hamiltonian approximation of general relativity in combination with a fitted merger and ringdown [57]. Several phenomenological models exist for BBH mergers,




FIG. 2. The distance our search can detect sources at a false alarm rate of 1/100 yr as a function of the primary and secondary masses, averaged over the possible sky locations and orientations of an isotropic source population, and averaged over the observation period. The horizon distance, the maximum distance a source could be found, is a factor of ~ 2.26 larger than the average range shown here.

however, they do not generalize to the high mass ratios we consider [58,59]. We assume our sources’ orbits have negligible eccentricity by the time of observation and that the component black holes are nonspinning. This choice is consistent with the prediction that PBHs have negligible component spin [60–63]. Because of the degeneracy between mass ratio and spin [64], we expect our search to be able to recover moderately spinning sources where $\chi_{1,2} \lesssim 0.1$ [65,66], which is well beyond the larger predictions at the percent level [67]. EOBNRv2 is too slow for use by our search directly. Instead, we use a straightforward interpolant based on $\sim 10^4$ pregenerated EOBNRv2 waveforms with different mass ratios that can be rapidly scaled to any point in parameter space. We crosscheck our model




FIG. 3. Comparison of the gravitational waveform for a $30 M_{\odot}$ - $0.01 M_{\odot}$ merger. The EOBNRv2 interpolant model used by our search is consistent with the EMRISur1dq1e4 surrogate model when the inclination of the source’s orbital plane is close to face-on or face-off. For sources with highly inclined orbital planes, higher order modes becomes increasingly important.

against the recent extreme mass ratio inspiral (EMRI) surrogate EMRISur1dq1e4 [68] and find that our interpolant, the base EOBNRv2 model, and the dominant-mode of EMRISur1dq1e4 have less than $< 1\%$ mismatch at all locations in our template bank, i.e., any of these models would recover $> 99\%$ of the SNR of a signal matching one of the other models. A visual comparison between a representative example of these models is in Fig. 3.

Gravitational waves are expressed in terms of spin-weighted spherical harmonic decomposed modes. Methods exist for incorporating higher order modes into searches at increased computation cost [69]. EOBNRv2 provides only the dominant mode of the gravitational waveform, and accurate models with higher order modes exist only for lower mass ratio sources [70–72], or short duration signals [68]. We compare our templates against these models to estimate the potential loss in search sensitivity. Neglecting higher order modes in our search imposes an source-orientation averaged loss in signal-to-noise ratio (SNR) of $\sim 5\%$, 10% , 15% , 22% for sources with a $20 M_{\odot}$, $40 M_{\odot}$, $60 M_{\odot}$, and $100 M_{\odot}$ primary mass, respectively. The most significant loss in sensitivity is for sources with near edge-on inclination of their orbital plane with respect to an observer, whereas higher order modes become negligible for sources with near face-on inclination. The search sensitivity and upper limits quoted in this Letter account for the detection rate reduction.

Observational results.—The most significant candidate from our search was observed at a false alarm rate of 3 per year (Additional details about the most significant candidates can be found at Ref. [73].), and if it were astrophysical, would be consistent with the merger of a $23 M_{\odot}$ primary black hole with a $0.012 M_{\odot}$ secondary. Considering the time searched, our results are consistent with a null observation.

We place upper limits at 90% confidence on the rate of mergers throughout the searched space using the loudest event method [74]. The upper limit R_{90} is given as

$$R_{90} = \frac{2.3}{VT}, \quad (1)$$

where V is the sensitive volume of our analysis at the false alarm rate of the most significant candidate, and T is the total time searched. We simulate a population of sources distributed isotropically in the sky and binary orientation, and uniform in volume, to measure the sensitive volume of our analysis as a function of the primary and secondary masses. Figure 4 shows the upper limit on the merger rate as a function of the secondary mass. Assuming a distribution of primary masses consistent with the black holes observed by LIGO and Virgo, we find that the rate of $0.01 M_{\odot}$ solar mass mergers is $< 1.7 \times 10^6 \text{ Gpc}^{-3} \text{ yr}^{-1}$ at 90% confidence.

Implications for primordial black holes.—Whereas stellar-mass black holes can be either the product of stellar evolution or PBHs, subsolar mass black holes only form in the primordial Universe given conventional stellar




FIG. 4. The 90% upper limit on the rate of mergers as a function of the mass of the secondary black hole, for a range of primary masses (various colors), and the average assuming a primary mass population consistent with observed BBH mergers from the 2-OGC catalog (black) [3]. The 1σ Monte Carlo statistical uncertainty is shown with shading.

evolution [39,40]. There are two possible origins for the high-mass-ratio binaries we considered; the first that only the secondary, lighter black hole is primordial and forms a binary with a stellar-origin black hole in the late Universe, and a second scenario where both black holes are primordial and formed a binary in the early Universe.

For the first scenario, binaries form in the galactic field through dynamical capture due to gravitational-wave bremsstrahlung. For PBHs with mass m_1 , stellar-origin black holes with mass m_2 and relative velocity v , the cross section for binary formation is given by Ref. [75] as

$$\sigma = 2\pi \left(\frac{85\pi}{6\sqrt{2}} \right)^{2/7} \frac{G^2 (m_1 + m_2)^{10/7} (m_1 m_2)^{2/7}}{c^{10/7} v^{18/7}}, \quad (2)$$

where G and c are the gravitational constant and speed of light, respectively. As shown by Ref. [22], binaries are expected to quickly merge after formation and disruption by other PBHs can be neglected.

To estimate the PBH distribution, we use a recent cosmological galaxy formation simulation *IllustrisTNG* [76]. In the redshift = 0 snapshot of the TNG-100 high resolution simulation, there are $\sim 10^5$ dark matter main subhalos with nonzero star formation within a ~ 100 Mpc size cube. For each main subhalo, we assume the dark matter number density follows the Navarro-Frenk-White (NFW) profile ρ_{NFW} [77], and that PBHs constitute a fraction of dark matter with mass fraction f_{PBH} .

Estimating the abundance of stellar-origin black holes is a challenge due to the lack of observation. Nevertheless, the synthesis population study of Ref. [78] shows that $\sim 0.006\%$ of the total galactic halo mass including dark matter is in the form of black holes. As an approximation, we take this value as the universal fraction over dark matter main subhalos to infer the mass density ρ_{BH} of stellar-origin black holes.

The rate density of dynamical captures between primordial and stellar-origin black holes is finally

$$R(m_1, m_2) = \sum_{\text{Halos}} \int_0^{\sqrt[3]{2}R_{\text{halfmass}}} \frac{\rho_{\text{BH}}}{m_1} \frac{f_{\text{PBH}} \rho_{\text{NFW}}(r)}{m_2} \sigma v d^3r. \quad (3)$$

Assuming a uniform spatial distribution of stellar-origin black holes, the radius r is integrated from the main subhalo center to $\sqrt[3]{2}$ times of the radius which contains half of the stellar mass, R_{halfmass} . The relative velocity v is approximated by the stellar dispersion velocity, provided by *IllustrisTNG*. We find that even for $f_{\text{PBH}} = 100\%$, this formation channel implies a merger rate $< 10^{-4} \text{ Gpc}^{-3} \text{ yr}^{-1}$ for $37 M_{\odot} - 0.01 M_{\odot}$ binaries.

The estimation of ρ_{BH} is a source of uncertainty, however, other variables in Eq. (3) such as dark matter halo abundance are not expected to change by orders of magnitude since they are extracted from the robust simulation. Given that the resultant rate is ~ 10 orders of

magnitude lower than that shown in Fig. 4, our conclusion that this scenario is unlikely is robust.

On the other hand, if both primary and secondary black holes are PBHs, a nearest pair may form a binary if decoupled from the Universe's expansion. References [25,79] consider a uniform spatial distribution when PBHs form and give the merger rate for a binary with mass m_1 and m_2 as

$$R(m_1, m_2) = 3.3 \times 10^6 \cdot f_{\text{PBH}}^2 (0.7 f_{\text{PBH}}^2 + \sigma_{\text{eq}}^2)^{-\frac{21}{74}} (m_1 m_2)^{\frac{3}{37}} \times (m_1 + m_2)^{\frac{36}{37}} \min \left(\frac{P(m_1)}{m_1}, \frac{P(m_2)}{m_2} \right) \left(\frac{P(m_1)}{m_1} + \frac{P(m_2)}{m_2} \right), \quad (4)$$

where mass m and merger rate R are in units of M_{\odot} and $\text{Gpc}^{-3} \text{ yr}^{-1}$, respectively. $P(m)$ is the normalized PBH mass distribution. The parameter σ_{eq} accounts for the variance of density perturbation from other dark matter aside from PBHs at the matter radiation equality epoch and is suggested to be 0.005 by Ref. [79].

The possibility that the currently observed stellar-mass BBH mergers were caused by PBHs is a topic of investigation [22–26,80]. In the most optimistic case, where the majority of LIGO and Virgo observed black holes are PBHs, $f_{\text{PBH}}^{\text{primary}} = 3 \times 10^{-3}$ by Ref. [80]. With this fixed fraction for the primary mass, we constrain the contribution of the secondary, subsolar mass black hole to dark matter. We assume a two-valued mass distribution, i.e., $P(m_1) + P(m_2) = 100\%$. Thus the fraction in dark matter for the primary and secondary black hole is $f_{\text{PBH}}^{\text{primary}} = P(m_1) f_{\text{PBH}}$ and $f_{\text{PBH}}^{\text{secondary}} = P(m_2) f_{\text{PBH}}$.

The upper limit for $f_{\text{PBH}}^{\text{secondary}}$ for a fixed fiducial primary mass $m_1 = 20/50/100 M_{\odot}$ and the average mass from the 2-OGC catalog ($\sim 37 M_{\odot}$) [3] are shown in Fig. 5. For the




FIG. 5. Upper limits on $f_{\text{PBH}}^{\text{secondary}}$ for the secondary, subsolar mass black hole assuming both primary and secondary black holes have primordial origin, where we choose the primary mass to be $20/50/100 M_{\odot}$ (blue/orange/green) or the average mass of the 2-OGC catalog ($\sim 37 M_{\odot}$) (black). The constraints from the LVC direct search for equal-mass PBHs [41] are plotted for comparison.

2-OGC average case, we find that $1(0.1) M_{\odot}$ PBHs cannot exceed 0.3(3)% of the total dark matter. In contrast, if we assume none of the LIGO/Virgo BBH detections are PBHs, our results cannot constrain f_{PBH} .

Our constraints can be directly compared with the targeted search for near equal-mass subsolar black holes [41,42] which used the same formation scenario as described by Eq. (4). Our results expand the range probed by direct merger observation down to $0.01 M_{\odot}$. The constraint on the abundance of PBHs is improved by an order of magnitude as we consider sources with higher total mass that emit stronger gravitational waves.

A significant source of uncertainty in this model derives from the fraction of binaries which survive to merger. Under active investigation is what fraction of PBH binaries would be disrupted after initial formation. If a significant fraction are disrupted, the event rate would be lowered and our constraints loosened. Reference [79] has shown analytically that the disruption can be neglected, however, recently Ref. [81] argues for a higher disruption rate. Further, if PBHs exhibit substantial clustering at formation, the event rate may be boosted and our constraints would be tighter [82,83]. As we consider the same model, both our results and LVC limits shown in Fig. 5 do not consider binary disruption and assume a uniform spatial distribution when PBHs form.

Stringent constraints for subsolar mass PBHs from pulsar timing arrays [45] have almost excluded the $0.001\text{--}1 M_{\odot}$ mass region, overlapping with our $0.01\text{--}1 M_{\odot}$ region. However, the scalar induced gravitational waves considered in Ref. [45] apply to *Gaussian* scalar curvature perturbation in the early Universe. References [84,85] have shown that non-Gaussianity can suppress the scalar induced gravitational waves by orders of magnitude depending on the detailed form of the perturbations. Positive results from a direct search for subsolar mass compact objects would imply large local non-Gaussianity of primordial perturbation to alleviate the tension. Null results may also have implications for early Universe non-Gaussianity, but a detailed analysis is beyond the scope of this work.

Conclusions.—We conduct a search for a previously unconsidered source of gravitational waves: the binary coalescence of high-mass-ratio sources, where the primary mass is $20\text{--}100 M_{\odot}$ and the secondary mass is $0.01\text{--}1 M_{\odot}$. We find no promising candidates, and thus place improved upper limits on the merger rate and the abundance of PBHs.

The merging of a PBH with a black hole formed through stellar evolution is extremely unlikely under the scenario of direct capture through gravitational-wave braking. A significantly more efficient binary formation mechanism would be required for this scenario to make a significant contribution. On the other hand, assuming both black holes are primordial in origin places constraints on the abundance of PBHs.

Advanced LIGO and Virgo are continually being upgraded [86], and the third generation of gravitational-wave detectors can further improve the horizon distance by an order of magnitude [87,88]. At that point, it will be possible to probe the redshift evolution of stellar-mass binaries to distinguish primordial and stellar-origin black hole distributions [80]. From our results, we expect the constraint on subsolar mass PBH abundance to be $10^{3\text{--}4}$ times tighter than the current search, assuming a null result. Combining the results of ground-based detectors, pulsar timing, and possibly space-based detectors in the future, can together probe the existence of PBHs and may investigate the structure of primordial perturbations in the early Universe [89].

The top candidates from our analysis along with the configuration files necessary to reproduce the search are available in Ref. [73].

This research has made use of data, software and/or web tools obtained from the Gravitational Wave Open Science Center [90], a service of LIGO Laboratory, the LIGO Scientific Collaboration and the Virgo Collaboration.

We acknowledge the Max Planck Gesellschaft and the Atlas cluster computing team at AEI Hannover for support. LIGO is funded by the U.S. National Science Foundation. Virgo is funded by the French Centre National de Recherche Scientifique (CNRS), the Italian Istituto Nazionale della Fisica Nucleare (INFN) and the Dutch Nikhef, with contributions by Polish and Hungarian institutes.

*alex.nitz@aei.mpg.de

- [1] B. P. Abbott *et al.* (Virgo and LIGO Scientific Collaborations), Observation of Gravitational Waves from a Binary Black Hole Merger, *Phys. Rev. Lett.* **116**, 061102 (2016).
- [2] A. H. Nitz, C. Capano, A. B. Nielsen, S. Reyes, R. White, D. A. Brown, and B. Krishnan, 1-OGC: The first open gravitational-wave catalog of binary mergers from analysis of public Advanced LIGO data, *Astrophys. J.* **872**, 195 (2019).
- [3] A. H. Nitz, T. Dent, G. S. Davies, S. Kumar, C. D. Capano, I. Harry, S. Mozzon, L. Nuttall, A. Lundgren, and M. Tpai, 2-OGC: Open gravitational-wave catalog of binary mergers from analysis of public Advanced LIGO and Virgo data, *Astrophys. J.* **891**, 123 (2019).
- [4] A. H. Nitz, T. Dent, G. S. Davies, and I. Harry, A search for gravitational waves from binary mergers with a single observatory, *Astrophys. J.* **897**, 169 (2020).
- [5] T. Venumadhav, B. Zackay, J. Roulet, L. Dai, and M. Zaldarriaga, New search pipeline for compact binary mergers: Results for binary black holes in the first observing run of Advanced LIGO, *Phys. Rev. D* **100**, 023011 (2019).
- [6] T. Venumadhav, B. Zackay, J. Roulet, L. Dai, and M. Zaldarriaga, New binary black hole mergers in the second observing run of Advanced LIGO and Advanced Virgo, *Phys. Rev. D* **101**, 083030 (2020).

- [7] B. Zackay, L. Dai, T. Venumadhav, J. Roulet, and M. Zaldarriaga, Detecting gravitational waves with disparate detector responses: Two new binary black hole mergers, [arXiv:1910.09528](#).
- [8] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), GWTC-1: A Gravitational-Wave Transient Catalog of Compact Binary Mergers Observed by LIGO and Virgo during the First and Second Observing Runs, *Phys. Rev. X* **9**, 031040 (2019).
- [9] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral, *Phys. Rev. Lett.* **119**, 161101 (2017).
- [10] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), GW190425: Observation of a compact binary coalescence with total mass $\sim 3.4 M_{\odot}$, *Astrophys. J. Lett.* **892**, L3 (2020).
- [11] J. Aasi *et al.* (LIGO Scientific Collaboration), Advanced LIGO, *Classical Quantum Gravity* **32**, 074001 (2015).
- [12] F. Acernese *et al.* (Virgo Collaboration), Advanced Virgo: A second-generation interferometric gravitational wave detector, *Classical Quantum Gravity* **32**, 024001 (2015).
- [13] R. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), GW190412: Observation of a Binary-Black-Hole Coalescence with Asymmetric Masses, *Phys. Rev. D* **102**, 043015 (2020).
- [14] R. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), GW190814: Gravitational waves from the coalescence of a 23 solar mass black hole with a 2.6 solar mass compact object, *Astrophys. J.* **896**, L44 (2020).
- [15] X. Cui *et al.* (PandaX-II Collaboration), Dark Matter Results from 54-Ton-Day Exposure of Pandax-II Experiment, *Phys. Rev. Lett.* **119**, 181302 (2017).
- [16] R. Agnese *et al.* (SuperCDMS Collaboration), Results from the Super Cryogenic Dark Matter Search Experiment at Soudan, *Phys. Rev. Lett.* **120**, 061802 (2018).
- [17] E. Aprile *et al.* (XENON Collaboration), Observation of excess electronic recoil events in XENON1T, *Phys. Rev. D* **102**, 072004 (2020).
- [18] S. Hawking, Gravitationally collapsed objects of very low mass, *Mon. Not. R. Astron. Soc.* **152**, 75 (1971).
- [19] B. J. Carr and S. W. Hawking, Black holes in the early Universe, *Mon. Not. R. Astron. Soc.* **168**, 399 (1974).
- [20] A. Dolgov and J. Silk, Baryon isocurvature fluctuations at small scales and baryonic dark matter, *Phys. Rev. D* **47**, 4244 (1993).
- [21] K. Jedamzik, Primordial black hole formation during the QCD epoch, *Phys. Rev. D* **55**, R5871 (1997).
- [22] S. Bird, I. Cholis, J. B. Muñoz, Y. Ali-Haïmoud, M. Kamionkowski, E. D. Kovetz, A. Raccanelli, and A. G. Riess, Did Ligo Detect Dark Matter?, *Phys. Rev. Lett.* **116**, 201301 (2016).
- [23] S. Clesse and J. García-Bellido, The clustering of massive primordial black holes as dark matter: Measuring their mass distribution with Advanced LIGO, *Phys. Dark Universe* **10**, 002 (2016).
- [24] M. Sasaki, T. Suyama, T. Tanaka, and S. Yokoyama, Primordial Black Hole Scenario for the Gravitational-Wave Event GW150914, *Phys. Rev. Lett.* **117**, 061101 (2016).
- [25] Z.-C. Chen and Q.-G. Huang, Merger rate distribution of primordial-black-hole binaries, *Astrophys. J.* **864**, 61 (2018).
- [26] V. De Luca, G. Franciolini, P. Pani, and A. Riotto, Primordial black holes confront LIGO/Virgo data: Current situation, *J. Cosmol. Astropart. Phys.* **06** (2020) 044.
- [27] C. L. Fryer, K. Belczynski, G. Wiktorowicz, M. Dominik, V. Kalogera, and D. E. Holz, Compact remnant mass function: Dependence on the explosion mechanism and metallicity, *Astrophys. J.* **749**, 91 (2012).
- [28] C. S. Kochanek, Failed Supernovae explain the compact remnant mass function, *Astrophys. J.* **785**, 28 (2014).
- [29] T. Ertl, H.-Th. Janka, S. E. Woosley, T. Sukhbold, and M. Ugliano, A two-parameter criterion for classifying the explodability of massive stars by the neutrino-driven mechanism, *Astrophys. J.* **818**, 124 (2016).
- [30] S. E. Woosley, Pulsational pair-instability Supernovae, *Astrophys. J.* **836**, 244 (2017).
- [31] P. Marchant, M. Renzo, R. Farmer, K. M. W. Pappas, R. E. Taam, S. E. de Mink, and V. Kalogera, Pulsational pair-instability supernovae in very close binaries, *Astrophys. J.* **882**, 36 (2019).
- [32] R. Farmer, M. Renzo, S. E. de Mink, P. Marchant, and S. Justham, Mind the gap: The location of the lower edge of the pair instability supernovae black hole mass gap, *Astrophys. J.* **887**, 53 (2019).
- [33] H. Yang, W. E. East, and L. Lehner, Can we distinguish low mass black holes in neutron star binaries?, *Astrophys. J.* **856**, 110 (2018); Erratum, *Astrophys. J.* **870**, 139 (2019).
- [34] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), Model comparison from LIGO–Virgo data on GW170817’s binary components and consequences for the merger remnant, *Classical Quantum Gravity* **37**, 045006 (2020).
- [35] B. P. Abbott *et al.* (LIGO Scientific, Virgo, Fermi-GBM, and INTEGRAL Collaborations), Gravitational waves and gamma-rays from a binary neutron star merger: GW170817 and GRB 170817A, *Astrophys. J. Lett.* **848**, L13 (2017).
- [36] B. P. Abbott *et al.* (GROND, SALT Group, OzGrav, DFN, INTEGRAL, Virgo, Insight-Hxmt, MAXI Team, Fermi-LAT, J-GEM, RATIR, IceCube, CAASTRO, LWA, ePESSTO, GRAWITA, RIMAS, SKA South Africa/MeerKAT, H.E.S.S., 1M2H Team, IKI-GW Follow-up, Fermi GBM, Pi of Sky, DWF (Deeper Wider Faster Program), Dark Energy Survey, MASTER, AstroSat Cadmium Zinc Telluride Imager Team, Swift, Pierre Auger, ASKAP, VINROUGE, JAGWAR, Chandra Team at McGill University, TTU-NRAO, GROWTH, AGILE Team, MWA, ATCA, AST3, TOROS, Pan-STARRS, NuSTAR, ATLAS Telescopes, BOOTES, CaltechNRAO, LIGO Scientific, High Time Resolution Universe Survey, Nordic Optical Telescope, Las Cumbres Observatory Group, TZAC Consortium, LOFAR, IPN, DLT40, Texas Tech University, HAWC, ANTARES, KU, Dark Energy Camera GW-EM, CALET, Euro VLBI Team, and ALMA Collaborations), Multimessenger Observations of a binary neutron star merger, *Astrophys. J.* **848**, L12 (2017).
- [37] M. Soares-Santos *et al.* (DES and Dark Energy Camera GW-EM Collaborations), The electromagnetic counterpart of the binary neutron star merger LIGO/Virgo GW170817.

- I. Discovery of the optical counterpart using the dark energy camera, *Astrophys. J.* **848**, L16 (2017).
- [38] D. A. Coulter *et al.*, Swope Supernova Survey 2017a (SSS17a), the optical counterpart to a gravitational wave source, *Science* **358**, 1556 (2017).
- [39] B. Carr and F. Kuhnel, Primordial black holes as dark matter, *Annu. Rev. Nucl. Part. Sci.* **70**, 355 (2020).
- [40] B. Carr, K. Kohri, Y. Sendouda, and J. Yokoyama, Constraints on primordial black holes, [arXiv:2002.12778](https://arxiv.org/abs/2002.12778).
- [41] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), Search for Substellar Mass Ultracompact Binaries in Advanced LIGO's Second Observing Run, *Phys. Rev. Lett.* **123**, 161102 (2019).
- [42] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), Search for Substellar-Mass Ultracompact Binaries in Advanced LIGO's First Observing Run, *Phys. Rev. Lett.* **121**, 231103 (2018).
- [43] S. Wang, Y.-F. Wang, Q.-G. Huang, and T. G. F. Li, Constraints on the Primordial Black Hole Abundance from the First Advanced LIGO Observation Run Using the Stochastic Gravitational-Wave Background, *Phys. Rev. Lett.* **120**, 191102 (2018).
- [44] Z. Arzoumanian *et al.* (NANOGrav Collaboration), The NANOGrav 11-year data set: High-precision timing of 45 millisecond pulsars, *Astrophys. J. Suppl. Ser.* **235**, 37 (2018).
- [45] Z.-C. Chen, C. Yuan, and Q.-G. Huang, Pulsar Timing Array Constraints on Primordial Black Holes with NANOGrav 11-Year Data Set, *Phys. Rev. Lett.* **124**, 251101 (2020).
- [46] M. Vallisneri, J. Kanner, R. Williams, A. Weinstein, and B. Stephens, The LIGO open science center, *J. Phys. Conf. Ser.* **610**, 012021 (2015).
- [47] R. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), Open data from the first and second observing runs of Advanced LIGO and Advanced Virgo, [arXiv:1912.11716](https://arxiv.org/abs/1912.11716).
- [48] S. A. Usman *et al.*, The PyCBC search for gravitational waves from compact binary coalescence, *Classical Quantum Gravity* **33**, 215004 (2016).
- [49] A. H. Nitz, I. W. Harry, J. L. Willis, C. M. Biwer, D. A. Brown, L. P. Pekowsky, T. Dal Canton, A. R. Williamson, T. Dent, C. D. Capano, T. J. Massinger, A. K. Lenon, A. B. Nielsen, and M. Cabero, PyCBC Software, <https://github.com/gwastro/pycbc> (2018).
- [50] B. Allen, A χ^2 time-frequency discriminator for gravitational wave detection, *Phys. Rev. D* **71**, 062001 (2005).
- [51] A. Harvey Nitz, Distinguishing short duration noise transients in LIGO data to improve the PyCBC search for gravitational waves from high mass binary black hole mergers, *Classical Quantum Gravity* **35**, 035016 (2018).
- [52] A. H. Nitz, T. Dent, T. Dal Canton, S. Fairhurst, and D. A. Brown, Detecting binary compact-object mergers with gravitational waves: Understanding and improving the sensitivity of the PyCBC search, *Astrophys. J.* **849**, 118 (2017).
- [53] G. S. Davies, T. Dent, M. Tápai, I. Harry, C. McIsaac, and A. H. Nitz, Extending the pycbc search for gravitational waves from compact binary mergers to a global network, *Phys. Rev. D* **102**, 022004 (2020).
- [54] S. Mozzon, L. K. Nuttall, A. Lundgren, S. Kumar, A. H. Nitz, and T. Dent, Dynamic normalization for compact binary coalescence searches in non-stationary noise, *Classical Quantum Gravity* **37**, 215014 (2020).
- [55] C. Messick *et al.*, Analysis framework for the prompt discovery of compact binary mergers in gravitational-wave data, *Phys. Rev. D* **95**, 042001 (2017).
- [56] I. W. Harry, B. Allen, and B. S. Sathyaprakash, A stochastic template placement algorithm for gravitational wave data analysis, *Phys. Rev. D* **80**, 104014 (2009).
- [57] Y. Pan, A. Buonanno, L. T. Buchman, T. Chu, L. E. Kidder, H. P. Pfeiffer, and M. A. Scheel, Effective-one-body waveforms calibrated to numerical relativity simulations: Coalescence of non-precessing, spinning, equal-mass black holes, *Phys. Rev. D* **81**, 084041 (2010).
- [58] G. Pratten *et al.*, Let's twist again: Computationally efficient models for the dominant and sub-dominant harmonic modes of precessing binary black holes, [arXiv:2004.06503](https://arxiv.org/abs/2004.06503).
- [59] S. Khan, S. Husa, M. Hannam, F. Ohme, M. Prerer, X. Jimnez Forteza, and A. Boh, Frequency-domain gravitational waves from non-precessing black-hole binaries. II. A phenomenological model for the advanced detector era, *Phys. Rev. D* **93**, 044007 (2016).
- [60] T. Chiba and S. Yokoyama, Spin distribution of primordial black holes, *Prog. Theor. Exp. Phys.* **2017**, 083E01 (2017).
- [61] V. De Luca, V. Desjacques, G. Franciolini, A. Malhotra, and A. Riotto, The initial spin probability distribution of primordial black holes, *J. Cosmol. Astropart. Phys.* **05** (2019) 018.
- [62] V. De Luca, G. Franciolini, P. Pani, and A. Riotto, The evolution of primordial black holes and their final observable spins, *J. Cosmol. Astropart. Phys.* **04** (2020) 052.
- [63] M. Mirbabayi, A. Gruzinov, and J. Norea, Spin of primordial black holes, *J. Cosmol. Astropart. Phys.* **03** (2020) 017.
- [64] E. Baird, S. Fairhurst, M. Hannam, and P. Murphy, Degeneracy between mass and spin in black-hole-binary waveforms, *Phys. Rev. D* **87**, 024035 (2013).
- [65] D. A. Brown, A. Lundgren, and R. O'Shaughnessy, Non-spinning searches for spinning binaries in ground-based detector data: Amplitude and mismatch predictions in the constant precession cone approximation, *Phys. Rev. D* **86**, 064020 (2012).
- [66] A. H. Nitz, A. Lundgren, D. A. Brown, E. Ochsner, D. Keppel, and I. W. Harry, Accuracy of gravitational waveform models for observing neutron-star-black-hole binaries in Advanced LIGO, *Phys. Rev. D* **88**, 124039 (2013).
- [67] K. Postnov, A. Kuranov, and N. Mitichkin, Spins of black holes in coalescing compact binaries, *Phys. Usp.* **62**, 1153 (2019).
- [68] N. E. M. Rifat, S. E. Field, G. Khanna, and V. Varma, Surrogate model for gravitational wave signals from comparable and large-mass-ratio black hole binaries, *Phys. Rev. D* **101**, 081502 (2020).
- [69] I. Harry, J. Calderon Bustillo, and A. Nitz, Searching for the full symphony of black hole binary mergers, *Phys. Rev. D* **97**, 023004 (2018).
- [70] Y. Pan, A. Buonanno, M. Boyle, L. T. Buchman, L. E. Kidder, H. P. Pfeiffer, and M. A. Scheel, Inspiral-merger-ringdown multipolar waveforms of nonspinning black-hole

- binaries using the effective-one-body formalism, *Phys. Rev. D* **84**, 124052 (2011).
- [71] L. London, S. Khan, E. Fauchon-Jones, C. Garca, M. Hannam, S. Husa, X. Jimnez-Forteza, C. Kalaghatgi, F. Ohme, and F. Pannarale, First Higher-Multipole Model of Gravitational Waves from Spinning and Coalescing Black-Hole Binaries, *Phys. Rev. Lett.* **120**, 161102 (2018).
- [72] R. Cotesta, S. Marsat, and M. Prrer, Frequency domain reduced order model of aligned-spin effective-one-body waveforms with higher-order modes, *Phys. Rev. D* **101**, 124040 (2020).
- [73] <https://github.com/gwastro/stellar-pbh-search>
- [74] R. Biswas, P. R. Brady, J. D. E. Creighton, and S. Fairhurst, The loudest event statistic: General formulation, properties and applications, *Classical Quantum Gravity* **26**, 175009 (2009).
- [75] H. Mouri and Y. Taniguchi, Runaway merging of black holes: Analytical constraint on the timescale, *Astrophys. J. Lett.* **566**, L17 (2002).
- [76] D. Nelson *et al.*, The IllustrisTNG Simulations: Public Data Release, [arXiv:1812.05609](https://arxiv.org/abs/1812.05609).
- [77] J. F. Navarro, C. S. Frenk, and S. D. M. White, The Structure of cold dark matter halos, *Astrophys. J.* **462**, 563 (1996).
- [78] A. Olejak, K. Belczynski, T. Bulik, and M. Sobolewska, Synthetic catalog of black holes in the Milky Way, *Astron. Astrophys.* **638**, A94 (2020).
- [79] Y. Ali-Haïmoud, E. D. Kovetz, and M. Kamionkowski, The merger rate of primordial-black-hole binaries, *Phys. Rev. D* **96**, 123523 (2017).
- [80] Z.-C. Chen and Q.-G. Huang, Distinguishing primordial black holes from astrophysical black holes by Einstein telescope and cosmic explorer, *J. Cosmol. Astropart. Phys.* **08** (2020) 039.
- [81] K. Jedamzik, Evidence for primordial black hole dark matter from LIGO/Virgo merger rates, [arXiv:2007.03565](https://arxiv.org/abs/2007.03565).
- [82] T. Bringmann, P. Frederik Depta, V. Domcke, and K. Schmidt-Hoberg, Towards closing the window of primordial black holes as dark matter: The case of large clustering, *Phys. Rev. D* **99**, 063532 (2019).
- [83] G. Ballesteros, P. D. Serpico, and M. Taoso, On the merger rate of primordial black holes: Effects of nearest neighbours distribution and clustering, *J. Cosmol. Astropart. Phys.* **10** (2018) 043.
- [84] T. Nakama, J. Silk, and M. Kamionkowski, Stochastic gravitational waves associated with the formation of primordial black holes, *Phys. Rev. D* **95**, 043511 (2017).
- [85] R.-G. Cai, S. Pi, S.-J. Wang, and X.-Y. Yang, Pulsar timing array constraints on the induced gravitational waves, *J. Cosmol. Astropart. Phys.* **10** (2019) 059.
- [86] B. P. Abbott *et al.* (Virgo, KAGRA, and LIGO Scientific Collaborations), Prospects for observing and localizing gravitational-wave transients with Advanced LIGO, Advanced Virgo and KAGRA, *Living Rev. Relativity* **21**, 3 (2018).
- [87] M. Punturo *et al.*, The einstein telescope: A third-generation gravitational wave observatory, *Classical Quantum Gravity* **27**, 194002 (2010).
- [88] D. Reitze *et al.*, Cosmic explorer: The U.S. contribution to gravitational-wave astronomy beyond LIGO, *Bull. Am. Astron. Soc.* **51**, 035 (2019), <https://baas.aas.org/pub/2020n7i035/release/1>.
- [89] A. M. Green and B. J. Kavanagh, Primordial Black Holes as a dark matter candidate, [arXiv:2007.10722](https://arxiv.org/abs/2007.10722).
- [90] <https://www.gw-openscience.org>.