

CLINICAL IMAGE

Gastric neuroendocrine tumor as a rare type of neoplasm in a kidney transplant recipient

Paulina Gołasa¹, Marcin Krzanowski¹, Przemysław Miarka¹,
Krzysztof Okoń², Marek Kuźniewski¹, Katarzyna Krzanowska¹

¹ Department of Nephrology, Jagiellonian University Medical College, Kraków, Poland

² Department of Clinical and Experimental Pathomorphology, Jagiellonian University Medical College, Kraków, Poland

Neuroendocrine neoplasms are rare tumors (2.89 to 6.98 cases per 100 000 people per year in the United States)^{1,2} originating from neuroendocrine cells (neural crest, neuroendocrine islets, and stem cells), with a rising incidence during several past decades. This type of tumors can be benign or malignant and may develop in various organs, particularly in the gastrointestinal and pulmonary systems.

One of their subtypes, gastric neuroendocrine tumors, account for 1 to 2 cases per 1 million persons per year and represent 8.7% of all gastrointestinal neuroendocrine tumors.^{3,4}

They originate from enterochromaffin-like cells of the gastric mucosa or, less commonly, from other types of endocrine cells (that secrete serotonin, gastrin, or adrenocorticotropic hormone).⁴ Patients may develop symptoms of uncontrolled hormone secretion or caused by tumor invasion.

Prognosis and survival depend on the type of tumor. There is a classification distinguishing 4 tumor types, based on their histological and morphological characteristics and pathogenesis.⁴ Well-differentiated type I and II gastric neuroendocrine neoplasms are mostly benign

tumors posing a low risk of malignancy, whereas type III neoplasm has a poor prognosis, and type IV neoplasm—an extremely poor prognosis with a mean survival time of several months.

We present a case of a 65-year-old female kidney transplant recipient treated with triple immunosuppressive therapy (prednisolone, cyclosporine, and mycophenolate mofetil), with end-stage renal disease of her own kidneys due to diabetic nephropathy, with hypertension, and a history of cytomegalovirus infection, who was incidentally diagnosed with asymptomatic neuroendocrine tumor of the gastric antrum 25 months after transplant. Routine endoscopy was performed and showed the granular, erythematous pattern of the mucous membrane in the antrum (**FIGURE 1A** and **1B**). Features of low-grade neuroendocrine tumor were found on histopathological examination, with a mitotic index below 1 mitosis / 10 high-power fields, and 70% of the section volume was involved, with infiltrated margins of the specimen slice. Additionally, traits of chronic atrophic gastritis with intestinal metaplasia were detected (**FIGURE 1C** and **1D**). The patient denied having any symptoms.




FIGURE 1 A, B – gastroscopy showing the granular, erythematous mucous membrane of the antrum

Correspondence to:

Katarzyna Krzanowska, MD, PhD,
Department of Nephrology,
Jagiellonian University Medical
College, ul. Jakubowskiego 2,
30-688 Kraków, Poland, phone:
+48 12 400 28 62, email:
kasiajanda@op.pl

Received: May 5, 2020.

Revision accepted: May 25, 2020.

Published online: June 3, 2020.

Pol Arch Intern Med. 2020;

130 (9): 801-802

doi:10.20452/pamw.15413

Copyright by the Author(s), 2020




FIGURE 1 C – hematoxylin-eosin staining, magnification $\times 400$: tumor cells forming cohesive nests (arrows). The nuclei are round or oval in shape and rather uniform, with finely dispersed chromatin. D – hematoxylin-eosin staining, magnification $\times 400$: the lamina propria infiltrated by rather uniform neuroendocrine cells (arrows)

Previous endoscopy (performed 15 months earlier) demonstrated only erosive gastritis in the antrum and polyps in the stomach body.

After establishing the diagnosis, a modification of the immunosuppressive therapy was recommended, ie, a switch from the calcineurin inhibitor to an mTOR inhibitor, but the patient disagreed with the suggestion and the previous therapy was therefore continued.

Concurrently, the patient was referred to the endocrinology outpatient clinic. She underwent 99m technetium sestamibi single-photon emission computed tomography of the abdomen and pelvis. No remarkable tracer concentration or increased tracer uptake were revealed. The case was discussed during a tumor board conference and a decision was made to closely monitor the patient.

Recent gastroscopy, performed 6 months after endoscopic tumor removal, showed signs of inactive chronic gastritis with foveolar hyperplasia in the antrum. No other abnormalities were detected.

This case is an interesting example of a very rare neoplasm developing in transplant recipients who are at increased risk of malignancies, but typically develop skin and renal malignancies or lymphomas. There are several cases of post-transplant neuroendocrine neoplasms, including mainly Merkel cell carcinoma or a single case of rectal neuroendocrine neoplasm.⁵ The presented case indicates the importance of accurate cancer screening in patients after organ transplants, as early detection makes it possible to implement proper treatment quickly—in this case, it was endoscopic tumor removal. In our clinical center, kidney transplant recipients have obligatory gastroscopy performed once a year and colonoscopy every 2 years or once a year in those with colon polyps. Close follow-up after cancer diagnosis is also crucial. However, the appropriate management of neuroendocrine neoplasms remains a challenge, as there are only scarce data available on this issue.

ARTICLE INFORMATION

CONFLICT OF INTEREST None declared.

OPEN ACCESS This is an Open Access article distributed under the terms of the Creative Commons AttributionNonCommercialShareAlike 4.0 International License (CC BY-NC-SA 4.0), allowing third parties to copy and redistribute the material in any medium or format and to remix, transform, and build upon the material, provided the original work is properly cited, distributed under the same license, and used for noncommercial purposes only. For commercial use, please contact the journal office at pamw@mp.pl.

HOW TO CITE Golasa P, Krzanowski M, Miarka P, et al. Gastric neuroendocrine tumor as a rare type of neoplasm in a kidney transplant recipient. *Pol Arch Intern Med.* 2020; 130: 801-802. doi:10.20452/pamw.15413

REFERENCES

- 1 Patel N, Benipal B. Incidence of neuroendocrine tumors in the United States from 2001-2015: a United States cancer statistics analysis of 50 states. *Cureus.* 2019; 11: e4322. [↗](#)
- 2 Wang R, Zheng-Pywell R, Chen HA, et al. Management of gastrointestinal neuroendocrine tumors. *Clin Med Insights Endocrinol Diabetes.* 2019; 12: 1-12. [↗](#)
- 3 Dias AR, Azevedo BC, Alban LBV, et al. Gastric neuroendocrine tumor: review and update. *Arq Bras Cir Dig.* 2017; 30: 150-154. [↗](#)
- 4 Li TT, Qiu F, Qian ZR, et al. Classification, clinicopathologic features and treatment of gastric neuroendocrine tumors. *World J Gastroenterol.* 2014; 20: 118-125. [↗](#)
- 5 Karunanithi S, Roy SG, Sharma P, et al. Metastatic neuroendocrine tumour in a renal transplant recipient: Dual-Tracer PET-CT with (18)F-FDG and (68)Ga-DOTANOC in this rare setting. *Nucl Med Mol Imaging* 2015; 49: 57-60. [↗](#)