

2-2021

Validation and Accuracy of Synovial Alpha Defensin in a Suburban Academic Center after Implantation as Part of the Workup for Periprosthetic Joint infections

Donald Willier, III

Kenneth J. McAlpine, MD

Anthony F. De Giacomo, MD

John S. Garfi, MS

Lawrence M. Specht, MD

See next page for additional authors

Follow this and additional works at: https://jdc.jefferson.edu/si_ctr_2023_phase1



Part of the [Translational Medical Research Commons](#)

[Let us know how access to this document benefits you](#)

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Phase 1 by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: JeffersonDigitalCommons@jefferson.edu.

Authors

Donald Willier, III; Kenneth J. McAlpine, MD; Anthony F. De Giacomo, MD; John S. Garfi, MS; Lawrence M. Specht, MD; and Michael S. Kain, MD



**Sidney Kimmel
Medical College™**
at Thomas Jefferson University

Validation and Accuracy of Synovial Alpha Defensin in a Suburban Academic Center after Implantation as Part of the Workup for Periprosthetic Joint Infections

Donald P. Willier III BS, Kenneth J. McAlpine MD, Anthony F. De Giacomo MD, John S. Garfi MS, Lawrence M. Specht MD, Michael S. Kain MD*

- The authors have no conflicts of interest in regards to this research study



- Periprosthetic joint infections increase morbidity and mortality after hip and knee arthroplasty^{1,2,3,4}
 - Revision arthroplasty and failure post revision
- Emergent problem that requires a clinical diagnosis
 - Introduction of MSIS criteria in 2011⁵



- Increasing development of biomarkers as diagnostic tests
 - Synovial fluid alpha-defensin
 - Incorporated into 2018 MSIS update⁶
- Prior literature has shown up to 100% for specificity and sensitivity^{7,8,9,10}
- Synovial alpha-defensin biomarker incorporated into our practice in 2014

Objectives & Hypothesis

- The purpose of this study was to evaluate our implementation of synovial alpha-defensin (Synovasure®) with comparison to the 2011 MSIS criteria for diagnosing periprosthetic joint infections after hip and knee arthroplasty
- We hypothesized that the sensitivity and specificity of the synovial alpha-defensin immunoassay will both be 100%

Approach & Results

- Single center, retrospective chart from 2014 – 2016
- 128 consecutive that came in for evaluation of potential hip / knee periprosthetic joint infection (PJI)
- All patients must have received synovial alpha-defensin immunoassay in addition to our standard PJI workup
- Data was collected from electronic medical records after IRB approval
- Rationale - Evaluate the potential diagnostic capability of synovial-alpha defensin

Approach & Results

- 28 patients had a positive synovial alpha defensin test vs. 100 patients with a negative test
 - There were no significant differences in demographics between groups

	Positive	Negative	P-Value
Age	69.5	67.2	0.381
Sex (M/F)	18/10	48/52	0.140
Hip / Knee	10/18	52/48	0.127
Tobacco Use	4	12	0.750
Diabetes	8	23	0.619

Approach & Results

- Results of our standard PJI workup:

	Positive Synovasure [®]	Negative Synovasure [®]
CRP Serum	112.4	20.9
Synovial WBC	42,723.5	739.3
Synovial PMN	87.2%	41.2%
Positive Culture	5 (17.9%)	7 (7%)
Gram Stain	8 (28.6%)	0
Crystals	9 (32.1%)	0

Approach & Results

- In comparison to 2011 MSIS as gold standard

	Value	95% Confidence Interval
Sensitivity	95.00%	75.13 – 99.87%
Specificity	91.67%	84.77 – 96.12%
Positive Predictive Value	67.86%	52.84 – 79.91%
Negative Predictive Value	99.00%	93.61 - 99.85%
Accuracy	92.19%	86.10 – 96.19%

Conclusions

- Our results demonstrated that there is a role for synovial-alpha defensin immunoassay in PJI workup
- However there was a low positive predictive value and a failure to achieve 100% in either sensitivity or specificity
 - Therefore it cannot be used as a standalone test
- Our findings support more recent literature of incorporating biomarkers into the clinical picture
 - 2018 MSIS criteria
- Analysis of our results has kept Synovasure® as part of our standard PJI workup

Future Directions

- Since the implementation of this study, potential sample size has grown
 - Possibility of re-evaluating with most up-to-date population
 - Possibility of utilizing the 2018 MSIS criteria for comparison

Acknowledgements

- Thank you to Drs. Kain, McAlpine, Specht and De Giacomo for bringing me onto this project in 2017
- Thank you to John Garfi for guiding the statistical analysis



References

1. Kamath AF, Ong KL, Lau E, Chan V, Vail TP, Rubash HE, et al. Quantifying the Burden of Revision Total Joint Arthroplasty for Periprosthetic Infection. *J Arthroplasty* 2015;30:1492–7. <https://doi.org/10.1016/j.arth.2015.03.035>.
2. Bozic KJ, Kurtz SM, Lau E, Ong K, Vail TP, Berry DJ. The epidemiology of revision total hip arthroplasty in the United States. *J Bone Joint Surg Am* 2009;91:128–33. <https://doi.org/10.2106/JBJS.H.00155>.
3. Jafari SM, Coyle C, Mortazavi SMJ, Sharkey PF, Parvizi J. Revision hip arthroplasty: infection is the most common cause of failure. *Clin Orthop Relat Res* 2010;468:2046–51. <https://doi.org/10.1007/s11999-010-1251-6>.
4. American Joint Replacement Registry (AJRR): 2019 Annual Report. Rosemont, IL: American Academy of Orthopaedic Surgeons (AAOS), 2019.
5. Parvizi J, Zmistowski B, Berbari EF, Bauer TW, Springer BD, Della Valle CJ, et al. New definition for periprosthetic joint infection: from the Workgroup of the Musculoskeletal Infection Society. *Clin Orthop Relat Res* 2011;469:2992–4. <https://doi.org/10.1007/s11999-011-2102-9>.
6. Parvizi J, Tan TL, Goswami K, Higuera C, Della Valle C, Chen AF, et al. The 2018 Definition of Periprosthetic Hip and Knee Infection: An Evidence-Based and Validated Criteria. *J Arthroplasty* 2018;33:1309-1314.e2. <https://doi.org/10.1016/j.arth.2018.02.078>.
7. Deirmengian C, Kardos K, Kilmartin P, Cameron A, Schiller K, Booth RE, et al. The alpha-defensin test for periprosthetic joint infection outperforms the leukocyte esterase test strip. *Clin Orthop Relat Res* 2015;473:198–203. <https://doi.org/10.1007/s11999-014-3722-7>.
8. Deirmengian C, Kardos K, Kilmartin P, Cameron A, Schiller K, Parvizi J. Diagnosing periprosthetic joint infection: has the era of the biomarker arrived? *Clin Orthop Relat Res* 2014;472:3254–62. <https://doi.org/10.1007/s11999-014-3543-8>.
9. Bingham J, Clarke H, Spangehl M, Schwartz A, Beauchamp C, Goldberg B. The alpha defensin-1 biomarker assay can be used to evaluate the potentially infected total joint arthroplasty. *Clin Orthop Relat Res* 2014;472:4006–9. <https://doi.org/10.1007/s11999-014-3900-7>.
10. Frangiamore SJ, Gajewski ND, Saleh A, Farias-Kovac M, Barsoum WK, Higuera CA. α -Defensin Accuracy to Diagnose Periprosthetic Joint Infection-Best Available Test? *J Arthroplasty* 2016;31:456–60. <https://doi.org/10.1016/j.arth.2015.09.035>.