brought to you by **CORE**

Aquaculture 536 (2021) 736510

Contents lists available at ScienceDirect

Aquaculture



journal homepage: www.elsevier.com/locate/aquaculture





Corrigendum to 'Dietary aflatoxin B1 (AFB1) reduces growth performance, impacting growth axis, metabolism, and tissue integrity in juvenile gilthead sea bream (*Sparus aurata*)'. Aquaculture, volume 533, 25 February 2021, 736189

A. Barany^{a,*}, M. Guilloto^a, J. Cosano^a, M. de Boevre^b, M. Oliva^a, S. de Saeger^b, J. Fuentes^c, G. Martínez-Rodriguez^d, J.M. Mancera^a

^a Department of Biology, Faculty of Marine and Environmental Sciences, Instituto Universitario de Investigación Marina (INMAR), Campus de Excelencia Internacional del Mar (CEI-MAR), University of Cádiz, 11510 Puerto Real, Cádiz, Spain

^b Centre of Excellence in Mycotoxicology & Public Health, MYTOX-SOUTH, Department of Bioanalysis, Ghent University, Ottergemsesteenweg 460, Ghent, Belgium

^c Centre of Marine Sciences (CCMar), Universidade do Algarve, Campus de Gambelas, 8005-139 Faro, Portugal

^d Instituto de Ciencias Marinas de Andalucía, Consejo Superior de Investigaciones Científicas (ICMAN-CSIC), E-11510 Puerto Real, Cádiz, Spain

The authors regret the errors in a few table references within the text. Specifically, it should reads as follows within the following subsections/ page:

3.2. Blood analysis (page 5)

Hematocrit showed a significant reduction in both groups fed with AFB1 (D1 and D2) compared to the CT group, F(2,24) = 6.5, P = 0.0054 (Table 3). No significant differences were detected in RBC, WBC, or SI (Table 3 and Table 4). Glucose and triglyceride levels were significantly lower in both groups fed with AFB1 (D1 and D2) compared to the CT group, F (2,22) = 8.6, P = 0.0017; F(2,22) = 48.6, P < 0.0001, respectively (Table 5). No significant differences were detected among experimental groups in neither lactate (Table 5) nor cortisol levels

(Fig. 2).

3.3. Liver (page 5)

HSI showed a significant reduction in both groups fed with AFB1 (D1 and D2) compared to the CT group, F(2,21) = 6.4, P = 0.0066 (Table 4).

In addition, gene nomenclature throughout the text should read: *eef1a* instead of *ef1a*.

pomca1 (proopiomelanocortin a1) instead of *pomca* (*proopiomelanocortin a*).

pomca2 (proopiomelanocortin a2) instead of pomcb (proopiomelanocortin b).

The authors would like to apologize for any inconvenience caused.

DOI of original article: https://doi.org/10.1016/j.aquaculture.2020.736189. * Corresponding author.

E-mail address: andre.barany@uca.es (A. Barany).

https://doi.org/10.1016/j.aquaculture.2021.736510

Available online 15 February 2021 0044-8486/© 2021 Elsevier B.V. All rights reserved.