

## Research Space

Journal article

**A qualitative screening tool to identify athletes with ‘high-risk’ movement mechanics during cutting: The cutting movement assessment score (CMAS)**

**Dos’Santos, T., McBurnie, A., Donelon, T., Thomas, C., Comfort, P. and Jones, P.A.**

<https://doi.org/10.1016/j.ptsp.2019.05.004>

## Manuscript Details

<b>Manuscript number</b>	YPTSP_2019_102_R1
<b>Title</b>	A QUALITATIVE SCREENING TOOL TO IDENTIFY ATHLETES WITH 'HIGH-RISK' MOVEMENT MECHANICS DURING CUTTING: THE CUTTING MOVEMENT ASSESSMENT SCORE (CMAS)
<b>Article type</b>	Research Paper

### Abstract

Objective: To assess the validity of the cutting movement assessment score (CMAS) to estimate the magnitude of peak knee abduction moments (KAM) against three-dimensional (3D) motion analysis, while comparing whole-body kinetics and kinematics between subjects of low (bottom 33%) and high CMASs (top 33%). Design: Cross-sectional study. Setting: Laboratory. Participants: Forty-one participants (soccer, rugby, netball, and cricket). Main outcome measures: Association between peak KAM and CMAS during a 90° cut. Comparison of 3D whole-body kinetics and kinematics between subjects with low (bottom 33%) and high CMASs (top 33%). Results: A very large significant relationship ( $p = 0.796$ ,  $p < 0.001$ ) between CMAS and peak KAM was observed. Subjects with higher CMASs displayed higher-risk cutting postures, including greater peak knee abduction angles, internal foot progression angles, and lateral foot plant distances ( $p \leq 0.032$ , effect size = 0.83-1.64). Additionally, greater cutting multiplanar knee joint loads (knee flexion, internal rotation, and abduction moments) were demonstrated by subjects with higher CMASs compared to lower ( $p \leq 0.047$ , effect size = 0.77-2.24). Conclusion: The CMAS is a valid qualitative screening tool for evaluating cutting movement quality and is therefore a potential method to identify athletes who generate high KAMs and "high-risk" side-step cutting mechanics.

<b>Keywords</b>	Anterior cruciate ligament; knee abduction moment; injury screening; injury-risk profile
<b>Corresponding Author</b>	Thomas Dos'Santos
<b>Order of Authors</b>	Thomas Dos'Santos, Alistair McBurnie, Thomas Donelon, Christopher Thomas, Paul Comfort, Paul Jones
<b>Suggested reviewers</b>	Anne Benjaminse, Gillian Weir

## Submission Files Included in this PDF

### File Name [File Type]

Cover Letter.docx [Cover Letter]

Comments from the editors and reviewers v3.docx [Response to Reviewers (without Author Details)]

Highlights-r1.docx [Highlights]

title pagev2.docx [Title Page (with Author Details)]

CMAS PTIS - FINAL- R1.docx [Manuscript (without Author Details)]

Conflict of Interest.docx [Conflict of Interest]

Ethical Statement.docx [Ethical Statement]

Supplement 1 - CMAS operational defsv3.pdf [e-Component]

Supplement 2 - Variables.docx [e-Component]

Supplement 3 - CMAS MANUALv4.pdf [e-Component]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.

## Research Data Related to this Submission

There are no linked research data sets for this submission. The following reason is given:  
Data will be made available on request

1  
2  
3 This manuscript is original and not previously published, nor is it being considered elsewhere  
4 until a decision is made as to its acceptability by the Physical Therapy in Sport Journal  
5 Review Board. All authors have made substantial contributions to all of the following: (1) the  
6 conception and design of the study, or acquisition of data, or analysis and interpretation of  
7 data, (2) drafting the article or revising it critically for important intellectual content, (3) final  
8 approval of the version to be submitted.  
9

10  
11 Thomas Dos'Santos

12  
13 Directorate of Sport, Exercise and Physiotherapy. University of Salford, Salford, Greater  
14 Manchester, UK  
15

16  
17 t.dossantos@edu.salford.ac.uk

18  
19 +44 7961744517  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59

1  
2  
3  
4 **-Reviewer 1**

5 **General:**

6 Thank you for reviewing this interesting manuscript. Overall, the development of the CMAS is very exciting.  
7 However, the test validity should be placed in its context and limitations (test is not executed in the field  
8 where complex interaction between athlete and environment plays a role) should be acknowledged. The  
9 authors have many places (e.g. lines 128, 198-199, 233, 316, 317, 1080, 1081, 1119, 1120) where many  
10 references are mentioned. Please limit the numbers of references.  
11

12 **Response:** Thank you for your comment. We have amended the number of citations in text throughout the  
13 manuscript, following your suggestions. Additionally, we have acknowledged the multifactorial and complex  
14 interaction of several factors regarding ACL injury. Consequently, paragraph 1 of the introduction has been  
15 amended.

16 **Introduction:**

17 Line 134-135: 'biomechanical and neuromuscular control deficits are critical factors that affect knee joint  
18 loading' > are these deficits causes or effects??  
19

20 **Response:** Thank you for your comment. We have amended this paragraph, and have subsequently removed  
21 this sentence.  
22  
23

24 Line 139-155: The authors forget to put ACL injury risk in a multifactorial perspective. The end-point of  
25 movement is an hazardous movement strategy with high knee joint load. The abnormal deficits mentioned are  
26 measured in lab situations, do athletes also show this in the field ? How about an environmental context, the  
27 anticipation strategies used by an athlete used milliseconds before getting injured ? Of course, evaluating  
28 movement quality is important, but please put this in a broader perspective before zooming in.  
29

30 **Response:** Thank you for your comment. We have now acknowledged that ACL injury risk factors are  
31 multifactorial, with a complex interaction of integral and external factors (i.e. anatomical, hormonal,  
32 environmental, shoe-surface interface, anticipation, and fatigue) during ACL injury inciting events. However,  
33 hazardous knee joint loads which are amplified with poor initial postures and mechanics, ultimately results in  
34 ACL rupture from a mechanism perspective. Yes, we agree anticipatory strategies, environment, fatigue, shoe-  
35 surface interface will also interact, but these biomechanical deficits are importantly modifiable, with the  
36 objective of qualitative screening tools to identify athletes who display these potentially 'high-risk' deficits, to  
37 subsequently enable individualised injury-risk mitigation training programmes to be developed. Therefore, this  
38 is why have concentrated focusing on the biomechanics and movement quality, because this is ultimately what  
39 the practitioner can modify.

40 We now state the following: "Although ACL injury-risk factors are multifactorial (74) and a complex interaction  
41 of internal and external factors (i.e. anatomical, hormonal, environmental, shoe-surface interface, anticipation,  
42 and fatigue) (8, 34, 54), a large proportion of ACL injuries are non-contact in nature during high velocity and  
43 impact sporting tasks, such as side-stepping (9, 39, 54). This occurrence can be attributed to the tendency to  
44 generate large multiplanar knee joint loading, such as knee abduction moments (KAM) and internal rotation  
45 moments (KIRM) (7, 19, 41), which increase ACL strain (4, 57, 76). These potentially hazardous knee joint loads  
46 are amplified when poor initial postures and movement is demonstrated (biomechanical and neuromuscular  
47 control deficits) during cutting (24, 34, 65), but importantly these deficits are modifiable (34, 71)."  
48

49 Line 158-174: Please limit wording here (shortening section) + try to shorten introduction in general.

50 **Response:** Thank you for your comment. We have removed the list of methodological issues: "(e.g. markers  
51 sets, marker placement, soft tissue artefact, body segment inertial parameters, modelling, reference frame,  
52 normalisation, low-pass filtering cut-frequencies)"  
53

54 We have done our best to be concise, but feel much of the content must remain in the introduction to help set  
55 the scene, gaps in the literature, and rationale for the study. Reviewer 2 has also specifically commented "Very  
56 nice and thorough introduction."  
57  
58  
59

60  
61  
62 Line 174-185: Not sure what the purpose is of this section, as the CMAS will neither be able to do this.  
63

64 Response: Thank you for your comment. We have removed the statements referring to ability to predict ACL  
65 injury. We have removed the following: "While the LESS has been validated against 3D motion analysis (65,  
66 69), its ability to predict ACL injury is questionable with conflicting observations found (67, 76). Similarly,  
67 although the TJA and QASLS demonstrate good face validity (28, 61), there is no empirical evidence to confirm  
68 these tools can identify athletes that display high knee joint loading or ability to predict non-contact ACL injury  
69 (23, 24)."

70 We hope this resolves this.  
71

72 Line 203: I would certainly not state this 'ecologically more valid'. Ecological validity means taking care of the  
73 interaction between person-task-environment. Even though addressing other, interesting aspects of  
74 movement in the light of ACL injury, the sidestep cutting technique as examined with the CMAS is still an  
75 isolated, preplanned movement. Please rewrite.

76 Response: Thank you for your comment. We have removed the term "ecologically more valid" and refer to  
77 specificity. The aim of paragraph 4 is to highlight the importance of specifically screening cutting mechanics,  
78 compared to inferring cutting mechanics from a jump-landing assessment.  
79

#### 80 81 **Methods:**

82 Line 446-447: Has this threshold been validated against visual observation?  
83

84 Response: Thank you for your comment. Yes, this threshold was used to account for the residual noise  
85 associated with the AMTI force plates in the lab and has been confirmed against visual observation in our lab.  
86 These vertical GRF thresholds have also been used by numerous research groups including Kristianslund et al.  
87 (Oslo research group), Robinson et al. (LJMU research group), and King et al. (Sports Surgery Clinic, Dublin).  
88

89 Line 504-505: How about the high-speed camera position and exit run? Was the exit run shorter than 3m. ?

90 Response: Thank you for your comment. As illustrated in Figure 1, the sagittal plane camera was position right  
91 of the timing gates, perpendicular to the PFC. Thus, athletes performed a 90° cut and exited 3-m from the final  
92 force plate.  
93

94 Line 593: Please write confidence instead of Confidence.

95 Response: Thank you for your comment. This has been amended to lower case.  
96

#### 97 **Discussion:**

98 Line 982-983: Please rewrite 'high CMAS' and 'low CMAS'.  
99

100 Response: Thank you for your comment. Apologies, but we do not fully understand the issue here? We have  
101 attempted to amend the sentence. We hope this resolves this, but any further clarification would be great.  
102

103 Line 993-1004: A large portion of this is repetition of what has been mentioned in the introduction. Please  
104 shorten/delete.

105 Response: Thank you for your comment. We have deleted the following: "Side-step cutting has been identified  
106 as a key action associated with non-contact ACL injury in multidirectional sports (6, 8, 11, 19, 37, 44, 46, 60, 65,  
107 79) due to the tendency to generate large multiplanar knee joint loading (KAMs and KIRMs) (5, 16, 39), which  
108 increases ACL strain (4, 55, 73). These potentially hazardous knee joint loads are amplified when  
109 biomechanical and neuromuscular control deficits (high-risk deficits), such as wide lateral foot plants (16, 27,  
110 38, 48), internally rotated foot postures (initial posture) (40, 75), internally rotated hip initial postures (27, 57,  
111 74, 75), knee abduction angles and motion (38, 40, 48, 57, 74), lateral trunk flexion (16, 26, 36, 38), and  
112 extended knee postures (13, 22, 46, 80) are demonstrated during side-step cutting."  
113

114 We hope this resolves this.  
115  
116  
117  
118

119  
120  
121 Line 1033-1038: Please rewrite and eliminate the 'hint' as if CMAS will be able to predict non-contact ACL  
122 injury.  
123

124  
125 **Response:** Thank you for your comment. We have removed the sentences relating to predicting ACL injury, but  
126 instead we focus on the drawbacks of evaluating landing mechanics to infer cutting mechanics, and we discuss  
127 that the LESS is the only other tool to be validated against 3D.  
128

129 Line 1071: Please add 'total' before 'CMAS'.

130  
131 **Response:** Thank you for your comment. We have added 'total'.  
132

133 Line 1119-1134: Do you really think the CMAS will be able to predict ACL injury ?? Is this the way such tests  
134 should be used at all ? I would be very careful with stating this. There is so many factors playing a rol in getting  
135 or not getting an ACL injury (in being at increased risk), which makes it too simple to say that one test could  
136 predict an injury. This section should be rewritten and the CMAS should but put in to a right perspective.  
137

138  
139 **Response:** Thank you for your comment. As stated earlier, we have removed the sentences relating to  
140 predicting ACL injury and the requirements to identify an "at-risk" threshold. We appreciate that ACL injury is  
141 multifactorial and influenced by both internal and external factors. The CMAS is simply a tool to assist in  
142 identification of poor movement quality cutting, and we appreciate that biomechanics and movement quality  
143 is only a contributing factor to a complex multifactorial mechanism of injury. However, with respect, athletes  
144 who demonstrate aberrant movement mechanics may have a potentially greater relative risk of sustaining an  
145 ACL injury, but we never state they will definitely sustain an injury.  
146

147 Line 1139: Isn't pre-rotation good for knee joint load (Dempsey MSSE 2007)?

148  
149 **Response:** Thank you for your comment. Our point regarding pre-rotation was regarding the potential  
150 difficulties in viewing lower-limb and trunk alignment because the athlete will not be perpendicular to the  
151 camera (i.e. parallax error). Research has shown that greater rotation of the foot (internal foot progression  
152 angles) increases peak KAMs (Sigward et al., 2007).  
153

154 Line 1141-1148: Please reword as I am not sure I understand this sentence.

155  
156 **Response:** Thank you for your comment. We have amended this. It now reads as follows: "It should be  
157 acknowledged that, due to the multiplanar nature of side-step cutting (7), some athletes pre-rotate towards  
158 the direction of travel during weight acceptance of the cut (77). This pre-rotation can potentially result in  
159 parallax error because the athlete is not perpendicular to the cameras which can restrict evaluations of  
160 particular CMAS criteria using the frontal plane and 45° cameras."  
161

162  
163 **-Reviewer 2**

#### 164 165 **General**

166 This is a really nice study and certainly something that can be readily utilized by community level athletes and  
167 coaches. I have a few minor comments below but my main curiosity was how your ran your statistics and I  
168 wonder if you can be more specific in your methods about how you accounted for subject variability in your  
169 correlations. See my specific comments below. It seems from what you describe in your methods and present  
170 in your results, that you ran correlations and group differences on the individual trials while not accounting for  
171 inter-subject variability which is a significant statistical flaw. While I definitely understand the importance of not  
172 averaging the trials as mechanically this can wash things out if you have one trial where they had high knee  
173 moments and a higher CMAS score and another trial with low knee moments and a low CMAS score and then  
174 you average the two and get a completely different result. However, there are correct ways of entering the  
175 individual trials into a general linear model for example and then entering the participants as random factors.  
176  
177

178  
179  
180 This needs to be further explained and/or amended in the methods/results section before it acceptable for  
181 publication  
182

183 **Response:** Thank you for your comment. In light of your comment regarding the pooling of trials for correlation  
184 analysis, we have consulted with a statistician and they agree with your point regarding inter-subject variability,  
185 and we have now removed the pooled correlations from the manuscript. Going forward, we have used the  
186 correlation value between the mean CMAS and mean peak KAMs for each subject (see figure 2), and we have  
187 made this clear in the statistical analyses section. Additionally, we have removed tables 3 and 4, and we have  
188 reperformed the analysis for comparisons in 3D mechanics between low and high CMAS. We now compare  
189 cutting mechanics between subjects with low and high CMASs (CMAS: top 33% and bottom 33% using subject  
190 mean data) with the results presented in Table 3. However, the primary findings are similar to the previous  
191 version of the manuscript, with subjects with high CMASs displaying higher-risk mechanics and multiplanar knee  
192 joint loads vs. subjects with lower CMASs. Finally, we did perform a general linear model which factors in CMAS,  
193 participant ID, and sex, and found it was statistically significant, with an R squared value of 0.889, and adjusted  
194 R squared of 0.763. However, in consultation with a statistician, we have decided not to include this in the  
195 manuscript because it violates the assumptions for general linear model testing because it is qualitative data  
196 and is non-parametric, and the pKAM and CMAS data were not symmetric.  
197

### 198 **Specific**

### 199 **Abstract**

200  
201 Page 2, line 87: Place  $p < 0.05$  within the brackets.  
202

203 **Response:** Thank you for your comment. We have added the p value in the abstract. ( $p \leq 0.047$ )  
204

### 205 **Introduction**

206  
207 Very nice and thorough introduction.  
208

209 **Response:** Thank you for your kind words.  
210

211  
212 Page 4, lines 198-202: No need to add this reference – but just for your interest and may add to your point. The  
213 authors compared single leg landing and unplanned sidestepping and had a similar conclusion to yours here.  
214

215 Chinasee, C., Weir, G., Sasimontokul, S., Alderson, J.A., & Donnelly, C.J. (2018). A Biomechanical Comparison of  
216 Single-Leg Landing and Unplanned Sidestepping. *International Journal of Sports Medicine*. 39: 1-10. doi:  
217 <https://doi.org/10.1055/a-0592-7422>  
218

219 **Response:** Thank you for your suggestion. We have added this reference because it further strengthens the  
220 requirement to specifically screen side-step mechanics rather than use a landing task.  
221

222 Page 4, line 221: *The p and d* values in brackets might be slightly misleading here as I read that sentence as  
223 talking about the predictions and not measured vs predicted. Consider rewording or remove the p and d values  
224 in brackets.  
225

226 **Response:** Thank you for your comment. Apologies for this coming across misleading. We have decided to  
227 remove the p and d values in the brackets.  
228

229 Page 4: I wonder whether it makes more sense to make the inter- and intra- rater reliability of your variables  
230 your first hypothesis as that is the order it is presented in your results.  
231

232 **Response:** Thank you for your comment. We have followed your suggestions and amended this section as  
233 follows: "Firstly, it was hypothesised that excellent inter- and intra-rater reliability would be demonstrated for  
234  
235  
236

237  
238  
239 CMAS items. Secondly, in line with Jones et al. (43), it was hypothesised that a strong relationship would be  
240 demonstrated between CMAS and peak KAM, and the CMAS would be able to discriminate between “low” and  
241 “high” CMASs in terms of “high-risk” whole-body kinetics and kinematics.”  
242

## 243 **Methods**

244  
245 Page 6, line 304: Were the cricket players in season at the same time as the winter sports?  
246

247 **Response:** Thank you for your comment. In England, the sports you may be referring to as Winter Sports (i.e.  
248 rugby and soccer) competitive seasons span approximately 9-10 months. Thus, when our testing took place, all  
249 athletes from all sports were still in-season (approx. April-May). We hope this clarifies this.  
250

251 Page 8, line 423-425: State that you measured completion time here and then place the times into the results  
252 section. Did athletes cut at their max speed or a pre-determined velocity? On line 441 you say that pre-contact  
253 velocity was 4.5m/s but it is not clear how you would have measured that as the timing gates are before and  
254 after the force platform. Instead state you measured velocity at foot strike or if you want to measure pre-contact  
255 velocity measure the average center of mass velocity from the penultimate toe off to cutting limb foot contact.  
256

257 **Response:** Thank you for your comment. Firstly, apologies, we have now stated that all cutting were trials were  
258 performed as fast as possible. Secondly, in terms of calculating approach velocity we stated: “as calculated by  
259 Jones et al, (41)”, but we understand this is not clear. Consequently, we have amended the section as follows:  
260 “Approach velocities were  $4.5 \pm 0.5$  m·s<sup>-1</sup> at initial contact (touch-down) of the PFC, by calculating the horizontal  
261 centre off mass velocity using the combined lower-limb and trunk model as recommended by Vanrenterghem  
262 et al. (77) and used previously in our laboratory (41).”  
263

264 Page 8, lines 451-452: Place the filtering and events part of your analysis before you talk about calculating  
265 external joint moments as if someone was going to replicate your study step by step. Also, state that you  
266 normalized joint moments by body mass here.  
267

268 **Response:** Thank you for your comment. Apologies for this error. We have taken on board your suggestion and  
269 moved the filtering events to the paragraph above.  
270

271 Page 9, lines 517-523: It is unclear how you ran your statistics...i.e. did you correlate mean KAM and mean CMAS  
272 for each subject, mean KAM and CMAS for one trial or did you correlate KAM and CMAS for each trial with the  
273 subject coded to each trial?  
274

275 **Response:** Thank you for your comment. We have provided a detailed response which is contained below your  
276 general comments section outlining our new statistical analyses and procedures. We have now correlated the  
277 mean CMAS and mean pKAM for each subject.  
278

## 279 **Results**

280 Figure 2 A-C: How did you account for the subject variance in the statistical analysis here? When you run a  
281 correlation on individual trials without accounting for intra-subject variability you violate some statistical  
282 assumptions.  
283

284 **Response:** Thank you for your comment. We have provided a detailed response which is contained below your  
285 general comments section outlining our new statistical analyses and procedures. We have now correlated the  
286 mean CMAS and mean pKAM for each subject.  
287

288 Page 831-847: Again here, how were the subject variances accounted for? I completely understand why you  
289 would evaluate individual trial biomechanics variables and CMAS rather than an average as to not lose resolution  
290 but inter and intra subject variance cannot be averaged together.  
291  
292  
293  
294  
295



296  
297  
298 Response: Thank you for your comment. We have provided a detailed response which is contained below your  
299 general comments section outlining our new statistical analyses and procedures. We have now correlated the  
300 mean CMAS and mean pKAM for each subject.  
301

302 Tables 3-4: There are a lot of variables here and you haven't justified why you would measure all of them in your  
303 introduction and methods. Place into the methods section what variables you measured and provide citations  
304 for each of them.  
305

306 Response: Thank you for your comment. We elude to some of these variables in paragraph 2 of the introduction.  
307 We have edited the methods (procedures section 2.4) following your suggestions. We provide a stronger  
308 justification for the additional kinetic and kinematic variables by stating the following: "Joint kinematics and GRF  
309 were also calculated using visual 3D, with Supplement 2 providing variables examined, definitions, and  
310 calculations. Briefly, the following kinetic and kinematics were examined to provide insight into potentially 'high-  
311 risk' cutting mechanics: vertical and horizontal GRF, knee flexion, rotation, and abduction angles and moments,  
312 hip rotation angle, trunk inclination angle, lateral foot plant distance, lateral trunk flexion, internal foot  
313 progression angle, and knee flexion. These kinetic and kinematics were evaluated because they have been  
314 shown to be associated with greater multiplanar knee joint loads (22, 48, 80), and have also been identified as  
315 visual characteristics of non-contact ACL injury during cutting (46, 52, 65). A more detailed rationale for  
316 investigating these variables is presented in Supplement 1. Supplement 1 contains citations for every variable  
317 examined.  
318

## 319 Discussion

320 Line 993-1004: This has already been stated in the introduction. Consider removing and focus upon placing your  
321 results into context with these studies.  
322

323 Response: Thank you for your comment. We have removed this section as Reviewer 1 also highlighted this.  
324

325 You present a lot of biomechanics results, to quantitatively measure the differences between CMAS groups  
326 based on the items (i.e. knee abduction angle) however you don't mention this at all in your discussion. Consider  
327 acknowledging that the scoring system and your raters were able to actually assess measurable differences in  
328 kinematics/kinetics between groups as it is a nice finding.  
329

330 Response: Thank you for your comment. We did discuss the differences in the kinetics and kinematics in the  
331 original manuscript in paragraph 3 of the discussion, but we do agree with your point of the raters being able to  
332 identify the abnormal movement mechanics. Consequently, we have added the following to the bottom of  
333 paragraph 3: "Krosshaug et al. (51) has highlighted the potential difficulties in estimating 3D joint kinematics  
334 based on 2D video evaluations of cutting mechanics. Conversely, the results indicate that the raters in the  
335 present study were capable of accurately evaluating and identifying aberrant lower-limb and trunk postures  
336 during cutting, as confirmed by the measurable difference in 3D kinetics and kinematics between subjects with  
337 "high" and "low" CMASs related to the CMAS scoring system (Table 3)."  
338

339 It is obviously a little harder to perform but multiple studies have compared planned vs unplanned sidestepping  
340 and identified that the mechanics are quite different and the KAM is much higher with much less muscle support  
341 to counter this load. I think you should identify this as a limitation of your study.  
342

343 Besier, T.F., Lloyd, D. G., Cochrane, J.L., & Ackland, T.R. (2001). External loading of the knee joint during running  
344 and cutting maneuvers. *Medicine & science in sports & exercise*, 33(7), 1168-1175.  
345

346 Besier, T. F., Lloyd, D. G., Ackland, T. R., & Cochrane, J. L. (2001). Anticipatory effects on knee joint loading during  
347 running and cutting maneuvers. *Medicine and science in sports and exercise*, 33(7), 1176-1181.  
348

349 Response: Thank you for your comment. We have acknowledged this as a limitation and stated the following:

350 "Finally, a pre-planned cutting task was used in the present study; however, results of previous research have  
351  
352  
353  
354

355  
356  
357 shown that unplanned side-stepping results in greater knee loads, more abnormal mechanics, and less muscle  
358 support to counteract the greater loads compared to pre-planned side-stepping (5, 6, 11).” We feel the CMAS  
359 could be applied to unanticipated cutting.  
360  
361

362 Were all of the raters biomechanists/grad students?  
363

364 Response: Thank you for your comment. Yes, they were biomechanists/ S&C/sports science graduates. We  
365 provide the specific information regarding the raters qualifications and experience in the statistical analysis  
366 section. We state the following: “The lead researcher, who has seven years’ strength and conditioning and  
367 biomechanics experience, viewed and graded each trial on two separate occasions separated by 7 days, in line  
368 with previous research (24, 72) to examine intra-rater reliability. Another researcher (experienced  
369 biomechanist; 17 years’ biomechanics and strength and conditioning experience), viewed and graded each trial  
370 once and these scores were compared to the lead researcher to establish inter-rater reliability. In addition, a  
371 recent sports science graduate also viewed and graded each trial once and these scores were compared to the  
372 lead researcher to establish inter-rater reliability.”  
373

374 Do you think your results would change if a coach did these measures?  
375

376 Response: Thank you for your comment. In response, possibly; however, we can not answer this based on our  
377 data, but it is indeed something to consider in future work which we will look to do. We state the following in  
378 the limitation’s sections: “Further work is required to establish agreements and reliability between different  
379 applied practitioners, such as sports rehabilitators, physiotherapists, sports coaches, in order to confirm its  
380 efficacy in the field.”  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413

1  
2  
3 **Highlights:**  
4

- 5 • CMAS is a valid and reliable screening tool for evaluating side-step cutting movement quality.
- 6 • **A very large significant** relationship was observed between CMAS and peak KAM.
- 7 • CMAS offers practitioners a cost-effective and easily applicable field-based screening tool to
- 8 identify athletes who generate high peak KAMs.
- 9 • CMAS allows practitioners to identify “high-risk” cutting mechanics in athletes
- 10 • CMAS can be used as a potential technical framework for coaching “safer” cutting.
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59

1  
2  
3 **A QUALITATIVE SCREENING TOOL TO IDENTIFY ATHLETES WITH ‘HIGH-RISK’**  
4 **MOVEMENT MECHANICS DURING CUTTING: THE CUTTING MOVEMENT ASSESSMENT**  
5 **SCORE (CMAS)**  
6

7 *Original Research*

8  
9 Thomas Dos’Santos<sup>1#</sup>, Alistair McBurnie<sup>1</sup>, Thomas Donelon<sup>2</sup>, Christopher Thomas<sup>1,3</sup>, Paul Comfort<sup>1</sup> and Paul  
10 A. Jones<sup>1</sup>  
11

12 Brief running head: Cutting Movement Assessment Score

13  
14 <sup>1</sup>Human Performance Laboratory, Directorate of Sport, Exercise, and Physiotherapy, University of Salford,  
15 Greater Manchester, United Kingdom

16 <sup>2</sup>Section of Sport and Exercise Sciences, Canterbury Christ Church University, Kent, United Kingdom

17 <sup>3</sup>Aspire Academy, Doha, Qatar  
18  
19

20 #Corresponding Author: Thomas Dos’Santos

21 Address: Allerton Building, Frederick Road Campus, Salford, Greater Manchester, United Kingdom, M6 6PU.

22 Telephone: +447961744517

23 Email: [t.dossantos@edu.salford.ac.uk](mailto:t.dossantos@edu.salford.ac.uk)  
24  
25

26  
27 Abstract word count: 200 words

28 Manuscript word count: 4489 words  
29  
30

31 Number of tables and figures: 3 Tables, 2 Figures

32 3 Supplementary materials  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59

---

**A QUALITATIVE SCREENING TOOL TO IDENTIFY ATHLETES WITH ‘HIGH-RISK’  
MOVEMENT MECHANICS DURING CUTTING: THE CUTTING MOVEMENT  
ASSESSMENT SCORE (CMAS)**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59

## **ABSTRACT**

**Objective:** To assess the validity of the cutting movement assessment score (CMAS) to estimate the magnitude of peak knee abduction moments (KAM) against three-dimensional (3D) motion analysis, while comparing whole-body kinetics and kinematics between subjects of **low (bottom 33%) and high CMASs (top 33%)**.

**Design:** Cross-sectional study.

**Setting:** Laboratory.

**Participants:** Forty-one participants (soccer, rugby, netball, and cricket).

**Main outcome measures:** Association between peak KAM and CMAS during a 90° cut. Comparison of 3D whole-body kinetics and kinematics between subjects **with low (bottom 33%) and high CMASs (top 33%)**.

**Results:** A **very large significant relationship** ( $\rho = 0.796, p < 0.001$ ) between CMAS and peak KAM was observed. **Subjects with higher CMASs displayed higher-risk cutting postures, including greater peak knee abduction angles, internal foot progression angles, and lateral foot plant distances** ( $p \leq 0.032$ , effect size = 0.83-1.64). **Additionally, greater cutting multiplanar knee joint loads (knee flexion, internal rotation, and abduction moments) were demonstrated by subjects with higher CMASs compared to lower** ( $p \leq 0.047$ , effect size = 0.77-2.24).

**Conclusion:** The CMAS is a valid qualitative screening tool for evaluating cutting movement quality and is therefore a potential method to identify athletes who generate high KAMs and “high-risk” side-step cutting mechanics.

**Keywords:** Anterior cruciate ligament; knee abduction moment; injury screening; injury-risk profile

### **Highlights:**

- CMAS is a valid and reliable screening tool for evaluating side-step cutting movement quality.
- **A very large significant** relationship was observed between CMAS and peak KAM.
- CMAS offers practitioners a cost-effective and easily applicable field-based screening tool to identify athletes who generate high peak KAMs.
- CMAS allows practitioners to identify “high-risk” cutting mechanics in athletes
- CMAS can be used as a potential technical framework for coaching “safer” cutting.

## 1. INTRODUCTION

Side-step lateral foot plant-and-cut actions are frequently performed movements in numerous sports (25, 84) and are also linked to decisive moments in matches, such as evading an opponent to penetrate the defensive line in rugby (tackle-break success in rugby) (84), or getting into to space to receive a pass in netball (25). Side-step cutting, however, are also actions associated with non-contact anterior cruciate ligament (ACL) injuries in sports (39, 48, 62). Although ACL injury-risk factors are multifactorial (74) and a complex interaction of internal and external factors (i.e. anatomical, hormonal, environmental, shoe-surface interface, anticipation, and fatigue) (8, 34, 54), a large proportion of ACL injuries are non-contact in nature during high velocity and impact sporting tasks, such as side-stepping (9, 39, 54). This occurrence can be attributed to the tendency to generate large multiplanar knee joint loading, such as knee abduction moments (KAM) and internal rotation moments (KIRM) (7, 19, 41), which increase ACL strain (4, 57, 76). These potentially hazardous knee joint loads are amplified when poor initial postures and movement is demonstrated (biomechanical and neuromuscular control deficits) during cutting (24, 34, 65), but importantly these deficits are modifiable (34, 71). As such, understanding the mechanics, interventions, and screening tools that can reduce ACL injury-risk factors is of critical importance.

The ability to identify athletes potentially at risk of injury is a critical step in effective ACL injury-risk reduction (26, 34). Although it is inconclusive whether screening tools can predict non-contact ACL injury (3, 27), evaluating movement quality and identifying biomechanical and neuromuscular control deficits (high-risk movement patterns) can provide important information regarding an athlete's "injury-risk profile" (33, 58, 61). These abnormal deficits include knee abduction angles (KAA) (40, 42, 50, 59, 77), lateral trunk flexion (19, 28, 38, 40), extended knee postures (16, 48, 83), and hip internal rotation (29, 59, 77, 78). This information from movement screening can subsequently be used to inform the future prescription of training and conditioning so specific deficits can be targeted through appropriate training interventions to decrease the relative risk of injury (33, 35, 61). Therefore, the inclusion of valid and reliable screening tools that assess movement quality are an important component of sports medicine and strength and conditioning testing batteries to provide an "injury-risk profile" for an athlete (33, 44).

Three-dimension (3D) motion analysis is considered the gold standard for evaluating movement kinetics and kinematics (27, 34); however, this method can be susceptible to errors, with a diverse range of data collection and analysis procedures available to practitioners which can impact outcome values, reliability, or subsequent evaluations of an athlete's biomechanical profile (12, 52). Given these methodological considerations and issues, and the fact the 3D motion analysis is expensive, time-consuming, requires expert and well trained assessors, and is usually restricted to testing one subject in laboratory setting, time- and cost-effective qualitative field-based screening tools have been developed, such as the landing error scoring system (LESS) (70, 72), tuck jump assessment (TJA) (32, 66), and

178  
179  
180 qualitative analysis of single leg loading (QASLS) (2, 31), to assess lower-limb and whole-body  
181 postures associated with increased potential risk of injury (high-risk movement patterns). **However, the**  
182 **LESS is the only screening tool of that has been validated against 3D motion analysis** (69, 72).  
183  
184

185 A fundamental shortcoming of the LESS, TJA, and QASLS are these assessments generally assess  
186 landing mechanics during a vertical-orientated task. Although screening landing mechanics is indeed  
187 applicable to jump-landing sports (netball, basketball, volleyball) where the primary action associated  
188 with non-contact ACL injury is landing manoeuvres (36, 54, 79), these aforementioned assessments  
189 may lack **specificity** to the unilateral, multiplanar plant-and-cut manoeuvres observed when changing  
190 direction (27, 44, 61). This is particularly important when aiming to screen athletes who participate in  
191 sports such as **soccer (82), handball (68), American football (39), badminton (46), and rugby (63)**, where  
192 directional changes are a primary action associated with non contact ACL injuries. Furthermore, there  
193 are mixed findings whether examination of landing mechanics can identify athletes with poor cutting  
194 mechanics (1, 13, 51, 67), with evidence suggesting an athlete's mechanics and "injury-risk profile" are  
195 task dependent (13, 45, 51, 64). **As such, screening side-step cutting technique, which is specific to the**  
196 **actions associated with non-contact ACL injuries in cutting sports (i.e. rugby, handball, soccer,**  
197 **American football), could be a more effective strategy for identify poor cutting movement quality in**  
198 **athletes, which can help inform future injury-risk mitigation training.**  
199  
200  
201  
202  
203  
204  
205  
206

207 Unfortunately, there is a paucity of field-based cutting screening tools available for practitioners.  
208 McLean et al. (60) initially evaluated two-dimensional (2D) estimates of frontal plane knee motion  
209 during cutting against the gold standard of 3D, and found 2D estimates correlated well with side-step  
210 ( $r^2 = 0.58$ ) and side-jump ( $r^2 = 0.64$ ) 3D valgus angles, but poorer associations were observed with 180°  
211 turn knee valgus angle ( $r^2 = 0.04$ ); thus, highlighting the difficulty in assessing 2D valgus motion in the  
212 frontal plane using a single camera during sharp CODs. Weir et al. (83) has recently demonstrated that  
213 2D measures of dynamic knee valgus angle, knee flexion angle at foot-strike and ROM, trunk flexion  
214 ROM, when inserted in regression equations, can be used to predict 3D peak knee flexor, KAM and  
215 KIRMs during unanticipated side-steps. Despite these promising relationships, such 2D side-step  
216 screening methods are not widely adopted by practitioners and clinicians. This lack of adoption could  
217 be attributed to the 2D method requiring additional time and software to measure joint kinematics, thus  
218 potentially limiting its applicability in field settings.  
219  
220  
221  
222  
223  
224

225 In light of the issues associated with 2D analysis, Jones et al. (44) have recently developed the  
226 cutting movement assessment score (CMAS), which is a qualitative screening tool that assesses cutting  
227 movement quality and specific lower-limb and trunk characteristics that are associated with (24, 50, 83)  
228 peak KAMs (Supplement 1), such as penultimate foot contact (PFC) braking strategy, and trunk, hip,  
229 knee, and foot positioning and motions. In this preliminary study, a strong relationship between CMAS  
230 and peak KAM ( $\rho = 0.633$ ;  $p < 0.001$ ) was demonstrated, while moderate to excellent intra-and inter-  
231  
232  
233  
234  
235  
236



rater agreements for all CMAS variables (Intra-rater:  $k = 0.60-1.00$ , 75-100% agreements; inter-rater:  $k = 0.71-1.00$ , 87.5-100% agreements) were observed, although lower inter-rater agreements for trunk positioning were observed ( $k = 0.40$ , 62.5% agreement). In light of these findings, the CMAS may have the potential to identify athletes displaying “high-risk” cutting mechanics but more importantly, could be used as a technical framework for coaching safer cutting mechanics. It should be noted, however, that the preliminary study contained a small sample size ( $n = 8$  subjects, 36 trials) and must be expanded with a greater sample size to confirm its validity and reliability. Furthermore, the authors recommended an additional camera to be placed at  $45^\circ$  relative to the COD and using a higher video capture rate ( $\geq 100$  Hz) to permit more accurate and reliable assessments for frontal and transverse plane technique deficits (i.e. trunk positioning, knee valgus).

The aim of this study, therefore, was to assess the validity of the CMAS tool to estimate the potential peak KAMs against the gold standard of 3D motion analysis, expanding on the work of Jones et al. (44) by examining a larger sample size and using an additional camera recording at a higher sampling rate. A further aim was to determine whether “higher-risk” movement mechanics were displayed by subjects with higher CMASs compared to subjects with lower CMASs. Firstly, it was hypothesised that excellent inter- and intra-rater reliability would be demonstrated for CMAS items. Secondly, in line with Jones et al. (44), it was hypothesised that a strong relationship would be demonstrated between CMAS and peak KAM, and the CMAS would be able to discriminate between “low” and “high” CMASs in terms of “high-risk” whole-body kinetics and kinematics.

## 2. METHODS

### 2.1 Experimental approach

This study used a cross-sectional design to determine the relationship between CMAS and peak KAMs during cutting over one session. Participants performed six  $90^\circ$  cuts ( $70-90^\circ$ ) whereby 3D motion and 2D video footage data were simultaneously captured to permit qualitative screening and comparisons to 3D motion data, similar to the procedures of previous research (44, 72).

### 2.2 Participants

Based on the work of Jones et al. (44) who determined the relationship between CMAS and peak KAM, a minimum sample size of 29 was determined from an *a priori* power analysis using G\*Power (Version 3.1, University of Dusseldorf, Germany) (22). This was based upon a correlation value of  $\rho = 0.633$ , a power of 0.95, and type 1 error or alpha level of 0.05. As such, 41 athletes (28 males/13 females) from multiple sports (soccer, rugby, netball, and cricket) (mean  $\pm$  SD; age:  $21.3 \pm 4.0$  years, height:  $1.75 \pm 0.08$  m, mass:  $72.8 \pm 11.8$  kg) participated in this study. For inclusion in the study, all athletes had played their respective sport for a minimum of 5 years and regularly participated in one game and performed two structured skill-based training sessions per week. All athletes were free from injury and had never suffered a prior traumatic knee injury such as an ACL injury. At the time of testing, players

were currently in-season (competition phase). The investigation was approved by the institutional ethics review board, and all participants were informed of the benefits and risks of the investigation prior to signing an institutionally approved consent and parental assent documents to participate in the study.

### 2.3 Cutting Movement Assessment Score

Table 1 presents the CMAS qualitative screening analysis tool to estimate the magnitude of KAMs during cutting, which has been slightly modified from the preliminary investigation by Jones et al. (44) (i.e. extra description provided to some criteria). The CMAS is based on research pertaining to technical determinants of peak KAMs during 30-90° side-step cutting (24, 50, 83) and visual observations of non-contact ACL injuries (39, 48, 68). Supplement 1 contains operation definitions and a biomechanical rationale of the CMAS. If an athlete exhibits any of the characteristics in Table 1 they are awarded a score, with a higher score representative of poorer technique and potentially greater peak KAM (44).

**Table 1. Cutting movement assessment score tool**

Camera	Variable	Observation	Score
<b>Penultimate contact</b>			
<i>Side / 45°</i>	<b>Clear PFC braking strategy</b> (at initial contact) <ul style="list-style-type: none"> <li>Backward inclination of the trunk</li> <li>Large COM to COP position – anterior placement of the foot</li> <li>Effective deceleration – heel contact PFC</li> </ul>	Y/N	Y=0/ N=1
<b>Final Contact</b>			
<i>Front / 45°</i>	<b>Wide lateral leg plant</b> (approx. > 0.35 m – dependent on subject anthropometrics) (at initial contact)	Y/N	Y=2/N=0
<i>Front / 45°</i>	<b>Hip in an initial internally rotated position</b> (at initial contact)	Y/N	Y=1/N=0
<i>Front / 45°</i>	<b>Initial knee ‘valgus’ position</b> (at initial contact)	Y/N	Y=1/N=0
<i>All 3</i>	<b>Foot not in neutral foot position</b> (at initial contact) Inwardly rotated foot position or externally rotated foot position (relative to original direction of travel)	Y/N	Y=1/N=0
<i>Front / 45°</i>	<b>Frontal plane trunk position relative to intended direction; Lateral or trunk rotated towards stance limb, Upright, or Medial</b> (at initial contact and over WA)	L/TR/U/M	L/TR=2/ U = 1, /M=0
<i>Side / 45°</i>	<b>Trunk upright or leaning back throughout contact</b> (not adequate trunk flexion displacement) (at initial contact and over WA)	Y/N	Y=1/N=0
<i>Side / 45°</i>	<b>Limited Knee flexion during final contact (stiff) <math>\leq 30^\circ</math></b> (over WA)	Y/N	Y=1/N=0
<i>Front / 45°</i>	<b>Excessive Knee ‘valgus’ motion during contact</b> (over WA)	Y/N	Y=1/N=0
		<b>Total Score</b>	<b>0 /11</b>

Key: PFC: Penultimate foot contact; COM: Centre of mass; COP: Centre of pressure; WA: weight acceptance; TR: Trunk rotation; Y: Yes; N: No; L: Lateral; TR: Trunk rotation; U: Upright; M: Medial.


## 2.4 Procedures

The warm up, 90° cut (21), marker placement (21, 41, 44), and 3D motion analysis (21, 41, 44), and CMAS (44) procedures were based on previously published methodologies (21, 44), thus a brief overview is provided here.

Participants performed six trials of a 90° cut **as fast as possible** (70-90°) (Figure 1). Completion time ( $2.11 \pm 0.14$  seconds, coefficient of variation = 2.71%) was measured to standardise performance between trials, and was assessed using two sets of Brower timing lights placed at hip height (Draper, UT, USA). Marker and force data were collected over the penultimate and final foot contact using ten Qualisys Oqus 7 (Gothenburg, Sweden) infrared cameras (240Hz) operating through Qualisys Track Manager software (Qualisys, version 2.16 (Build 3520), Gothenburg, Sweden) and GRF's were collected from two 600 mm × 900 mm AMTI (Advanced Mechanical Technology, Inc, Watertown, MA, USA) force platforms (Model number: 600900) embedded into the running track sampling at 1200Hz, respectively.

Using the pipeline function in visual 3D, joint coordinate (marker) and force data were smoothed using a Butterworth low-pass digital filter with cut-off frequencies of 15 and 25 Hz, based on *a priori* residual analysis (86), visual inspection of motion data, and recommendations by Roewer et al. (75). Lower limb joint moments were calculated using an inverse dynamics approach (85) through Visual 3D software (C-motion, version 6.01.12, Germantown, USA) and were defined as external moments and normalised to body mass. Joint kinematics and GRF were also calculated using visual 3D, with Supplement 2 providing the variables examined, definitions, and calculations. Briefly, the following kinetic and kinematics were examined to provide insight into potentially “high-risk” cutting mechanics: vertical and horizontal GRF, knee flexion, rotation, and abduction angles and moments, hip rotation angle, trunk inclination angle, lateral foot plant distance, lateral trunk flexion, initial foot progression angle, and knee flexion angle. These aforementioned kinetic and kinematics were evaluated because they have been shown to be associated with greater multiplanar knee joint loads (24, 50, 83), and have also been identified as visual characteristics of non-contact ACL injury during cutting (39, 48, 68). A more detailed rationale for investigation of these variables is presented in Supplement 1.

The trials were time normalised for each subject to 101 data points with each point representing 1% of the weight acceptance (WA) phase (0 to 100% of WA) of the cutting task. Initial contact was defined as the instant after ground contact that the vertical GRF was higher than 20 N, and end of contact was defined as the point where the vertical GRF subsided past 20 N (42, 50, 52). The WA phase was defined as the instant of initial contact to the point of maximum knee flexion (29, 40, 41). Approach velocities were  $4.5 \pm 0.5$  m·s<sup>-1</sup> at initial contact (touch-down) of the PFC, by calculating the horizontal centre of mass velocity using the combined lower-limb and trunk model, as recommended by Vanrenterghem et al. (80) and used previously in our laboratory (43).



**Figure 1. Plan view of the experimental set-up. The task involved subjects approaching 5-m towards turning point on 2<sup>nd</sup> force platform. At the turning point, subjects cut to the left 90° using their right limb between timing gates placed 3-m away. Marker, GRF, and 2D camera data were collected simultaneously.**

### 2.5 Qualitative assessment: CMAS

While marker and GRF data were collected, three Panasonic Lumix FZ-200 high speed cameras sampling at 100 Hz simultaneously filmed the cutting trials. These cameras were positioned on tripods 3-m away from the force plates at a height of 0.60 m and were placed in the sagittal and frontal plane, with a camera also placed 45° relative the cut, in accordance with previous recommendations (44) (Figure 1). Video footage was subsequently viewed in Kinovea software (0.8.15 for Windows), which is free, and was used for qualitative screening using the CMAS (Table 1). This software allowed videos to be played at various speeds and frame-by-frame. The three raters were allowed to independently watch the videos as many times as necessary (23, 69), at whatever speeds they needed to score each test, and could also pause footage for evaluative purposes (23). On average, qualitative screening of one trial took ~3 minutes.

Prior to qualitative screening, all raters attended a one-hour training session outlining how to grade the cutting trials using the CMAS, and to establish and uniformly agree on low-risk and high-risk movement patterns using pilot video footage. Subsequently, the lead researcher created a manual for all raters which contained guidelines, operational definitions (Supplement 1 and 3), and example images of low-risk and high-risk motions of each screening criteria to assist CMAS screening.

## 2.6. Statistical analyses

Thirty-two trials were discarded due to technical issues with camera footage, 3D data, or subjects slid or missed the platform that went unnoticed during data collection, thus resulting in 214 trials (minimum 4 trials from 41 athletes) screened and used for further analysis. All statistical analyses were performed in SPSS v 24 (SPSS Inc., Chicago, IL, USA) and Microsoft Excel (version 2016, Microsoft Corp., Redmond, WA, USA). To determine inter- and intra-rater reliability, 41 trials (one trial from each subject) were randomly selected by the lead researcher, similar to the procedures of previous research (44). The lead researcher, who has seven years' strength and conditioning and biomechanics experience, viewed and graded each trial on two separate occasions separated by 7 days, in line with previous research (23, 72) to examine intra-rater reliability. Another researcher (experienced biomechanist; 17 years' biomechanics and strength and conditioning experience), viewed and graded each trial once and these scores were compared to the lead researcher to establish inter-rater reliability. In addition, a recent sports science graduate also viewed and graded each trial once and these scores were compared to the lead researcher to establish inter-rater reliability.

Intra-class correlation coefficients (ICC) (two-way mixed effects, average measures, absolute agreement) for total score were determined. Intraclass correlations were interpreted based on the following scale presented by Koo and Li (49): poor ( $< 0.50$ ), moderate (0.50-0.75), good (0.75-0.90), and excellent ( $> 0.90$ ). For each item within the CMAS (Table 1), percentage agreements (agreements /agreements + disagreements  $\times 100$ ) and Kappa co-efficients were calculated. Kappa co-efficients were calculated using the formula;  $k = \text{Pr}(a) - \text{Pr}(e) / 1 - \text{Pr}(e)$ , where  $\text{Pr}(a)$  = relative observed agreement between raters;  $\text{Pr}(e)$  = hypothetical probability of chance agreement, which describes the proportion of agreement between the two methods after any agreement by chance has been removed (81). The kappa co-efficient was interpreted based on the following scale of Landis and Koch (55): slight (0.01-0.20), fair (0.21-0.40), moderate (0.41-0.60), good (0.61-0.80), and excellent (0.81-1.00). Percentage agreements were interpreted in line with previous research (15, 69) and the scale was as follows: excellent ( $>80\%$ ), moderate (51-79%), and poor ( $< 50\%$ ) (15, 69).

The relationship between CMAS and the “gold standard” determination of peak KAM during the final foot contact (FFC) of the cutting task from 3D motion analysis using the **means of each subject** was explored using Spearman's rank correlation, **with 95% confidence intervals (CI)**, due to the non-parametric nature of the qualitative data. Correlations were evaluated as follows: trivial (0.00-0.09), small (0.10–0.29), moderate (0.30–0.49), large (0.50–0.69), very large (0.70–0.89), nearly perfect (0.90–0.99), and perfect (1.00) (37). This analysis was performed using the 214 trials screened by the lead researcher.

**Subjects were classified into low CMAS (bottom 33%,  $n = 14$ ) and high CMAS (top 33%,  $n = 14$ ) groups based on their mean CMASs. Subsequently, cutting 3D kinetics and kinematics were compared between the two groups (subject mean data) using independent sample t tests for parametric**

532  
533  
534 **data and Mann-Whitney U tests for non-parametric data.** To explore the magnitude of differences  
535  
536 between groups, **mean differences with 95% CIs and** Hedges' *g* effect sizes with 95% **CIs** were also  
537 calculated as described previously (30), and interpreted as trivial (< 0.19), small (0.20–0.59), moderate  
538 (0.60–1.19), large (1.20–1.99), very large (2.0–3.99), and extremely large ( $\geq 4.00$ ) (37). Statistical  
539 significance was defined  $p \leq 0.05$  for all tests.  
540  
541

### 542 **3. RESULTS**

#### 543 **3.1 Intra- and inter-rater reliability**

544  
545 Excellent intra-rater reliability was observed for CMAS total score (ICC = 0.946). Intra- and inter-rater  
546 percentage agreements and Kappa coefficients are presented in Table 2. Excellent intra-rater  
547 percentage-agreements and kappa-coefficients were demonstrated for all CMAS variables (Table 2),  
548 with two variables scoring 100% agreement. For inter-rater reliability, most items displayed moderate  
549 to excellent percentage agreements (Table 2), while most items displayed moderate to good kappa  
550 coefficients between the lead researcher and experienced biomechanist. Conversely, kappa coefficients  
551 ranged from slight to good between the lead researcher and recent graduate, and most items displayed  
552 moderate to excellent percentage agreements (Table 2). Moderate inter-rater reliability was observed  
553 for CMAS total score between raters (ICC = 0.690)  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590


Table 2. Intra- and inter-rater reliability for CMAS criteria and total score

Variable/ CMAS tool criteria	Intra-rater reliability (Lead researcher)		Inter-rater reliability - Lead research vs experienced biomechanist		Inter-rater reliability - Lead researcher vs recent graduate	
	% agreement	<i>k</i>	% agreement	<i>k</i>	% agreement	<i>k</i>
Clear PFC braking	97.6	0.940	82.9	0.633	82.9	0.633
Wide lateral leg plant	95.1	0.900	82.9	0.629	87.8	0.747
Hip in an initial internally rotated position	100.0	1.000	63.4	0.194	43.9	0.067
Initial knee 'valgus' position	90.2	0.805	75.6	0.512	75.6	0.512
Inwardly rotated foot position	100.0	1.000	80.5	0.599	90.2	0.784
Frontal plane trunk position relative to intended direction	90.2	0.805	73.2	0.551	87.8	0.767
Trunk upright or leaning back throughout contact	100.0	1.000	90.2	0.554	78.0	0.220
Limited Knee Flexion during final contact	97.6	0.932	80.5	0.431	80.5	0.381
Excessive Knee 'valgus' motion during contact	95.1	0.898	80.5	0.605	70.7	0.376
Average	96.2	0.920	78.9	0.52	77.5	0.50

Key: CMAS: Cutting movement assessment score; PFC: Penultimate foot contact

### 3.2 Relationships between CMAS and peak KAM

Mean  $\pm$  SD from each trial of the 41 subjects were  $5.1 \pm 1.8$  CMAS and peak KAM  $1.00 \pm 0.44$  Nm/kg. CMASs and KAMs for males and females were  $5.1 \pm 1.7$ ,  $1.07 \pm 0.45$  Nm/kg and  $5.2 \pm 2.1$ , KAM  $0.81 \pm 0.35$  Nm/kg, respectively. Figure 2 shows a linear and positive relationship between CMAS and peak KAMs. Spearman's correlation revealed a significant and very large ( $\rho = 0.796$ , 95% CI = 0.647-0.887,  $p < 0.001$ ) association between CMAS and peak KAMs.



**Figure 2. Relationship between CMAS and peak KAMs (pKAM) subject mean data.**

### 3.3 Comparisons in cutting 3D kinetics and kinematics between subjects with low and high CMASs

Descriptive statistics,  $p$  values, and effect sizes for kinetic and kinematic measures for subjects with low and high CMASs are presented in Table 3. Subjects with higher CMASs displayed significantly greater FFC mean VBFs, HBFs, and mean HBF ratios, and greater peak knee abduction angles, internal foot progression angles, and lateral foot plant distances (Table 3), with moderate to large effect sizes. Additionally, significantly greater cutting multiplanar knee joint loads (KFMs, KIRMs, and KAMs) were demonstrated by subjects with higher CMASs compared to lower (Table 3), with moderate to very large effect sizes.



Table 3. Comparisons in 3D cutting mechanics between subjects with lower and higher CMAS containing *p* values and effect size

Variable	Foot contact	Low CMAS ( <i>n</i> = 14)		High CMAS ( <i>n</i> = 14)		<i>p</i>	<i>g</i>	95% <i>g</i>		Mean difference	Mean difference 95% CI			
		Mean	SD	Mean	SD			LB	UB		LB	UB		
CMAS		3.34	0.70	6.95	0.63	<b>&lt;0.001</b>	-5.29	-6.87	-3.72	-3.61	-4.13	-3.10		
GRF	peak VBF (BW)	PFC	2.67	0.55	2.72	0.63	0.855	-0.07	-0.81	0.67	-0.04	-0.50	0.42	
	mean VBF (BW)	PFC	0.95	0.16	0.97	0.20	0.879	-0.06	-0.80	0.68	-0.01	-0.15	0.13	
	peak HBF /BW)	PFC	-1.53	0.52	-1.50	0.48	0.872	-0.06	-0.80	0.68	-0.03	-0.42	0.36	
	mean HBF /BW)	PFC	-0.56	0.12	-0.53	0.14	0.617	-0.18	-0.92	0.56	-0.02	-0.12	0.07	
	peak VBF (BW)	FFC	2.55	0.53	2.64	0.46	0.632	-0.18	-0.92	0.56	-0.09	-0.48	0.30	
	mean VBF (BW)	FFC	1.54	0.18	1.71	0.21	<b>0.029</b>	-0.84	-1.61	-0.07	-0.17	-0.33	-0.02	
	peak HBF (BW)	FFC	-1.44	0.35	-1.45	0.24	0.975	0.02	-0.73	0.76	0.00	-0.23	0.23	
	mean HBF (BW)	FFC	-0.78	0.16	-0.94	0.13	<b>0.009</b>	1.03	0.24	1.82	0.16	0.04	0.27	
	peak HBF ratio	both	1.03	0.35	1.06	0.39	<i>0.909</i>	-0.09	-0.83	0.66	-0.03	-0.32	0.26	
	mean HBF ratio	both	1.42	0.29	1.88	0.65	<b><i>0.018</i></b>	-0.88	-1.66	-0.10	-0.45	-0.84	-0.06	
	Joint kinematics	peak KFA (°)	FFC	66.6	9.0	62.5	7.5	0.209	0.47	-0.28	1.22	4.0	-2.4	10.5
		KFA - IC (°)	FFC	23.1	5.1	23.6	4.9	0.766	-0.11	-0.85	0.63	-0.6	-4.5	3.3
KFA ROM (°)		FFC	43.5	7.3	38.9	5.9	0.080	0.67	-0.09	1.43	4.6	-0.6	9.8	
peak KAA (°) (- abduction, + adduction)		FFC	-7.8	6.5	-13.4	6.6	<b>0.032</b>	0.83	0.06	1.60	5.6	0.5	10.7	
KAA - IC (°) (- abduction, + adduction)		FFC	4.3	4.8	0.6	4.7	0.052	0.75	-0.02	1.51	3.7	0.0	7.4	
KAA ROM (°)		FFC	-12.1	4.9	-14.0	5.4	0.321	0.37	-0.38	1.12	2.0	-2.0	5.9	
KRA - IC (°) (- internal, + external)		FFC	-10.7	6.9	-4.5	6.2	<b>0.020</b>	-0.91	-1.69	-0.13	-6.2	-11.3	-1.1	
peak KRA (°) (- internal, + external)		FFC	-9.6	7.4	-1.0	8.6	<b>0.009</b>	-1.04	-1.83	-0.25	-8.6	-14.8	-2.3	
Hip rotation angle - IC (°) (- internal, + external)		FFC	11.0	7.1	7.9	10.6	0.377	0.33	-0.42	1.08	3.1	-3.9	10.1	
Technique	Trunk inclination angle - IC (°) (relative to vertical line, + forward, - backward)	PFC	6.8	3.9	8.1	3.4	0.361	-0.34	-1.09	0.41	-1.3	-4.1	1.6	
	Trunk inclination angle - IC (°) (relative to vertical line, + forward, - backward)	FFC	17.2	31.3	10.4	6.0	0.437	0.29	-0.46	1.03	6.7	-10.8	24.2	
	IFPA - IC (°) (- internal, + external)	FFC	9.0	10.2	25.5	9.3	<b>&lt;0.001</b>	-1.64	-2.49	-0.78	-16.5	-24.1	-8.9	
	Lateral trunk flexion - IC (°) (- over stance leg, + direction of travel)	FFC	-18.4	8.0	-17.6	7.3	0.794	-0.10	-0.84	0.64	-0.8	-6.7	5.2	
	Lateral foot plant distance - IC (m)	FFC	-0.299	0.041	-0.336	0.044	<b>0.028</b>	0.85	0.08	1.63	0.038	0.004	0.071	
Joint moment	peak KFM (Nm/kg)	FFC	3.06	0.60	3.64	0.72	<b>0.027</b>	-0.86	-1.64	-0.09	-0.59	-1.10	-0.07	
	peak KRM (Nm/kg) (- internal, + external)	FFC	-0.69	0.39	-1.10	0.61	<b>0.047</b>	0.77	0.01	1.54	0.41	0.01	0.81	
	peak KAM (Nm/kg) (+ abduction, - adduction)	FFC	0.73	0.27	1.37	0.28	<b>&lt;0.001</b>	2.24	-3.18	-1.29	-0.63	-0.85	-0.42	

Key: VBF: Vertical braking force; HBF: Horizontal braking force; FFC: Final foot contact; PFC: Penultimate foot contact; IC: Initial contact; BW: Body weight; KFA: Knee flexion angle; ROM: Range of motion; KAA: Knee abduction angle; KRA: Knee rotation angle; IFPA: Initial foot progression angle; KFM: Knee flexor moment; KRM: Knee rotation moment; KAM: Knee abduction moment; ES: Effect size; CMAS: Cutting movement assessment scores; Sag: Sagittal. CI: Confidence interval; LB: Lower bound; UB: Upper bounds; ES: Effect size. Note: Bold denotes statistically significant difference ( $p < 0.05$ ) and italic denotes non-parametric.

#### 4. DISCUSSION

The primary aim of this study was to examine the validity and relationship between the CMAS attained from a qualitative screening tool and peak KAM quantified via 3D motion analysis. This study expanded on the preliminary work of Jones et al. (44) by using an additional camera filming at a higher sampling rate, and also investigating a larger sample size. In line with the study hypotheses, and substantiating Jones et al. (44), a very large ( $\rho = 0.796, p < 0.001$ ) (Figure 2) relationship was observed between CMAS and peak KAM. Moreover, “higher-risk” cutting mechanics associated with greater knee joint loading, thus ACL injury-risk, were displayed by subjects with higher CMASs (~7) compared to subjects with lower CMASs (~3) (Table 3). The CMAS also demonstrated excellent intra-rater reliability (Table 2), and generally moderate-to-excellent inter-rater reliability (Table 2). Therefore, these findings indicate that the CMAS qualitative screening tool can be considered a reliable and valid method to identify athletes who generate high KAMs and “high-risk” cutting mechanics. This tool offers practitioners a field-based screening method which can be included in testing and screening batteries for cutting sports so “high-risk” cutting deficits can be identified and “injury-risk profiles” can be created for athletes.

In light of kinetic and kinematics (high-risk) cutting deficits associated with greater knee joint loads during side-step cutting (Supplement 1), Jones et al. (44) developed the CMAS screening tool and reported a large relationship CMAS and peak KAM ( $\rho = 0.633; p < 0.001$ ). Expanding on the preliminary investigation by Jones et al. (44), the present study observed a stronger relationship between CMAS and peak KAMs ( $\rho = 0.796, p < 0.001$ , Figure 2), in a substantially greater sample size (41 vs. 8 subjects). The stronger relationships observed in the present study, compared to Jones et al. (44), could be attributed to the additional camera placed at 45° and increased sampling rate of the cameras (100 vs. 30 Hz). These additions may have permitted more accurate screening and evaluations of frontal and transverse plane deficits, such as trunk positioning and knee valgus. Nevertheless, these findings confirm that the CMAS is able to identify athletes who generate high peak KAMs, which offers practitioners a cheaper, time-efficient, and field-based applicable screening tool compared to 3D motion analysis using only three high-speed cameras and free video-analysis software.

While screening tools such as the LESS (70, 72), TJA (32, 66), and QASLS (2, 31) are useful for identifying abnormal and “high-risk” jump-landing mechanics, there is mixed evidence whether the examination of landing mechanics can identify athletes with poor cutting mechanics (1, 13, 51, 67). This issue is pertinent for practitioners who work with athletes who participate cutting dominant sports. In addition, the LESS is the only screening tool to have been validated and assessed against 3D motion analysis (69, 72), with no evidence to suggest that the TJA and QASLS is capable of identifying athletes who generate greater multiplanar knee joint loads. Conversely, in the present study, “higher-risk” cutting mechanics and greater multiplanar knee joint loads (Table 3) were demonstrated by subjects

with high CMASs compared to subjects with low CMASs. These “higher-risk” mechanics included greater mean VBF and HBFs, greater KAAs, greater lateral foot plant distances, greater internal foot progression angles, and lower knee flexion ROM (Table 3), with moderate to large effect sizes. Moreover, greater multiplanar knee joint loads (knee flexion, abduction, and internal rotation moments) were also demonstrated by subjects with high CMASs compared to low, with moderate to very large effect sizes (Table 3). This finding is important because combined multiplanar loads strain the ACL to a greater extent compared to uniplanar loading (4, 57, 76). Krosshaug et al. (53) has highlighted the potential difficulties in estimating 3D joint kinematics based on 2D video evaluations of cutting mechanics. Conversely, the results indicate that the raters in the present study were capable of accurately evaluating and identifying aberrant lower-limb and trunk postures during cutting, as confirmed by the measurable difference in 3D kinetics and kinematics between subjects with “high” and “low” CMASs related to the CMAS scoring system (Table 3).

Supporting Jones et al. (44), higher CMASs were associated with greater peak KAMs (Figure 2), and “higher-risk” cutting mechanics were displayed by subjects with high CMASs (Table 3). These findings indicate that higher scores are representative of, in general, poorer cutting technique. The CMAS tool can therefore be useful for practitioners who want to screen and evaluate cutting movement quality to identify potentially “high-risk” athletes (33, 35, 58, 61), so these athletes can be targeted with biomechanical and neuromuscular informed training interventions to reduce potential injury-risk (33, 35, 61). Qualitative screening tools such as the JTA (47), LESS (20, 73), and QASLS (17) have been used to monitor the effectiveness of training interventions on jump-landing or single leg control mechanics; therefore, the CMAS could be used to monitor pre-to-post changes in cutting movement quality in response to training interventions, and is subsequently a recommended future direction of research. However, it is emphasised that lower CMASs do not necessarily equate to optimal or “safe” technique, and practitioners should not only focus on total score, but focus on the CMAS criteria where athletes scored deficits (27, 44). For example, an athlete who scores 2-3 points may still display “high-risk” cutting deficits such as knee valgus, lateral trunk flexion, limited knee flexion, or hip internal rotation and thus, would still warrant specific injury-risk mitigation training and conditioning. As such, practitioners should be cautious and are advised to look beyond the total CMAS score and use the CMAS tool to assist in the identification of potentially “high-risk” cutting deficits. The information attained from the CMAS may help inform the future prescription of training and conditioning to correct these deficits, and thus potential injury risk (33, 35, 61).

Although a plethora of investigations have focused on COD biomechanics associated with increased risk of injury and have identified a range of factors linked to knee joint loading (Table 1) (19, 28, 29, 38, 40, 41, 50, 59, 77, 78, 83), technical guidelines for coaching safer side-step cutting are limited. A unique aspect of the CMAS is that the criteria (Table 1) can be used as a technical framework for coaching safer side-step cutting which practitioners can use when working with their athletes (44).

COD technique modification has been shown to be an effective modality for reducing high-risk mechanics and knee joint loading during COD (16, 18). Consequently, using the CMAS as a screening tool and a technical framework for safer cutting could be a viable strategy which coaches and practitioners could use to identify specific “high-risk” cutting deficits (i.e. lateral trunk flexion, knee valgus) to help inform preventative COD technique modification training.

It is worth noting, however, that some of the “high-risk” cutting deficits may be needed for faster cutting performance (16, 24, 29). For example, a wide lateral foot plant is needed to generate medio-lateral propulsive force and impulse (29, 40), thus subsequent exit velocity; however, this technique concurrently elevates peak KAMs (19, 40, 50). Limited knee flexion and motion is associated with potentially shorter GCTs (16, 24), but this posture increases KAMs (50, 83), knee flexor joint loads and GRFs (16, 87), thus potential injury-risk (24). Moreover, lateral trunk flexion, from an attacking and evasive perspective, may be performed to feint and deceive opponents (10), but is a critical factor that augments potentially hazardous KAMs (19, 36). Consequently, practitioners should acknowledge the trade-off between knee joint loading (injury-risk) and performance when screening cutting mechanics, because some of the high-risk deficits demonstrated could be effective for performance. Nonetheless, practitioners should ensure that their athletes’ have the physical capacity (i.e. neuromuscular control, co-contraction, and rapid force production) to tolerate the knee joint loading demands of side-steps (40, 56, 71). Further research is required to improve our understanding of the potential performance-injury conflict during cutting (24).

## **5. LIMITATIONS**

It should be acknowledged that, due to the multiplanar nature of side-step cutting (7), some athletes pre-rotate towards the direction of travel during weight acceptance of the cut (77). **This pre-rotation can potentially result in parallax error because the athlete is not perpendicular to the cameras which can restrict evaluations of particular CMAS criteria using the frontal plane and 45° cameras.** Additionally, the current study only investigated a side-step cutting action; thus, the CMAS screening tool is specific to side-step cutting only. Specific screening tools must be developed and validated for assessing other COD actions, such as crossover cuts and pivots, which are also performed and associated with injury in multidirectional sport (14, 39). However, side-step cutting appears to be the predominant COD action associated with non-contact ACL injury (14, 62); therefore, highlighting the importance and inclusion of side-step cutting screening tools (CMAS) in testing batteries for athletes who participate in cutting sports, such as soccer, rugby, handball, American football, and badminton. **Furthermore, the intra- and inter-rater reliability, generally, was moderate to excellent (Table 2), but limited to biomechanists and strength and conditioning coaches. Further work is required to establish agreements and reliability between different applied practitioners, such as sports rehabilitators, physiotherapists, AND sports coaches, in order to confirm its efficacy in the field. Finally, a pre-planned cutting task was used in the**

present study; however, results of previous research have shown that unplanned side-stepping results in greater knee joint loads, more abnormal mechanics, and less muscle support to counteract the greater loads compared to pre-planned side-stepping (5, 6, 11).

## 6. CONCLUSION

In conclusion, a very large significant relationship was observed between CMAS and peak KAM, and “higher-risk” cutting mechanics associated with greater knee joint loading were displayed by subjects with “high” CMASs (~7) compared to subjects with “low” CMASs (~3). As such, the CMAS is a valid and reliable screening tool for evaluating side-step cutting movement quality and offers practitioners a cost-effective and easily applicable field-based screening tool to identify athletes who generate high peak KAMs during side-step cutting. Practitioners should therefore consider including the CMAS in their fitness and testing batteries when screening and profiling athletes who participate in multidirectional sports. Equally, the CMAS allows practitioners to identify “high-risk” cutting deficits in athletes and subsequently create an “injury-risk profile”. These identified deficits can be targeted and addressed through biomechanical and neuromuscular informed training interventions. Finally, the CMAS can be used as a potential technical framework for coaching “safer” cutting.

## REFERENCES

1. Alenezi F, Herrington L, Jones P, and Jones R. Relationships between lower limb biomechanics during single leg squat with running and cutting tasks: a preliminary investigation. *Br J Sports Med* 48: 560-561, 2014.
2. Almangoush A, Herrington L, and Jones R. A preliminary reliability study of a qualitative scoring system of limb alignment during single leg squat. *Physical Therapy and Rehabilitation* 1: 2, 2014.
3. Bahr R. Why screening tests to predict injury do not work and probably never will: a critical review. *Br J Sports Med* 50: 776-780, 2016.
4. Bates NA, Myer GD, Shearn JT, and Hewett TE. Anterior cruciate ligament biomechanics during robotic and mechanical simulations of physiologic and clinical motion tasks: a systematic review and meta-analysis. *Clin Biomech* 30: 1-13, 2015.
5. Besier TF, Lloyd DG, and Ackland TR. Muscle activation strategies at the knee during running and cutting maneuvers. *Med Sci Sports Exerc* 35: 119-127, 2003.
6. Besier TF, Lloyd DG, Ackland TR, and Cochrane JL. Anticipatory effects on knee joint loading during running and cutting maneuvers. *Med Sci Sports Exerc* 33: 1176-1181, 2001.
7. Besier TF, Lloyd DG, Cochrane JL, and Ackland TR. External loading of the knee joint during running and cutting maneuvers. *Med Sci Sports Exerc* 33: 1168-1175, 2001.
8. Bittencourt NFN, Meeuwisse WH, Mendonça LD, Nettel-Aguirre A, Ocarino JM, and Fonseca ST. Complex systems approach for sports injuries: moving from risk factor identification to injury pattern recognition-narrative review and new concept. *Br J Sports Med* 50: 1309-1314, 2016.
9. Boden BP, Dean GS, Feagin JA, and Garrett WE. Mechanisms of anterior cruciate ligament injury. *Orthopedics* 23: 573-578, 2000.
10. Brault S, Bideau B, Kulpa R, and Craig CM. Detecting deception in movement: the case of the side-step in rugby. *PloS one* 7: e37494, 2012.
11. Brown SR, Brughelli M, and Hume PA. Knee mechanics during planned and unplanned sidestepping: a systematic review and meta-analysis. *Sports Med* 44: 1573-1588, 2014.

12. Camomilla V, Cereatti A, Cutti AG, Fantozzi S, Stagni R, and Vannozzi G. Methodological factors affecting joint moments estimation in clinical gait analysis: a systematic review. *Biomedical engineering online* 16: 106, 2017.
13. Chinnasee C, Weir G, Sasimontonkul S, Alderson J, and Donnelly C. A Biomechanical Comparison of Single-Leg Landing and Unplanned Sidestepping. *Int J Sports Med* 39: 636-645, 2018.
14. Cochrane JL, Lloyd DG, Buttfeld A, Seward H, and McGivern J. Characteristics of anterior cruciate ligament injuries in Australian football. *J Sci Med Sport* 10: 96-104, 2007.
15. Cortes N and Onate J. Clinical assessment of drop-jump landing for determination of risk for knee injury. *International Journal of Athletic Therapy and Training* 18: 10-13, 2013.
16. Dai B, William EG, Michael TG, Darin AP, Robin MQ, and Bing Y. The Effects of 2 Landing Techniques on Knee Kinematics, Kinetics, and Performance During Stop-Jump and Side-Cutting Tasks. *Am J Sport Med* 43: 466-474, 2014.
17. Dawson SJ and Herrington L. Improving Single-Legged "Squat Performance: Comparing 2 Training Methods With Potential Implications for Injury Prevention. *J Athl Training* 50: 921-929, 2015.
18. Dempsey AR, Lloyd DG, Elliott BC, Steele JR, and Munro BJ. Changing sidestep cutting technique reduces knee valgus loading. *Am J Sport Med* 37: 2194-2200, 2009.
19. Dempsey AR, Lloyd DG, Elliott BC, Steele JR, Munro BJ, and Russo KA. The effect of technique change on knee loads during sidestep cutting. *Med Sci Sports Exerc* 39: 1765-1773, 2007.
20. Distefano LJ, Marshall SW, Padua DA, Peck KY, Beutler AI, de la Motte SJ, Frank BS, Martinez JC, and Cameron KL. The effects of an injury prevention program on landing biomechanics over time. *Am J Sport Med* 44: 767-776, 2016.
21. Dos'Santos T, Comfort P, and Jones PA. Average of trial peaks versus peak of average profile: impact on change of direction biomechanics. *Sport Biomech*: 1-10, 2018.
22. Faul F, Erdfelder E, Buchner A, and Lang A-G. Statistical power analyses using G\* Power 3.1: Tests for correlation and regression analyses. *Behav Res Methods* 41: 1149-1160, 2009.
23. Fort-Vanmeerhaeghe A, Montalvo AM, Lloyd RS, Read P, and Myer GD. Intra-and inter-rater reliability of the modified tuck jump assessment. *J Sport Sci Med* 16: 117, 2017.
24. Fox A. Change-of-Direction Biomechanics: Is What's Best for Anterior Cruciate Ligament Injury Prevention Also Best for Performance? *Sports Med* 48: 1799-1807, 2018.
25. Fox A, Spittle M, Otago L, and Saunders N. Offensive agility techniques performed during international netball competition. *Int J Sports Sci Coach* 9: 543-552, 2014.
26. Fox AS, Bonacci J, McLean SG, and Saunders N. Efficacy of ACL injury risk screening methods in identifying high-risk landing patterns during a sport-specific task. *Scand J Med Sci Spor* 27: 525-534, 2017.
27. Fox AS, Bonacci J, McLean SG, Spittle M, and Saunders N. A Systematic Evaluation of Field-Based Screening Methods for the Assessment of Anterior Cruciate Ligament (ACL) Injury Risk. *Sports Med*: 1-21, 2015.
28. Frank B, Bell DR, Norcross MF, Blackburn JT, Goerger BM, and Padua DA. Trunk and hip biomechanics influence anterior cruciate loading mechanisms in physically active participants. *Am J Sport Med* 41: 2676-2683, 2013.
29. Havens K and Sigward SM. Cutting mechanics: relation to performance and anterior cruciate ligament injury risk. *Med Sci Sports Exerc* 47: 818-824, 2015.
30. Hedges L and Olkin I. *Statistical Methods for Meta-Analysis*. New York: Academic Press, 1985.
31. Herrington L, Myer G, and Horsley I. Task based rehabilitation protocol for elite athletes following anterior cruciate ligament reconstruction: a clinical commentary. *Phys Ther Sport* 14: 188-198, 2013.
32. Herrington L, Myer GD, and Munro A. Intra and inter-tester reliability of the tuck jump assessment. *Phys Ther Sport* 14: 152-155, 2013.

- 1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103
33. Herrington LC, Munro AG, and Jones PA. Assessment of factors associated with injury risk. *Performance Assessment in Strength and Conditioning*: 53-95, 2018.
  34. Hewett T. Preventive biomechanics: A paradigm shift with a translational approach to biomechanics. *Am J Sports Med* 45: 2654-2664, 2017.
  35. Hewett TE. Response to: 'Why screening tests to predict injury do not work -and probably never will...: a critical review'. *Br J Sports Med* 50: 1353-1353, 2016.
  36. Hewett TE, Torg JS, and Boden BP. Video analysis of trunk and knee motion during non-contact anterior cruciate ligament injury in female athletes: lateral trunk and knee abduction motion are combined components of the injury mechanism. *Br J Sports Med* 43: 417-422, 2009.
  37. Hopkins WG. A scale of magnitudes for effect statistics. *A new view of statistics*: Retrieved from <http://sportsci.org/resource/stats/effectmag.html>, 2002.
  38. Jamison ST, Pan X, and Chaudhari AMW. Knee moments during run-to-cut maneuvers are associated with lateral trunk positioning. *J Biomech* 45: 1881-1885, 2012.
  39. Johnston JT, Mandelbaum BR, Schub D, Rodeo SA, Matava MJ, Silvers HJ, Cole BJ, ElAttrache NS, McAdams TR, and Brophy RH. Video analysis of anterior cruciate ligament tears in professional American football athletes. *Am J Sport Med* 46: 862-868, 2018.
  40. Jones P, Herrington L, and Graham-Smith P. Technique determinants of knee joint loads during cutting in female soccer players. *Hum Movement Sci* 42: 203-211, 2015.
  41. Jones P, Herrington L, and Graham-Smith P. Braking characteristics during cutting and pivoting in female soccer players. *J Electromyogr Kines* 30: 46-54, 2016.
  42. Jones P, Herrington L, and Graham-Smith P. Technique determinants of knee abduction moments during pivoting in female soccer players. *Clin Biomech* 31: 107-112, 2016.
  43. Jones P, Thomas C, Dos'Santos T, McMahon J, and Graham-Smith P. The Role of Eccentric Strength in 180° Turns in Female Soccer Players. *Sports* 5: 42, 2017.
  44. Jones PA, Donelon T, and Dos' Santos T. A preliminary investigation into a qualitative assessment tool to identify athletes with high knee abduction moments during cutting: Cutting Movement Assessment Score (CMAS). *Professional Strength and Conditioning*, 2017.
  45. Jones PA, Herrington LC, Munro AG, and Graham-Smith P. Is there a relationship between landing, cutting, and pivoting tasks in terms of the characteristics of dynamic valgus? *Am J Sport Med* 42: 2095-2102, 2014.
  46. Kimura Y, Ishibashi Y, Tsuda E, Yamamoto Y, Tsukada H, and Toh S. Mechanisms for anterior cruciate ligament injuries in badminton. *Br J Sports Med* 44: 1124-1127, 2010.
  47. Klugman MF, Brent JL, Myer GD, Ford KR, and Hewett TE. Does an in-season only neuromuscular training protocol reduce deficits quantified by the tuck jump assessment? *Clinics in sports medicine* 30: 825-840, 2011.
  48. Koga H, Nakamae A, Shima Y, Iwasa J, Myklebust G, Engebretsen L, Bahr R, and Krosshaug T. Mechanisms for noncontact anterior cruciate ligament injuries knee joint kinematics in 10 injury situations from female team handball and basketball. *Am J Sport Med* 38: 2218-2225, 2010.
  49. Koo TK and Li MY. A guideline of selecting and reporting intraclass correlation coefficients for reliability research. *Journal of chiropractic medicine* 15: 155-163, 2016.
  50. Kristianslund E, Faul O, Bahr R, Myklebust G, and Krosshaug T. Sidestep cutting technique and knee abduction loading: implications for ACL prevention exercises. *Br J Sports Med* 48: 779-783, 2014.
  51. Kristianslund E and Krosshaug T. Comparison of drop jumps and sport-specific sidestep cutting: implications for anterior cruciate ligament injury risk screening. *Am J Sport Med* 41: 684-688, 2013.
  52. Kristianslund E, Krosshaug T, and van den Bogert AJ. Effect of low pass filtering on joint moments from inverse dynamics: implications for injury prevention. *J Biomech* 45: 666-671, 2012.

- 1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162
53. Krosshaug T, Nakamae A, Boden B, Engebretsen L, Smith G, Slauterbeck J, Hewett TE, and Bahr R. Estimating 3D joint kinematics from video sequences of running and cutting maneuvers—assessing the accuracy of simple visual inspection. *Gait Posture* 26: 378-385, 2007.
  54. Krosshaug T, Nakamae A, Boden BP, Engebretsen L, Smith G, Slauterbeck JR, Hewett TE, and Bahr R. Mechanisms of anterior cruciate ligament injury in basketball video analysis of 39 cases. *Am J Sport Med* 35: 359-367, 2007.
  55. Landis JR and Koch GG. The measurement of observer agreement for categorical data. *biometrics*: 159-174, 1977.
  56. Lloyd DG and Buchanan TS. Strategies of muscular support of varus and valgus isometric loads at the human knee. *J Biomech* 34: 1257-1267, 2001.
  57. Markolf KL, Burchfield DM, Shapiro MM, Shepard MF, Finerman GAM, and Slauterbeck JL. Combined knee loading states that generate high anterior cruciate ligament forces. *J Orthop Res* 13: 930-935, 1995.
  58. McCunn R and Meyer T. Screening for risk factors: if you liked it then you should have put a number on it. *Br J Sports Med* 50: 1354, 2016.
  59. McLean SG, Huang X, and van den Bogert AJ. Association between lower extremity posture at contact and peak knee valgus moment during sidestepping: implications for ACL injury. *Clin Biomech* 20: 863-870, 2005.
  60. McLean SG, Walker K, Ford KR, Myer GD, Hewett TE, and van den Bogert AJ. Evaluation of a two dimensional analysis method as a screening and evaluation tool for anterior cruciate ligament injury. *Br J Sports Med* 39: 355-362, 2005.
  61. Mok K-M and Leow R-S. Measurement of movement patterns to enhance ACL injury prevention-A dead end? *J Sports Med Arthrosc Rehabil Techno* 5: 13-16, 2016.
  62. Montgomery C, Blackburn J, Withers D, Tierney G, Moran C, and Simms C. Mechanisms of ACL injury in professional rugby union: a systematic video analysis of 36 cases. *British Journal of Sports Medicine*: bjsports-2016-096425, 2016.
  63. Montgomery C, Blackburn J, Withers D, Tierney G, Moran C, and Simms C. Mechanisms of ACL injury in professional rugby union: a systematic video analysis of 36 cases. *Br J Sports Med* 52: 944-1001, 2018.
  64. Munro AG, Herrington L, and Comfort P. The Relationship Between 2-Dimensional Knee-Valgus Angles During Single-Leg Squat, Single-Leg-Land, and Drop-Jump Screening Tests. *Journal of Sports Rehabilitation* 26: 72-77, 2017.
  65. Myer GD, Brent JL, Ford KR, and Hewett TE. Real-time assessment and neuromuscular training feedback techniques to prevent ACL injury in female athletes. *Strength Cond J* 33: 21, 2011.
  66. Myer GD, Ford KR, and Hewett TE. Tuck jump assessment for reducing anterior cruciate ligament injury risk. *Athletic therapy today: the journal for sports health care professionals* 13: 39, 2008.
  67. O'Connor KM, Monteiro SK, and Hoelker IA. Comparison of selected lateral cutting activities used to assess ACL injury risk. *Journal of applied biomechanics* 25: 9-21, 2009.
  68. Olsen O-E, Myklebust G, Engebretsen L, and Bahr R. Injury mechanisms for anterior cruciate ligament injuries in team handball a systematic video analysis. *Am J Sport Med* 32: 1002-1012, 2004.
  69. Onate J, Cortes N, Welch C, and Van Lunen B. Expert versus novice interrater reliability and criterion validity of the landing error scoring system. *Journal of Sport Rehabilitation* 19: 41-56, 2010.
  70. Padua DA, Boling MC, DiStefano LJ, Onate JA, Beutler AI, and Marshall SW. Reliability of the landing error scoring system-real time, a clinical assessment tool of jump-landing biomechanics. *Journal of Sport Rehabilitation* 20: 145-156, 2011.



- 1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221
71. Padua DA, DiStefano LJ, Hewett TE, Garrett WE, Marshall SW, Golden GM, Shultz SJ, and Sigward SM. National Athletic Trainers' Association Position Statement: Prevention of Anterior Cruciate Ligament Injury. *J Athl Training* 53: 5-19, 2018.
  72. Padua DA, Marshall SW, Boling MC, Thigpen CA, Garrett WE, and Beutler AI. The landing error scoring system (LESS) is a valid and reliable clinical assessment tool of jump-landing biomechanics the JUMP-ACL study. *Am J Sport Med* 37: 1996-2002, 2009.
  73. Parsons JL, Sylvester R, and Porter MM. The effect of strength training on the jump-landing biomechanics of young female athletes: results of a randomized controlled trial. *Clin J Sport Med* 27: 127-132, 2017.
  74. Quatman CE, Quatman-Yates CC, and Hewett TE. A 'Plane'Explanation of Anterior Cruciate Ligament Injury Mechanisms. *Sports Med* 40: 729-746, 2010.
  75. Roewer BD, Ford KR, Myer GD, and Hewett TE. The 'impact' of force filtering cut-off frequency on the peak knee abduction moment during landing: artefact or 'artifiction'? *Br J Sports Med* 48: 464-468, 2014.
  76. Shin CS, Chaudhari AM, and Andriacchi TP. Valgus plus internal rotation moments increase anterior cruciate ligament strain more than either alone. *Med Sci Sports Exerc* 43: 1484-1491, 2011.
  77. Sigward SM, Cesar GM, and Havens KL. Predictors of frontal plane knee moments during side-step cutting to 45 and 110 degrees in men and women: implications for anterior cruciate ligament injury. *Clin J Sport Med* 25: 529-534, 2015.
  78. Sigward SM and Powers CM. Loading characteristics of females exhibiting excessive valgus moments during cutting. *Clin Biomech* 22: 827-833, 2007.
  79. Stuelcken MC, Mellifont DB, Gorman AD, and Sayers MG. Mechanisms of anterior cruciate ligament injuries in elite women's netball: a systematic video analysis. *J Sports Sci* 34: 1516-1522, 2016.
  80. Vanrenterghem J, Gormley D, Robinson M, and Lees A. Solutions for representing the whole-body centre of mass in side cutting manoeuvres based on data that is typically available for lower limb kinematics. *Gait Posture* 31: 517-521, 2010.
  81. Viera AJ and Garrett JM. Understanding interobserver agreement: the kappa statistic. *Fam Med* 37: 360-363, 2005.
  82. Walden M, Krosshaug T, Bjerneboe J, Andersen TE, Faul O, and Hagglund M. Three distinct mechanisms predominate in non-contact anterior cruciate ligament injuries in male professional football players: a systematic video analysis of 39 cases. *Br J Sports Med* 49: 1452-1460, 2015.
  83. Weir G, Alderson J, Smailes N, Elliott B, and Donnelly C. A Reliable Video-based ACL Injury Screening Tool for Female Team Sport Athletes. *Int J Sports Med*, 2019.
  84. Wheeler KW, Askew CD, and Sayers MG. Effective attacking strategies in rugby union. *Eur J Sports Sci* 10: 237-242, 2010.
  85. Winter DA. *Biomechanics and motor control of human motion*. New York: Wiley-Interscience, 1990.
  86. Winter DA. *Biomechanics and motor control of human movement*. John Wiley & Sons, 2009.
  87. Zhang S-N, Bates BT, and Dufek JS. Contributions of lower extremity joints to energy dissipation during landings. *Med Sci Sports Exerc* 32: 812-819, 2000.

1  
2  
3 **Declarations of interest**  
4

5  
6 None declared  
7

8  
9 **Conflict of Interest**  
10

11  
12 None declared.  
13

14 **Funding**  
15

16 This research did not receive any specific grant from funding  
17 agencies in the public, commercial, or not-for-profit sectors.  
18  
19

20 **Ethical approval**  
21

22 Ethical Approval for the study was provided by the Ethics committee at the University of X  
23 (HSR1617-02). The work described has been carried out in accordance with The Code of Ethics  
24 of the World Medical Association (Declaration of Helsinki). All subjects provided written  
25 informed consent prior to participating in the study.  
26  
27  
28  
29  
30

31 **Acknowledgements**  
32

33 The authors would like to thank all participants for partaking in this study.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59

1  
2  
3 **Ethical statement**  
4

5 Ethical Approval for the study was provided by the Ethics committee at the University of X  
6  
7 (HSR1617-02). The work described has been carried out in accordance with The Code of Ethics  
8  
9 of the World Medical Association (Declaration of Helsinki). All subjects provided written  
10  
11 informed consent prior to participating in the study.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59

**Supplement 1. Cutting movement assessment score: CMAS operational definitions and biomechanical rationale**

Suggested viewing camera	CMAS variable	Observation and score	Operational Definition	Biomechanical rationale
<u>Penultimate contact</u>				
<i>Side / 45°</i>	Clear PFC braking strategy (at initial contact) <ul style="list-style-type: none"> <li>• Backward inclination of the trunk</li> <li>• Large COM to COP position - anterior placement of the foot</li> <li>• Effective deceleration – heel contact PFC</li> </ul>	Y/N  Y=0/ N=1	If the subject does not demonstrate a clear PFC braking strategy that emphasises large anterior placement of the foot relative to the centre of mass and does not demonstrate backwards trunk inclination (relative to a vertical straight line), then a score (+1) is awarded  Practitioners may consider referring to a vertical straight line for evaluating trunk inclination.	COD is multistep action with evidence to suggest that the PFC is involved in deceleration prior to directional change, and is a 'preparatory step' (1, 20). A 'large anterior placement of the foot relative to COM and backward inclination of the trunk relative to planted foot' is considered to increase horizontal braking forces during PFC, based on research demonstrating a relationship between average horizontal GRF during PFC and peak KAMs during FFC (20). Reducing the majority of momentum during PFC, will reduce the braking requirements of the FFC, which may result in lower knee joint loads and protect against injury (19-21).
<u>Final Contact</u>				
<i>Front / 45°</i>	Wide lateral leg plant (approx. > 35 cm – dependent on subject anthropometrics) (at initial contact)	Y/N  Y=2/N=0	If the subject demonstrates a wide lateral foot leg plant, then a score (+2) is awarded. A wide foot plant is considered as a distance > 0.35 m) between the hip and plant foot contact IC; however, this is dependent on subject anthropometrics. Practitioners assessing athletes of small stature and leg length may change this accordingly (i.e. > 0.25 m for youth athletes)	A 'wide lateral leg plant' is a major determinant of peak KAM (8, 13, 19, 24). A wide foot plant creates a GRF vector acting laterally relative to the knee, whereby greater foot plant distances creating a greater moment arm (relative to knee joint centre) and thus, KAM.  Abducted hip positions are also commonly observed characteristics displayed during visual inspection of non-contact ACL injuries during COD actions (5, 12, 18, 22).
<i>Front / 45°</i>	Hip in an initial internally rotated position (at initial contact)	Y/N  Y=1/N=0	If the subject's femur is in an internally rotated position at initial contact, then a score (+1) is awarded.	Internal hip rotation can lead to a more medially positioned knee relative to the GRF vector, thus increase moment arm distance and subsequent KAM (13, 28, 39, 41).
<i>Front / 45°</i>	Initial knee 'valgus' position (at initial contact)	Y/N  Y=1/N=0	If the subject's knee is in a valgus (medial) position at initial contact, then a score (+1) is awarded.	'Initial knee valgus position' has been shown to be associated with peak KAM (19, 21, 24, 28, 39). An increased knee abduction angle at initial contact has an effect of placing the knee more medial to the resultant GRF vector and thus, increases the lever arm of the resultant GRF vector relative to the knee joint leading to an increased KAM. Prospective research showed greater valgus angles were associated with increased risk of non-contact ACL injury (14). Increase in knee valgus angle of 2° can lead to a 40 Nm change in valgus moment (29).
<i>All 3</i>	Foot not in neutral foot position (at initial contact)	Y/N	If the subject's foot is not in a neutral position (i.e. approx. 0°) and is inwardly or externally rotated at initial contact (relative to approach direction of travel), then a score (+1) is awarded	Initial foot progression angle is associated with KAM, with a neutral foot position considered the safest strategy (8, 21, 41). Internally rotated foot positions during weight acceptance can lead to a more medially positioned knee relative to the GRF vector, thus increase moment arm distance and subsequent KAM (21, 41). A

42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82

	Inwardly rotated foot position or externally rotated foot position (relative to approach direction of travel)	Y=1/N=0		neutral foot position would most likely result in forces being absorbed in the sagittal plane utilising the large knee and hip extensor musculature, which is potentially a safer strategy (21). Excessive foot external rotation increases susceptibility to eversion and pronation which could lead to knee valgus and tibial internal rotation (10, 26, 33), thus ACL loading. External rotation of the foot has also been stated as characteristics during visual inspection of non-contact ACL injuries during change of direction (18).
<i>Front / 45°</i>	Frontal plane trunk position relative to intended direction; Lateral or trunk rotated towards stance limb (L/TR), Upright (U) or Medial (M) (at initial contact and over WA)  (use shoulder positioning as guide)	L/TR/U/M  L=2/ TR = 2/  U = 1,  /M=0	If the subject's trunk is laterally flexed over the stance (push-off) limb or rotated towards the stance limb at initial contact and over WA, then a score (+2) is awarded.  If the subject's trunk is upright (vertical relative to straight line), then a score (+1) is awarded.  If the subject's trunk is medial (leaning towards the intended direction of travel), then no score is awarded.  Practitioners may consider referring to a vertical straight line and use shoulder position as an indicator.	The trunk contains approx. half of the body's mass, and during cutting the entire body's mass must be balanced and supported on one leg, thus trunk control and positioning is a critical factor influencing knee joint loads (15, 16, 30). Lateral trunk flexion (8, 17, 19) or trunk rotation (8, 11) towards stance limb are major determinants of peak KAM. A laterally flexed trunk or rotated trunk towards the planted leg side shifts the athlete's weight laterally creating a laterally directed force vector, increasing the moment arm relative to the knee joint and thus, KAMs. Prospective research has shown deficits in trunk control and proprioception are associated with increased risk of non-contact ACL injury (48, 49). Lateral trunk flexion over plant leg also a commonly observed visual characteristic of non-contact ACL injuries during plant-and cut manoeuvres and landing (16, 22, 42).
<i>Side / 45°</i>	Trunk upright or leaning back throughout contact over whole contact (not adequate trunk flexion displacement) (at initial contact and over WA)	Y/N  Y=1/N=0	If the subject's trunk is upright or leaning back throughout weight (i.e. appears limited hip flexion) acceptance and push-off during the FFC and does not go through an adequate range of trunk-flexion displacement, then a score (+1) is awarded.	Trunk inclination (leaning back or upright) with minimal trunk flexion displacement during weight acceptance may increase the overall knee joint load due an increased lever arm of the trunk relative to the knee and increasing the COM distance from the base of support (37). Some trunk flexion allows generation of hip moments to help absorb the GRF during weight acceptance and thus, may lower KAMs (21).  Increasing hip flexion and promoting a hip dominant strategy are involved GRF attenuation (38, 40, 47), energy dissipation (35, 46), reducing loading rates (40) and reducing knee joint loads (32, 35, 36, 40) during high impact tasks. Increasing hip flexion increases the moment arm distance at the hip which creates a greater hip flexor moment (utilising hip extensor musculature). This can have the effect of unloading the knee by more evenly distributing loading proximally up the lower-limb chain (21, 35, 36, 40), thus reducing the demands for the knee.
<i>All 3</i>	Limited knee flexion during final contact (stiff) $\leq 30^\circ$	Y/N  Y=1/N=0	If the subject's knee goes through limited knee flexion (approximately $\leq 30^\circ$ ) over weight acceptance appears 'stiff', then a score (+1) is awarded.	Stiffer weight acceptance strategies can increase impact GRFs (7, 9, 50), and greater GRFs are associated with increased KAMs (39, 41). Less knee flexion is also associated with greater KAMs (24, 44). Furthermore, extended knee positions with high anterior tibial loading and shear force can also increase ACL strain (2, 3, 27, 45). Extended knee positions are also commonly observed characteristics of non-contact ACL injury during directional changes and landing (4-6, 12, 18, 22, 23, 25, 31, 34, 43).

Front / 45°	Excessive knee 'valgus' motion during WA	Y/N  Y=1/N=0	If the subject's knee demonstrates visible valgus motion during weight acceptance, then a score (+1) is awarded.	Knee valgus motion during FFC is considered because it is a key indicator of ACL injury risk (14), and can contribute to large front plane knee joint loading. Dynamic knee valgus positions are also commonly observed characteristics of non-contact ACL injury during directional changes and landing (5, 6, 12, 18, 22, 23, 25, 34, 42, 43).
Key: PFC: Penultimate foot contact; FFC: Final foot contact; KAM: peak Knee abduction moment; ACL: Anterior cruciate ligament; IC: Initial contact; WA: Weight acceptance; GRF: Ground reaction force. Y: Yes; N: No; L: Lateral; TR: Trunk rotation; U: Upright; M: Medial.				

## References

- Andrews JR, McLeod WD, Ward T, and Howard K. The cutting mechanism. *Am J Sport Med* 5: 111-121, 1977.
- Beynon B, Howe J, Pope MH, Johnson RJ, and Fleming B. The measurement of anterior cruciate ligament strain in vivo. *International orthopaedics* 16: 1-12, 1992.
- Beynon BD, Fleming BC, Johnson RJ, Nichols CE, Renström PA, and Pope MH. Anterior cruciate ligament strain behavior during rehabilitation exercises in vivo. *Am J Sport Med* 23: 24-34, 1995.
- Boden BP, Dean GS, Feagin JA, and Garrett WE. Mechanisms of anterior cruciate ligament injury. *Orthopedics* 23: 573-578, 2000.
- Brophy RH, Stepan JG, Silvers HJ, and Mandelbaum BR. Defending puts the anterior cruciate ligament at risk during soccer: a gender-based analysis. *Sports health* 7: 244-249, 2015.
- Cochrane JL, Lloyd DG, Buttfield A, Seward H, and McGivern J. Characteristics of anterior cruciate ligament injuries in Australian football. *J Sci Med Sport* 10: 96-104, 2007.
- Dai B, William EG, Michael TG, Darin AP, Robin MQ, and Bing Y. The Effects of 2 Landing Techniques on Knee Kinematics, Kinetics, and Performance During Stop-Jump and Side-Cutting Tasks. *Am J Sport Med* 43: 466-474, 2014.
- Dempsey AR, Lloyd DG, Elliott BC, Steele JR, Munro BJ, and Russo KA. The effect of technique change on knee loads during sidestep cutting. *Med Sci Sports Exerc* 39: 1765-1773, 2007.
- Devita P and Skelly WA. Effect of landing stiffness on joint kinetics and energetics in the lower extremity. *Med Sci Sports Exerc* 24: 108-115, 1992.
- Ford KR, Myer GD, Toms HE, and Hewett TE. Gender differences in the kinematics of unanticipated cutting in young athletes. *Med Sci Sports Exerc* 37: 124-129, 2005.
- Frank B, Bell DR, Norcross MF, Blackburn JT, Goerger BM, and Padua DA. Trunk and hip biomechanics influence anterior cruciate loading mechanisms in physically active participants. *Am J Sport Med* 41: 2676-2683, 2013.
- Grassi A, Smiley SP, Roberti di Sarsina T, Signorelli C, Marcheggiani Muccioli GM, Bondi A, Romagnoli M, Agostini A, and Zaffagnini S. Mechanisms and situations of anterior cruciate ligament injuries in professional male soccer players: a YouTube-based video analysis. *Eur J Orthop Surg Traumatol* 27: 697-981, 2017.
- Havens K and Sigward SM. Cutting mechanics: relation to performance and anterior cruciate ligament injury risk. *Med Sci Sports Exerc* 47: 818-824, 2015.
- Hewett T, Myer GD, Ford KR, Heidt RS, Colosimo AJ, McLean SG, Van den Bogert AJ, Paterno MV, and Succop P. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sport Med* 33: 492-501, 2005.
- Hewett TE and Myer GD. The mechanistic connection between the trunk, knee, and anterior cruciate ligament injury. *Exercise and sport sciences reviews* 39: 161, 2011.
- Hewett TE, Torg JS, and Boden BP. Video analysis of trunk and knee motion during non-contact anterior cruciate ligament injury in female athletes: lateral trunk and knee abduction motion are combined components of the injury mechanism. *Br J Sports Med* 43: 417-422, 2009.
- Jamison ST, Pan X, and Chaudhari AMW. Knee moments during run-to-cut maneuvers are associated with lateral trunk positioning. *J Biomech* 45: 1881-1885, 2012.
- Johnston JT, Mandelbaum BR, Schub D, Rodeo SA, Matava MJ, Silvers HJ, Cole BJ, ElAttrache NS, McAdams TR, and Brophy RH. Video analysis of anterior cruciate ligament tears in professional American football athletes. *Am J Sport Med* 46: 862-868, 2018.
- Jones P, Herrington L, and Graham-Smith P. Technique determinants of knee joint loads during cutting in female soccer players. *Hum Movement Sci* 42: 203-211, 2015.
- Jones P, Herrington L, and Graham-Smith P. Braking characteristics during cutting and pivoting in female soccer players. *J Electromyogr Kines* 30: 46-54, 2016.
- Jones P, Herrington L, and Graham-Smith P. Technique determinants of knee abduction moments during pivoting in female soccer players. *Clin Biomech* 31: 107-112, 2016.
- Kimura Y, Ishibashi Y, Tsuda E, Yamamoto Y, Tsukada H, and Toh S. Mechanisms for anterior cruciate ligament injuries in badminton. *Br J Sports Med* 44: 1124-1127, 2010.
- Koga H, Nakamae A, Shima Y, Iwasa J, Myklebust G, Engebretsen L, Bahr R, and Krosshaug T. Mechanisms for noncontact anterior cruciate ligament injuries knee joint kinematics in 10 injury situations from female team handball and basketball. *Am J Sport Med* 38: 2218-2225, 2010.
- Kristianslund E, Faul O, Bahr R, Myklebust G, and Krosshaug T. Sidestep cutting technique and knee abduction loading: implications for ACL prevention exercises. *Br J Sports Med* 48: 779-783, 2014.
- Krosshaug T, Nakamae A, Boden BP, Engebretsen L, Smith G, Slauterbeck JR, Hewett TE, and Bahr R. Mechanisms of anterior cruciate ligament injury in basketball video analysis of 39 cases. *Am J Sport Med* 35: 359-367, 2007.
- Loudon JK, Jenkins W, and Loudon KL. The relationship between static posture and ACL injury in female athletes. *J Orthop Sport Phys* 24: 91-97, 1996.

124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164

27. Markolf KL, Gorek JF, Kabo JM, and Shapiro MS. Direct measurement of resultant forces in the anterior cruciate ligament. An in vitro study performed with a new experimental technique. *J Bone Joint Surg Am* 72: 557-567, 1990.
28. McLean SG, Huang X, and van den Bogert AJ. Association between lower extremity posture at contact and peak knee valgus moment during sidestepping: implications for ACL injury. *Clin Biomech* 20: 863-870, 2005.
29. McLean SG, Lipfert SW, and Van den Bogert AJ. Effect of gender and defensive opponent on the biomechanics of sidestep cutting. *Med Sci Sports Exerc* 36: 1008-1016, 2004.
30. Mendiguchia J, Ford KR, Quatman CE, Alentorn-Geli E, and Hewett TE. Sex differences in proximal control of the knee joint. *Sports Med* 41: 541-557, 2011.
31. Montgomery C, Blackburn J, Withers D, Tierney G, Moran C, and Simms C. Mechanisms of ACL injury in professional rugby union: a systematic video analysis of 36 cases. *British Journal of Sports Medicine: bjsports-2016-096425*, 2016.
32. Nguyen A-D, Taylor JB, Wimbish TG, Keith JL, and Ford KR. Preferred Hip Strategy During Landing Reduces Knee Abduction Moment in Collegiate Female Soccer Players. *Journal of Sport Rehabilitation* 20: 1-5, 2018.
33. Nyland J, Caborn DNM, Shapiro R, Johnson DL, and Fang H. Hamstring extensibility and transverse plane knee control relationship in athletic women. *Knee Surgery, Sports Traumatology, Arthroscopy* 7: 257-261, 1999.
34. Olsen O-E, Myklebust G, Engebretsen L, and Bahr R. Injury mechanisms for anterior cruciate ligament injuries in team handball a systematic video analysis. *Am J Sport Med* 32: 1002-1012, 2004.
35. Pollard CD, Sigward SM, and Powers CM. Limited hip and knee flexion during landing is associated with increased frontal plane knee motion and moments. *Clin Biomech* 25: 142-146, 2010.
36. Powers CM. The influence of abnormal hip mechanics on knee injury: a biomechanical perspective. *J Orthop Sport Phys* 40: 42-51, 2010.
37. Sheehan FT, Sipprell Iii WH, and Boden BP. Dynamic sagittal plane trunk control during anterior cruciate ligament injury. *Am J Sport Med* 40: 1068-1074, 2012.
38. Shimokochi Y, Ambegaonkar JP, Meyer EG, Lee SY, and Shultz SJ. Changing sagittal plane body position during single-leg landings influences the risk of non-contact anterior cruciate ligament injury. *Knee Surgery, Sports Traumatology, Arthroscopy* 21: 888-897, 2013.
39. Sigward SM, Cesar GM, and Havens KL. Predictors of frontal plane knee moments during side-step cutting to 45 and 110 degrees in men and women: implications for anterior cruciate ligament injury. *Clin J Sport Med* 25: 529-534, 2015.
40. Sigward SM and Pollard CD. Proximal Risk Factors for ACL Injury: Role of the Hip Joint and Musculature, in: *ACL Injuries in the Female Athlete*. Springer, 2018, pp 207-223.
41. Sigward SM and Powers CM. Loading characteristics of females exhibiting excessive valgus moments during cutting. *Clin Biomech* 22: 827-833, 2007.
42. Stuelcken MC, Mellifont DB, Gorman AD, and Sayers MG. Mechanisms of anterior cruciate ligament injuries in elite women's netball: a systematic video analysis. *J Sports Sci* 34: 1516-1522, 2016.
43. Walden M, Krosshaug T, Bjerneboe J, Andersen TE, Faul O, and Hagglund M. Three distinct mechanisms predominate in non-contact anterior cruciate ligament injuries in male professional football players: a systematic video analysis of 39 cases. *Br J Sports Med* 49: 1452-1460, 2015.
44. Weir G, Alderson J, Smailes N, Elliott B, and Donnelly C. A Reliable Video-based ACL Injury Screening Tool for Female Team Sport Athletes. *Int J Sports Med*, 2019.
45. Withrow TJ, Huston LJ, Wojtys EM, and Ashton-Miller JA. The relationship between quadriceps muscle force, knee flexion, and anterior cruciate ligament strain in an in vitro simulated jump landing. *Am J Sport Med* 34: 269-274, 2006.
46. Yeow CH, Lee PVS, and Goh JCH. Non-linear flexion relationships of the knee with the hip and ankle, and their relative postures during landing. *Knee* 18: 323-328, 2011.
47. Yu B, Lin C-F, and Garrett WE. Lower extremity biomechanics during the landing of a stop-jump task. *Clin Biomech* 21: 297-305, 2006.
48. Zazulak BT, Hewett TE, Reeves NP, Goldberg B, and Cholewicki J. Deficits in Neuromuscular Control of the Trunk Predict Knee Injury Risk: Prospective Biomechanical-Epidemiologic Study. *Am J Sport Med* 35: 1123-1130, 2007.
49. Zazulak BT, Hewett TE, Reeves NP, Goldberg B, and Cholewicki J. The effects of core proprioception on knee injury: a prospective biomechanical-epidemiological study. *Am J Sports Med* 35: 368-373, 2007.
50. Zhang S-N, Bates BT, and Dufek JS. Contributions of lower extremity joints to energy dissipation during landings. *Med Sci Sports Exerc* 32: 812-819, 2000.

**Supplement 2. Definitions and calculations for kinetic and kinematic variables examined during cutting**

	<b>Variable</b>	<b>Foot contact</b>	<b>Abbreviation</b>	<b>Definition or calculation</b>
Sagittal plane knee joint moments	Peak knee flexor moment	FFC	HFM	Peak external joint moments during weight acceptance of FFC using inverse dynamics.
Sagittal plane joint angles	knee flexion angle	FFC	KFA	Derived from the following order of rotations: flexion (+)/extension (-)
Frontal plane injury risk parameters	Peak knee abduction moments	FFC	KAM	Peak external knee abduction moment (+ abduction/- adduction) during weight acceptance phase of FFC using inverse dynamics.
	Knee abduction angle	FFC	KAA	Knee abduction angle (-) during weight acceptance phase of final contact /adduction (+)
Transverse plane injury risk parameters	Peak knee rotation moment	FFC	KRM	Peak external knee rotation moment (+ external/- internal) during weight acceptance phase of final contact using inverse dynamics
	Knee rotation angle	FFC	KRA	Knee rotation angle (- internal/ + external) during weight acceptance phase of FFC
GRF	Peak vertical braking force (Fz)	PFC and FFC	VBF	Peak normalised vGRF (Fz) value during weight acceptance
	Mean vertical braking force (Fz)	PFC and FFC	Mean VBF	Average normalised vGRF (Fz) during weight acceptance
	Peak horizontal braking force (Fx)	PFC and FFC	HBF	Peak normalised hGRF (Fx) value during weight acceptance
	Mean horizontal braking force (Fx)	PFC and FFC	Mean HBF	Average normalised hGRF (Fx) during weight acceptance
	Braking force ratio	Between the two contacts	-	FFC braking force / PFC braking force
Trunk variables	Lateral Trunk flexion	FFC	-	Angle of trunk relative to vertical line perpendicular to the pelvis: (0°) upright / (+) trunk flexion away from plant leg/ (-) trunk flexion towards plant leg
	Trunk inclination angle	PFC and FFC	-	Angle of trunk relative to a vertical straight-line straight line, (+) forward trunk lean/ (-) backward trunk lean
	Hip rotation angle	FFC	-	Femur internally rotated (-)/ external rotation (+)
Hip, pelvis, and foot	Lateral foot plant distance	FFC	-	Lateral distance from initial foot contact of foot COM to proximal end of pelvis – for cutting
	Initial foot progression angle	FFC	IFPA	Angle of foot progression relative to lab coordinate system configuration/ original direction: straight (0°)/inward rotation (+)/outward rotation (-) angle (°)

Key: PFC: Penultimate foot contact; FFC: Final foot contact; COM: Centre of mass; COD: Change of direction; IC: Initial contact; GRF: Ground reaction force; vGRF: Vertical GRF; hGRF: Horizontal GRF;



42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82



# Supplement 2. CMAS MANUAL

Table 1. Cutting movement assessment score tool

Camera	Variable	Observation	Score
<b><u>Penultimate contact</u></b>			
<i>Side / 45°</i>	<b>Clear PFC braking strategy</b> (at initial contact) <ul style="list-style-type: none"> <li>• Backward inclination of the trunk</li> <li>• Large COM to COP position – anterior placement of the foot</li> <li>• Effective deceleration – heel contact PFC</li> </ul>	Y/N	Y=0/ N=1
<b><u>Final Contact</u></b>			
<i>Front / 45°</i>	<b>Wide lateral leg plant</b> (approx. >0.35 m – dependent on subject anthropometrics) (at initial contact)	Y/N	Y=2/N=0
<i>Front / 45°</i>	<b>Hip in an initial internally rotated position</b> (at initial contact)	Y/N	Y=1/N=0
<i>Front / 45°</i>	<b>Initial knee ‘valgus’ position</b> (at initial contact)	Y/N	Y=1/N=0
<i>All 3</i>	<b>Foot not in neutral foot position</b> (at initial contact) Inwardly rotated foot position or externally rotated foot position (relative to original direction of travel)	Y/N	Y=1/N=0
<i>Front / 45°</i>	<b>Frontal plane trunk position relative to intended direction; Lateral or trunk rotated towards stance limb, Upright or Medial.</b> (at initial contact and over WA)	L/TR/U/M	L/TR=2/ U = 1, /M=0
<i>Side / 45°</i>	<b>Trunk upright or leaning back throughout contact</b> (not adequate trunk flexion displacement) - (at initial contact and over WA)	Y/N	Y=1/N=0
<i>Side / 45°</i>	<b>Limited Knee Flexion during final contact (stiff) <math>\leq 30^\circ</math></b> (over WA)	Y/N	Y=1/N=0
<i>Front / 45°</i>	<b>Excessive Knee ‘valgus’ motion during contact</b> (over WA)	Y/N	Y=1/N=0
		<b>Total Score</b>	<b>0 /11</b>

Key: PFC: Penultimate foot contact; COM: Centre of mass; COP: Centre of pressure; WA: weight acceptance; TR: Trunk rotation; Y: Yes; N: No; L: Lateral; TR: Trunk rotation; U: Upright; M: Medial.

Table 1. Cutting movement assessment score tool

Camera	Variable	Observation	Score
<b><u>Penultimate contact</u></b>			
<i>Side / 45°</i>	<b>Clear PFC braking strategy (at initial contact)</b> <ul style="list-style-type: none"> <li>• Backward inclination of the trunk</li> <li>• Large COM to COP position – anterior placement of the foot</li> <li>• Effective deceleration – heel contact PFC</li> </ul>	Y/N	Y=0/ N=1

**Yes**



**No**



37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72

Table 1. Cutting movement assessment score tool

Camera	Variable	Observation	Score
<u>Final Contact</u>			
Front / 45°	Wide lateral leg plant (approx. >0.35 m – dependent on subject anthropometrics) (at initial contact)	Y/N	Y=2/N=0

Yes



No



73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108

Table 1. Cutting movement assessment score tool

Camera	Variable	Observation	Score
<u>Final Contact</u>			
Front / 45°	Hip in an initial internally rotated position (at initial contact)	Y/N	Y=1/N=0

Yes

No



109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144

Table 1. Cutting movement assessment score tool

Camera	Variable	Observation	Score
<u>Final Contact</u>			
Front / 45°	Initial knee 'valgus' position (at initial contact)	Y/N	Y=1/N=0

Yes

No



145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180

Table 1. Cutting movement assessment score tool

Camera	Variable	Observation	Score
<u>Final Contact</u>			
All 3	<b>Foot not in neutral foot position</b> (@ initial contact) Inwardly rotated foot position or externally rotated foot position (relative to original direction of travel)	Y/N	Y=1/N=0

**Yes**

**No**



181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216

Table 1. Cutting movement assessment score tool

Camera	Variable	Observation	Score
<u>Final Contact</u>			
<i>Front / 45°</i>	Frontal plane trunk position relative to intended direction; Lateral or trunk rotated towards stance limb, Upright or Medial. (at initial contact and over WA)	L/TR/U/M	L/TR=2/ U = 1, /M=0

L

TR

U

M



217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252



Table 1. Cutting movement assessment score tool

Camera	Variable	Observation	Score
<u>Final Contact</u>			
Side / 45°	Trunk upright or leaning back throughout contact (not adequate trunk flexion displacement) - (at initial contact and over WA)	Y/N	Y=1/N=0

Yes

No



253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288

Table 1. Cutting movement assessment score tool

Camera	Variable	Observation	Score
<u>Final Contact</u>			
<i>Side / 45°</i>	Limited Knee Flexion during final contact (stiff) $\leq 30^\circ$ (over WA)	Y/N	Y=1/N=0

Yes



No



289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324

Table 1. Cutting movement assessment score tool

Camera	Variable	Observation	Score
<u>Final Contact</u>			
<i>Front / 45°</i>	Excessive Knee 'valgus' motion during contact (over WA)	Y/N	Y=1/N=0

Yes

No



325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360