



University of Dundee

Normal spirometry equates to normal impulse oscillometry in healthy subjects

Lipworth, Brian; Chan, Rory

Published in:
Respiratory Research

DOI:
[10.1186/s12931-021-01693-0](https://doi.org/10.1186/s12931-021-01693-0)

Publication date:
2021

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Lipworth, B., & Chan, R. (2021). Normal spirometry equates to normal impulse oscillometry in healthy subjects. *Respiratory Research*, 22, [96]. <https://doi.org/10.1186/s12931-021-01693-0>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

LETTER TO THE EDITOR

Open Access



Normal spirometry equates to normal impulse oscillometry in healthy subjects

Brian Lipworth*  and Rory Chan

Background

Impulse oscillometry (IOS) is an effort independent test performed with tidal breathing which can ascertain the presence of small airways dysfunction by measuring the heterogeneity of resistance between 5 and 20 Hz (R5–R20) or the area under the reactance curve (AX). The article of Li et al. reported on impulse oscillometry (IOS) values in young non smoking symptomatic otherwise healthy subjects with preserved spirometry without a diagnosis of chronic respiratory disease [1]. In subjects who had forced expiratory flow rates <65% predicted they reported poor area under curve values <0.70 with associated low sensitivity and specificity values on the whole being less than 80%.

In our clinic we would consider pragmatic abnormal IOS values as $R5 > 0.5$ kPa/l.s, $R5-R20 > 0.1$ kPa/l/s and $AX > 1.0$ kPa/l, in terms of being indicative of a clinically relevant degree of airflow obstruction [2]. We would also like to point out that in persistent asthma patients with preserved spirometry who have $FEV1 > 80\%$ predicted, the presence of raised values for R5–R20 was associated with a greater likelihood of long term poor disease control [3]. Moreover small airways dysfunction as R5–R20 and AX is related to type 2 inflammation in asthma [4].

Li et al. have demonstrated that otherwise healthy individuals with symptoms who have abnormal forced expiratory flows also have normal IOS values. Moreover it is important to note that forced expiratory rates are rather volume dependent such that false positive low values are commonly seen when subjects do not exhale fully to

residual volume. We would therefore have concerns that clinicians might end up erroneously interpreting what are essentially normal IOS values which would lead to unnecessary further investigation and cause potential unwarranted anxiety to patients with normal tests.

Conclusions

Impulse oscillometry should be interpreted in the light of abnormal values as being indicative of small airways dysfunction.

Abbreviations

R5: Resistance at 5 Hz (total resistance); R20: Resistance at 20 Hz (central resistance); R5–R20: Resistance heterogeneity between 5 and 20 Hz (peripheral resistance); AX: Area under reactance curve; FEV1: Forced expiratory volume in 1 s; IOS: Impulse oscillometry.

Acknowledgements

Not applicable.

Authors' contributions

BJL and RC were responsible for writing the letter and read and approved the manuscript. All contributors who do not meet the criteria for authorship should be listed in an acknowledgements section. Both authors read and approved the final manuscript.

Funding

Not applicable.

Availability of data and materials

Not applicable.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

*Correspondence: bj.lipworth@dundee.ac.uk
Scottish Centre for Respiratory Research, Ninewells Hospital,
Dundee DD19SY, UK



Competing interests

Drs. Lipworth and Chan have no conflicts of interest to report.

Received: 10 March 2021 Accepted: 23 March 2021

Published online: 31 March 2021

References

1. Li LY, Yan TS, Yang J, Li YQ, Fu LX, Lan L, et al. Impulse oscillometry for detection of small airway dysfunction in subjects with chronic respiratory symptoms and preserved pulmonary function. *Respir Res*. 2021;22:68.
2. Lipworth B, Manoharan A, Anderson W. Unlocking the quiet zone: the small airway asthma phenotype. *Lancet Respir Med*. 2014;2:497–506.
3. Manoharan A, Anderson WJ, Lipworth J, Ibrahim I, Lipworth BJ. Small airway dysfunction is associated with poorer asthma control. *EurRespir J*. 2014;44:1353–5.
4. Kuo CR, Jabbal S, Lipworth B. Is small airways dysfunction related to asthma control and type 2 inflammation? *Ann Allergy Asthma Immunol*. 2018;121:631–2.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

