Body temperature and physical activity correlates of the menstrual cycle in female chacma baboons (*Papio hamadryas ursinus*).

Trevor Tapiwa Nyakudya

A dissertation submitted to the Faculty of Health Sciences, University of the Witwatersrand, in fulfillment of the requirements for the degree of Master of Science in Medicine.

Johannesburg, South Africa, 2010

DECLARATION

I, Trevor Tapiwa Nyakudya, declare that the work contained in this dissertation is my own, except where acknowledged as otherwise. This dissertation is being submitted for the Degree of Master of Science in Medicine in the Faculty of Health Sciences at the University of the Witwatersrand. The work herein has not been submitted before for any degree or examination at any other university.

Trevor Tapiwa Nyakudya

Signed on the______ day of______, 2010

I certify that all the experimental procedures used in this dissertation were approved by the Animals Ethics Screening Committee of the University of the Witwatersrand (AESC number: 2007/60/5).

CONFERENCE PROCEEDINGS AND PRESENTATIONS

The following oral and poster presentations are offered in support of this dissertation.

1) **Nyakudya TT**, Fuller A, Meyer LCR and Mitchell D. Body temperature correlates of the menstrual cycle in baboons, *Papio hamadryas*. International Brain Research Organisation (IBRO) IBRO/UNESCO/ISN African Neuroscience School of Chronobiology. Theme "Brain and Clocks: Neurochemical, Pharmacological and Therapeutic Approaches" Ourzazate-Zagora, Morocco, 20-27 October 2008. (Oral presentation).

2) **Nyakudya TT**, Fuller A, Meyer LCR, Maloney SK and Mitchell D. Body temperature and activity correlates of the menstrual cycle in baboons, *Papio hamadryas*. 3rd International Symposium on Physiology and Pharmacology of Temperature Regulation 2009 (PPTR2009) Matsue, Izumo, Japan, 23-26 July 2009. (Poster presentation).

3) **Nyakudya TT**, Fuller A, Meyer LCR, Maloney SK and Mitchell D. Body temperature and activity correlates of the menstrual cycle in baboons, *Papio hamadryas.* The 36th International Congress of Physiological Sciences (IUPS) "Function of Life: Elements

and Integration" Kyoto International Conference Center (ICC Kyoto). Takaragaike, Sakyo-ka, Kyoto, Japan, 27 July to 01 August 2009. (Poster presentation).

4) **Nyakudya TT**, Fuller A, Meyer LCR, Maloney SK and Mitchell D. Correlates of the menstrual cycle in non-human primates. The 20th IBRO/ARC School in Neuroscience in Africa. Nemba Training Hospital, Gekenke, Rwanda, 31 August-05 September 2009. (Oral presentation).

5) **Nyakudya TT**, Fuller A, Meyer LCR, Maloney SK and Mitchell D. Body temperature and activity correlates of the menstrual cycle in baboons, *Papio hamadryas ursinus*. The 37th Annual Congress of the Physiological Society of Southern Africa. University of Stellenbosch, Stias Conference Centre, South Africa, 09-12 September 2009. (Oral presentation).

6) **Nyakudya TT**, Fuller A, Meyer LCR, Maloney SK and Mitchell D. Body temperature and activity correlates of the menstrual cycle in baboons, *Papio hamadryas ursinus*. The 2nd Cross Faculty Postgraduate Symposium. University of the Witwatersrand, South Africa, 20-21 October 2009. (Poster presentation).

ABSTRACT

I investigated the relationship between abdominal temperature, physical activity, anogenital swellings, and faecal and urine ovarian steroid hormonal concentrations over the menstrual cycle in baboons in an attempt to devise a reliable non-hormonal physiological indicator to detect ovulation. Using a miniature thermometric data logger surgically implanted in the abdominal cavity and an activity data logger implanted subcutaneously on the trunk, I measured, continuously over six months at a 10 min interval, abdominal temperature and physical activity patterns in four female adult baboons, Papio hamadryas ursinus (12.9-19.9 kg), unrestrained in cages in an indoor animal facility (22-25°C). I monitored menstrual bleeding, and anogenital swelling changes using digital photography, and collected urine and faeces, daily, to ascertain the stage and length of the menstrual cycle. The length of the menstrual cycle, determined from daily observations of menstrual bleeding and anogenital swellings, was 36 ± 2 days (mean \pm SD). Baboons exhibited a cyclic change in anogenital swellings, abdominal temperature, physical activity, urine and faecal steroid hormones over the menstrual cycle. Mean 24-h abdominal temperature during the luteal phase was significantly higher (ANOVA, p = 0.04; $F_{(2,9)} = 4.7$) than during the ovulatory phase, but not different to the follicular phase. Physical activity also followed a similar pattern, with mean 24 h physical activity almost twice as high in the luteal than in the ovulatory phase (ANOVA, p = 0.58; F_(2,12) = 5.8). As expected, urine and faecal oestradiol was higher in the follicular than in the luteal phase, while progesterone was higher in the luteal than the follicular phase. Cortisol in both urine and faecal samples did not show any

recognisable menstrual cycle related pattern. I have characterised correlates of the menstrual cycle in baboons and shown, for the first time, a rhythm of physical activity over the baboon menstrual cycle. I have also shown, from the measurements of abdominal temperature, physical activity, ovarian steroid hormonal concentrations and anogenital swellings, that ovulation in captive unrestrained baboons, and probably also free-living baboons, can be estimated from anogenital swellings or possibly abdominal temperature and physical activity, without the need for hormone measurements.

ACKNOWLEDGEMENTS

My dissertation is a result of many hours of hard work and the invaluable assistance of several people. I would therefore like to extend my sincere gratitude to my supervisors, Professor Duncan Mitchell, Professor Andrea Fuller and Dr. Leith Meyer for their professional guidance and continued support over the entire period that I was pursuing my studies. My sincere gratitude also goes to Professor Peter Henzi for suggesting the idea of identifying ovulation remotely in order to establish paternity.

I would also like to thank the staff of the Central Animal Services for their husbandry of the baboons and assistance during surgeries. Special thanks go to Professor Shane Maloney for the statistical advice, encouragement and setting up the telemetry system, Mrs. Marie Lawson for doing the radioimmunoassays for the samples, and Mr. Floyd Olsen of the National Health Laboratory Services for doing the urine creatinine assays. Finally I would like to thank my family and friends who have been there for me through the challenging times. I will always appreciate the support, patience and love that they gave me all the way. Many thanks go to Lavender Chaparadza for being there for me always and keeping me sane.

To God our heavenly Father I give glory and honor for the gift of life and all the other blessings.

vii

The work in this dissertation was funded by grants from Medical Faculty Research Endowment Fund of the University of the Witwatersrand and National Research Foundation Grants to Professor A. Fuller and Professor D. Mitchell.

TABLE OF CONTENTS

DECLARATION	ii
CONFERENCE PROCEEDINGS AND PRESENTATIONSi	ii
ABSTRACT	V
ACKNOWLEDGEMENTSv	ii
LIST OF FIGURESxi	ii
LIST OF TABLES	ii
LIST OF ABBREVIATIONS xvi	ii
CHAPTER 1- INTRODUCTION	1
1.1 Introduction	2
1.2. Menstrual cycle	4
1.2.1. Introduction	4
1.2.2. Events of the menstrual cycle	5
1.3. Hormonal control of the menstrual cycle	7
1.3.1. Introduction	7
1.3.2. Oestradiol	3
1.3.3. Progesterone 10	C
1.3.4. Measurement of steroid hormones in non-human primates1	1
1.4. Cortisol and the menstrual cycle1	3
1.5. Anogenital swellings	5

PAGE

1.5.1. Effect of oestradiol and progesterone on the anogenital area	16
1.5.2. Estimation of ovulation using anogenital swelling changes	17
1.6. Menstrual cycle related body temperature regulation	18
1.7. Physical activity patterns	24
1.8. Aims of the dissertation	27

2. 1. Experimental animals and housing 2	29
2.2. Surgery	3
2.2.1. Anaesthesia protocol 3	3
2.2.2. Abdominal logger and telemeter implantation	3
2.2.3. Activity logger implantation 3	\$4
2.2.4. Post-surgical care3	6
2.3. Recovery of data loggers and radiotelemeters	6
2.4. Measurement of abdominal temperature	\$7
2.5 Measurement of physical activity	9
2.6. External reproductive changes 4	1
2.6.1. Menstrual bleeding 4	1
2.6.2. Anogenital swellings 4	1
2.7. Hormone analysis 4	2
2.7.1. Urine sample collection and processing 4	2
2.7.1.1. Extraction of urine steroid hormones	4

2.7.1.2. Measurement of creatinine	44
2.7.2. Faecal sample collection and processing	45
2.7.2.1. Drying	46
2.7.2.2. Extraction of faecal steroid hormones	46
2.7.3. Radioimmunoassays	47
2.7.3.1.Cortisol assays	48
2.7.3.2. Progesterone assays	49
2.7.3.3. Oestradiol assays	50
2.8 Data analysis	52

PTER 3- RESULTS

3.1. Menstrual cycle record 50
3.1.1. Menstrual bleeding50
3.2. Hormone analysis
3.2.1. Urine progesterone and oestradiol over the menstrual cycle 50
3.2.2. Comparison of urine and faecal steroid hormones
3.3. Anogenital swelling changes over the menstrual cycle
3.4. Nychthemeral rhythms of abdominal temperature and physical activity 6
3.5. Abdominal temperature over the menstrual cycle7
3.6. Physical activity over the menstrual cycle
3.7. Correlation of abdominal temperature and physical with theday of
ovulation

CHAPTER 4- DISCUSSION	
CHAPTER 5- CONLUSIONS	
REFERENCES	

LIST OF FIGURES

Page

Figure 1.1	Serum hormone concentrations during the baboon, Papio	
	hamadryas cynocephalus menstrual cycle.	9
Figure 1.2	Hormonal and anogenital profiles in captive chimpanzee, Pan	
	troglodytes, over the menstrual cycle.	19
Figure 1.3	Variation of rectal temperature and ovarian hormones over the	
	menstrual cycle in women.	21
Figure 1.4	Nocturnal body temperature (mean \pm SEM) in 12	
	chimpanzees.	22
Figure 2.1	An adult female chacma baboon, Papio hamadryas ursinus.	30
Figure 2.2	Photographs showing A) Three adjacent cages (1-3) and B)	
	modified sluice under the cages to trap urine samples.	32
Figure 2.3	Implantation of an activity logger.	35
Figure 2.4	Thermometric data logger before waxing (left) and after	
	after waxing (right).	38
Figure 2.5	Activity data logger after waxing (left) and before waxing	
	(right)	40

Figure 2.6	Different stages of the anogenital swellings during the	
	menstrual cycle in baboons, complete anogenital	
	deturgescence (left) and maximum swelling (right).	43
Figure 3.1	Urine steroid hormonal profiles, expressed per mmol of	
	creatinine (Cr) in the same sample for baboon B, cycle 2.	58
Figure 3.2	Urine steroid hormonal profiles, expressed per mmol of	
	creatinine (Cr) in the same sample for two consecutive cycles	
	in baboon C.	59
Figure 3.3	Urine steroid hormonal profiles, expressed per mmol of	
	creatinine (Cr) in the same sample for two consecutive cycles	
	in baboon D.	60
Figure 3.4	Oestradiol (mean \pm SD) and progesterone (mean \pm SD)	
	concentrations, expressed per mmol of creatinine in the	
	same sample in different phases of the menstrual cycle.	61
Figure 3.5	Urine and faecal steroid hormonal profiles during the	
	menstrual cycles of two baboons.	63
Figure 3.6	Correlation of progesterone and oestradiol concentrations	
	in urine and faecal samples for two baboons.	64

Figure 3.7	Digital photographs showing the different stages of the	
	anogenital swellings during the menstrual cycle.	66
Figure 3.8	Anogenital swelling changes over one menstrual cycle in	
	one of the study animals.	68
Figure 3.9	Cyclic anogenital swellings over menstrual cycles in three	
	of the study animals as a function of time.	69
Figure 3.10	Nychthemeral rhythms of 10-min recordings of abdominal	
	temperature (mean \pm SD) and physical activity (mean \pm SD)	
	as a function of time from one baboon.	70
Figure 3.11	A typical example of mean \pm SD, minimum and maximum	
	24-h abdominal temperature changes over one menstrual	
	cycle from one of the animal.	72
Figure 3.12	Mean 24-h abdominal temperatures calculated from 10-min	
	recordings over menstrual cycles in four baboons.	73
Figure 3.13	Mean \pm SD, maximum \pm SD and minimum \pm SD 24-h	
	abdominal temperature in different phases of the menstrual	
	cycle.	75
Figure 3.14	Typical mean \pm SD 24-h physical activity changes over one	
	menstrual cycle from one of the animal.	77
Figure 3.15	Mean 24-h physical activity over menstrual cycles in three	

xv

	baboons over a period four months.	78
Figure 3.16	Mean \pm SD 24-h physical activity in different phases of the	
	menstrual cycle.	79
Figure 3.17	Estimation of the time ovulation for one cycle in	
	baboon C.	81
Figure 4.1	Double plots of the nychthemeral rhythm of body core	
	temperature of baboons and a woman.	89
Figure 4.2	Progesterone, anogenital swellings, abdominal temperature	
	and physical activity changes over one menstrual cycle in one	
	of the study animals.	91
Figure 4.3	The physical activity (percentage of maximum recorded	
	activity for each woman, mean \pm SD) of the women with a	
	history of dysmenorrhoea and women without a history of	
	dysmenorrhoea when not menstruating.	96

LIST OF TABLES

Table 3.1	Duration of the menstrual cycles (days) in baboons.	57
Table 3.2	Duration of different stages of anogenital swellings	67
Table 3.3	Abdominal temperature data for all the animals .	74
Table 3.4	Estimation of the day of ovulation from abdominal	
	temperature and physical activity data as days prior to	
	anogenital deturgescence .	82

LIST OF ABBREVIATIONS

ANOVA	Analysis of variance
Cr	Creatinine
DNA	Deoxyribonucleic acid
FSH	Follicle stimulating hormone
h	Hours
I ¹²⁵	Radioisotope of iodine
kg	Kilogram
LH	Luteinising hormone
min	Minutes
ml	Millilitres
RIA	Radioimmunoassay
SD	Standard deviation
°C	Degrees Celsius