

THE POPULATION DYNAMICS OF SOME WOODY SPECIES
IN THE KALAHARI SAND VEGETATION OF HWANGE
NATIONAL PARK

Susan Lawley Childes

A Dissertation Submitted to the Faculty of Science
University of the Witwatersrand, Johannesburg
for the Degree of Master of Science

Hwange 1984

ABSTRACT

THE POPULATION DYNAMICS OF SOME WOODY SPECIES IN THE
KALAHARI SAND VEGETATION OF HWANGE NATIONAL PARK

CHILDES, Susan Lawley, M.Sc. University of the
Witwatersrand, Johannesburg, 1984.

The population dynamics and principal controlling factors of six woody species found in the Kalahari Sand vegetation of Hwange National Park are investigated. The populations from woodland and scrub areas of known human disturbance (logging) are compared with undisturbed sites. The effects of edaphic factors, frost, fire and elephant damage on the woody species are examined.

The results show the populations from the disturbed areas to be generally unstable with a decline in the woodlands. The estimated recruitment of Baikiaea plurijuga is 0,2 of the rate needed to maintain the woodlands in the present state. The influence of soil moisture is of major importance in determining vegetation physiognomy and species composition. Frost and fire also have a strong modifying influence on the vegetation through the differing sensitivities of each species. Frost, fire and elephant account for 57%, 24% and 19% of the damage (excluding unknown factors).

In conclusion a hypothesis showing the interaction of abiotic and biotic factors on the three stages of vegetation cycling : woodland, scrub and grassland, is presented.

DECLARATION

I declare that this dissertation is my own, unaided work. It is being submitted for the degree of Master of Science in the University of the Witwatersrand, Johannesburg. It has not been submitted before for any degree or examination in any other University.

Susan Lawley Childes

Susan Lawley Childes

20 day of March, 1984.

	Page
PREFACE	iii
Chapter	
1. INTRODUCTION	1
2. THE STUDY AREA	5
2.1. Situation	5
2.2. Climate	5
2.3. Topography and Geology	9
2.4. Vegetation	9
2.5. Biotic Influences	14
3. LITERATURE REVIEW	18
3.1. Geology	18
3.1.1. Geological History of the Kalahari Sands	18
3.1.2. Pan Formation	19
3.1.3. Characteristics of Kalahari Sand	20
3.2. Vegetation	21
3.2.1. Forestry Activities	21
3.2.2. Composition and Classification	24
3.2.3. Succession	27
4. VEGETATION PHYSIOGNOMY, GROWTH AND MORTALITY.	36
4.1. Introduction	36
4.2. Methods	38
4.2.1. Vegetation Size Structure and Composition	38
4.2.2. Growth Rates	41
4.2.3. Mortality	43
4.2.4. Coppice Competition Experiment.	43

Chapter	Page
4.3. Results	43
4.3.1.1. Vegetation Composition	43
4.3.1.2. Size Structure	46
4.3.1.3. Quotients per Size Class ..	51
4.3.1.4. Disturbance Index	59
4.3.2. Growth Rates	59
4.3.3. Mortality	67
4.3.4. Competition	73
4.4. Discussion	73
5. DETERMINANTS OF VEGETATION CHANGE	76
5.1. Introduction	76
5.1.1. Literature Review	76
5.1.1.1. Edaphic Factors	76
5.1.1.2. Frost	78
5.1.1.3. Fire	79
5.1.1.4. Elephant	80
5.1.1.5. Logging Activities	80
5.2. Methods	81
5.2.1. Edaphic Factors	81
Soil Analyses	81
Soil Moisture	81
Hard Layer	88
Root Systems	88
Phenology	89
5.2.2. Frost	89
5.2.3. Fire	93
5.2.4. Frost, Fire and Elephant Damage	93
5.3. Results	93
5.3.1. Edaphic Factors	93
Soil Analyses	93
Soil Moisture	98
Hard Layer Testing	106
Root Systems	110
Phenology	116

Chapter	Page
5.3.2. Frost	125
5.3.3. Fire	135
5.3.4. Frost, Fire and Elephant Damage Assessments	138
5.3.4.1. Damage Assessments for the six main species	138
5.3.4.2. Damage Assessments for other species	143
5.3.4.3. Heights at which damage occurs	143
5.3.5. Logging Activities	146
5.4. Discussion	149
5.4.1. Edaphic Factors	149
Soil Moisture	149
Root Systems	150
Phenology	150
5.4.2. Frost Damage	151
5.4.3. Fire Damage	152
5.4.4. Elephant Damage	152
5.4.5. Logging Effects	154
5.4.6. Hypothesis	155
6. GENERAL DISCUSSION AND CONCLUSIONS	158
6.1. <u>Baikiaea plurijuga</u> Dynamics	158
6.1.1. Introduction	158
6.1.2. Methods	164
6.1.3. Results	166
6.1.4. Discussion	169
6.2. General Vegetation Dynamics	171
6.2.1. Introduction	171
6.2.2. Dynamics and Succession	172
6.2.3. Factors affecting Dynamics ...	172
6.2.4. Stability and Resilience	178
6.3. Critique of Study	179
6.4. Importance of Study	181
6.4.1. Management Guide	181
LIST OF REFERENCES	184

PREFACE

This dissertation was undertaken as a part-time study while I was employed as an Ecologist by the Department of National Parks and Wildlife Management. The study was also listed as a Departmental project for Hwange National Park.

The purpose of the study was to test and develop earlier work by my supervisor, Prof. B.H. Walker, and my predecessor, Mr J.E. Rushworth, on the pattern and dynamics of Kalahari Sand vegetation. A further understanding of the Kalahari Sand ecosystem would provide a firmer base for Park management.

The study was limited to the population dynamics of six of the main woody species found in woodland and scrub communities on the Sands. It continued through three dry seasons and two wet seasons from 1980 - 1982.

I wish to thank Ms W. Knoop for her invaluable help with the computer program, Mr G.M. Calvert and the Forestry Commission for making data and reports available, Mrs J. Hussein and the Department of Research and Specialist Services for the loan of the soil moisture meter and the soil analyses, and Dr T. Smith for testing the Baikiaea population model. Assistance in the field from the Research Scouts at Main Camp is gratefully acknowledged. Finally, I am thankful to my supervisor for his stimulating discussions and guidance.

LIST OF TABLES

	Page
Chapter	
2. THE STUDY AREA	
Table 2.1. Meteorological data for Main Camp, Hwange National Park	7
Table 2.2. Frequency of severe frost at Main Camp	10
Table 2.3. Area of Kalahari Sands covered by different vegetation communities ...	10
Table 2.4. Population estimates for the main herbivore species in Hwange National Park	16
4. VEGETATION PHYSIOGNOMY, GROWTH AND MORTALITY	
Table 4.1. Stands sampled for vegetation structure, composition and damage assessments	40
Table 4.2. Vegetation composition of the five stands	45
Table 4.3. Disturbance indices for the six species in the five stands	61
Table 4.4. Growth rates of the five main species	65
5. DETERMINANTS OF VEGETATION CHANGE	
Table 5.1. Depth and siting of soil moisture blocks	86
Table 5.2. Tree and shrub species marked for phenology records	90
Table 5.3. Heights of thermometers placed in nine vegetation sites near Main Camp.	92
Table 5.4. Laboratory analyses of Kalahari Sand samples	94

Chapter	Page
Table 5.5. Summary of mean monthly minimum temperatures at Livingi and Malindi sites	126
Table 5.6. Mean monthly minimum temperatures at Dopi	131
Table 5.7. Annual rainfall and the proportion of Kalahari Sand vegetation burnt	136
Table 5.8. Percentage weight damage of the six major species	140
Table 5.9. Percentage weighted damage for other woody species	144
 6. GENERAL DISCUSSION AND CONCLUSIONS	
Table 6.1. The life stages and corresponding sizes of <u>Baikiaea plurijuga</u>	160
Table 6.2. Seedling survivorship of <u>Baikiaea plurijuga</u>	160
Table 6.3. Life table for <u>Baikiaea plurijuga</u> .	163
Table 6.4. Numbers of trees in each life stage after running a matrix population model for 50 years	167
Table 6.5. Approximate time taken for <u>Baikiaea plurijuga</u> trees to pass through each life stage	168
Table 6.6. Densities of <u>Baikiaea plurijuga</u> in the five life stages in woodland and scrub	168
Table 6.7. Key to management types of scrub vegetation	182

LIST OF FIGURES

	Page
Chapter	
2. THE STUDY AREA	
Fig.2.1. Map of Hwange National Park showing situation and study sites	6
Fig.2.2. Annual rainfall and potential evapotranspiration for Main Camp	8
Fig.2.3. Parallel arrangement of dunes and vegetation on the Kalahari Sands in western Hwange National Park	11
Fig.2.4. Geology map of Hwange National Park adapted from the Geology of Rhodesia 1:1 000 000 map, Surveyor General, 1971	12
Fig.2.5. Vegetation map at Hwange National Park	13
Fig.2.6. Photograph taken from the bottom of a fossil drainage line near Manga 3 Pan, Hwange National Park	15
3. LITERATURE REVIEW	
Fig.3.1. Map of Forestry concessions in Hwange National Park and adjacent areas	23
Fig.3.2. Photograph of <u>Baikiaea plurijuga</u> and <u>Colophospermum mopane</u> growing in close proximity, indicating the abrupt soil changes in the dunes	29
Fig.3.3. Boughey's (1963) hypothesis for the derivation of various secondary communities from <u>Baikiaea</u> woodland	31

Chapter	Page
4. VEGETATION PHYSIOGNOMY, GROWTH AND MORTALITY	
Fig.4.1. Photograph of dry season <u>Baikiaea plurijuga</u> woodland near Mitswiri Pan, Hwange National Park	39
Fig.4.2. Diagram showing positions of circumference measurements above and below ground	42
Fig.4.3. Photograph showing the use of weather resistant tapes for measuring the growth rates of trees.	42
Fig.4.4. Photograph of regenerating (after logging) <u>Baikiaea plurijuga</u> woodland near Caterpillar Pan, Hwange National Park	44
Fig.4.5(a) Above and below ground size structures of <u>Baikiaea plurijuga</u> populations in two woodland stands .	46 A
4.5(b) Above and below ground size structures of <u>Baikiaea plurijuga</u> populations in one woodland and one scrub stand.....	47
4.5(c) Above and below ground size structures of <u>Terminalia sericea</u> populations in two woodland and two scrub stands.....	49
4.5(d) Above and below ground size structures of <u>Guibourtia coleosperma</u> populations in two woodland and one scrub stand	50
4.5(e) Above and below ground size structures of <u>Burkea africana</u> populations in two woodland and two scrub stands	52

Chapter	Page
4.	
4.5(f) Above and below ground size structures of <u>Erythrophleum africanum</u> populations in two woodland and one scrub stand.....	53
4.5(g) Above and below ground size structures of <u>Ochna pulchra</u> populations in two woodland and two scrub stands	54
Fig.4.6(a) Quotients for successive 2000cm ² size classes for three woodland stands of <u>Baikiaea plurijuga</u>	55
4.6(b) Quotients for successive 2000cm ² size classes for one woodland and one scrub stand of <u>Guibourtia coleosperma</u> and two woodland and two scrub stands of <u>Burkea africana</u>	57
4.6(c) Quotients for successive 2000cm ² size classes for two woodland and one scrub stand of <u>Erythrophleum africanum</u>	58
4.6(d) Quotients for successive 2000cm ² size classes for two woodland and two scrub stands of <u>Terminalia sericea</u> and <u>Ochna pulchra</u>	60
Fig.4.7(a) Growth of <u>Baikiaea plurijuga</u> using data from this study and Forestry Commission data	63
4.7(b) Growth of <u>Guibourtia coleosperma</u> using data from this study and Forestry Commission data	64

Chapter	Page
4.	
4.7(c)Growth of <u>Burkea africana</u> using data from this study and Forestry Commission data	68
4.7(d)Growth of <u>Erythrophleum africanum</u> using data from this study and Forestry Commission data	69
4.7(e)Growth of <u>Terminalia sericea</u> and <u>Ochna pulchra</u> using data from this study and Forestry Commission data .	70
Fig.4.8. Growth of coppiced individuals of <u>Baikiaea plurijuca</u> , <u>Guibourtia coleosperma</u> , <u>Burkea africana</u> and <u>Erythrophleum africanum</u>	71
Fig.4.9. Percentage of dead trees per 10cm circumference size class for the six species in six different sites	72
Fig.4.10. A comparison between the growth of pruned and unpruned (control) coppiced stems of <u>Baikiaea plurijuca</u> .	74
5. DETERMINANTS OF VEGETATION CHANGE	
Fig.5.1. Photograph of a degraded sand dune near Dopu Pan	83
Fig.5.2. Photographs of the three main vegetation communities along the dune	84
Fig.5.3. Positions of the six soil moisture recording sites in the three communities along the dune	85
Fig.5.4. Diagram showing the positions of auger holes in a soil pit for the installation of soil moisture blocks	87

Chapter	Page
5.	
Fig.5.5. Positioning of thermometers for the recording of minimum temperatures at different heights along the dune near Dopi Pan	91
5.6(a) Mean monthly changes in soil moisture in two <u>Baikiaea</u> woodland sites (nos. 3 and 4) on a dune ridge near Dopi Pan	99
5.6(b) Mean monthly changes in soil moisture in two <u>Terminalia</u> scrub sites (nos. 2 and 6) on a dune ridge near Dopi Pan	100
5.6(c) Mean monthly changes in soil moisture in two <u>Burkea</u> open woodland sites (nos. 1 and 5) on a dune ridge near Dopi Pan	101
Fig.5.7. The different drying rates at 1,0m depth of the three vegetation communities along a dune near Dopi Pan	102
Fig.5.8(a) Mean monthly changes in soil moisture in a <u>Baikiaea</u> woodland site near Livingi Pan where there was no obvious topographical changes	104
5.8(b) Mean monthly changes in soil moisture in a mixed scrub site near Livingi Pan where there was no obvious topographical changes	105
Fig.5.9. Photograph of 10m deep soil pit dug in <u>Baikiaea</u> woodland near Livingi Pan	107
Fig.5.10(a) Mean monthly changes in soil moisture in a <u>Baikiaea</u> scrub site near Malindi, where no hard layer was detected	108

Chapter	Page
5.	
5.10(b) Mean monthly changes in soil moisture in a mixed scrub near Livingi Pan where a hard layer was present at 2,3m depth	109
Fig.5.11	
(a) - (i) Photographs of excavated root systems of the six species in woodland and scrub sites	111-115
Fig.5.12	
(a) Phenological changes of two well-developed woodland species, <u>Baikiaea plurijuga</u> and <u>Guibourtia coleosperma</u>	117
(b) Phenological changes of two disturbed woodland species, <u>Burkea africana</u> and <u>Erythrophleum africanum</u>	119
(c) Phenological changes of two scrub species, <u>Terminalia sericea</u> and <u>Ochna pulchra</u>	120
(d) Phenological changes of two woodland shrub species, <u>Croton pseudopulchellus</u> & <u>C. gratissimus</u>	122
Fig.5.13. Photograph of <u>Baikiaea plurijuga</u> flowers	121
Fig.5.14. Photograph of <u>Erythrophleum africanum</u> flushing after a drought and late rains in April 1982	123
Fig.5.15. Photograph of <u>Terminalia sericea</u> flushing along a road verge	124
Fig.5.16. Mean monthly minimum temperature profiles in scrub and woodland sites near Main Camp for the period May - August 1982	127

Chapter	Page
5.	
Fig.5.17. Absolute minimum temperature profiles in six scrub sites for July 1982	128
Fig.5.18. Absolute minimum temperature profiles along the dune study site for the period May - August 1982	130
Fig.5.19. Absolute minimum temperatures at different heights along the dune study site for July 1982	132
Fig.5.20. The relationship between mean weekly dewpoint and minimum temperature recorded at Main Camp from May - August 1982	133
Fig.5.21. Annual rainfall and absolute minimum temperature (May-Aug.) measured at Main Camp from 1951-1982	134
Fig.5.22. Monthly rainfall (histogram) and percentage area of Kalahari Sand vegetation burnt (x --- x) from 1969-1982	137
Fig.5.23. Fire frequency map of the Kalahari Sand area in Hwange National Park for the period 1968-1982	139
Fig.5.24(a) Percentage total damage and the height distribution of <u>Baikiaea plurijuga</u> populations in three woodland and one scrub stands	145
5.24(b) Percentage total damage and the height distribution of <u>Terminalia sericea</u> populations in two woodland and two scrub stands	147

Chapter	Page
5.	
Fig.5.24(c) Percentage total damage and the height distribution of <u>Burkea africana</u> , <u>Erythrophleum africanum</u> , <u>Ochna pulchra</u> and <u>Guibourtia coleosperma</u> in woodland and scrub stands	148
Fig.5.25. Photograph illustrating facilitatory effect of <u>Terminalia sericea</u> trees protecting <u>Baikiaea plurijuga</u> seedlings and coppice from frost	153
Fig.5.26. An hypothesis to explain the role of edaphic features in controlling the vegetation structure and composition	156
6. GENERAL DISCUSSION AND CONCLUSIONS	
Fig.6.1. Life cycle graphs of a <u>Baikiaea plurijuga</u> population classified by size/stage	162
Fig.6.2. Diagram illustrating the hypothetical interactions of abiotic and biotic factors on the successional phases of vegetation on deep, well-drained Kalahari Sands	173
Fig.6.3. The dynamics or relative rates of change between the various stages in the three vegetation phases, woodland, scrub and grassland	174
Fig.6.4. Dry season photograph of area around Nyamandhlovu Pan, Hwange National Park showing the effects of herbivore pressure on grass and woody vegetation	177

Chapter

6.

Fig.6.5. Photograph of a 1m² grassland enclosure plot at Ngamo, Hwange National Park showing the intensity of grazing by wildebeest 177

disorders, particularly in the dry, flat inland land country. A brief review of past disturbances and the problems associated with the establishment of the Park are given below. A more detailed historical perspective appears in Chapter 7.

At the beginning of this century the only inhabitants of the Hwange district were the Bushmen. European influence was initially confined around the river which separated them from the Kalahari Desert to the west during the dry months. Later, the value of indigenous wild game for the hunting and exploitation of the bush by colonial (British) hunters increased, with overall development of the country. Much of the time, extraction was uncontrolled and without any regard to the future of the wild game areas. The direct and indirect effects of this disturbance are of major importance in the modification of Hwange's land vegetation. A major factor, responsible for much of the fire damage to the vegetation, was the construction of a railway line from Bulawayo to Victoria Falls. This locomotive, and still is, had a coal-burning boiler and a tall chimney, a constant source of fire.

As with most protected wildlife areas, there arose a conflict between wildlife and human activities. In 1940 a fence was erected to prevent the spread of foot and mouth disease to cattle by the adjacent farming areas. The fence ran from the Zambezi river south to the Victoria Falls. Since it also habitually blocks the migration routes of wildebeest, it caused controversy between the

CHAPTER 1. INTRODUCTION

Hwange (formerly Wankie) National Park is the largest National Park in Zimbabwe and the third largest in Africa.

Despite its recent history of human interference, much of Hwange still remains a comparatively untouched wilderness, particularly in the dry, flat Kalahari Sand country. A brief review of past disturbances and the problems associated with the establishment of the Park are given below. A more detailed historical perspective appears in Chapter 2.

At the beginning of this century the only inhabitants of the Kalahari Sands were nomadic Bushmen. European influence was initially centred around hunting the game which migrated from the waterless Sands to the Gwaai river during the dry months. Later, the value of indigenous timber was realised and exploitation of the teak or 'mkusi' (Baikiaea plurijuga) commenced, with overall development of the country. Much of the timber extraction was uncontrolled and without any regard to the future of the slow growing forests. The direct and indirect effects of this disturbance are of major importance in the modification of Kalahari Sand vegetation. A second factor, responsible for much of the fire damage to the vegetation, was the construction of a railway line from Bulawayo to Victoria Falls. Steam locomotives were, and still are, a continual source of fire.

As with most protected wildlife areas, there arose a conflict between wildlife and domestic stock. In 1960 a fence was erected to prevent the spread of foot and mouth disease to cattle in the adjacent farming areas. The fence ran from the Gwaai river south to the Botswana border. Since it was habitually broken by elephant, buffalo and wildebeest, it became Veterinary Department and

National Parks policy to shoot all game within a mile of the fence. This caused a heavy drain on the game population (Davison, 1977). The effects of this fence protection must have improved the balance between herbivore and predator populations, as it had been the practice for the previous 30 years to reduce lion, hyaena and wild dog numbers. In 1981/82 a 'game-proof' fence was erected by Botswana authorities along the south western boundary of the Park. The repercussions of this barrier on the movements of buffalo, elephant and other species remain to be seen, but will probably cause a further compression of game into the Park.

Another major problem in the Kalahari Sand region is the presence of widespread artificially supplied water during the dry season. Boreholes were first sunk in the 1940's in order to attract the game away from the farming areas. The permanent water supplies soon began to have an effect on game migrations and by 1945 elephant and buffalo appeared in greater numbers (Davison, 1977). Wankie Game Reserve was officially designated as a National Park in 1950 and following this, tourism became an important consideration in the Park. Pumped waterholes became popular game viewing spots, but because of game pressure the vegetation was heavily damaged. Further boreholes were then drilled in an attempt to alleviate the pressures around the older waterholes. The provision of additional water during the critical dry season did little to reduce vegetation damage but allowed herbivore populations, especially elephant, to increase. Levels of elephant damage became unacceptable and in 1970 a population reduction exercise began, and continued annually.

To summarize, the Kalahari Sand vegetation of Hwange has been subjected to a past eighty years of

disturbance through timber extraction and an increase in game populations and fire frequency. As most of the vegetation is dominated by woody plants (for explanation see Chapter 6), the effects of disturbance would be reflected in these species. The vegetation falls into two main physiognomic categories: woodland, dominated by Baikiaea plurijuga, and scrub, dominated by Terminalia sericea. The questions underlying this project were firstly, are the woodland and scrub areas inherently the same, i.e.: stages in a succession? Secondly, what factors cause changes in the vegetation, and what is the effect of past disturbance? The study focussed on the two dominant species B. plurijuga and T. sericea. Four other woody species were also considered as they are common components of woodland and scrub communities. Those were Guibourtia coleosperma, Burkea africana, Erythrophleum africanum and Ochna pulchra.

The objectives of the study were to determine:

- (i) The population status and dynamics of these six woody plants.
- (ii) What factors influence the vegetation structure, composition and dynamics. This would then provide an insight as to whether management was feasible or necessary to counter the effects of these factors.

The following is a résumé of the work, outlining the arrangement and themes of the chapters. Chapter 2 introduces the study area, with details of past history, climate, soils, vegetation and animals. Chapter 3 provides the literary background with a discussion of the work of early botanists and forestry officers on Kalahari Sand vegetation, together with more recent research and ideas. In Chapter 4 the theme of a comparison between woodland

Author Childes S L

Name of thesis The Population dynamics of some woody species in the Kalahari Sand Vegetation of Hwange National Park
1984

PUBLISHER:

University of the Witwatersrand, Johannesburg

©2013

LEGAL NOTICES:

Copyright Notice: All materials on the University of the Witwatersrand, Johannesburg Library website are protected by South African copyright law and may not be distributed, transmitted, displayed, or otherwise published in any format, without the prior written permission of the copyright owner.

Disclaimer and Terms of Use: Provided that you maintain all copyright and other notices contained therein, you may download material (one machine readable copy and one print copy per page) for your personal and/or educational non-commercial use only.

The University of the Witwatersrand, Johannesburg, is not responsible for any errors or omissions and excludes any and all liability for any errors in or omissions from the information on the Library website.