

HIGH INHERITANCE ELONGATE STROMATOLITIC MOUNDS FROM THE TRANSVAAL DOLOMITE

by

K. A. Eriksson and J. F. Truswell

Department of Geology, University of the Witwatersrand, Johannesburg.

ABSTRACT

Elongate mound-like structures up to 10 metres across and 40 metres in length are described. Characteristic features of the mounds are their high inheritance and considerable relief which, along with the absence of sub-aerial exposure features, the fine-grained nature of the carbonate, and their vertical persistence, are taken to suggest a subtidal origin. Marine currents are considered adequate to account for their elongation.

INTRODUCTION

Study of the environmental significance of stromatolites owes much to the work of Black at Andros Island in the Bahamas. There, in 1933, he recorded contemporary algal mats in the intertidal and supratidal zones, and showed that specific growth forms developed in distinct geographic

settings. He went on to draw an analogy between these mats and the stromatolitic structures so prevalent in many Palaeozoic and Precambrian carbonate sequences. Early corroboration of this analogy was provided from the Transvaal Dolomite in the northern Cape by Young (1934). Young also sought to relate stromatolitic and other structures




Fig. 1. Outcrops of the Transvaal Dolomite and Locality of the Section (marked by arrow).

dimensions. While the dominant minor structure remains the above low-amplitude crinkled bedding, other structures were also recorded. These include a linked columnar form with variable spacing (Fig. 7) and a height of 15–40 cm and diameter of 4–8 cm; and in one horizon an unusual fan-shaped domical structure up to 12 cm high (Young 1940) (Fig. 8).

AN ENVIRONMENTAL INTERPRETATION

A survey of the literature reveals no obvious analogy with the Hol River stromatolitic mounds although a number of structures with a similar external geometry have been reported. Both thrombolites, as discussed by Aitken (1967), and loferites (Fischer, 1964) are characterized by broad contiguous arched bedding structures. However the former has a spongy or clotted internal fabric, while the latter by definition contains small ragged pores filled by clear blocky calcite; and neither of these contain internal lamination. Hoffman (in press) has described large features from the platform edge in the Pethei Group at the Great Slave Lake in Canada. These do have a similar relief to the mounds on the upper fall in the Groot Boetsap River but contain branched stromatolites in a specific vertical sequence. Knight (1968) described both large elongate domical structures and undulatory stromatolites in the Medicine Bow mountains in Wyoming, but recorded no association of these two types in the mapped bioherms.

Among the features that are characteristic of the mounds in the Groot Boetsap are their high inheritance and relief. There is a lack of any sub-aerial exposure features. All these criteria suggest that the mounds are subtidal phenomena but within the realms of light penetration. The fine-grained carbonate developed through the upper fall has been shown elsewhere to be dominant in deeper water carbonate environments (Irwin 1965, Laporte 1967, 1969, and Kepper 1972). The vertical persistence through a thickness of 45 metres in the gorge is also most noticeable.

In a discussion of Palaeozoic carbonate facies Laporte (1971) noted that vertical facies variations, particularly in the deep subtidal environment, are very gradual and infrequent. The marked vertical persistence of the stromatolite mounds in the Boetsap River would suggest an analogy with Laporte's observations; specifically of the formation of these structures in a deep-water environment that may well have been subtidal.

It has been noted earlier that the dominant minor structure is a low amplitude crinkled bedding, with which linked columnar and small domical forms may be associated in the smaller mounds. Heath et al. (1967) have shown that within the subtidal environment an increase in energy occurs towards the land, and in the Groot Boetsap River the minor structure is regarded as a reflection of this, with the smaller mounds and their greater potential variation in minor structure being developed landward of the larger mounds. At

Shark Bay Logan (1961) noted that outwards in the intertidal zone flat-bedding gave way successively to dome-like stromatolites, linked columnar forms, and unlinked columnar stromatolites with energy increase towards the surf zone. In the Groot Boetsap section small changes in the minor structures are taken to suggest lesser changes of energy.

A further distinctive, and unusual, feature of the mounds is their consistent east-west elongation, i.e. at right angles to the depositional shoreline (see also Visser and Grobler (in press)). This orientation must be related to persistent, if weak, currents within the subtidal environment. Irwin (1965) considered that in the subtidal parts of an epeiric sea marine currents would constitute the only form of hydraulic energy. Shaw's comment (1964) that such currents were small in size compared with oceanic currents, but if they were steady would have exerted a marked effect, is regarded as being most pertinent in the development of the mounds in the Groot Boetsap River. The stromatolite mounds from the Pethei Group in Canada (Hoffman, in press) have been shown to have formed on a shelf edge, basinwards of which greywackes, marlstones and shales were deposited. To date only cratonic shelf sedimentation in an epeiric sea is indicated as the depositional conditions of the Transvaal Dolomite. There is no evidence of an off-shelf facies, or that the elongate mounds developed on a shelf-edge.

REFERENCES

- AITKEN, J.D. (1967). Classification and environmental significance of cryptalgal limestones and dolomites, with illustrations from the Cambrian and Ordovician of S. W. Alberta. *Jour. Sed. Petrology*, 37, 1163–1178.
- BLACK, M. (1934). Algal sediments of Andros Island, Bahamas. *Phil. Trans. Roy. Soc. London, Ser. B*, 222, 165–192.
- FISCHER, A. G. (1965). Fossils, early life and atmosphere history. *Proc. Nat. Acad. Sci. United States*, 53, 1205–1215.
- HEATH, C. P., LUMSDEN, D. N., AND CAROZZI, A. (1967). Petrography of a carbonate transgressive/regressive sequence: the Bird Spring Group (Pennsylvanian), Arrow Canyon Range, Clark County, Nevada. *Jour. Sed. Petrology*, 37, 377–400.
- HOFFMAN, H. J. (1969a). Attributes of Stromatolites. *Geol. Surv. Canada. Paper 69–35*, pp. 58.
- (1969b). Stromatolites from the Proterozoic Animikie and Sibley Groups, Ontario, Canada. *Geol. Surv. Canada. Paper 68–69*, pp. 77.
- HOFFMAN, P. (In press). Proterozoic carbonate platform-to-basin facies zonation, Pethei Group, Great Slave Lake, North-west Territories, Canada.

- IRWIN, M. L. (1965). General theory of epeiric clear water sedimentation. *Amer. Assoc. Petroleum Geologists Bull.*, 49, 445-459.
- KEPPER, J. C. (1972). Palaeoenvironmental patterns in Middle to Lower Upper Cambrian Interval in Eastern Great Basin. *Amer. Assoc. Petroleum Geologists Bull.*, 56, 503-528.
- KNIGHT, S. H. (1968). Precambrian stromatolites, bioherms and reefs in the lower half of the Nash Formation, Medicine Bow Mountains, Wyoming. *Wyoming Univ., Contrib. to Geol.*, 7, 73-116.
- KRUMBEIN, W. C. and SLOSS, L. L. (1963). Stratigraphy and Sedimentation. Freeman, 2nd Ed., pp. 660.
- LAPORTE, L. F. (1967). Deposition near the mean sea-level and resultant facies mosaic: Manlius Formation (Lower Devonian) of New York State. *Amer. Assoc. Petroleum Geologists Bull.*, 51, 73-101.
- (1969). Recognition of a transgressive carbonate sequence within an epeiric sea; Helderberg Group (Lower Devonian) of New York State. In G. Friedman, ed. *Depositional Environments in Carbonate Rocks, Soc. Econ. Pal. and Min. Spec. Pub.* 14.
- (1971). Palaeozoic carbonate facies of the Central Appalachian Shield. *Jour. Sed. Petrology*, 41, 724-741.
- LOGAN, B. W. (1961). Cryptozoon and associated stromatolites from the Recent, Shark Bay, Western Australia. *J. Geol.*, 69, 517-533.
- SHAW, A. B. (1964). Time in Stratigraphy. McGraw Hill, New York, 365 p.
- TOENS, P. D. (1966). Precambrian limestone and dolomite in the northern Cape Province. *Mem. Geol. Surv. S. Afr.*, 57, pp. 109.
- TRUSWELL, J. F., and ERIKSSON, K. A. (1972). The morphology of stromatolites from the Transvaal Dolomite north-west of Johannesburg, South Africa. *Trans. geol. Soc. S. Afr.*, 75, 99-111.
- VISSER, J. N. J. and GROBLER, N. J. (In press). The transition beds at the base of the Dolomite Series in the northern Cape Province. *Trans. geol. Soc. S. Afr.*
- YOUNG, R. B. (1928). Pressure phenomena in the dolomitic limestones of the Campbell Rand Series in Griqualand West. *Trans. Geol. Soc. S. Afr.*, 31, 157-165.
- (1932). The occurrence of stromatolitic or algal limestones in the Campbell Rand Series, Griqualand West. *Trans. Geol. Soc. S. Afr.*, 35, 29-36.
- (1933). Conditions of deposition of the Dolomite Series. *Trans. geol. Soc. S. Afr.*, 36, 121-135.
- (1934). A comparison of certain stromatolitic rocks in the Dolomite Series of South Africa with marine algal sediments in the Bahamas. *Trans. geol. Soc. S. Afr.*, 37, 153-162.
- (1940). Further notes on algal structures in the Dolomite Series. *Trans. geol. Soc. S. Afr.*, 43, 17-21.
- (1943). The domical-columnar structures and other minor deformations in the Dolomite Series. *Trans. geol. Soc. S. Afr.*, 46, 91-107.



Fig. 3. Contiguous tops to large elongate mounds, elongated east-west. (Scale : 60 cm in length)



Fig. 4. High inheritance and relief of elongate mounds. (Scale : 60 cm in length)



Fig. 5. Axial zone domical structure. (Scale : 60 cm in length)




Fig. 6. Low amplitude stromatolitic bedding on elongate mounds. (Scale : 15 cm in length)



Fig. 7. Linked columnar minor structure on elongate mounds. (Scale : 15 cm in length)



Fig. 8. Fan-shaped domical structure on elongate mounds. (Scale : 3,5 cm in width)