



The Contribution of Electrophysiology to Entomology

Paul-André Calatayud*

Research Institute for Development (IRD), French Institute, France

By definition, electrophysiology “is the study of the electrical properties of biological cells and tissues”. Thereby the flow of ions in biological tissues (living organisms, excised tissues / organs and cells) is assessed. This tool has been initially developed for human health (e.g., electrocardiography, electroencephalography...). In the mid 20th century, electrophysiology started to be used in entomology.

In 1960 an electronic method was designed to monitor electrically the biting and feeding behaviour of piercing-sucking phytophagous and hematophagous insects [1,2]. This technique was first used to record the probing, penetration, salivation, engorgement and withdrawal of mosquitoes in relation to their hosts [3]. It was thereafter extensively used to discern hemipteran stylet activity within plant tissues [4,5]. This technique, now universally termed electrical penetration graph (EPG), is used to monitor routinely insects feeding in order to understand the mouthpart pathway and activity into host tissues. More than 200 papers have been published, mostly on aphid and leafhopper feeding [5,6]. EPG has also been largely used to understand the mode of virus transmission by hemipteran in plant tissues [6]. It can also be used to monitor chewing insects (e.g. caterpillar [7]), and more generally to monitor the organ insertion of insects into hosts such as ovipositor insertion of parasitic wasps into the host body (Backus E., personal communication).

The peripheral nervous system in insects has been extensively examined by electrophysiology. In 1957, the German biologist Schneider invented the electroantennogram (EAG) [8], which ables to “measure the average voltage output of the insect antennae to the brain by an odour”. Since then, this technique has been largely used for a comprehensive identification of the presence of receptors in insect antennae to given odours. Since [9], a blend of odours can now be screened by coupling EAG with GC (gas chromatography) and to identify the compound(s) from the blend that is/are detected by the insect. Although these techniques have been developed initially for pheromone detection, they are now largely used for semiochemicals (kairomones and allomones) detection in general. At the end of the last century, single cell recording (SCR) was developed to record the action potential in single olfactory receptor neurons [10]. By SCR, the specificity in each neuron can be revealed. Simultaneously, another tool has been also developed to record the action potential in single taste receptor neurons [11,12], allowing screening the detection of hydrosoluble compounds of single taste receptors in insects. More recently, Hiroi et al. [13] adapted this technique to enable the screening of hydrophobic substances. The patch clamp technique, which allows studying single or multiple ions channels in cells, was also applied in entomology [14]. The patch clamp technique was initially used to study the effect of insecticides on insect cell membranes and it is now used to study the neuronal function of the insect brain [15] and, in general, the mode of action of molecules in the cell physiology of insects.

The patch clamp technique allowed examining the physiology of the central nervous system in insects. Associated to that, the development of a new class of voltage-sensitive dyes [16] has offered “the possibility to optically image the functionality of neuronal circuits at both the single neuron and whole brain level”. The development of calcium-sensitive dyes [17] has provided another universal and sensitive method

to “study distinct information processing pathways in whole neural networks in particular when used in combination with two-photon laser scanning microscopy [18]”. In the last years, these developments have been increasingly used in entomology and tremendous insights into the insect’s olfactory system have been gained.

The current trend of entomology is to use routinely electrophysiology. This trend is likely to continue and combined with other studies like genomics/transcriptomics can give deeper physiological insights into the insect’s chemosensory system, which is the crucial system interacting with the environment.

Acknowledgements

Thanks are given to Fritz Schulthess for revising the manuscript and his English corrections.

References

- McLean DL, Kinsey MG (1964) A technique for electronically recording aphid feeding and salivation. *Nature* (London) 202: 1358-1359.
- Kashin P, Wakeley HG (1965) An insect ‘Biometer’. *Nature* (London) 208: 462-464.
- Kashin P (1966) Electronic recording of the mosquito bite. *J Insect Physiol* 12: 281-286.
- Tjallingii WF (1978) Electronic recording of penetration behaviour by aphids. *Entomol Exp Appl* 24: 721-730.
- Backus EA (1994) History, development, and applications of the AC electronic monitoring system for insect feeding, pp. 1-51. In: M. M. Ellsbury, E. A. Backus, and D. L. Ullman [eds.], History, development, and application of AC electronic insect feeding monitors. Thomas Say Publ. Entomol. Entomological Society of America, Lanham, MD.
- Walker GP, Backus EA [eds.] (2000) Principles and applications of electronic monitoring and other techniques in the study of homopteran feeding behavior. Thomas Say Publ Entomol Entomol Soc Am, Lanham, MD.
- Saxena RC, Khan ZR (1991) Electronic recording of feeding behaviour of *Cnaphalocrocis medinalis* (Lepidoptera: Pyralidae) on resistant and susceptible rice cultivars. *Ann Entomol Soc Am* 84: 316-318.
- Schneider D (1957) Elektrophysiologische untersuchungen von chemo- und mechanorezeptoren der antenne des seidenspinners *Bombyx mori*. *Z Vergl Physiol* 40: 8-41.
- Arn H, Städler E, Rausfcher S (1975) The electroantennographic detector: a selective and sensitive tool in the gas chromatographic analysis of insect pheromones. *Z. Naturforsch* 30C: 722-725.
- Masson C, Mustaparta H (1990) Chemical information processing in the olfactory system of insects. *Physiol Rev* 70: 199-245.
- Avenet P, Lindemann B (1991) Noninvasive recording of receptor cell action

*Corresponding author: Paul-André Calatayud, Senior Research Scientist, Research Institute for Development (IRD), French Institute, France, Tel: + (254) 720 957 877; E-mail: pcalatayud@icipe.org

Received February 20, 2014; Accepted February 23, 2014; Published February 26, 2014

Citation: Calatayud PA (2014) The Contribution of Electrophysiology to Entomology. *Entomol Ornithol Herpetol* 3: e108. doi:[10.4172/2161-0983.1000e108](https://doi.org/10.4172/2161-0983.1000e108)

Copyright: © 2014 Calatayud PA. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

- potentials and sustained currents from single taste buds maintained in the tongue: the response to mucosal NaCl and amiloride. *J Membr Biol* 124: 33-41.
12. Marion-Poll FC, Guillaumin D, Masson C (1992) Sexual dimorphism of tarsal receptors and sensory equipment of the ovipositor in the European corn borer, *Ostrinia nubilalis*. *Cell Tissue Res* 267: 507-518.
 13. Hiroi M, Tanimura T, Marion-Poll F (2008) Hedonic taste in *Drosophila* revealed by olfactory receptors expressed in taste neurons. *PLoS One* 3: e2610.
 14. Amar M, Pichon Y (1989) Effects of two sodium channel 'agonists': veratridine and deltamethrin on single channel activity in inexcitable embryonic cockroach neurones. *Biophys J* 55: 315a.
 15. Palmer M, Connolly C (2013) Patch-clamp recording from Kenyon cells in acutely-isolated bee brain. *Protocol Exchange* doi:[10.1038/protex.2013.038](https://doi.org/10.1038/protex.2013.038).
 16. Salzberg BM, Davila HV, Cohen LB (1973) Optical recording of impulses in individual neurones of an invertebrate central nervous system. *Nature* 246: 508-509.
 17. Grynkiewicz G, Poenie M, Tsien RY (1985) A new generation of Ca²⁺ indicators with greatly improved fluorescence properties. *J. Biol. Chem* 260: 3440-3450.
 18. Denk W1, Strickler JH, Webb WW (1990) Two-photon laser scanning fluorescence microscopy. *Science* 248: 73-76.

Citation: Calatayud PA (2014) The Contribution of Electrophysiology to Entomology. Entomol Ornithol Herpetol 3: e108. doi:[10.4172/2161-0983.1000e108](https://doi.org/10.4172/2161-0983.1000e108)

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 300 Open Access Journals
- 25,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: www.omicsonline.org/submission/

