

Agamermis catadecaudata n. sp. (Nematoda : Mermithidae), a parasitoid of Orthoptera in south-eastern Australia

Graeme L. BAKER * and George O. POINAR Jr. **

* Biological and Chemical Research Institute, NSW Agriculture, Private Mail Bag 10, Rydalmere, NSW 2116, Australia, and

** Division of Entomology and Parasitology, University of California, Berkeley, CA 94720, USA.

Accepted for publication 22 April 1994.

Summary – *Agamermis catadecaudata* n. sp. is a parasitoid of the acridids *Phaulacridium vittatum* (Sjöstedt), *Praxibulus exsculptus* Rehn and *Brachyexarna lobipennis* Sjöstedt, the tetrigid *Paratettix australicus* Walker, the katydid *Conocephalus* sp. and the cricket *Bobilla victoriae* Otte & Alexander in the Northern Tablelands of New South Wales. In the laboratory *A. catadecaudata* n. sp. infected and developed in the acridid *Chortoicetes terminifera* (Walker). *A. catadecaudata* n. sp. is distinguished from all previously described species by the length of the infective juvenile and the position of the node. The infective is decaudate. Adult *A. catadecaudata* n. sp. are similar to *A. decaudata* but the egg is smaller and the male has a shorter spicule. Throughout its range there are two forms of parasitic juvenile the relative abundances of which are variable. An undescribed species, *Agamermis* sp. aff. *sinuosa* Kaiser, was recorded in the Southern, Central and Northern Tablelands and was sympatric with *A. catadecaudata* n. sp. in the Northern Tablelands. The host of *A. sp. aff. sinuosa* is unknown.

Résumé – *Agamermis catadecaudata* n. sp. (Nematoda : Mermithidae), un parasitoïde des Orthoptères du sud-est de l'Australie – *Agamermis catadecaudata* n. sp. est un parasitoïde des acridiens *Phaulacridium vittatum* (Sjöstedt), *Praxibulus exsculptus* Rehn et *Brachyexarna lobipennis* Sjöstedt, du Tétrigide *Paratettix australicus* Walker, du Katydid *Conocephalus* sp. et du grillon *Bobilla victoriae* Otte & Alexander dans le nord des Tablelands en Nouvelle Galle du Sud. Au laboratoire, *Agamermis catadecaudata* n. sp. infeste et se développe sur l'acridien *Chortoicetes terminifera* (Walker). *Agamermis catadecaudata* n. sp. se distingue de toutes les espèces déjà connues du genre par la longueur des juvéniles infestants et la position du nœud. Ces juvéniles infestants perdent leur queue. Les adultes de *Agamermis catadecaudata* n. sp. sont semblables à ceux de *A. decaudata*, mais les œufs sont plus petits et le spicule plus court. Il existe deux types de juvéniles préinfestants dont l'abondance relative est variable. Une espèce non encore décrite, *Agamermis* sp. aff. *sinuosa* Kaiser, a été rencontrée dans les régions sud, centrale et nord des Tablelands; elle est sympatrique d'*Agamermis catadecaudata* n. sp. dans la région nord; ses hôtes sont inconnus.

Key-words : Entomophilic nematodes, Mermithidae, *Agamermis*, Acrididae.

During sampling of terrestrial mermithid nematodes as part of a study of nematode parasitism in acridids (Orthoptera) in the tablelands of New South Wales by G. L. B., mermithid adults, eggs and infectives of an *Agamermis* type similar to *A. sinuosa* Kaiser, 1977, were occasionally recorded. The species had a wide distribution from latitude 28-32° S. Attempted infection of third instar stage Australian plague locust nymphs, *Chortoicetes terminifera* (Walker), using L 2 larvae hatched from field collected eggs failed. No decaudate parasitic juveniles were recovered from acridids during an extensive parasite sampling program conducted between 1979-1983 (Baker, 1986) and the host of the species remained unknown.

In March 1985 A. J. Campbell, NSW Agriculture, Armidale, collected a large series of conspecific adult material of a second species of *Agamermis* from soil at Beaumont's Road, Hernani (32° 22' S 152° 28' E). Orthopteran insects collected from the pasture at the sampling site were heavily parasitised by parasitic juveniles

of an *Agamermis* species. Infection of *C. terminifera* in the laboratory using L2 larvae hatched from eggs was successful.

The species is new to science and is herein described. The description is based solely on material from the Hernani area. However, a detailed comparison is made of eggs and L2 infectives from other locations. Specimens were heat killed, fixed in 3% formaldehyde and then processed to glycerine. In the description the first figure is the dimension of the holotype (female) and allotype (male) and the figures in parentheses the range of paratype specimens. In addition to those specimens used in the description a further large series of females, eggs and progeny were measured.

Morphometric data were analysed using regression analysis and analysis of variance.

The species of *Agamermis* described in this paper has been the subject of studies on behaviour (Robinson *et al.*, 1990) and parasitic development (Baker *et al.*, 1988).

Agamermis Cobb, Steiner & Christie, 1923

DIAGNOSIS (Kaiser, 1991, *emend.*).

Medium to very large nematodes, 10–465 mm in length. Mouth opening terminal. Amphids small. Six head papillae, lateral and sub-medial in position. Cuticle with criss-cross fibres. In mid-body six hypodermal chords. Tail tip in both sexes bluntly rounded. Male : with two short spicules, genital papillae in 4–6 irregular rows. Female : vagina S-shaped, vulvar cone well developed, cuticularised. Infectious stage : amputates tail, i.e. 75 % of body, at a preformed node just before penetrating the host. Postparasitic and parasitic juveniles with craterlike appendage from the decaudate tail. Eggs : large without byssii.

TYPE SPECIES

- Agamermis decaudata* Cobb, Steiner & Christie, 1923
- = *A. decaudata* var. *paraguayensis* Steiner, 1924 n. syn.
- = *A. angusticephala* Steiner, 1924 n. syn.
- = *A. pachycephala* Steiner, 1928 n. syn.

OTHER SPECIES

- A. unka* Kaburaki & Imamura, 1932
- A. cobbi* Schuurmans-Stekhoven & Mawson, 1955
- A. sinuosa* Kaiser, 1977
- A. changshaensis* Bao, Lou & Lou, 1992
- A. catadecaudata* n. sp.

SPECIES INQUIRENDA

- A. parva* Rubtsov, 1976

SPECIES FORMERLY PLACED IN THE GENUS

- Agamermis dubia* (Steiner, 1924) n. comb.
- = *Agamermis dubia* Steiner, 1924
- Agamermis hangaica* (Rubtsov, 1976) n. comb.
- = *Agamermis hangaica* Rubtsov, 1976
- Agamermis saldulae* (Rubtsov, 1969) n. comb.
- = *Agamermis saldulae* Rubtsov, 1969
- Hexamermis microamphidis* Steiner, 1925
- = *Agamermis paradecaudata* Steiner, 1925 n. syn. (male)
- Oesophagomermis terricola* (Hagmeier, 1912) Artyukhovski, 1969
- = *Agamermis terricola* (Hagmeier, 1912) Kiryanova, Karavaeva & Romanenko, 1959
- = *Mermis terricola* Hagmeier, 1912
- Schuurmanimermis couturieri* (Schuurmans Steckhoven & Mawson, 1955) Rubtsov, 1978
- = *Agamermis couturieri* Schuurmans Steckhoven & Mawson, 1955

REMARKS

A. decaudata var. *paraguayensis*, *A. angusticephala* and *A. pachycephala* are here synonymised with *A. decaudata*. *A. decaudata* var. *paraguayensis* is distinguished from *A. decaudata* in having a ventral cleft between the ventral

sub-medial head papillae (Steiner, 1924). This cleft is not a good diagnostic character being also present in *A. decaudata* as figured in Kaiser (1991) and is also present in *A. catadecaudata* n. sp. between both the dorsal and ventral submedial head papillae. The description of *A. decaudata* var. *paraguayensis* by Steiner (1924) is regarded as the first description of *A. decaudata*, which was subsequently illustrated by Nickle (1972), Camino *et al.* (1986) and Kaiser (1991). *A. angusticephala* is most probably the postparasitic juvenile of *A. decaudata* var. *paraguayensis*, having the same host and locality data. *A. pachycephala* is described from juvenile specimens with a head which is holocephalic while *A. decaudata* is described as hemicephalic (Steiner, 1928) as is the head of *A. decaudata* var. *paraguayensis* (Steiner, 1924) and the differences could be attributed to their being different stages of a single species. The fact that Leidy's material, on which the description of *A. pachycephala* is based, was from the grasshopper *Dissosteira carolina* (L.) and apparently common throughout Philadelphia would further indicate the species to be *A. decaudata*.

A. parva is declared *species inquirenda* on the grounds that the adults were inadequately described for assignment to the genus *Agamermis* being described from an immature female and the egg and preparasitic juvenile are unknown (Rubtsov, 1976 *b*). The unusually shaped vagina is not characteristic of *Agamermis* and it may represent a new genus.

A. terricola was designated by Artyukhovski (1969) as the type species of the genus *Oesophagomermis*. *A. couturieri* was considered to be a species of *Amphimermis* by Kaiser (1977) and was subsequently designated by Rubtsov (1978) as the type species of the genus *Schuurmanimermis*. *A. dubia*, *A. saldulae* and *A. (?) hangaica* were described from juvenile material and are assigned to the genus *Agamermis* on the basis of criteria outlined by Poinar and Welch (1981). *A. saldulae* has a scab on the terminus of the tail and is probably *Agamermis*. However, *A. dubia* and *A. (?) hangaica* do not have a scab on the terminus of the tail (Steiner, 1924; Rubtsov, 1976 *a*) and do not belong to the genus *Agamermis*.

A. paradecaudata is considered a junior synonym of *Hexamermis microamphidis* because of the dubious assignment of material from the same host. Of the 28 nematode specimens ex *Helopeltis antonii* Sign (Hemiptera : Miridae), all the parasitic juveniles and females were assigned to a new species *H. microamphidis*, and all the males and a parasitic juvenile based on a sketch, to *A. paradecaudata*. The authors consider it probable that Steiner's material was the male and female of a single species of *Hexamermis*. Steiner (1925) gives no reason for linking a sketch of the parasitic juvenile, which is a typical *Agamermis* parasitic juvenile, with all the male material. The meagre description of *H. microamphidis* by Steiner (1925) is accepted as valid as convention would otherwise name the species *Hexamermis parade-*

caudata (Steiner, 1925) n. comb which would unfortunately provide a species of *Hexameremis* with a specific name indicating a decaudate preparasitic juvenile.

***Agameremis catadecaudata* n. sp.**

(Figs 1-4)

MEASUREMENTS

Female (n = 16) : L = 154 (56-200) mm; diam. mid-body = 326 (242-326) μm ; head diam. (at level of cephalic papillae) = 62 (55-72) μm , at neck = 72 (65-85) μm ; body diam. at nerve ring = 148 (120-150) μm ; cuticle width at nerve ring = 21 (5-21) μm , at mid-body = 12 (6-21) μm ; width hypodermis, at mid-body = 10 (7-20) μm ; amphid aperture (residual); amphid pouch (residual); distance of nerve ring from mouth = 319 (300-380) μm ; V = 51.29 (51.2-58.0); length of vagina = 318 (187-318) μm ; diameter of vagina = 156 (80-162) μm ; width of lateral hypodermal chord = 32 (17-45) μm ; distance of vestigial anus from tail = 199 (150-232) μm ; diameter of eggs, in uterus = 118 (110-130) μm , laid = 130 (110-130) μm .

Male : (n = 12) : L = 16.2 (5.4-24.0) mm; diam., mid-body = 121 (88-185) μm ; head diam. at level of cephalic papillae = 54 (42-58) μm , at neck = 52 (52-62) μm ; body diam. at nerve ring = 80 (67-105) μm ; cuticle width at nerve ring = 12 (3-12) μm , mid-body = 14 (3-30) μm ; hypodermis width mid-body = 10 (5-10) μm ; amphid aperture = 1 μm ; amphid pouch 3 \times 5 μm ; distance of nerve ring from mouth = 275 (187-280) μm ; spicule length = 97 (77-137) μm ; spicule head width = 13 (7-20) μm ; spicule width mid-shaft = 9 (5-10) μm ; tail length = 124 (102-150) μm ; tail width = 103 (75-124) μm ; position of proximal papillae anterior to cloaca = 84 (45-120) μm ; number of genital papillae = 34 (34-48).

Preparasitic juvenile (n = 25) : L = 2.6 mm (2.3-3.3 mm); mid-body diam. = 15 μm (11-18); head diam. = 7 μm (7-8); buccal cavity length = 30 μm (30-32); stylet length = 28 μm (25-30); distance of nerve ring from mouth = 98 μm (98-140); body width at nerve ring = 12 μm (12-19); position of stichosome = 150 μm (150-210) to 220 μm (220-400); position of node as per cent of body length = 12.6 (12.2-17.2).

DESCRIPTION

General : Body length exhibiting extreme sexual dimorphism, females being 2-20 \times the length of males. This is greater than for any other species of *Agameremis* for which both sexes have been described. Mouth terminal. Amphids poorly developed and making no incursion into the lateral cephalic papillae; simply forming void in cuticle with very indistinct connection to outer cuticle.

Female : Amphids less distinct than in males, reduced to rudimentary duct in close association with second errant nerve ending in neck region. Amphid not visible




Fig. 1. *Agameremis catadecaudata* n. sp., female. A : Head, dorsal view; B : Head, lateral view; C : Tail, lateral view; D : Vagina, ventral view; E : Vagina, lateral view; F : Egg.

in *en face* view nor in lateral view. Vagina, though horn-shaped, closely opposed to uterus with junction in same transverse plane. Vaginal opening oblique slit. Vestigial anus well developed with cone-shaped external pore and filamentous tubules within hyaline cuticular body, base of which lies in depression or pocket in hypodermis. Eggs undeveloped when laid.

Male : Head rounded, slightly set off from body (neck diameter sometimes less than width at level of cephalic papillae). Bilateral symmetry of head pronounced, large dorsal and ventral cuticular incursions separating paired submedial papillae. Lateral papillae broader than submedial papillae. Amphids indistinct, set posterior (approx. 10 μm) and slightly lateral to sub-medial papillae. Spicule relatively narrow, evenly curved, head slightly flared due to prolongation of ventral wall; ventral wall slightly greater length than dorsal wall. Spicule length approximately equal to body width at cloaca. Tail length greater than spicule length (\times 1.1-1.5). Proximal genital papillae less than spicule length anterior to cloaca (\times 0.6-0.9). Genital papillae in three rows of unequal length, medial row bifurcate in region of cloaca. Medial and




Fig. 2. *Agameremis catadecaudata* n. sp., male. A: Head, dorsal view; B: Head, lateral view; C: Tail, lateral view; D: Tail, schematic view; E: Tail, ventral view; F: Cross section, mid-body; G: Head, en face.

sub-lateral rows with reduced number of papillae (two to four) as compared with nine to twelve papillae in sub-medial and lateral rows.

Juvenile st. 2 (preparasitic): Long, slender larvae. Proximal 10-20% of body broader than remainder. Large stylet with broad head, barbed on one side. Paired penetration glands immediately anterior to stichosome. Stichosome short in relation to body length. Stichocytes undifferentiated. Node, between primordial (vestigial) gonads and trophosome, distinct and composed of two to seven transverse plates. Trophosome interspaces distinct, rectangular and number approximately 30. Tail attenuated to a fine point.

Parasitic and postparasitic juveniles: Preparasitic juvenile sheds trophosome on entering the host. Body of uniform width along entire length. Tail with raised concave stub formed from scar of decaudate tail. Stichocytes distinct during early development, either paired or single, numbering fourteen to fifteen.

Postparasitic juveniles with rounded tail. Stub retained. Pharynx thickened at proximal end. Double moult to adult stage.

TYPE HOST AND LOCALITY

Praxibulus exsculptus Rehn (Orthoptera: Acrididae). Beaumont Road, Hernani (32° 22' S 152° 28' E), New South Wales, Australia.

HOST RANGE

In addition to the type host, *P. exsculptus*, *A. catadecaudata* n. sp. has been reared from *Phaulacridium vittatum* (Sjöstedt), *Brachyexarna lobipennis* Sjöstedt (Orthoptera: Acrididae), *Paratettix australicus* Walker (Orthoptera: Tetrigidae), *Conocephalus* sp. (Orthoptera: Tettigoniidae) and *Bobilla victoriae* Otte & Alexander (Orthoptera: Gryllidae). In the laboratory *A. catadecaudata* n. sp. readily infected and developed in *Chortoicetes terminifera* (Walker) (Orthoptera: Acrididae).

TYPE SPECIMENS

Holotype (female) and **allotype** (male) in Department of Nematology, University of California, Davis, USA (UCD). **Paratypes** (one male and one female deposited in South Australian Museum, Adelaide, Australia (acquisition numbers AHC 18807-18809) and Laboratoire de Biologie Parasitaire, Protistologie, Helminthologie, Muséum National d'Histoire Naturelle, Paris, France.

DIAGNOSIS AND RELATIONSHIPS

A. catadecaudata n. sp. differs from all described species of *Agameremis* in regard to the diameter of the egg in relation to body length in the female; body length and position of the node in the preparasitic juvenile; and spicule length in relation to body length in the male.

The original description of *A. decaudata* contained no dimensions and in the following study of relationships the dimensions are from a redescription by Nickle (1972). Further dimensions given by Camino *et al.* (1986) fall within the range given by Nickle (1972) unless otherwise indicated.

Female *A. catadecaudata* n. sp. differ from *A. decaudata* in the generally shorter body length: 56-200 *vs* 50-465 mm (30-465 mm in Camino *et al.*, 1986) and smaller eggs, especially in relation to body length (110-130 *vs* 150-180 µm) and narrower mid-body width (approx. 300 *vs* 500 µm); *A. unka*, *A. sinuosa* and *A. changshaensis* in the greater body length (56-200 *vs* 24-34, 30-37 and 23-41 mm respectively) and relatively larger eggs in relation to female length (110-130 *vs* 70-75, 95-110 and 105-140 µm respectively) and *A. cobbi* in the shorter body length (56-200 *vs* 213-323 mm) and smaller eggs (110-130 *vs* 150-200 µm).

Male *A. catadecaudata* n. sp. differ from *A. decaudata* in having a generally smaller body length (5.4-24 *vs* 10-120 mm in Nickle, 1972) and smaller spicule length,




Fig. 3. *Agameremis catadecaudata* n. sp., egg and preparasitic juvenile. A : Egg; B : Head of preparasitic juvenile with skin of st. 1 attached; C : Node; D : Whole body; E : Anterior region. (Bar equivalents : A, D, E = 50 μ m; B, C = 10 μ m.)




Fig. 4. *Agameremis catadecaudata* n. sp., parasitic and post-parasitic juvenile. **A**: Six day-old parasitic juvenile (stichosome ruptured); **B**: Parasitic, juvenile, scab of residual node; **C**: Head, post-parasitic juvenile undergoing double moulting; **D**: Tail, post parasitic juvenile undergoing double moulting. (Bar equivalents: A, C, D = 100 μ m; B = 10 μ m.)

especially in relation to body length (77-137 vs 150-180 μ m); from *A. unka* and *A. changshaensis* in much smaller spicule length (77-137 vs 190-230 and 170-267 μ m respectively), especially significant given the similarity of the body length of the three species (5.4-24 vs 11-22 and 13-28 mm, respectively); from *A. sinuosa*

in arrangement of the genital papillae (three rows, median row bifurcate in region of cloaca vs six rows). The male of *A. cobbi* is unknown.

Preparasitic juveniles differ from *A. decaudata* (based on paralectotype material from Falls Creek, Virginia collected by Christie and held in UCD), in generally short-




Fig. 5. *Agameremis catadecaudata* n. sp. Frequency distribution histograms of female length, egg diameter, length of infective and position of node as a per cent of body length.

er length (2.3-3.3 vs 2.0-5.6 mm for *A. decaudata*); and in the mean lengths of two forms (2.6 and 3.1 vs 2.1 and 5.6 mm); position of node of the two forms (12 and 17 vs 15%) with little variation; length in relation to egg

size (130 and 140 vs 120 and 163 µm); from *A. sinuosa* in the greater length (2.3-3.3 vs 1.6 mm), position of the node (12-17 vs 17-21%) and the greater length of the stichosome in relation to length of body anterior to node (35 vs 26%) and from *A. unka* and *A. changshaensis* in greater length (2.3-3.3 vs 0.6 and 1.4-1.6 mm respectively) and *A. cobbi* in the shorter body length (2.3-3.3 vs 4-5.3 mm).

BIOLOGY AND ECOLOGY

Extensive collections of *Agameremis* spp. were made at two sites in the Northern Tablelands of New South Wales, Hernani and Black Mountain between 1983-1985.

There were no apparent differences between adults from both sites. However, the preparasitic juveniles were highly variable both at the same site and between sites. *A. catadecaudata* n. sp. collected at Hernani were readily divided into two forms based on the bimodal nature of the relationship between body length (L) and position of node (N): *form one*, short (2.6 mm) with relatively short body anterior to node (13%), *form two*, long (3.0 mm) with a relatively long body anterior to the node (17%). The relationship between body length and position of node (L/N) within each form was negative and between forms was positive (Fig. 6).

Specimens from Black Mountain consisted of three forms; a low incidence of the two forms of *A. catadecaudata* n. sp. found at Hernani, and a third, more abundant form which was very short (1.8 mm) with a relatively long body anterior to the node (20%) (Fig. 6). The latter form had the shortest body length yet the proportionally longest body length anterior to the node. This trend is at variance with that exhibited by the two forms of *A. catadecaudata* n. sp. i.e. the shorter specimens have the proportionally shortest body length anterior to the node and the material is considered a distinct species, with close affinity to *Agameremis sinuosa* Kaiser, 1977.

The slope of the relationship L/N for *Agameremis* sp. *aff. sinuosa* from Black Mountain is negative. This is the same relationship that occurs in both forms one and two of *A. catadecaudata* n. sp. This is in contrast with the positive relationship exhibited between forms one and two of *A. catadecaudata* n. sp. and substantiates the status of *A. sp. aff. sinuosa* as a separate species to *A. catadecaudata* n. sp.

The host of *A. sp. aff. sinuosa* is unknown. The host of *A. sinuosa* in Europe is Cicadellidae (Kaiser, 1977). No Cicadellidae were dissected during the course of this study. No parasitism of acridids (*P. vittatum* and *B. lobipennis*) was recorded at the Black Mountain site where *A. sp. aff. sinuosa* predominated and conversely no *A. sp. aff. sinuosa* was found at Hernani where parasitism of acridids was common. It is not known if parasitism of acridids by *A. catadecaudata* n. sp. at the Hernani site was caused by form one or two or both but it




Fig. 6. The relationship between female egg diameter and length of infective and position of node. A, B : *Agameremis catadecaudata* n. sp. population from Hernani; C, D : Mixed population of *A. catadecaudata* n. sp. (●) and *A. sp. aff. sinuosa* (▲) from Black Mountain.

appears the host range of *A. sp. aff. sinuosa* is different to that of *A. catadecaudata* n. sp.

A. catadecaudata n. sp. was also present at Black Mountain, but had a very low incidence. Given the abundance of acridid hosts, the low incidence of adults in soil samples and apparent absence of parasitism in acridids is unexplained.

A bimodal frequency distribution in both egg size and infective length has been recorded for *A. decaudata* in North America and both forms were accepted as being of the same species (Christie, 1936). Similarly, and in the absence of evidence of reproductive isolation, the two forms of *A. catadecaudata* n. sp. are considered to represent a single species.

Discussion

The hosts of the described species vary considerably: *A. decaudata* is recorded from grasshoppers (Orthoptera: Acrididae), *A. sinuosa* is apparently specific to cica-dellids (Kaiser, 1977), *A. unka* and *A. changsaensis* to the brown rice planthopper *Nilaparvata lugens* Stal and whitebacked planthopper *Sogatella furcifera* (Horvath) (Hemiptera: Delphacidae) (Bao *et al.*, 1992; Choo & Kaya, 1993) and *A. catadecaudata* n. sp. to a range of orthopteran insects. The host of *A. cobbi* is unknown.

A. decaudata and *A. catadecaudata* n. sp. share many features including the same extreme sexual dimorphism, bimodal distribution of egg diameter, infective length and position of node and the same host range (Orthoptera) and are obviously very closely related. *A. unka*, *A. changsaensis* and *A. sinuosa* do not exhibit sexual dimorphism to the same extent as *A. decaudata* and *A. catadecaudata* n. sp.

Cobb *et al.* (1923) and Christie (1936) state there is a positive relationship between female length and egg diameter in *A. decaudata*. In *A. catadecaudata* n. sp. there is no significant relationship between egg diameter and female length [P (egg diam.) = 0.2; correlation coefficient < 0.1].

There is a strong positive relationship between egg diameter and length of the infective (correlation coefficient > 0.4) (Fig. 6 A). The data are evenly distributed (unimodal with low SE) which is again at variance with the bimodal distribution displayed by *A. decaudata*. Poinar (1979) suggested the difference in length of infectives probably reflected differences in the availability of nutrients during egg development.

The fact that the relationship L/N is both negative and of similar slope within the two forms but is positive between forms, possibly indicates incipient differentiation into sibling species. The differentiation into sibling species is not as advanced as that apparent in *A. decaudata* where in addition to infective length, egg diameter and female length also have a bimodal distribution.

The failure to record a decaudate parasitic stage during the survey of parasites of acridids throughout the tablelands of New South Wales contrasts with the high incidence of *A. catadecaudata* n. sp. in acridids at Hernani. This indicates differences in host susceptibility to infection between Hernani and the remainder of the State. No eggs or infectives of *A. sp. aff. sinuosa* have been recorded from the Hernani area, indicating there may be some spatial separation of the two species. Mixed populations were recorded at Springmount (Black Mountain) with *A. sp. aff. sinuosa* predominating, whereas only *A. sp. aff. sinuosa* has been recorded in the central and southern tablelands. There also appears to be a regional variation in the ability of *A. catadecaudata* n. sp. to infect acridids for, despite its occurrence in districts other than Hernani, it has only been recorded in acridids in the Hernani area. This may be a

consequence of unsatisfied environmental conditions during host infection in other districts, at least as these conditions relate to the infection of acridids. Perhaps, significantly, Hernani has the highest annual rainfall of any district in New South Wales.

Acknowledgments

The authors would like to thank Dr A. Campbell, NSW Agriculture, Armidale for providing the material from Hernani on which the description is based, Ms H. M. Holmes, formerly NSW Agriculture, Rydalmere, for assistance with processing samples and Dr A. Clift, NSW Agriculture, Rydalmere, for regression analysis of morphometric data. The data reported in this paper were obtained during studies of the population dynamics of *P. vittatum* supported by grants from the Tablelands Wingless Grasshopper Committee and Australian Meat Research Council (DAN 19 S).

References

- ARTYUKHOVSKI, A. K. (1969). [New genera of the family Mermithidae (Enoplida, Nematoda)]. *Zool. Zh.*, 48 : 1309-1319.
- BAKER, G. L. (1986). The ecology of mermithid nematode parasites of grasshoppers and locusts in south-eastern Australia. In: Samson, R. A., Vlask, J. M. & Peters, D. (Eds). *Fundamental and applied aspects of invertebrate pathology*. Wageningen, The Netherlands, Foundation 4th intern. Colloq. Invert. Pathol. : 277-280.
- BAKER, G. L., DAVISON, M. & HOLMES, H. M. (1988). The parasitic development of *Amphimermis* sp. and *Agameremis* sp. In: *Research Report 1986-1988, BCRI*. Rydalmere, NSW Agriculture & Fisheries : 41.
- BAO, X. C., LOU, Q. G. & LUO, D. M. (1992). [A new species of *Agameremis* from China (Nematoda : Mermithidae).] *Acta zootaxon. Sinica*, 17 : 20-25.
- CAMINO, N. B., BELLUSCI DE MIRALLES, D. A. & MARCHISSIO, S. L. (1986). Aportes al conocimiento de la fauna Argentina de mermithidos. 1. Sobre *Agameremis decaudata* Cobb, Steiner and Christie 1923. *Neotropica*, 32 : 67-70.
- CHOO, Y. K. & KAYA, H. K. (1993). Parasitism of brown planthopper by the mermithid nematode *Agameremis* in rice. In: Bedding, R., Akhurst, R. & Kaya, H. (Eds). *Nematodes and the biological control of insect pests*. Melbourne, CSIRO : 21-26.
- CHRISTIE, J. R. (1936). Life history of *Agameremis decaudata*, a nematode parasite of grasshoppers and other insects. *J. agric. Res.*, 52 : 161-198.
- COBB, N. A., STEINER, G. & CHRISTIE, J. R. (1923). *Agameremis decaudata* : a nema parasite of grasshoppers and other insects. *J. Agric. Res.*, 11 : 921-926.
- HAGMEIER, A. (1912). Beiträge zur Kenntnis der Mermithiden 1. Biologische Notizen und systematische Beschreibung einiger alten und neuen Arten. *Zool. Jahrb. Syst.*, 32 : 521-612.
- KABURAKI, T. & IMAMURA, S. (1932). Mermithid worm parasitic in leaf hoppers, with notes on its life history and habits. *Proc. Imp. Acad. Japan*, 8 : 136-141.

- KAISER, H. (1977). [Studies on the morphology, biometry, biology and systematics of mermithids : a contribution to the problem of separating morphologically closely related species.] *Zool. Jahrb. Syst.*, 104 : 20-71.
- KAISER, H. (1991). Terrestrial and semiterrestrial Mermithidae. In : Nickle, W. R. (Ed.). *Manual of agricultural nematology*. New York, Marcel Dekker : 899-965.
- KIRYANOVA, E., KARAVAEVA, R. & ROMANENKO, K. (1959). [Mermithids (Nematodes)- parasites of *Hyponomeuta malinella* and *H. padella* in Southern Kirgiz. S.S.R.]. *Trudi Kirgizskoi Lesnoi Opitnoi Stantsii*, 2 : 195-240.
- NICKLE, W. R. (1972). A contribution to our knowledge of the Mermithidae. *J. Nematol.*, 4 : 113-146.
- POINAR, G. O., JR. (1979). *Nematodes for biological control of insects*. Boca Raton, Florida, CRC Press : 277 p.
- POINAR, G. O., JR. & WELCH, H.E. (1981). Parasites of invertebrates in the terrestrial environments. In : Slusarski, W. (Ed.). *Review of advances in parasitology*. Warszawa, PWN-Polish Scientific Publ. : 947-954.
- ROBINSON, A. F., BAKER, G. L. & HEALD, C. M. (1990). Transverse phototaxis by infective juveniles of *Agamermis* sp. and *Hexamermis* sp. *J. Parasitol.*, 76 : 147-152.
- RUBTSOV, I. A. (1969). [On a new species of *Agamermis* (Mermithidae) of a bug in Poland]. *Acta parasit. polon.*, 16 : 97-100.
- RUBTSOV, I. A. (1976 a). [Mermithids (Nematoda, Mermithidae) parasites of insects in Mongolia]. In : *Nasekomye Mongolii, Akademiya Nauk SSR, Zoologicheskii Institut, No. 4*. Leningrad, Izdatel'stvo " Nauka ", Leningradskoe Otdelenie : 596-614.
- RUBTSOV, I. A. (1976 b). [Mermithids from Kirgizia]. In : *Entomologicheskie issledovaniya v Kirgizii. "Ilim" Frunze*, 11 : 87-100.
- RUBTSOV, I. A. (1978). [Mermithids : Classification, importance, use]. Leningrad, " Nauka " : 207 p.
- SCHUURMANS STEKHOVEN, J. H. & MAWSON, P. M. (1955). Mermithidés d'Alsace. *Annales Parasit.*, 30 : 69-82.
- STEINER, G. (1924). Beiträge zur Kenntnis der Mermithiden. 2. Teil. Mermithiden aus Paraguay in der Sammlung des zoologischen Museums zu Berlin. *Zentralbl. Bakt. Parasit.-Kde.*, 62 : 90-110.
- STEINER, G. (1925). Mermithids parasitic in the tea bug (*Helopeltis antonii* Sign.). *Meded. Proefst. Thee Batavia*, 94 : 3-16.
- STEINER, G. (1928). A revision of Leidy's collection of mermithids. *Proc. Acad. nat. Sci. Philadelphia*, 80 : 547-552.