



0040-4020(94)E0101-X

Euryspongiols: Ten New Highly Hydroxylated 9,11-Secosteroids with Antihistaminic Activity from the Sponge *Euryspongia* sp. Stereochemistry and Reduction.

José Dopeso, Emilio Quiñoá and Ricardo Riguera*

Departamento de Química Orgánica, Facultad de Química, Universidad de Santiago de C., Santiago de Compostela, 15706, Spain

Cécile Debitus

Centre ORSTOM, B. P. A5, Nouméa Cedex, New Caledonia

Patricia R. Bergquist

School of Biological Sciences, University of Auckland, Auckland, New Zealand

Abstract: Ten novel polyhydroxylated 9,11-secosterols, 1-10, have been isolated from the New Caledonian sponge *Euryspongia* sp. and their structures elucidated by spectroscopic studies, selective acetylation and reduction, leading to the complete assignment of all the NMR signals. Euryspongiols A1-A5 differ in their side chains but have the same $2\alpha,3\beta,4\alpha,6\beta,11,19$ hexahydroxy-9,11-secocholestane skeleton. Euryspongiols B1-B5 are the corresponding 3α epimers. Compounds 1-10 are the most highly hydroxylated secosteroids isolated so far, and are the first hydroxylated at C-4. NaBH_4 reduction of the carbonyl group at C-9 of 1 and 2 is stereospecific, producing the heptahydroxylated secosteroids 1a and 2b with a β C-9 hydroxyl group. Compounds 1 and 2 have been found to strongly inhibit the release of histamine from rat mastocysts.

INTRODUCTION

Steroids, as components of all living organisms, have been a frequent subject of marine studies. Apart from the large number of novel structures disclosed, interesting biogenetic and synthetic work have been described. In the last few years, advances in this field have coincided with advances in the study of water-soluble marine natural products. Specifically, there is now a well established methodology for studying the intact structure of steroid (and terpenoid) saponins.

The known water-soluble marine steroids are the polyhydroxysteroids found in sponges¹ and equinoderms² and the polyhydroxysecosteroids from sponges, soft corals and gorgonians.³

Among this latter class of compounds, the 9,11-secosteroids are the most abundant. Structurally, all 9,11-secosteroids have a keto group at C-9 and a side chain like those usually found in "normal" steroids;⁴ differences reside in the A-B ring fusion (*cis/trans* or Δ^5) and the number and position of the hydroxyl (or epoxide) groups, which are always located on rings A and B.

Fonds Documentaire IRD



010026189

3813

Fonds Documentaire IRD

Cote : Bx 26189 Ex :

Trans decalin 9,11-secosteroids have been found in the sponges *Aplysilla glacialis*⁵ (glaciasterols A and B) and *Spongia officinalis*⁶ both *trans* and Δ^5 decalin 9,11-secosteroids in the soft coral *Sclerophyllum* sp.⁷ and Δ^5 -secosteroids in the gorgonian *Pseudopterogorgia americana*³ and the soft coral *Sinularia* sp.^{8,9} To date, the sole *cis* decalin system reported is herbasterol, found by Faulkner *et al.* from *Dysidea herbacea*.¹⁰

According to the degree and pattern of hydroxylation, the simplest compounds of this class are the 3,11-dihydroxy and the 3,6,11-trihydroxy-9,11-secosteroids.^{3,6,7} Glaciasterols⁵ have four oxygenated sites (3,5,6 and 11), with hydroxyl groups at C-3 and C-11 and an epoxide bridge at C-5, C-6. Herbasterol¹⁰ which is pentahydroxylated at 2,3,6,11 and 19, has the highest degree of oxygenation hitherto reported in the secosteroid literature.

RESULTS AND DISCUSSION

In the course of our investigations on marine invertebrates we have now isolated ten hexahydroxylated 9,11-secosterols from the polar extracts of a new species of a marine sponge *Euryspongia* sp. (Porifera, Demospongiae, family Dysideidae), which was collected on the coast of New Caledonia and selected for study because of the cytotoxicity and strong antihistaminic activity of its extracts.

The structure of the new compounds which we have named euryspongiols (1-10, Fig. 1), were elucidated by spectral analysis and chemical transformations.

The freshly collected sponge (556.30 g) was extracted, defatted and chromatographed to afford a fraction containing the secosteroids, which were separated by reversed phase HPLC to yield compounds 1 (30 mg), 2 (32 mg), 3 (1.5 mg), 4 (1 mg), 5 (7 mg), 6 (17 mg), 7 (5 mg), 8 (1.5 mg), 9 (1.5 mg) and 10 (6 mg).




Figure 1. Structures of the 9,11-secosterols from *Euryspongia* sp.

These secosteroids can be classified in two series, euryspongiols A (compounds 1-5) and euryspongiols B (compounds 6-10), according to their stereochemistry at C-3. For the sake of simplicity, their structural

091
123

features will be presented in accordance with this classification.

Eurysongiol A1 (1). The molecular formula of the major component **1** was determined by high resolution EIMS as $C_{27}H_{46}O_7$ (5 unsaturations). EIMS showed main fragments at m/z 452 corresponding to $C_{26}H_{44}O_6$ $[M-CH_2O]^+$; 434 $[M-CH_2O-H_2O]^+$; 416 $[M-CH_2O-2H_2O]^+$ and 398 $[M-CH_2O-3H_2O]^+$. The presence of an unsaturated C_8H_{15} steroidal side chain was indicated by an ion at m/z 341 $[M-CH_2O-C_8H_{15}]^+$. Some of the most relevant fragments are shown in Figure 2.

Negative ion mode FAB MS produced a prominent fragment at m/z 451 $[M-H-CH_2O]^-$, while positive ion mode FAB MS in a glycerol matrix showed the expected $[M+H]^+$ ion at m/z 483 and also the loss of a molecule of water at m/z 465 $[M+H-H_2O]^+$, loss of five molecules of water at m/z 393 $[M+H-5H_2O]^+$, and loss of the side chain at m/z 354 $[M+H-H_2O-C_8H_{15}]^+$.




Figure 2. Fragments observed in EIMS of compound **1**.

The ^{13}C NMR spectrum of **1** contained signals for all twenty seven carbon atoms. A ketone signal at δ 215.4 (C-9) and a pair of sp^2 carbons at δ 138.4 and δ 128.4 (C-22/23) indicated the presence of a carbonyl group and a disubstituted double bond, so only three rings (as against four in a normal steroid) are present to justify the unsaturation number, suggesting a secosteroid structure. Six resonances between δ 82.8 and 59.2 supported the existence of six sites with a heteroatom (oxygen) substituent. On the basis of DEPT experiments, two of these signals (δ 63.9 and 59.2) were assigned to hydroxymethylene carbons and the other four (δ 82.8, 71.1, 70.5 and 64.7) to hydroxymethines (see Table 1).

Due to the polyhydroxylated structure of **1**, pyridine-induced deshielding¹¹ was prominent and the 1H NMR pattern varied greatly as compared with the spectrum taken in CD_3OD (see Table 2). In pyridine- d_5 , compound **1** showed signals for four of the five methyl groups of a C-27 sterol at δ 0.84 (6H, d, H-26/27), 0.91 (3H, s, H-18), 1.03 (3H, d, H-21).

The singlet that one would expect for the C-19 methyl protons is absent from the spectrum of **1**, and instead doublets were observed at δ 5.73 and 4.40 ppm (1H each, $J = 11.7$ Hz) corresponding to an isolated C-19 hydroxymethylene group. A second hydroxymethylene group gave rise to signals at δ 4.20 (1H) and 4.10 (1H) that agree well with the C-11 protons of a 9,11-secosterol.

In spite of the information provided by the pyridine-induced effects, the overlapping of signals was too extensive for complete analysis of the relevant spin systems and identification of signals by 2D correlations. However, a combination of the NMR data (COSY and HETCOR) for **1**, and various derivatives allowed complete signal assignment and stereochemical analysis, as follows.

The mutually coupled signals observed in the COSY spectrum at δ 4.48 (1H, dd, $J = 8.7, 4.7$ Hz), 3.95

(1H, t, $J = 8.7$ Hz) and 4.58 (1H, dd, $J = 11.2, 8.7$ Hz) were taken to correspond to the three axial hydroxymethine protons of a $2\alpha,3\beta,4\alpha$ -trihydroxysterol.

Table 1. ^{13}C NMR Spectral data for the secosterols **1**, **1a** and **7**

Carbon No.	1 δ , DEPT (CD ₃ OD)	1a δ , DEPT (CD ₃ OD)	7 δ , DEPT (CD ₃ OD)
1	35.5, t	37.0, t	30.6, t
2	71.1, d	70.8, d	68.4, d
3	82.8, d	83.4 ^b , d	74.3, d
4	70.5, d	70.6, d	68.2, d
5	56.3, d		50.2, d
6	64.7, d	64.8, d	65.1, d
7	41.3, t	41.3 ^a , t	41.4, t
8	39.7, d	39.7, d	39.9, d
9	215.4, s	83.2 ^b , d	215.0, s
10	56.2, s		56.1, s
11	59.2, t	59.1, t	59.2, t
12	41.3, t		41.2, t
13	46.6, s	47.0, s	46.7, s
14	43.1, d		42.9, d
15	23.9, t	22.7, t	23.8, t
16	26.7, t	27.3, t	26.8, t
17	51.0, d		50.8, d
18	18.1, q	18.6, q	17.8, q
19	63.9, t	63.6, t	63.5, t
20	39.7, d	40.0 ^a , d	35.6, d
21	22.3, q	22.7, q	19.8, q
22	138.4, d	138.6, d	36.7, t
23	128.4, d	128.3, d	25.6, t
24	43.2, t		40.6, t
25	29.7, d	29.7, d	29.1, d
26	22.7, q	22.2, q	22.9, q
27	22.7, q	22.2, q	22.9, q

^{a,b} Assignments may be interchanged

The HETCOR spectrum showed correlation of the peak at δ 82.8 attributed to C-3 with the triplet at 3.95 (H-3) and correlation of the signals at δ 71.1 and 70.5 with the multiplets at δ 4.48 and 4.58 assigned to H-2 and H-4. The remaining signal at δ 5.05 (1H) was then attributed to an equatorial hydroxymethine proton α H-6 by analysis of the C-4/C-5/C-6 spin system (*vide infra*).


Table 2. ¹H NMR Spectral data for the secosterol **1**, **1a** and **7**

Proton No.	1 δ , mult, J(Hz) (C ₅ D ₅ N)	1 δ , mult, J(Hz) (CD ₃ OD)	1a δ , mult, J(Hz) (CD ₃ OD)	7 δ , mult, J(Hz) (CD ₃ OD)
1 α	2.20	1.45	0.99 t, 12.7	1.64
1 β	3.12 dd, 13.3, 4.7	2.34 dd, 13.7, 4.8	2.65 dd, 12.7, 4.6	2.00 dd, 13.2, 4.15
2	4.48 dd, 8.7, 4.7,	3.69	3.66	3.82
3	3.95 t, 8.7	3.10 t, 8.9	3.14 t, 8.9	3.93 dd, 2.9, 2.4
4	4.58 dd, 11.2, 8.7	3.64	3.60	3.80 dd, 11.4, 2.4
5	2.05 dd, 11.2, 2.7	1.49 dd, 11.3, 2.3	1.20 dd, 11.3, 2.3	1.86 dd, 11.4, 2.4
6	5.05	4.32 dd, 2.4, 2.4	4.15 dd, 2.4, 2.4	4.28 dd, 2.4, 2.4
7 α	2.45	2.32	1.92	2.28
7 β	1.70	1.56	1.29	1.52
8	4.20	3.46 ddd, 13.6, 4.6, 4.1	1.65	3.47 ddd, 13.2, 4.6, 4.1
9			2.98 d, 10.8	
11	4.10	3.66	3.70	3.60
11'	4.20	3.66	3.70	3.60
12	2.00	1.64	1.78	1.67
14	3.00	2.49	2.32	2.47
15	1.55	1.37	1.48	1.37
16	1.48	1.36	1.44	1.37
17	1.70	1.60	1.59	1.53
18	0.91 s	0.81 s	0.82 s	0.77 s
19	4.40 d, 11.7	3.70	3.64 d, 12.1	3.70 d, 11.5
19'	5.73 d, 11.7	4.85	4.43 d, 12.1	4.78 d, 11.5
20	2.10	2.24	2.16	1.42
21	1.03 d, 6.7	1.07 d, 6.8	1.06 d, 6.8	1.00 d, 6.4
22	5.28	5.36	5.33	
23	5.28	5.36	5.33	
24	1.80	1.88 dd, 9.4, 3.3	1.86	1.10
25	1.50	1.53	1.61	1.54
26	0.84 d, 6.6	0.91 ^a d, 6.6	0.87 d, 6.6	0.87 d, 6.7
27	0.84 d, 6.6	0.89 ^a d, 6.6	0.87 d, 6.6	0.87 d, 6.7

^a Assignments may be interchanged

The coupling constants corresponding to H-5 (δ 2.05, dd, J = 11.2, 2.7 Hz) are indicative of its *trans* diaxial relationship with H-4 and axial-equatorial relationship with H-6. The *trans* A/B ring fusion in **1** can be deduced by comparison of the C-13 chemical shifts, (especially that of the C-19 hydroxymethyl group) with the calculated values for *cis* and *trans* decalines. Addition of the standard -OH group α effect (+49 ppm) to the chemical shift of the methyl group in *trans* (15.7 ppm) and *cis* (28.2 ppm) 9-methyldecalin predicts shifts of

64.7 ppm and 77.2 ppm for the corresponding *trans* and *cis* 9-hydroxymethyl decalines respectively; thus the experimental value of 63.9 ppm for **1** clearly implies *trans* fusion.¹² Furthermore, direct experimental evidence of *trans* fusion came from a NOE experiment run on derivative **1a** (Fig. 3), which showed NOEs among H-5 (ax) and H-3 (ax) and H-9 (ax) and H-7 (ax) (Table 3).



- 1a** R = OH R' = OH R'' = a
1b R = OAc R' = OAc R'' = a
2a R = OAc R' = O R'' = b
2b R = OAc R' = OH R'' = b
2c R = OAc R' = OAc R'' = b

Figure 3

The signal due to H-8 is especially difficult to identify in this type of secosteroids, not only because of the complexity of the spectra and the rotation around the C-8/C-14 bond, but also because these compounds are not structurally stable under basic treatment,¹¹ which in consequence cannot be used to simplify the spectra by deuterium exchange. In C_5D_5N , and due to overlapping, the coupling constants of H-8 with H-7/7' and H-14 could not be measured. In CD_3OD , they could be obtained ($J = 13.6, 4.6, 4.1$ Hz), and were assigned taking into account the *trans* diaxial relationship between H-8 and H-9 (10.8 Hz in **1a**). Thus H-8 lies in axial position, with coupling constants of 13.6 Hz with α -H-7 (ax) and 4.6 and 4.1 Hz with β -H-7 (eq) and H-14 respectively.

Comparison of the 1H and ^{13}C NMR data for the side chain with those of the literature¹³ suggested the presence of an E Δ^{22} double bond in **1**.

Taken together, the above evidence indicates that compound **1**, named eurysspongiol A1, is 2 $\alpha,3\beta,4\alpha,6\beta,11,19$ -hexahydroxy-9,11-secocholest-22(E)-en-9-one.

Table 3. NOEs observed in compound **1a**

Irradiated proton	NOE				
	H-5	H-1 α	-----	-----	-----
H-3	H-5	H-1 α	-----	-----	-----
H-9	H-14	H-5	H-1 α	-----	-----
H-5	H-6	H-3	H-9	H-7 α	H-1 α
H-1 α	H-3	H-9	H-1 β	H-5	-----
H-6	H-7 α	H-5	-----	-----	-----
H-19'	H-19	-----	-----	-----	-----

Eurysspongiols A2-A5 (2-5). The molecular formula of compound **2**, eurysspongiol A2, was deduced as $C_{27}H_{48}O_7$ from HREIMS, FAB MS (negative ion mode) and ^{13}C NMR/DEPT spectral data, ^{13}C and 1H chemical shift values in compound **2** were virtually identical to those of compound **1**, the absence of the double bond being the only difference and suggesting that **2** is the saturated counterpart of **1**. An EIMS fragment at m/z 323 [$M-CH_2O-H_2O-C_8H_{17}$]⁺ confirmed the presence of a saturated C_8H_{17} side chain in **2**. The ^{13}C NMR spectra of **3** and **4**, eurysspongiols A3 and A4 respectively, contained signals for twenty eight carbon atoms,

while compound **5**, eurysspongiol A5, had twenty six. Comparison of their ^1H and ^{13}C -NMR data with those of **1** and **2** indicate that all the secosteroids of the A series share the same skeleton, substitution pattern and regio; and stereochemistry, and differ only in their side chain structures.

Both **3** and **4** have the same molecular formula, $\text{C}_{28}\text{H}_{48}\text{O}_7$, showing the presence of a disubstituted double bond (δ 137.7, C-23; δ 135.2, C-22) and one more methyl group than in **1** and **2** (**3**, δ 0.94, d, $J = 6.9$ Hz, Me-28; **4**, δ 0.96, d, $J = 6.9$ Hz, Me-28). Both these features are located in their side chains, and make them (22E)-24-methyl- Δ^{22} -sterols.¹⁴ Under (-)FAB MS, both compounds had the same $[\text{M}-\text{H}]^-$ ion (m/z 495) and the same fragmentation pattern, suggesting that these two isomers are epimers at C-24. Their EIMS showed ions at m/z 466 $[\text{M}-\text{CH}_2\text{O}]^+$ and m/z 341 $[\text{M}-\text{CH}_2\text{O}-\text{C}_9\text{H}_{17}]^+$, confirming the presence of the unsaturated C_9H_{17} side chains. Their stereochemistry at C-24 was deduced by comparison with known 24R and 24S epimers of (22E)-24-methyl- Δ^{22} -sterols, using the proton chemical shift of the Me-21 protons as diagnostic signal (the 24R epimer signal lies downfield of the 24S signal.^{14,15} Since Me-21 resonates at δ 1.06 (d, $J = 6.9$ Hz) in **3** at δ 1.07 (d, $J = 6.8$ Hz) in **4**, compound **3** is $2\alpha,3\beta,4\alpha,6\beta,11,19$ -hexahydroxy-9,11-secosteroid-(22E,24S)-24-methyl-en-9-one and compound **4** $2\alpha,3\beta,4\alpha,6\beta,11,19$ -hexahydroxy-9,11-secosteroid-(22E,24R)-24-methyl-en-9-one.

Secosterol **5**, eurysspongiol A5, has the molecular formula $\text{C}_{26}\text{H}_{44}\text{O}_7$, corresponding to a norsecosteroid. A (+)FAB MS experiment in a NaCl+glycerol matrix produced an $[\text{M}+\text{Na}]^+$ ion at m/z 491 and a fragment at m/z 394 $[\text{M}+\text{Na}-\text{C}_7\text{H}_{13}]^+$ due the loss of the unsaturated C_7H_{13} side chain. The ^1H NMR spectrum of **5** contained resonances that could be assigned to one tertiary methyl group (δ 0.78, s, 3H, Me-18) and three secondary methyl groups (δ 0.96, 6H, d, $J = 6.7$ Hz, Me-25/26 and 1.03, 3H, d, $J = 6.8$ Hz, Me-21). The ^{13}C NMR spectrum showed two sp^2 carbons at δ 136.0 and 134.0 ppm identified as C-22 and C-23. These data, together with comparison with the literature,¹⁶ indicate that **5** is $2\alpha,3\beta,4\alpha,6\beta,11,19$ -hexahydroxy-9,11-secosteroid-(22E)-en-9-one.

Eurysspongiols B1-B5 (6-10). Comparison of their spectral data showed that secosteroids **6-10** (eurysspongiols B) are the C-3 epimers of secosteroids **1-5** (eurysspongiols A). Their structures were determined by exhaustive spectroscopic analysis, which includes HMBC, HMQC, COSY and TOCSY experiments on **7** and **8**.

The ^1H NMR data of **7**, eurysspongiol B2, showed important A-ring differences from secosteroids **1-5** in the 3-4 ppm region. The ^{13}C NMR spectrum confirmed these differences, the hydroxylated carbons now resonating at δ 74.3, 68.4, 68.2, 65.1, 63.5 and 59.2. HMQC and HMBC experiments allowed assignment of all the carbons with their hydrogens (see Tables 1 and 2). ^1H NMR COSY and TOCSY experiments were used to analyse all the relevant spin systems and to measure the vicinal coupling constants (J) at the three hydroxylated carbons in ring A: H-4 (δ 3.80, dd, $J = 11.4$ and 2.4 Hz) was shown to have *trans*-diaxial coupling to H-5 and an ax-eq relationship with H-3 (δ 3.93 dd, $J = 2.9$ and 2.4 Hz), which in turn is coupled to H-2 at 3.82 ppm with a small constant ($J = 2.9$ Hz) indicative of an ax-eq orientation. Thus both H-2 and H-4 are axial, while H-3 is equatorial and H-5 is axial.

The *trans* ring fusion of the A/B rings was deduced as before by comparison with models and by ROESY. Figure 4 shows relevant NOEs that corroborated the above stereochemical results.

Thus compound **7**, named eurysspongiol B2 is $2\alpha,3\alpha,4\alpha,6\beta,11,19$ -hexahydroxy-9,11-secosteroid-en-9-one, and compounds **6**, **8**, **9** and **10** are the C-3 epimers of **1**, **3**, **4** and **5** respectively. Compound **6**, eurysspongiol B1 is $2\alpha,3\alpha,4\alpha,6\beta,11,19$ -hexahydroxy-9,11-secosteroid-(22E)-en-9-one. Compound **8**,

euryspongiol B3 is $2\alpha,3\alpha,4\alpha,6\beta,11,19$ -hexahydroxy-9,11-secocholest-(22E,24S)-24-methyl-en-9-one. Compound **9**, euryspongiol B4 is $2\alpha,3\alpha,4\alpha,6\beta,11,19$ -hexahydroxy-9,11-secocholest-(22E,24R)-24-methyl-en-9-one. Compound **10**, euryspongiol B5 is $2\alpha,3\alpha,4\alpha,6\beta,11,19$ -hexahydroxy-9,11-seconorcholest-(22E)-en-9-one.




Figure 4. Selected NOEs for compound **7**.

Conformation and absolute stereochemistry. In addition to coupling constant values, ROESY experiments gave information on rotation about the C-8/C-14 bond and the presence of a conformational preference. The coupling constant between H-8 and H-14 about 4 Hz, (see Table 2) suggested a gauche-like arrangement of these hydrogens, and this was confirmed by the ROESY spectrum. Strong NOEs were observed between H-8 and H-14; H-8 and H-7(β , eq); H-14 and H-7(β , eq); and between Me-18 and H-7(β , eq). Thus, the preferred conformation is as depicted in Figure 5.

Molecular Mechanics calculations (MM2 Force Field) confirmed the existence of an energy minimum when the dihedral angle between H-8 and H-14 was 50° , in which conformation their calculated coupling constant was 4.55 Hz. MM2 calculations also confirmed the dihedral angles and general geometry of the A/B decalone system.




Figure 5

Finally, the absolute stereochemistry of compounds **1-10** was investigated by circular dichroism measurements. The CD spectrum of **2** showed a negative Cotton effect in the 295 nm carbonyl absorption region. This band coincides with the one observed for similar 9,11-secoosterols with an equatorial substituent at C(8),¹⁰ which have the configuration shown in the Figure 1. As compounds **1-10** all had very similar negative optical rotation values, it can safely be concluded that they all belong to the enantiomeric series shown in the figures.

Hydride reduction of the 9 keto group of 9,11-secosteroids. The reduction of the C-9 carbonyl group of **1** with NaBH₄/MeOH yielded exclusively the heptahydroxylated 9,11-secosteroid **1a** (δ 2.98, d, J = 10.8 Hz, H-9 α), which was further transformed into hepta-acetate **1b**. The reduction reaction was shown to be stereospecific, giving exclusively the β C-9 equatorial alcohol. This result is noteworthy and rather unexpected because no intramolecular assistance by any of the hydroxyl groups seems to occur in spite of the apparently favourable spatial location of some of them (notably the secondary axial hydroxyl at C-6 and the primary OH at C-19)(Figure 6). To get further insight into this behaviour, we proceeded to transform the -OH groups into non-hydroxyl groups before reduction. Secosteroid **2** was acetylated to the keto-hexa-acetate **2a**, and the keto group was then reduced with NaBH₄/MeOH to afford **2b** (δ 3.12, d, J = 10.6 Hz, H-9 α), which was further acetylated to **2c**. Changing the hydroxyl groups to acetyl groups was not observed to influence the stereoselectivity of the process.

The formation of the 9 β -ol can be explained in terms of the mechanism suggested by Wigfield,¹⁷ in which the transition state includes one molecule of solvent alcohol, one molecule of alkoxide, one borohydride ion and the ketone, with the borohydride attacking the carbonyl carbon at an angle of 126° and not at \approx 90°. In this long acyclic transition state, β attack (leading to the 9 α -ol) would be impeded by interactions between the borohydride ion and both the axial 19-CH₂OH and, more severely, the axial 6 β -ol (Fig. 7). However, α attack, which yields the 9 β -ol, lacks similar impediments.




Figure 6




Figure 7

Antihistaminic activity. Euryssongiols **1** and **2** were tested for *in vitro* antiallergic activity by non immunological stimulus on rat mast cells. We used as chemical inducer for this assay a low molecular weight mixed polymer of N-Methyl-p-methoxyfenetilamine cross linked with formaldehyde, marketed as compound 48/80® and considered the prototype of polycations that release histamine.¹⁸ Compounds **1** and **2** were found to reduce the release of histamine induced on rat mast cells by the polymer 48/80 to about two thirds (26 %) and one half (15 %) respectively of the amount released by action of the control 48/80 (35 %). Similar values of inhibition of histamine release, were observed when GRF (growth releasing factor) was used as stimulus instead of the polymer 48/80.

Furthermore, the response of the rat mast cells to **1** and **2** is dose-dependent, principally in the case of **1** (Fig 8). We found also that **1** and **2** act as inhibitors only if there is a preincubation time before the stimulus is added to the cells. Simultaneous addition of the stimulus and the chemical showed no inhibitory effect.




Figure 8

The similarity of action of **1** and **2** with that of disodium cromoglycate (DSCG),¹⁹ a well known antihistaminic agent, suggest a common mode of action.

CONCLUSIONS

The genus *Euryspongia* is known to be an important source of furan and pyran-based terpenoids but this is the first report of secosteroids in that genus. Structurally, euryspogiols A (**1-5**) and euryspogiols B (**6-10**) constitute two series epimeric at C-3 and represent the most highly hydroxylated secosterols isolated so far and the first hydroxylated at C-4. The use of **1** as a model for the study of the stereoselective hydride reduction of the carbonyl group is noteworthy due to the high number of hydroxylic groups and their location to produce intramolecular assistance.

The antihistaminic activity of these compounds is interesting because it might be an indication of their potential use as anti-allergic and in the treatment of asthma. Certain polyhydroxylated steroids have been recently found²⁰ to be strong inhibitors of histamine release induced by anti-IgE. The structural analogy of those compounds with euryspogiols, may indicate some pharmacological similarity in the mechanism of action.

EXPERIMENTAL SECTION

General: NMR spectra were recorded on a Bruker WM-250 and a Varian XL500; chemical shifts are reported in δ units relative to tetramethylsilane ($\delta = 0$) as internal standard, with C_5D_5N and CD_3OD as solvents; J values are given in Hz. 1H NMR and ^{13}C NMR assignments were made using 2D COSY, 2D HETCOR, DEPT, TOCSY, ROESY, HMBC and HMQC sequences. EIMS were obtained at 70 eV on a Hewlett-Packard HP 59970 MS mass spectrometer. High resolution mass spectra and FAB MS (positive and negative ion modes) were recorded on a KRATOS MS 50 spectrometer using a sample dissolved in a glycerol matrix. Infrared spectra were recorded on a Perkin-Elmer 1420 spectrophotometer. Optical rotations were measured on

a JASCO DIP-370 digital polarimeter. Circular dichroism spectra were recorded on a Jobin-Ivon instrument in absolute MeOH. Column chromatography was performed on Amberlite XAD-2 from Sigma (20-60 mesh) and Sephadex LH-20 (20-100 mm). Thin layer chromatography (TLC) analyses were performed on Merck GF-254 precoated silicagel. Reversed phase HPLC was performed with a Waters Associates 590 pump, a μ -Bondapak C₁₈ column (7.8 mm x 30 cm) and a differential refractometer as detector. Compound 48/80 was purchased from Sigma Chemical Co.

Eluents: compounds **1**, **2** and **7**, 7:3 MeOH/H₂O at 3 mL min⁻¹; compound **5**, 7:3 MeOH/H₂O at 2 mL min⁻¹; compounds **3**, **4**, **6**, **8**, **9** and **10**, 6:4 MeOH/H₂O at 4 mL min⁻¹.

Collection and extraction of the sponge: *Euryspongia* sp. (reference R1225), was collected by divers using scuba from St. 184, Point Bovis, New Caledonia on 3/12/1979 at a depth of 22 m on coral abutting fine sand. Other sponges and soft coral were in association. A voucher specimen will be deposited in the Museum of Natural History, Paris and in Museum Orstom in Nouméa.

The specimen is 10 cm high, 15 cm wide, thrown in to low lobes along the crest of which large oscules lie flush with the surface. The colour is a brilliant lilac purple (Munsell RP⁶/6) mottled with whitish conules which mark the point of intersection of primary fibres with the translucent, finely reticulated pinacoderm. The skeleton is a strong reticulum in which the primary, ascending fibres are clear of debris but contain scattered bacterial-type particles. The secondary connecting fibres form an irregular, relatively dense reticulum. The choanocyte chambers are large, oval and eurypylous. The sponge, given these characteristics, falls into the genus *Euryspongia* (Family Dysideiidae, Order Dictyoceratida) and is distinguished from other members of that genus in colour, and in the peculiar nature of the coring material in the fibres. Should subsequent electron microscopy study on appropriately fixed material confirm the nature of the inclusions, it could be that a new genus within the Family Dysideiidae will have to be established for the species. A full description will be included in a forthcoming publication on the Dictyoceratida of New Caledonia.

The freshly collected sponge was cut into pieces and extracted at room temperature with methanol. The solvent was removed from the methanolic extracts by evaporation in vacuo, and the residue was partitioned successively between hexane and 10 % aqueous methanol, between Cl₂CH₂ and 20 % aqueous methanol, and between n-BuOH and 40 % aqueous methanol; evaporation of the solvents gave a hexane extract (9.3 g), a Cl₂CH₂ extract (9.9 g) and an n-BuOH extract (5.0 g). The n-BuOH extract was added to a column of Amberlite XAD-2, and the column was washed with H₂O (3L at 2 mL min⁻¹) and then eluted with MeOH (2L at 10 mL min⁻¹).

The MeOH eluates were concentrated in vacuo to give a residue (2.2 g) that was chromatographed on a Sephadex LH-20 column eluted with 2:1 MeOH/H₂O. Fractions of 10 mL each were collected and analysed by TLC on SiO₂ in 12:3:5 n-BuOH/AcOH/H₂O.

Fractions D (208 mg) and E (125 mg), which contained the secosterols, were separated by semipreparative reversed phase HPLC on a C₁₈ μ -Bondapak column, eluted with MeOH/H₂O. This afforded pure samples of **1** (30 mg, R_t 44.01 min), **2** (32 mg, R_t 70.00 min), **3** (1.5 mg, R_t 120.11 min), **4** (1 mg, R_t 120.90 min), **5** (7 mg, R_t 30.00 min), **6** (17 mg, R_t 60.37 min), **7** (5 mg, R_t 60.22 min), **8** (1.5 mg, R_t 120.00 min), **9** (1.5 mg, R_t 120.50 min) and **10** (6 mg, R_t 46.00 min).

Euryspongiol A1 (1): (30 mg); [α]_D²¹ -42° (c 0.001, MeOH); IR spectrum: ν_{\max} (KBr disc) 3450 (br), 2950, 1700 cm⁻¹; ¹H NMR (CD₃OD, C₅D₅N, 250 Mz) see Table 2; ¹³C NMR (CD₃OD, 250 Mz) see Table 1;

HR EIMS: obsd 482.3217, calcd for $C_{27}H_{46}O_7$ 482.3243; EIMS m/z (%) 482 (1), 452 (4), 434 (3), 416 (2), 408 (2), 390 (2), 299 (12), 281 (7), 218 (5), 147 (8), 125 (76), 67 (34), 54 (100 rel); FAB MS (+, glycerol matrix) m/z 483, 465, 393, 354, 316, 224; FAB MS (-, glycerol matrix) m/z 451, 387, 275, 257, 183, 127, 91.

Euryspongiol A2 (2): (32 mg); $[\alpha]_D^{25} -22^\circ$ (*c* 0.001, MeOH); CD (MeOH) $(\theta)_{295} - 6796^\circ$; UV λ_{max} (MeOH) 208 nm; HR EIMS: obsd 484.3419, calcd for $C_{27}H_{48}O_7$ 484.3399; EIMS m/z (%) 484 [M^+ , 1], 454 [$M-CH_2O^+$, 1], 436 [$M-CH_2O-H_2O^+$, 5], 418 [$M-CH_2O-2H_2O^+$, 3], 410 [$M-CH_2O-C_3H_8^+$, 1], 400 [$M-CH_2O-3H_2O^+$, 1], 382 [$M-CH_2O-4H_2O^+$, 1], 364 [$M-CH_2O-5H_2O^+$, 1], 323 [$M-CH_2O-H_2O-C_8H_{17}^+$, 2]; FAB MS (-, glycerol matrix) m/z : 483, 465, 453, 433, 386, 350, 325, 311, 273, 243, 219, 181, 165, 151, 121, 97, 93; 1H NMR (C_5D_5N , 250 Mz) δ 0.91 (3H, s, H-18), 0.98 (3H, d, *J* = 6.4 Hz, H-21), 1.74 (1H, H-7 β), 2.31 (1H, m, H-1 α), 2.51 (1H, H-7 α), 3.34 (1H, dd, *J* = 13.2, 4.7 Hz, H-1 β), 3.97 (1H, t, *J* = 8.8 Hz, H-3 α), 4.23 (1H, H-8), 4.44 (1H, d, *J* = 11.2 Hz, H-19), 4.58 (1H, m, H-2 α), 4.62 (1H, m, H-4 α), 5.13 (1H, s, H-6 β), 5.78 (1H, d, *J* = 11.2 Hz, H-11'); ^{13}C NMR (C_5D_5N , 250 Mz) δ 17.4 (C-18), 19.4 (C-21), 22.5^b (C-26), 22.7^b (C-26), 23.0 (C-15), 24.5 (C-23), 26.1 (C-16), 27.9 (C-25), 34.5 (C-20), 35.8^a (C-22), 35.6^a (C-1), 38.7 (C-8), 39.5 (C-24), 41.1 (C-12), 41.9 (C-14), 42.0 (C-7), 45.9 (C-13), 49.7 (C-17), 55.9 (C-10), 56.3 (C-5), 58.1 (C-11), 63.7 (C-19), 64.0 (C-6), 70.1 (C-4), 70.7 (C-2), 83.2 (C-3), 215.4 (C-9).

Euryspongiol A3 (3): (1.5 mg); HR EIMS obsd 496.3401, calcd for $C_{28}H_{48}O_7$ 496.3400; EIMS m/z (%) 496 [M^+ , 1], 466 [$M-CH_2O^+$, 7]; FAB MS (-, glycerol matrix) m/z 495 [$M-H^-$]; 1H NMR (CD_3OD , 250 Mz) δ 0.79 (3H, s, H-18), 0.87 (6H, d, *J* = 6.9 Hz, H-26, 27), 0.94 (3H, d, *J* = 6.9 Hz, H-28), 1.06 (3H, d, *J* = 6.9 Hz, H-21), 1.34 (2H, m, H-16), 1.36 (2H, m, H-15), 1.46 (1H, H-1 α), 1.49 (1H, H-5), 1.57 (1H, H-7 β), 2.20 (1H, H-7 α), 2.34 (1H, dd, *J* = 13.6, 4.9 Hz, H-1 β), 2.49 (1H, H-14), 3.07 (1H, t, *J* = 9.0 Hz, H-3), 3.47 (1H, H-8), 3.62 (1H, H-4), 3.68 (2H, H-11), 3.68 (1H, d, *J* = 11.5 Hz, H-19), 3.70 (1H, H-2), 4.30 (1H, H-6), 4.85 (1H, d, *J* = 11.5 Hz, H-19), 5.25 (2H, m, H-22, 23); ^{13}C NMR (CD_3OD , 250 Mz) δ 18.1 (C-18), 19.6 (C-28), 23.5 (C-21), 23.9 (C-15), 24.2 (C-26, 27), 26.7 (C-16), 35.5 (C-1), 35.7 (C-25), 39.7 (C-8), 41.3 (C-7), 41.3 (C-12), 41.7 (C-20), 43.1 (C-14), 45.8 (C-24), 46.6 (C-13), 51.0 (C-17), 56.2 (C-10), 56.3 (C-5), 59.2 (C-11), 63.9 (C-19), 64.7 (C-6), 70.5 (C-4), 71.1 (C-2), 82.8 (C-3), 135.2 (C-22), 137.7 (C-23), 215.4 (C-9).

Euryspongiol A4 (4): (1 mg); HR EIMS obsd 496.3401, calcd for $C_{28}H_{48}O_7$ 496.3400; EIMS m/z (%) 496 [M^+ , 1], 466 [$M-CH_2O^+$, 7]; FAB MS (-, glycerol matrix) m/z 495 [$M-H^-$]; 1H NMR (CD_3OD , 250 Mz) δ 0.81 (3H, s, H-18), 0.89 (6H, d, *J* = 6.6 Hz, H-26, 27), 0.96 (3H, d, *J* = 6.9 Hz, H-28), 1.07 (3H, d, *J* = 6.8 Hz, H-21), 1.36 (2H, m, H-16), 1.37 (2H, m, H-15), 1.46 (1H, H-1 α), 1.49 (1H, H-5), 1.58 (1H, H-7 β), 1.67 (2H, H-12), 2.24 (1H, H-7 α), 2.32 (1H, dd, *J* = 13.6, 4.8 Hz, H-1 β), 2.47 (1H, H-14), 3.10 (1H, t, *J* = 8.9 Hz, H-3), 3.46 (1H, H-8), 3.64 (1H, H-4), 3.64 (2H, m, H-11), 3.66 (1H, H-2), 3.75 (1H, d, *J* = 11.5 Hz, H-19), 4.33 (1H, H-6), 4.85 (1H, d, *J* = 11.5 Hz, H-19), 5.20 (2H, m, H-22, 23); ^{13}C NMR (CD_3OD , 250 Mz) δ 18.1 (C-18), 19.6 (C-28), 23.5 (C-21), 23.9 (C-15), 24.2 (C-26, 27), 26.7 (C-16), 35.5 (C-1), 35.7 (C-25), 39.7 (C-8, C-20), 41.3 (C-7), 41.3 (C-12), 43.1 (C-14), 45.8 (C-24), 46.6 (C-13), 51.0 (C-17), 56.2 (C-10), 56.3 (C-5), 59.2 (C-11), 63.9 (C-19), 64.7 (C-6), 70.5 (C-4), 71.1 (C-2), 82.8 (C-3), 135.2 (C-22), 137.7 (C-23), 215.4 (C-9).

Eurysspongiol A5 (5): (7 mg); $[\alpha]^{21}_{\text{D}} -48^{\circ}$ (*c* 0.0001, MeOH); HR EIMS: obsd 468.3090, calcd for $\text{C}_{26}\text{H}_{44}\text{O}_7$ 468.3087; FAB MS (+, NaCl+glycerol matrix) *m/z* 491 $[\text{M}+\text{Na}]^+$, 394 $[\text{M}+\text{Na}-\text{C}_7\text{H}_{13}]^+$; ^1H NMR (CD_3OD , 250 Mz) δ 0.78 (3H, s, H-18), 0.96 (6H, d, *J* = 6.7 Hz, H-25, 26), 1.03 (3H, d, *J* = 6.8 Hz, H-21), 1.34 (2H, m, H-16), 1.38 (2H, m, H-15), 1.45 (1H, H-1 α), 1.45 (1H, dd, *J* = 11.7, 2.5 Hz, H-5), 1.53 (1H, H-7 β), 1.64 (2H, m, H-12), 2.21 (1H, H-24), 2.22 (1H, H-7 α), 2.30 (1H, dd, *J* = 13.8, 5.1 Hz, H-1 β), 2.44 (1H, H-14), 3.07 (1H, t, *J* = 9.0 Hz, H-3), 3.46 (1H, H-8), 3.61 (2H, m, H-11), 3.65 (1H, H-4), 3.69 (1H, H-2), 3.72 (1H, d, *J* = 11.6 Hz, H-19), 4.31 (1H, d, *J* = 2.5 Hz, H-6), 4.87 (1H, d, *J* = 11.8 Hz, H-19), 5.30 (2H, m, H-22, 23); ^{13}C NMR (CD_3OD , 250 Mz) δ 18.1 (C-18), 22.2 (C-21), 23.9 (C-15), 26.3 (C-16), 32.3 (C-24), 35.5 (C-1), 39.4 (C-20), 39.8 (C-8), 41.3 (C-7,12), 43.1 (C-14), 46.7 (C-13), 51.0 (C-17), 56.2 (C-10), 56.3 (C-5), 59.2 (C-11), 64.0 (C-19), 64.7 (C-6), 70.5 (C-4), 71.1 (C-2), 82.8 (C-3), 134.0 (C-23), 136.0 (C-22), 214.8 (C-9).

Eurysspongiol B1 (6): (17 mg); $[\alpha]^{21}_{\text{D}} -39^{\circ}$ (*c* 0.003, MeOH); HR EIMS: obsd 482.3239, calcd for $\text{C}_{27}\text{H}_{46}\text{O}_7$ 482.3243; ^1H NMR (CD_3OD , 500 Mz) δ 0.84 (3H, s, H-18), 0.91 (6H, d, *J* = 6.6 Hz, H-26, 27), 1.10 (3H, d, *J* = 6.8 Hz, H-21), 1.32 (2H, m, H-16), 1.34 (2H, m, H-15), 1.55 (1H, H-25), 1.55 (1H, H-7 β), 1.60 (1H, H-17), 1.63 (2H, H-12), 1.66 (1H, H-1 α), 1.75 (1H, H-5), 1.88 (1H, dd, *J* = 8.9, 3.0 Hz, H-24), 2.01 (1H, dd, *J* = 13.6, 4.5 Hz, H-1 β), 2.26 (1H, H-20), 2.26 (1H, H-7 α), 2.55 (1H, H-14), 3.47 (1H, H-8), 3.65 (2H, H-11), 3.78 (1H, d, *J* = 12.8 Hz, H-19), 3.84 (1H, dd, *J* = 11.3, 2.6 Hz, H-4), 3.89 (1H, dd, *J* = 11.6, 4.5 Hz, H-2), 3.99 (1H, dd, *J* = 4.5, 2.7 Hz, H-3), 4.33 (1H, d, *J* = 2.8 Hz, H-6), 4.86 (1H, d, *J* = 11.3 Hz, H-19), 5.36 (2H, m, H-22, 23); ^{13}C NMR (CD_3OD , 500 Mz) δ 18.1 (C-18), 22.3 (C-21), 22.7 (C-26, 27), 24.0 (C-15), 26.7 (C-16), 29.7 (C-25), 30.6 (C-1), 39.7 (C-8), 40.0 (C-20), 41.3 (C-12), 41.5 (C-7), 43.1 (C-14), 43.2 (C-24), 46.6 (C-13), 51.0 (C-17), 56.2 (C-10), 59.2 (C-11), 63.5 (C-19), 65.1 (C-6), 68.2 (C-4), 68.4 (C-2), 74.3 (C-3), 128.4 (C-23), 138.4 (C-22), 216.8 (C-9).

Eurysspongiol B2 (7): (5 mg); $[\alpha]^{21}_{\text{D}} -50^{\circ}$ (*c* 0.0006, MeOH); HR EIMS obsd 484.3403, calcd for $\text{C}_{27}\text{H}_{48}\text{O}_7$ 484.3400; EIMS (%) *m/z* 484 $[\text{M}^+$, 1], 454 $[\text{M}-\text{CH}_2\text{O}^+$, 5], 436 $[\text{M}-\text{CH}_2\text{O}-\text{H}_2\text{O}^+$, 23], 418 $[\text{M}-\text{CH}_2\text{O}-2\text{H}_2\text{O}^+$, 5], 400 $[\text{M}-\text{CH}_2\text{O}-3\text{H}_2\text{O}^+$, 4], 305 $[\text{M}-\text{CH}_2\text{O}-\text{H}_2\text{O}-\text{C}_9\text{H}_{17}^+$, 8]; FAB MS (+, NaCl+glycerol matrix) *m/z* 507 $[\text{M}+\text{Na}]^+$, 394 $[\text{M}+\text{Na}-\text{C}_8\text{H}_{17}]^+$; ^1H NMR (CD_3OD , 500 Mz) δ see table 2; ^{13}C NMR (CD_3OD , 500 Mz) δ see table 1.

Eurysspongiol B3 (8): (1.5 mg); HR EIMS: obsd 496.3405, calcd for $\text{C}_{28}\text{H}_{48}\text{O}_7$ 496.3400; EIMS *m/z* (%) 496 $[\text{M}^+$, 1] 466 $[\text{M}-\text{CH}_2\text{O}^+$, 6], 305 $[\text{M}-\text{CH}_2\text{O}-2\text{H}_2\text{O}-\text{C}_9\text{H}_{17}^+$, 8]; FAB MS (+, NaCl+KCl+glycerol matrix) *m/z* 535 $[\text{M}+\text{K}]^+$, 519 $[\text{M}+\text{Na}]^+$, 371 $[\text{M}-\text{C}_9\text{H}_{17}]^+$; ^1H NMR (CD_3OD , 500 Mz) δ 0.79 (3H, s, H-18), 0.89 (6H, d, *J* = 7.0 Hz, H-26, 27), 0.93 (3H, d, *J* = 7 Hz, H-28), 1.05 (3H, d, *J* = 7.0 Hz, H-21), 1.38 (2H, m, H-16), 1.51 (1H, H-7 β), 1.52 (2H, m, H-15), 1.58 (2H, H-12), 1.64 (1H, H-25), 1.65 (1H, H-1 α), 1.85 (1H, dd, *J* = 11.9, 3.0 Hz, H-5), 2.00 (1H, dd, *J* = 13.5, 3.9 Hz, H-1 β), 2.18 (1H, H-20), 2.23 (1H, H-7 α), 2.48 (1H, H-14), 3.48 (1H, H-8), 3.59 (2H, H-11), 3.72 (1H, d, *J* = 11.5 Hz, H-19), 3.80 (1H, dd, *J* = 11.9, 3.0 Hz, H-4), 3.82 (1H, H-2), 3.93 (1H, t, *J* = 3.0 Hz, H-3), 4.28 (1H, H-6), 4.85 (1H, d, *J* = 11.5 Hz, H-19), 5.24 (2H, dd, *J* = 15.0, 7.5 Hz, H-22), 5.28 (1H, d, *J* = 15.0, 8.0 Hz, H-23); ^{13}C NMR (CD_3OD , 500 Mz) δ 19.0 (C-18), 19.6 (C-28), 23.5 (C-21), 24.2 (C-26, 27), 25.1 (C-15), 28.1 (C-16), 31.7 (C-1), 35.7 (C-25), 41.0 (C-8), 41.1 (C-20), 42.2 (C-12), 42.8 (C-7), 44.1 (C-14), 45.8 (C-24), 47.9 (C-13), 51.3 (C-5), 57.3 (C-10), 60.5 (C-11), 64.5 (C-19), 66.2 (C-6), 69.6 (C-4), 69.6 (C-2), 75.4 (C-3),

135.2 (C-22), 137.7 (C-23), 215.4 (C-9).

Euryspongiol B4 (9): (1.5 mg); HR EIMS: obsd 496.3405, calcd for $C_{28}H_{48}O_7$ 496.3400; EIMS m/z (%) 496 $[M^+$, 1] 466 $[M-CH_2O^+$, 6], 305 $[M-CH_2O-2H_2O-C_9H_{17}^+$, 8]; FAB MS (+, NaCl+KCl+glycerol matrix) m/z 535 $[M+K]^+$, 519 $[M+Na]^+$, 371 $[M-C_9H_{17}]^+$; 1H NMR (CD_3OD , 500 Mz) δ 0.79 (3H, s, H-18), 0.87 (6H, d, $J = 6.5$ Hz, H-26, 27), 0.93 (3H, d, $J = 6.5$ Hz, H-28), 1.06 (3H, d, $J = 7.0$ Hz, H-21), 1.38 (2H, m, H-16), 1.51 (1H, H-7 β), 1.52 (2H, m, H-15), 1.58 (2H, H-12), 1.64 (1H, H-25), 1.65 (1H, H-1 α), 1.85 (1H, dd, $J = 11.9, 3.0$ Hz, H-5), 2.00 (1H, dd, $J = 13.5, 3.9$ Hz, H-1 β), 2.18 (1H, H-20), 2.23 (1H, H-7 α), 2.48 (1H, H-14), 3.48 (1H, H-8), 3.59 (2H, H-11), 3.72 (1H, d, $J = 11.5$ Hz, H-19), 3.80 (1H, dd, $J = 11.9, 3.0$ Hz, H-4), 3.82 (1H, H-2), 3.93 (1H, t, $J = 3.0$ Hz, H-3), 4.28 (1H, H-6), 4.85 (1H, d, $J = 11.5$ Hz, H-19), 5.24 (2H, dd, $J = 15.0, 7.5$ Hz, H-22), 5.28 (1H, dd, $J = 15.0, 8.0$ Hz, H-23); ^{13}C NMR (CD_3OD , 500 Mz) δ 19.0 (C-18), 19.6 (C-28), 23.5 (C-21), 24.2 (C-26, 27), 25.1 (C-15), 28.1 (C-16), 31.7 (C-1), 35.7 (C-25), 41.0 (C-8), 41.1 (C-20), 42.2 (C-12), 42.8 (C-7), 44.1 (C-14), 45.8 (C-24), 47.9 (C-13), 51.3 (C-5), 57.3 (C-10), 60.5 (C-11), 64.5 (C-19), 66.2 (C-6), 69.6 (C-4), 69.6 (C-2), 75.4 (C-3), 135.2 (C-22), 137.7 (C-23), 215.0 (C-9).

Euryspongiol B5 (10): (6 mg); $[\alpha]_D^{21}$ -45° (c 0.002, MeOH); HR EIMS: obsd 468.3092, calcd for $C_{26}H_{44}O_7$ 468.3087; FAB MS (+, NaCl+glycerol matrix) m/z 491 $[M+Na]^+$; 1H NMR (CD_3OD , 250 Mz) δ 0.78 (3H, s, H-18), 0.97 (6H, d, $J = 6.7$ Hz, H-25, 26), 1.03 (3H, d, $J = 6.8$ Hz, H-21), 1.54 (1H, H-7 β), 1.61 (2H, H-12), 1.65 (1H, H-1 α), 1.83 (1H, dd, $J = 11.5, 2.8$ Hz, H-5), 2.02 (1H, dd, $J = 13.3, 4.6$ Hz, H-1 β), 2.22 (1H, H-7 α), 2.47 (1H, H-14), 3.43 (1H, H-8), 3.63 (2H, H-11), 3.69 (1H, d, $J = 11.8$ Hz, H-19), 3.83 (1H, dd, $J = 11.5, 2.6$ Hz, H-4), 3.85 (1H, H-2), 3.86 (1H, dd, $J = 4.2, 2.6$ Hz, H-3), 4.28 (1H, H-6), 4.83 (1H, d, $J = 11.8$ Hz, H-19), 5.29 (2H, m, H-22, 23); ^{13}C NMR (CD_3OD , 250 Mz) δ 18.1 (C-18), 22.2 (C-21), 22.9 (C-26, 27), 30.7 (C-1), 32.3 (C-24), 39.3 (C-20), 40.1 (C-8), 41.5 (C-7), 41.5 (C-12), 43.2 (C-14), 46.7 (C-13), 51.2 (C-5), 56.1 (C-10), 59.2 (C-11), 63.5 (C-19), 65.1 (C-6), 68.2 (C-4), 68.5 (C-2), 74.3 (C-3), 134.0 (C-23), 137.0 (C-22), 215.0 (C-9).

NaBH₄ Reduction of euryspongiol A1 (1). NaBH₄ (2 mg) was added to **1** (12 mg) in MeOH (2 mL). The mixture was stirred 7 h, dilute HCl (10 %, 1 mL) was added, and the solvent was evaporated. Purification of the product by preparative HPLC on μ -Bondapak C₁₈ (6:4 MeOH/H₂O) yielded **1a** (7.2 mg, R_t 80.09 min): $[\alpha]_D^{21}$ -35° (c 0.001, MeOH); UV λ_{max} (MeOH) 206 nm; HR EIMS: obsd 484.3410, calcd for $C_{27}H_{48}O_7$ 484.3400; EIMS m/z (%) 484 (1), 454 (3), 422 (4), 355 (5), 338 (9), 320 (10), 306 (19), 289 (11), 235 (36), 191 (16), 147 (27), 95 (61), 69 (100); 1H NMR (CD_3OD) δ see Table 2; ^{13}C NMR (CD_3OD) δ see Table 1.

Acetylation of 1a to 1b. A mixture of **1a** (2 mg) and excess Ac₂O in 0.5 ml of dry pyridine was kept at room temperature for 12 h, after which removal of the excess reagents in vacuo yielded **1b** (4 mg): $[\alpha]_D^{21}$ -50° (c 0.0001, Cl₂CH₂); HR EIMS: obsd 778.4150, calcd for $C_{41}H_{62}O_{14}$ 778.4140; 1H NMR (CD_3OD , 250 Mz) δ 0.77 (3H, s, H-18), 0.90 (6H, d, $J = 6.6$ Hz, H-26, 27), 1.08 (3H, d, $J = 6.8$ Hz, H-21), 2.01, 2.01, 2.03, 2.07, 2.10, 2.18 and 2.18 (each 3H, s, OAc).

Acetylation of 2 to 2a. A crystal of 4-dimethylaminopyridine was left overnight at room temperature

in a mixture of **2** (13 mg) and excess of Ac₂O in 1 mL of dry pyridine. Removal of excess reagents in vacuo afforded the acetylated compound **2a** (23 mg): [α]²¹_D -29° (*c* 0.002, Cl₂CH₂); UV λ_{\max} (Cl₃CH) 244 nm; HR EIMS: obsd 736.4029, calcd for C₃₉H₆₀O₁₃ 736.4034; EIMS *m/z* (%) 736 (1), 368 (100), 353 (24), 326 (4), 260 (21), 247 (20), 213 (13), 147 (41), 105 (24), 81 (21); ¹H NMR (Cl₃CD, 250 Mz) δ 0.71 (3H, s, H-18), 0.87 (6H, d, *J* = 6.5 Hz, H-26, 27), 0.98 (3H, d, *J* = 6.7 Hz, H-21), 2.00, 2.01, 2.02, 2.03, 2.09, and 2.10 (each 3H, s, OAc), 2.33 (1H, H-1 β), 2.48 (1H, m, H-14), 3.37 (1H, m, H-8), 4.11 (2H, m, H-11), 4.57 (1H, d, *J* = 12.6 Hz, H-19), 5.13 (1H, m, H-2), 5.38 (1H, d, *J* = 12.6 Hz, H-19).

Reduction of 2a to 2b. NaBH₄ (3 mg) was added to **2a** in MeOH (2 mL). The mixture was stirred for two hours at 0 °C, 0.5 % HCl was added, and the solvent was evaporated. Purification by partition between Cl₂CH₂ and water and concentration and drying of the Cl₂CH₂ eluates gave 23 mg of a product that was subjected to preparative HPLC on a μ -Bondapak C₁₈ column (95:5 MeOH/H₂O at 3 mL.min⁻¹) to yield **2b** (11 mg, R_t 5.10 min): [α]²¹_D -87° (*c* 0.001, Cl₂CH₂); HR EIMS: obsd 738.4198, calcd for C₃₉H₆₂O₁₃ 738.4190; ¹H NMR (CD₃OD, 250 Mz) δ 0.71 (3H, s, H-18), 0.90 (6H, d, *J* = 6.5 Hz, H-26, 27), 1.01 (3H, d, *J* = 6.4 Hz, H-21), 1.45 (1H, H-1 α), 1.51 (1H, H-7 β), 1.80 (2H, H-12), 1.91 (1H, dd, *J* = 10.7, 2.4 Hz, H-5), 1.95 (1H, H-7 α), 2.02 (1H, H-8), 2.65 (1H, dd, *J* = 12.9, 5.2 Hz, H-1 β), 2.33 (1H, H-14), 3.10 (1H, d, *J* = 10.6 Hz, H-9), 4.18 (2H, H-11), 4.43 (1H, d, *J* = 12.3 Hz, H-19), 4.69 (1H, d, *J* = 12.3 Hz, H-19), 5.02 (1H, H-6), 5.09 (1H, t, *J* = 9.4 Hz, H-3), 5.34 (1H, H-2), 5.49 (1H, H-4); ¹³C NMR (CD₃OD, 250 Mz) δ 17.6 (C-18), 18.5 (C-21), 22.4^a (C-26), 22.6^a (C-27), 45.7 (C-13), 48.6 (C-17), 61.5 (C-11), 63.2 (C-19), 65.6 (C-6), 68.6 (C-4), 69.8 (C-2), 75.9 (C-9), 80.6 (C-3), 169.9, 169.9, 170.0, 170.3, 171.2, 171.9 (OAc).

Acetylation of 2b to 2c: A mixture of **2b** (1 mg) and excess Ac₂O in 0.3 mL of pyridine was kept at room temperature for twelve hours. Removal of excess reagents in vacuo yielded **2c** (2 mg): [α]²¹_D -50° (*c* 0.0002, Cl₂CH₂); HR EIMS: obsd 780.4291, calcd for C₄₁H₆₄O₁₄ 780.4296; ¹H NMR (CD₃OD, 250 Mz) δ 0.77 (3H, s, H-18), 0.91 (6H, d, *J* = 6.6 Hz, H-26, 27), 1.03 (3H, d, *J* = 6.2 Hz, H-21), 2.01, 2.01, 2.04, 2.07, 2.09, 2.18 and 2.18 (each 3H, s, OAc).

ACKNOWLEDGEMENTS

This work was supported by CICYT (SAF 92-1023) and Xunta de Galicia (XUGA 20906B91 and XUGA 20906B92). J. D. acknowledges a collaboration from the Instituto Farmacológico Español. We are grateful to prof. L. Botana and M. Estévez of the Department of Pharmacology for the antihistaminic tests.

REFERENCES

- 1 See for instance Milkova, Ts. S.; Mikhova, B. P.; Nikolov, N. M.; Popov, S. S. *J. Nat. Prod.* **1992**, *55*, 974.
- 2 Minale, L.; Pizza, C.; Riccio, R.; Greco, O. S.; Zollo, F. *J. Nat. Prod.* **1984**, *47*, 784.
- 3 Enwall, E. L.; van der Helm, D.; Hsu, I. N.; Pattabhiraman, T.; Schmitz, F. J.; Spraggins, R. L.; Weinheimer, A. *J. C. S. Chem. Comm.* **1972**, 215.

- 4 West, R. R.; Cardellina II, J. H. *J. Org. Chem.* **1988**, *53*, 2782.
- 5 Pika, J.; Tischler, M.; Andersen, R. J. *Can. J. Chem.* **1992**, *70*.
- 6 Migliuolo, A.; Piccialli, V.; Sica, D. *Tetrahedron* **1991**, *47*, 7937 and Migliuolo, A.; Piccialli, V.; Sica, D. *Steroids* **1992**, *57*, 344.
- 7 Kobayashi, M.; Kobayashi, K.; Ramana, K. V.; Lakshmana Rao, Ch. V.; Vencata Rao, D.; Bheemasankara Rao, Ch. *J. Chem. Soc. Perkin Trans. I* **1991**, 493.
- 8 Kazlauskas, R.; Murphy, P. T.; Ravi, B. N.; Sanders, R. L.; Wells, R. J. *Aust. J. Chem.* **1982**, *35*, 69.
- 9 Bonini, C.; Cooper, C. B.; Kazlauskas, R.; Wells, R. J.; Djerassi, C. *J. Org. Chem.* **1983**, *48*, 2108.
- 10 Capon, R. J.; Faulkner, D. J. *J. Org. Chem.* **1985**, *50*, 4771.
- 11 Demarco, P. V.; Farkas, E.; Doddrell, D.; Mylari, B. L.; Wenkert, E. *J. Am. Chem. Soc.* **1968**, *90*, 5480.
- 12 Breitmaier, E.; Voelter, W. *Carbon-13 NMR Spectroscopy*, Weinheim Ed; New York, **1987**, pp. 190.
- 13 Itoh, T.; Sica, D.; Djerassi, C. *J. Chem. Soc. Perkin Trans. I* **1983**, 147.
- 14 Rubistein I.; Goad, L. J.; Clague, A. D. H.; Mulheirn, L. J. *Phytochemistry* **1976**, *15*, 195.
- 15 Nes, W. R.; Kreritz, K.; Behzadam, S. *Lipids* **1976**, *11*, 118.
- 16 Madaio, A.; Piccialli, V.; Sica, D. *J. Nat. Prod.* **1989**, *52*, 952.
- 17 Wigfield, D. C.; Gowland, F. W. *Tetrahedron Letters* **1976**, *38*, 3373.
- 18 Lagunoff, D.; Martin, T. W.; Read, G. *Ann. Rev. Pharmacol.* **1983**, *23*, 331.
- 19 Khandwala, A.; van Inwegen, R.; Coutts, S.; Dally-Meade, V.; Youssefyeh, R. D. *Int. J. Immunopharmac.* **1983**, *5*, 491.
- 20 Noboru, S.; Akemi, U.; Kogyoku, S.; Katsuko, T.; Shigenobu, A.; Kobayashi, J.; Takei, M. *J. Org. Chem.* **1992**, *57*, 2996.

(Received in UK 25 October 1993; revised 17 January 1994; accepted 21 January 1994)