

Bionematicides as an alternative to methyl bromide fumigation

B. B. Westerdahl^{1,*}, J. Hasey²,
J. Grant³ and L. W. Beem⁴

¹Department of Entomology
and Nematology, University of
California, Davis, CA 95616.

²University of California Cooperative
Extension, Yuba City, CA 95991.

³University of California Cooperative
Extension, Stockton, CA 95206.

⁴Beem Consulting, Sacramento,
CA 95827.

*E-mail: bbwesterdahl@ucdavis.edu

This paper was edited by
Shaun D. Berry.

Received for publication
October 2, 2020.

Abstract

Fumigant use in perennial crops can be reduced through prolonging the life of existing orchards. The longer an orchard remains healthy and productive, the less often it will be terminated, fumigated, and replanted. Two trials were conducted to determine the effectiveness of DiTera, a toxin produced by the fungus (*Myrothecium verrucaria*) and Nema-Q, an extract of the soap bark tree (*Quillaja saponaria*) for management of root-lesion (*Pratylenchus vulnus*) and ring (*Mesocriconema xenoplax*) nematodes on walnuts. In the first trial, spring and fall treatments of DiTera were applied each year for four years to variety 'Chandler' scion on seedling 'Paradox' rootstock, and to own-rooted 'Chandler' trees. On 'Paradox' rootstock, both DiTera and Nema-Q increased walnut yields ($P \leq 0.05$) and produced more vigorous trees ($P \leq 0.05$) without significant reductions in nematode populations ($P \leq 0.05$). A second trial was conducted with three rates of DiTera and four rates of Nema-Q, combinations of the two products, and an untreated control on 'Chandler' scion on 'Paradox' rootstock. The highest rate of Nema-Q ($P \leq 0.05$), and a combination treatment of DiTera plus Nema-Q ($P \leq 0.05$) increased trunk circumference. Several treatments showed reductions in root-lesion and ring nematodes ($P \leq 0.05$). Bionematicides can improve yield, growth, and vigor in walnut orchards infested with plant-parasitic nematodes.

Keywords

Biological Control, DiTera, Management, *Mesocriconema Xenoplax*, *Myrothecium Verrucaria*, Nema-Q, Nematode, *Pratylenchus Vulnus*, *Quillaja Saponaria*, Walnut.

Root-lesion, *Pratylenchus vulnus* Allen and Jensen; and ring, *Mesocriconema xenoplax* (Raski, 1952) Loof & De Grisse, 1989 nematodes reduce walnut (*Juglans* sp.) yields through root damage from direct feeding and by placing trees under stress (Lownsbery, 1956, 1959; Lownsbery et al., 1978). Root-lesion nematodes are likely to be found within roots as well as in soil, while ring nematodes are external parasites of roots. Recently, two biological nematicides achieved registration in California for use on walnuts, DiTera, a toxin produced by the fungus *Myrothecium verrucaria* (Alb. & Schwein.) Ditmar

(1813) and Nema-Q, an extract of the soap bark tree *Quillaja saponaria* Molina.

In recent years, extensive research has been conducted to find replacements for methyl bromide, widely used as a pre-plant soil fumigant before being implicated in the depletion of ozone in the stratosphere (United States Environmental Protection Agency, 1993). A way to reduce the frequency of fumigant use in perennial crops is through prolonging the life of existing orchards. The longer an orchard remains healthy and productive, the less often it will be terminated, fumigated, and replanted.

Over time, this reduces the amount of fumigant used. In addition to fumigation, a variety of approaches have been researched for management of plant-parasitic nematodes including development of pre-plant hot water treatments of rootstocks, evaluation of rootstock susceptibility, and evaluation of biological products (Buzo et al., 2009; Giraud et al., 2011; Hasey et al., 2004; Westerdahl and Radewald, 2011).

Only about a dozen nematicidal active ingredients have ever achieved registration in California, and several of the most effective of these lost their registrations owing to groundwater contamination, air pollution, or carcinogenicity (Ferris, 2021). The loss of use of the nematicide dibromochloropropane (DBCP) in 1977, that had been widely used post-plant on bearing fruit and nut tree crops in California, created a tremendous need for replacement products (United States Environmental Protection Agency, 2014).

The goal of this study was to evaluate the potential of DiTera and Nema-Q for post-plant management of root-lesion and ring nematodes in commercial walnut orchards.

Materials and methods

Two field trials were conducted to evaluate the potential of two bionematicides for nematode management on walnuts. Because the actual active ingredients in both products tested are uncertain, all rates are expressed in terms of amount of product per ha.

Sutter County Trial: The first trial was conducted in an orchard in Sutter County, CA on Hollilipah loamy sand. This orchard was previously used for an own-rooted 'Chandler' compared to 'Chandler' on 'Paradox' rootstock trial (Hasey et al., 2004). In that trial that was planted in 1991, two rootstocks, micropropagated 'Chandler' (*Juglans regia* L.) on its own-roots and nursery grafted 'Chandler' on seedling 'Paradox,' *J. hindsii* (Jeps.) Jeps. Ex R.E. Sm., \times *J. regia*, rootstock were spaced at 7.62 m \times 7.62 m in a randomized complete block design with 20 individual tree replicates. In 1998, the orchard was found to be infested with plant-parasitic nematodes. From this previous trial, 15 trees of each rootstock were selected for a trial in a randomized complete block design with five replicates of three treatments: Untreated Control (UC), DiTera (DT), and Nema-Q (NQ). DiTera[®] DF (Valent BioSciences Corporation, Libertyville, IL) at 56 kg/ha, and Nema-Q[®] (Monterey AgResources, Fresno, CA) at 23.4 L/ha, were applied in a 50% banded spray, so that actual amounts applied were 28 kg DT and 11.7 L NQ/treated ha. One day prior to treatment, 1.25 cm of irrigation was applied to the

orchard from 39.8 Lph microsprinklers (Antelco Rotor Spray Mini Sprinkler, Antelco, Longwood, FL). Each tree was treated individually with 5.7 L of solution from a 7.6-L watering can (Bloem Classic 2 Gal. Blue Plastic Watering Can, Bloem, Hudsonville, MI) twice yearly for four years. Treatment was immediately followed by an additional 1.25 cm of irrigation. Treatments were applied in 2003 (April 11 and October 24), 2004 (April 6 and September 29), 2005 (April 11 and October 13), and 2006 (May 10 and October 30). The orchard was irrigated by the grower as needed following California Irrigation Management Information System (CIMIS) guidelines (CIMIS station #84) (CIMIS, 2021). Weather data for the trial area is available from CIMIS station #84 (UCIPM, 2021). The orchard was managed by the grower and treated with standard practices with respect to fertilization, insecticides, and fungicides.

Soil samples were taken prior to each treatment date using a 5-cm diameter bucket auger to a depth of 60-cm midway between the dripline of the tree canopy and the tree trunk, in the fall and spring of each year. Nematodes were extracted from a 400-cm³ soil sub-sample with a modified semiautomatic elutriator and sucrose centrifugation technique (Byrd et al., 1976). Extracted nematodes were identified and counted at \times 45 magnification under a stereoscopic dissecting microscope (Bausch & Lomb, Bridgewater, NJ). The sampling method used did not yield sufficient roots to permit extraction of nematodes. Yields and trunk circumference were evaluated each year from 2002 (pretreatment baseline) to 2006. Circumference of each tree trunk was measured at 60 cm above the ground and trunk cross-sectional area was calculated from the circumference measurements (Retzlaff et al., 1992). In addition, tree vigor was visually evaluated in 2005 and 2006 using a rating scale developed by the authors: 0=Dead, 1=Very low vigor, dieback, 2=Early yellowing, 3=No new shoot growth, 4=Some new shoot growth, and 5=Most vigorous.

San Joaquin County Trial: The second trial was conducted in San Joaquin County, CA with 'Chandler' scion on 'Paradox' rootstock. The randomized complete block trial with 6 replicates of 14 treatments evaluated three rates of DT, four rates of NQ and six combinations of DT and NQ for their effectiveness in controlling root-lesion and ring nematodes. Treatments included an UC. Treatments were applied to the soil surface in a 50% banded spray. One day prior to treatment, 1.25 cm of irrigation was applied to the orchard with 30.2 Lph microsprinklers (Rain Bird Micro-Quick Spray Assembly, Rain Bird, Azusa, CA). Each tree was treated individually with 5.7 L of solution from a 7.6-liter watering can (Bloem Classic 2 Gal. Blue Plastic Watering Can, Bloem,

Hudsonville, MI). Treatment was immediately followed by an additional 1.25 cm of irrigation. Orchards were irrigated by the grower as needed following CIMIS guidelines (CIMIS station #70) (CIMIS, 2021). Weather data for the trial area is available from CIMIS station #70 (UCIPM, 2021). The orchard was managed by the grower and treated with standard practices with respect to fertilization, insecticides, and fungicides.

Treatment effectiveness was evaluated via soil and root sampling, and trunk circumference measurements. Nematodes were extracted from soil as described for the Sutter County Trial. Nematodes were also extracted from roots that were weighed

and placed in an intermittent mist chamber for 72h (Ayoub, 1977). Treatments were applied in the spring of 2005 following measurement of trunk circumference. Additional nematode sampling followed by repeated treatments were conducted in October 2005, April 2006, and October 2006. Post-treatment trunk circumference measurements were done in October 2005, April 2006, and October 2006.

Data analysis: Both trials were conducted in randomized complete block design. There were five replicates per treatment in the Sutter County Trial and data were analyzed using Repeated Measures Analysis of Variance (ANOVA) ($P \leq 0.05$), followed by

Table 1. Yield data for Sutter County trial.

Year	Treatment	Yield (kg/tree)					
		Combined		Paradox		Own-rooted	
2002	UC	23.4		26.3		20.5	
	DT	24.0		25.2		22.8	
	NQ	28.8		32.4		25.0	
2003	UC	32.0		37.4		26.5	
	DT	36.9		40.8		32.9	
	NQ	42.1	0.0027 ^a	47.0	0.0338	37.1	0.0383
2004	UC	22.8		25.45		20.2	
	DT	27.1		31.1		23.0	
	NQ	26.5		33.1		20.0	
2005	UC	33.0		33.8		32.2	
	DT	39.3		43.4	0.0325	35.3	
	NQ	38.3		49.2	0.0008	27.4	
2006	UC	33.0		37.7		28.3	
	DT	42.3	0.0068	49.1	0.0114	35.5	
	NQ	42.6	0.0041	51.2	0.0031	33.9	
2007	UC	27.0		31.2		22.8	
	DT	39.4	0.0004	37.4		41.4	0.0005
	NQ	35.8	0.0080	41.1	0.0279	30.6	
Yield difference (kg/tree) between UC in 2002, and UC, DT, and NQ in 2007							
	UC	3.6		4.9		2.3	
	DT	16.0	0.0001	11.1	0.0142	20.9	0.0001
	NQ	12.5	0.0003	14.9	0.0013	10.1	0.0494

Notes: Data are means of 5 replicates. ^aFigures indicate the probability of that treatment being different from the UC.

independent contrasts for mean separation ($P \leq 0.05$) (Super Anova, Abacus Concepts, Berkeley, CA); and by linear regression and correlation analysis ($P \leq 0.05$) (Prism 7, GraphPad Software, Inc., San Diego, CA). There were six replicates per treatment in the San Joaquin County Trial and data were analyzed with Repeated Measures Analysis of Variance ANOVA ($P \leq 0.05$), followed by independent contrasts for mean separation ($P \leq 0.05$) (SuperAnova, Abacus Concepts, Berkeley, CA).

Results

Sutter County Trial: Pretreatment yield data taken in 2002 indicated that there were no significant differences at the beginning of the Sutter County trial ($P > 0.05$) (Table 1). During the course of the trial, combined data for 'Paradox' and own-rooted treatments indicated yields for the trees treated

with NQ treatment were greater than for UC trees in 2003 ($P \leq 0.0027$), 2006 ($P \leq 0.0041$), and 2007 ($P \leq 0.0080$). For trees treated with DT, yields were greater than for UC trees in 2006 ($P \leq 0.0068$) and 2007 ($P \leq 0.0004$). For 'Paradox' rootstock, yields for trees treated with NQ were greater than UC trees for 2003 ($P \leq 0.0338$), 2005 ($P \leq 0.0008$), 2006 ($P \leq 0.0031$), and 2007 ($P \leq 0.0279$). Yields for trees treated with DT were greater than UC trees for 2005 ($P \leq 0.0325$) and 2006 ($P \leq 0.0114$). For own-rooted trees, yields for trees treated with NQ were greater than UC trees in 2003 ($P \leq 0.0383$), as were yields for trees treated with DT in 2007 ($P \leq 0.0005$). Overall, from 2002 to 2007, yields for UC trees did not increase. In contrast, compared to UC trees, yields increased for trees treated with DT ($P \leq 0.0142$) or with NQ ($P \leq 0.0013$) on 'Paradox'; for trees treated with DT ($P \leq 0.0001$) or with NQ ($P \leq 0.0494$) on own-rooted; and for trees treated

Table 2. Trunk cross-sectional area for Sutter County trial.

Year	Treatment	Trunk cross-sectional area (cm)			
		Combined	Paradox	Own-rooted	
2002	UC	354	221	488	
	DT	352	279	425	
	NQ	386	324	448	0.0396
2003	UC	422	292	552	
	DT	396	363	428	0.0396
	NQ	483	449	516	0.0001
2004	UC	448	324	572	
	DT	468	400	535	0.0302
	NQ	526	509	542	0.0001
2005	UC	460	354	567	
	DT	476	425	526	0.0420
	NQ	535	542	528	0.0001
2006	UC	503	386	620	
	DT	555	487	623	0.0044
	NQ	605	613	597	0.0001
2007	UC	533	421	647	
	DT	571	525	619	0.0037
	NQ	648	677	620	0.0001

Notes: Data are means of 5 replicates. ^aFigures indicate the probability of that treatment being different from the UC.

Table 3. Tree rating ranging from 0 (dead) to 5 (most vigorous) for Sutter County trial^a.

Year	Treatment	___Paradox___	___Ownroot___
2005	UC	3.6	3.0
	DT	3.8	3.6
	NQ	4.4	3.8
2006	UC	3.9	3.5
	DT	4.6	4.2
	NQ	4.6	4.1

Notes: Data are means of 5 replicates. ^aRating Scale: 0=Dead, 1= Very low vigor, dieback, 2=Early yellowing, 3=No new shoot growth, 4=Some new shoot growth, 5=Most vigorous. ^bFigures indicate the probability of that treatment being different from the UC.

with DT ($P \leq 0.0001$) or with NQ ($P \leq 0.0003$) on both rootstocks combined.

Pretreatment, in 2002, there were no differences in trunk cross sectional area for both rootstocks combined or for own-rooted trees ($P > 0.05$) (Table 2). For ‘Paradox,’ initial trunk cross sectional area for trees treated with NQ was greater than that for UC trees ($P \leq 0.0396$). For both rootstocks combined, trunk cross sectional area for trees treated with NQ was greater than UC trees for 2003 ($P \leq 0.0064$), 2004 ($P \leq 0.0009$), 2005 ($P \leq 0.0012$), 2006 ($P \leq 0.0001$), and 2007 ($P \leq 0.0001$), and for trees treated with DT in 2006 ($P \leq 0.0428$). For ‘Paradox’ rootstock, trunk cross sectional area was greater than the UC trees for trees treated with DT in 2003 ($P \leq 0.0396$), 2004 ($P \leq 0.0302$), 2005 ($P \leq 0.0420$), 2006 ($P \leq 0.0044$), and 2007 ($P \leq 0.0037$), and for trees treated with NQ in 2003 ($P \leq 0.0001$), 2004 ($P \leq 0.0001$), 2005 ($P \leq 0.0001$), 2006 ($P \leq 0.0001$), and 2007 ($P \leq 0.0001$). For own-rooted trees, trunk cross sectional area of trees treated with DT was greater than UC trees in 2003 ($P \leq 0.0003$). Overall, differences in trunk cross sectional area between when the trial was initiated in 2002 and terminated in 2007 were greater ($P \leq 0.05$) for all rootstock and treatment combinations (data not shown).

Visual rating of tree vigor conducted in 2005 indicated ‘Paradox’ trees treated with NQ were more vigorous than UC trees ($P \leq 0.02$) (Table 3). ‘Paradox’ trees treated with DT were more vigorous than UC trees in 2006 ($P \leq 0.04$). Own-rooted trees treated with NQ were more vigorous than UC trees in 2005 ($P \leq 0.04$).

Regression and correlation analysis support the results discussed above. Slopes of the lines are not

significantly different, but the y intercepts for yield are significantly different between trees treated with DT and UC trees. Linear regression analysis over time demonstrated positive but not significant slopes for yield ($P > 0.05$) for all treatments except for own-root UC trees that had a negative slope. The elevation of the line for own-root trees treated with DT ($yield = 0.6822 + 6.8978 \cdot time$) was greater ($P \leq 0.025$) than that for UC trees ($yield = -29.5084 - 1.5548 \cdot time$).

For trunk cross sectional area (TCSA), linear regression demonstrated positive slopes for all treatments that were significant at $P \leq 0.05$. For trees treated with NQ, y intercepts were significantly different from UC trees for both ‘Paradox’ and own-rooted trees. Line elevation for own-rooted trees treated with NQ ($TCSA = 9.8716 + 3.8494 \cdot time$, $P \leq 0.0158$) was greater than that for UC trees ($TCSA = 7.3885 + 2.9901 \cdot time$). The same relationship was true for trees treated with NQ that were on ‘Paradox’ rootstock ($TCSA = 23.0473 + 5.9105 \cdot time$, $P \leq 0.0009$) compared to UC trees ($TCSA = 20.221 + 5.7692 \cdot time$).

Prior to treatment on the first sampling date, there were no differences in numbers of root-lesion or ring nematode for either rootstock ($P > 0.05$) (Table 4). Throughout the course of the trial, reductions were not observed in nematode populations in soil ($P > 0.05$). For trees treated with DT, increases in the population of ring nematode in soil were observed on ‘Paradox’ in October of 2006 ($P \leq 0.050$), and on own-rooted trees in April 2004 ($P \leq 0.004$), October 2004 ($P \leq 0.002$), and April 2005 ($P \leq 0.04$). An increase in the population of root-lesion nematode for trees treated with NQ was observed on own-rooted trees in October 2005 ($P \leq 0.006$).

Table 4. Effect of treatments on densities of nematodes per 1,000 cm³ soil in Sutter County Trial.

Sample	Date	Treatment	Paradox		Own-rooted		
			Lesion	Ring	Lesion	Ring	
October 2003		UC	3,030	0	4,380	10	
		DT	3,200	0	5,740	100	
		NQ	3,820	0	4,410	0	
April 2004		UC	1,440	40	2,560	110	
		DT	1,330	540	2,020	3,270	0.004
		NQ	1,320	300	1,870	0	
October 2004		UC	1,130	20	1,230	450	
		DT	2,250	30	2,470	3,830	0.002
		NQ	1,210	420	1,580	0	
April 2005		UC	870	40	610	300	
		DT	1,410	0	580	2,480	0.040
		NQ	2,290	260	1,350	10	
October 2005		UC	1,470	10	1,890	1,380	
		DT	2,060	30	2,580	2,340	
		NQ	2,850	540	4,970	30	0.006
April 2006		UC	1,380	1,610	1,640	590	
		DT	2,650	980	1,300	2,560	
		NQ	1,310	2,020	1,680	830	
October 2006		UC	470	280	740	470	
		DT	320	1,340	960	290	0.050 ^a
		NQ	620	1,030	2,080	660	

Notes: Data are means of 5 replicates. ^aFigures indicate the probability of that treatment being different from the UC.

San Joaquin County Trial: In the San Joaquin County trial, in October 2005, six months after treatment, numerically, 10 out of 13 treatments showed an increase in trunk circumference ranging from 7.13 to 12.05% compared to a 7.1% increase for the UC trees (Table 5). The largest increases were 12.05% for trees treated with NQ at 56L/ha ($P \leq 0.0105$) followed by 11.36% for trees treated with DT 28kg/ha + NQ 37L/ha ($P \leq 0.0269$). These same two treatments continued to show significant increases in trunk circumference on subsequent sampling dates ($P \leq 0.05$). Trees treated with DT at 14kg/ha was the only treatment that failed to show a numerical increase in trunk circumference compared to UC on at least one occasion ($P > 0.05$).

At the time of application in April 2005, although the number of root-lesion nematode in soil was numerically greater than the UC trees for all but one treatment, there were no significant differences between treatments ($P > 0.05$) (Table 6). When sampled in October 2005, trees treated with DT at 14kg/ha + NQ 23.4L/ha was the only treatment to show a reduction in the soil population of nematodes ($P \leq 0.0005$). When sampled in October 2006, trees treated with DT at 28kg/ha + NQ 23.4L/ha ($P \leq 0.05$) was the only treatment to show a reduction in nematodes in soil.

At the time of application in April 2005, roots from trees treated with DT at 14kg/ha ($P \leq 0.02$) and with NQ at 37.4L/ha ($P \leq 0.05$) had more root-lesion nematode per gram of root than the UC

Table 5. Circumference of trees in San Joaquin County trial.

Treatment (L/ha)	Initial trunk Circumference (cm)	Percent increase from initial circumference			
		October 2005	April 2006	October 2006	
UC	68.3	7.1	7.2	10.6	
DT 56	66.5	7.0	7.7	10.6	
DT 28	61.4	7.2	8.0	10.9	
DT 14	67.1	5.3	5.7	7.9	
DT 56+NQ 37.4	71.2	8.4	8.9	11.7	
DT 56+NQ 23.4	69.7	9.2	9.9	12.8	
DT 28+NQ 37.4	61.5	11.4	12.9	16.3	0.0269 ^a 0.0052 0.0091
DT 28+NQ 23.4	68.2	8.5	9.5	12.8	
DT 14+NQ 37.4	69.0	7.3	8.1	11.1	
DT 14+NQ 23.4	71.7	7.1	5.9	10.0	
NQ 56	70.9	12.1	12.8	15.8	0.0105 0.0062 0.0176
NQ 37.4	64.0	7.6	8.6	11.3	
NQ 23.4	72.3	6.9	7.7	10.4	
NQ 12	61.6	7.2	8.1	10.8	

Notes: Data are means of 6 replicates. ^aFigures indicate the probability of that treatment being different from the UC.

(Table 7). In October 2005, roots from trees treated with NQ at 12L/ha had a greater number of root-lesion nematode per gram of root than UC trees ($P \leq 0.01$). In April 2006, there were no differences in number of root-lesion nematode per gram of root.

At the time of application in April 2005, there were no differences in number of ring nematode per liter of soil ($P > 0.05$) (Table 8). In October 2005, trees treated with DT at 14 kg/ha + NQ 23.4L/ha and with NQ 56L/ha had fewer ring nematode in soil than UC trees ($P \leq 0.01$). In April 2006, trees in five treatments had fewer ring nematode in soil than UC: DT 56 kg/ha + NQ 23.4L/ha ($P \leq 0.03$), DT 28 kg/ha + NQ 37.4L/ha ($P \leq 0.003$), DT 14 kg/ha + NQ 23.4L/ha ($P \leq 0.003$), NQ 56L/ha ($P \leq 0.003$), and NQ 23.4L/ha ($P \leq 0.03$).

Discussion

Bionematicides were evaluated on two walnut rootstocks, own-rooted 'English' ('Chandler') and 'Paradox.' Own-rooted 'English' walnut trees can be used in areas where commonly used rootstocks such as 'Paradox' (*J. hindsii* and *J. hindsii* x *J. regia*) are

undesirable because of hypersensitivity to cherry leaf roll virus. This disease is known as walnut blackline because a black line forms at the graft union in infected trees (Mircetich et al., 1998). The disease can be overcome by using either 'English' walnut rootstocks (*J. regia*) or 'English' walnut cultivars growing on their own roots. Micropropagation techniques can be used to produce own-rooted 'English' walnut cultivars (McGranahan et al., 1988).

DT is a killed-microbial product of the fungus *Myrothecium verucaria*. The mode of action of DT is due primarily to the presence of many, relatively low-molecular-mass, water-soluble, compounds, which act synergistically (Wilson and Jackson, 2013). It has been shown to kill nematodes via contact as well as to inhibit hatching and development of eggs, cause muscle paralysis, feeding inhibition, depletion of lipids, and changes in sensory perception affecting activities such as host and mate-finding (Twomey et al., 2000, 2002; Rehberger et al., 2002). In addition to activity to nematodes, increased plant health, shoot and root weights, greening, and root proliferation have also been observed in trials by others (Spence and Lewis, 2010). Additional work has shown that it

Table 6. Effect of treatments on densities of lesion nematodes per 1,000 cm³ soil in San Joaquin County trial.

Treatment (L/ha)	Date sampled		
	April 2005	October 2005	April 2006
UC	2,783	4,208	2,950
DT 56	6,708	4,167	2,375
DT 28	4,825	8,075	6,075
DT 14	4,458	4,883	3,875
DT 56+NQ 37.4	4,233	7,417	1,967
DT 56+NQ 23.4	3,708	6,625	2,350
DT 28+NQ 37.4	8,733	4,583	1,525
DT 28+NQ 23.4	3,492	5,325	1,575
DT 14+NQ 37.4	5,625	8,050	2,200
DT 14+NQ 23.4	8,405	2,333	3,150
NQ 56	4,667	4,850	4,475
NQ 37.4	2,817	6,600	2,650
NQ 23.4	3,875	5,433	1,633
NQ 12	2,517	3,175	1,583

Notes: Data are means of 6 replicates. ^aFigures indicate the probability of that treatment being different from the UC.

enhances microbial antagonism towards nematodes. This antagonism was associated with structural and functional changes of the rhizosphere bacterial and fungal community (Fernandez et al., 2001). DT applied at planting to strawberries in a greenhouse trial decreased populations of *Pratylenchus penetrans* and stimulated root and crown growth compared to infested controls (Pinkerton and Kitner, 2006).

NQ is an extract of *Quillaja saponaria* a tree endemic to Chile that is rich in secondary plant metabolites including saponins, glycosides, polyphenols, and tannins that are found in the cortex, leaves and flowers (Insunza et al., 2001). Aqueous extracts have been shown to have nematicidal effects against a variety of nematode species and to increase root growth (Martin and Magunacelaya, 2005).

Our trials have demonstrated that bionematicides can improve yield, growth, and vigor in walnut orchards infested with plant-parasitic nematodes. This prolongs the viable life of an orchard and reduces the frequency of pre-plant fumigations. This research contributed to the registration of these organic nematicides (OMRI certified) in California.

As we move away from traditional fumigant and nonfumigant nematicides towards natural products with different modes of action, the most effective application methods, rates, and timing, and interpretation of results become less straight forward. For example, in the trial conducted in San Joaquin County, increasing yields were associated with an increase in populations of nematodes. This could be an indication of the development of a more vigorous root system that is capable of supporting greater populations of nematodes.

For more than 30 years we have searched for a product that would replace DBCP. What we have found after years of believing that “the only good plant-parasitic nematode is a dead nematode” is that products are available that will increase yields in the presence of plant-parasitic nematodes and, may actually permit populations to increase. This opens the door to additional research on how to best utilize the new tools that we have to maximize yields for growers. It also raises questions for additional long-term research on perennial crops. Will yields continue to increase as demonstrated in these trials, will yields

Table 7. Effects on lesion nematode per gram of roots in San Joaquin County trial.

Treatment (L/ha)	Date sampled			
	April 2005	October 2005	April 2006	
UC	8	9	0	
DT 56	142	9	2	
DT 28	121	18	4	
DT 14	203	0.02 ^a	109	4
DT 56+NQ 37.4	11	12	5	
DT 56+NQ 23.4	18	15	2	
DT 28+NQ 37.4	8	81	1	
DT 28+NQ 23.4	45	4	0	
DT 14+NQ 37.4	36	74	1	
DT 14+NQ 23.4	1	22	2	
NQ 56	45	94	13	
NQ 37.4	130	0.05	13	16
NQ 23.4	165	25	5	
NQ 12	11	229	0.01	1

Notes: Data are means of 6 replicates. ^aFigures indicate the probability of that treatment being different from the UC.

stabilize, or will yields crash at some point in the future? Research with bionematicides on annual crops has also shown increases in yield without a reduction in nematode populations (Westerdahl and Radewald, 2011). The current research is also a challenge to others to take another look at data they may have set aside because yields increased, but effects on nematode populations did not match expectations.

Acknowledgments

This work was supported by the California Walnut Marketing Board, Desert King, and Valent Laboratories. The authors are grateful to the grower

Table 8. Effect of treatments on densities of nematodes per 1,000 cm³ soil in San Joaquin County trial.

Treatment (L/ha)	Ring nematode			
	April 2005	October 2005	April 2006	
UC	975	317	1,575	
DT 56	1,133	7,300	2,150	
DT 28	1,500	950	525	
DT 14	83	1,117	900	
DT 56+NQ 37.4	583	450	2,000	
DT 56+NQ 23.4	3,283	2,500	350	0.03
DT 28+NQ 37.4	83	300	0	0.003
DT 28+NQ 23.4	900	1,650	1,600	
DT 14+NQ 37.4	575	1,183	825	
DT 14+NQ 23.4	63	0	0.01 ^a	0 0.003
NQ 56	0	0	0.01	200 0.003
NQ 37.4	150	800	1,275	
NQ 23.4	750	4,883	475	0.03
NQ 12	2,900	950	1,575	

Notes: Data are means of 6 replicates. ^aFigures indicate the probability of that treatment being different from the UC.

cooperator Joe Conant of the Whitney Warren Ranch for his assistance and long-term cooperation, and to Barton Ranch. We are also grateful to Cindy Anderson, Samuel Metcalf, and Claudia Negron for field and laboratory assistance.

References

Ayoub, S. M. 1977. Plant nematology an agricultural training aid California Department of Food and Agriculture, Division of Plant Industry, Sacramento, CA.
 Buzo, T., McKenna, J., Kaku, S., Anwar, S. A. and Mckenry, M. V. 2009. VX211, A vigorous new walnut

hybrid clone with nematode tolerance and a useful resistance mechanism. *Journal of Nematology* 41: 211–6.

Byrd, D. W. Jr, Barker, K. R., Ferris, H., Nusbaum, C. J., Griffin, W. E., Small, R. H. and Stone, C. A. 1976. Two semi-automatic elutriators for extracting nematodes and certain fungi from soil. *Journal of Nematology* 8:206–12.

CIMIS 2021. California Irrigation Management Information System. State of California, available at: <https://cimis.water.ca.gov>.

Fernandez, C. R., Rodriguez-Kabana, R., Warrior, P. and Kloepper, J. W. 2001. Induced soil suppressiveness to a root-knot nematode species by a nematicide. *Biological Control* 22:103–14.

Ferris, H. 2021. Chemical methods of nematode management, available at: <http://nemaplex.ucdavis.edu/Mangmnt/Chemical.htm>.

Giraud, D., Riddle, L. J., Anderson, C. E. and Westerdahl, B. B. 2011. New products to improve growth of field grown Easter lily bulbs. *Acta Horticulturae* 900:333–7.

Hasey, J. K., Westerdahl, B. B. and Lampinen, B. 2004. Long-term performance of own-rooted ‘Chandler’ walnut compared to ‘Chandler’ walnut on Paradox rootstock. *Acta Horticulturae* 636:83–6.

Insunza, V., Aballay, E. and Macaya, J. 2001. *In vitro* nematicidal activity of aqueous plant extracts on Chilean populations of *Xiphinema americanum sensu lato*. *Nematropica* 31:47–54.

Lownsbery, B. F. 1956. *Pratylenchus vulnus*, primary cause of the root-lesion disease of walnuts. *Phytopathology* 46:375–9.

Lownsbery, B. F. 1959. Studies of the nematode, *Criconemoides xenoplax*, on peach. *Plant Disease Reporter* 43:913–7.

Lownsbery, B. F., Moody, E. H., Moretto, A., Noel, G. R. and Burlando, T. M. 1978. Pathogenicity of *Macroposthonia xenoplax* to Walnut. *Journal of Nematology* 10:232–6.

Martin, R. S. and Magunacelaya, J. C. 2005. Control of plant-parasitic nematodes with extracts of *Quillaja saponaria*. *Nematology* 7:577–85.

McGranahan, G., Leslie, C. A. and Driver, J. A. 1988. *In vitro* propagation of mature Persian walnut cultivars. *HortScience* 23:220.

Mircetich, S. M. J., Rowhani, A., Civerolo, E. L. and Ramos, D. E. 1998. “Blackline disease”, In Ramos, D. E. (Ed.), *Walnut Production Manual*, UC ANR Publication 3373, Oakland, pp. 233–41.

Pinkerton, J. N. and Kitner, M. L. C. 2006. Effects of biologically-derived products on mobility and reproduction of the root-lesion nematode *Pratylenchus penetrans*, on strawberry. *Nematropica* 36:181–96.

Raski, D. 1952. On the morphology of *Criconemoides* Taylor, 1936, with descriptions of six new species. *Proceedings of the Helminthological Society* 19:85–99.

Rehberger, L., Shammo, B. and Warrior, P. 2002. Examination and disclosure of the multifaceted modes of action of DiTera®, a biological nematicide. *Biological Control* 4:290.

Retzlaff, W. A., Williams, L. E. and DeJong, T. M. 1992. Photosynthesis, growth, and yield response of ‘Casselman’ plum to various ozone partial pressures during orchard establishment. *Journal of the American Society for Horticultural Science* 117:703–10.

Spence, K. O. and Lewis, E. E. 2010. Biopesticides with complex modes of action: direct and indirect effects of DiTera on *Meloidogyne incognita*. *Nematology* 12:835–46.

Twomey, U., Warrior, P., Kerry, B. R. and Perry, R. N. 2000. Effects of the biological nematicide, DiTera®, on hatching of *Globodera rostochiensis* and *G. pallida*. *Nematology* 2:355–62.

Twomey, U., Rolfe, R. N., Warrior, P. and Perry, R. N. 2002. Effects of the biological nematicide, DiTera®, on movement and sensory responses of second stage juveniles of *Globodera rostochiensis*, and stylet activity of *G. rostochiensis* and fourth stage juveniles of *Ditylenchus dipsaci*. *Nematology* 4:909–15.

UCIPM 2021. Weather models and degree-days,” Agriculture and Natural Resources University of California, Oakland, CA, available at: <http://ipm.ucanr.edu/WEATHER/index.html>.

United States Environmental Protection Agency 1993. Protection of Stratospheric Ozone: Final rule. *Federal Register*, 58, 236, December 10, pp. 65018–62.

United States Environmental Protection Agency 2014. Hazard Summary 1,2-Dibromo-3-Chloropropane (DBCP) United States Environmental Protection Agency Technology Transfer Network – Air Toxics Web Site, Washington, DC, available at: <https://www.epa.gov/sites/production/files/2016-09/documents/1-2-dibromo-3-chloropropane.pdf>.

Westerdahl, B. B. and Radewald, J. D. 2011. Natural products for nematode management in tomatoes. *Acta Horticulturae* 914:275–80.

Wilson, M. J. and Jackson, T. A. 2013. Progress in the commercialization of bionematicides. *Biocontrol* 58:715–22.